

Supporting Information for Algorithmic Generation and Exploitation of Topology-Aware Models for Semiexperimental Structure Refinement

Vincenzo Barone

S1 Global Benchmark Diagnostics

Tables S1 and S2 collect the numerical values underlying the graphical benchmark summaries in the main text.

S2 Atom Numbering Used in the Tables

Figure S1 reproduces the atom numbering shown in the main text for use with the structural tables and diagnostic comparisons. The same Cartesian numbering is used by the topology perception, coordinate generation, and least-squares reporting steps. The p-EBN structure is included as a legacy MSR-import companion case for numbering consistency.

S3 Glycine I C_s Constraint Benchmark

Table S5 reports the Gly-I benchmark run from the QC reference Cartesian geometry with no stored structural constraints. The fit uses moments of inertia as observables and all three rotational components because the C_s structure is not a planar rotor in the inertial sense: the two methylene and two amino hydrogens lie outside the heavy-atom plane. The ground-state constants and statistical weights are from Godfrey and Brown, J. Am. Chem. Soc. 117, 2019-2023 (1995), Table 1. The corrected constants are $B_e = B_0 + \Delta B_{\text{vib}}$ with the Gly-Ip zero-point corrections of Kasalová et al. The MORPHEUS advisor generated the following four hard constraints from the QC reference geometry and isotope-availability pattern:

```
HNH(Frozen,Value=105.634453352622)=A(3,2,10)
NH2WAG(Frozen,Value=55.901476176288)=U(1,2,3,10)
ROH(Frozen,Value=0.965159162788)=R(6,7)
ACOH(Frozen,Value=106.584216097335)=A(4,6,7)
```

The numerical data entering the fit are collected in Table S4 so that the corrected constants can be reproduced without consulting the original tables.

Table S1: One-page audit of the representative MORPHEUS semiexperimental refinements. “Model” is the working-coordinate model used for the main row. “Fit” gives the independent rotational components entering the normal equations; for moment-based fits, AB denotes (I_a, I_b) . “GIC/A” gives the algorithmically generated non-redundant coordinate count and the totally symmetric active count before constraint projection. “Eff./rank” gives the independent variables after primitive-constraint projection and the final numerical rank. Rotational residuals are RMS/maximum absolute values in MHz over all reported diagnostic constants.

System	PG	Model	Fit	Iso.	GIC/A	Eff./rank	Constraints	Steps	RMS/max
Glycolaldehyde	C_s	GIC	ABC	10	18/12	12/12	none	3/0; min	0.5640/1.7961
Cyclopentadiene	C_{2v}	Cart.	ABC	14	27/10	10/10	none	10/0; min	0.0635/0.2289
Nitrobenzene	C_{2v}	GIC	AB	9	36/13	13/13	8 primitive	4/0; pos.	0.1480/0.2605
p-EBN	C_{2v}	GIC	AB	11	39/14	6/6	22 frozen	4/20; min	0.1190/0.5674
Azulene	C_{2v}	GIC	AB	12	48/17	10/10	8 primitive	8/28; min	0.0731/0.2900
Norcamphor	C_1	GIC	ABC	9	48/48	18/18	30 H primitive	8/0; min	0.0710/0.2091
Glycine I	C_s	GIC	ABC	5	24/15	11/11	4 function	3/17; min	0.0269/0.0570
Glycine II	C_1	GIC	ABC	10	24/24	19/19	5 function	26/8; min	0.2491/0.6707

Table S2: Planar-component diagnostics for the planar benchmarks. Only two principal moments are independent for a planar molecule; the selected pair marked by an asterisk is used in the normal equations, while all three rotational constants are recomputed after convergence. The residuals are RMS/maximum absolute MHz values over the complete diagnostic A , B , and C constant set.

System	Pair	Fitted moments	Rank	Cond.	Diagnostic RMS/max
Nitrobenzene	AB*	(I_a, I_b)	13	27464.4	0.1480/0.2605
Nitrobenzene	AC	(I_a, I_c)	13	56007.7	0.2593/0.4597
Nitrobenzene	BC	(I_b, I_c)	13	10623.3	2.4602/4.3693
Azulene	AB*	(I_a, I_b)	10	956.7	0.0731/0.2900
Azulene	AC	(I_a, I_c)	10	1780.0	0.1086/0.4222
Azulene	BC	(I_b, I_c)	10	950.2	0.3232/0.7541

Table S3: Comparison of MORPHEUS working-coordinate models. “Work.” is the number of non-redundant GICs or symmetry-adapted Cartesian vibrational directions, “Sym.” is the number of totally symmetric working coordinates before primitive-constraint projection, and “Eff.” is the number of independent least-squares variables after projection. Residuals are RMS/maximum absolute values in MHz.

System	Model	Work.	Sym.	Eff.	Rank	Cond.	Residuals
Cyclopentadiene	GICs	27	10	10	10	454.5	0.0635/0.2289
Cyclopentadiene	Sym. Cart.	27	10	10	10	635.4	0.0635/0.2289
p-EBN	GICs	39	14	6	6	2440.2	0.1190/0.5674
Azulene	GICs	48	17	10	10	956.7	0.0731/0.2900
Azulene	Sym. Cart.	48	17	10	10	744.8	0.0731/0.2900
Norcamphor	GICs	48	48	18	18	3700.3	0.0710/0.2091
Norcamphor	Sym. Cart.	48	48	18	18	1615.8	0.0710/0.2091




Figure S1: Atom numbering used for the benchmark and companion structures: glycolaldehyde, cyclopentadiene, nitrobenzene, p-EBN, azulene, norcamphor, glycine I, glycine II, and the maleic, phthalic, and succinic anhydride series. Testosterone is included as the 49-atom fused-ring scale check.

Table S4: Glycine I rotational input data. B_0 , $\sigma(B_0)$, ΔB_{vib} , and $B_e = B_0 + \Delta B_{\text{vib}}$ are in MHz. The $^{13}\text{C}(1)$ and $^{13}\text{C}(2)$ labels of Godfrey and Brown correspond here to C13_carboxyl and C13_methylene, respectively.

Isotopologue	Constant	B_0	$\sigma(B_0)$	ΔB_{vib}	B_e
parent	A	10341.5210	0.0890	78.6700	10420.1910
parent	B	3876.1785	0.0012	31.3000	3907.4785
parent	C	2912.3509	0.0010	22.3300	2934.6809
C13_carboxyl	A	10340.9430	0.0290	77.8000	10418.7430
C13_carboxyl	B	3869.1142	0.0057	30.9500	3900.0642
C13_carboxyl	C	2908.3166	0.0026	22.0600	2930.3766
C13_methylene	A	10227.0840	0.0880	76.7300	10303.8140
C13_methylene	B	3859.0510	0.0210	30.9500	3890.0010
C13_methylene	C	2893.5960	0.0200	21.9200	2915.5160
CD2	A	9318.4520	0.0230	69.8400	9388.2920
CD2	B	3799.2353	0.0023	31.2100	3830.4453
CD2	C	2832.3222	0.0015	21.4800	2853.8022
N15	A	10341.4500	0.1200	78.7600	10420.2100
N15	B	3762.4570	0.0530	30.1300	3792.5870
N15	C	2847.6390	0.0290	21.7100	2869.3490

Table S5: Complete rotational-constant comparison for glycine I. Corrected experimental constants, calculated constants, and differences are in MHz.

Isotopologue	Constant	Corrected exp.	Calculated	Difference
parent	A	10420.1910	10420.2481	-0.0570
parent	B	3907.4785	3907.4786	-0.0001
parent	C	2934.6809	2934.6808	+0.0001
C13_carboxyl	A	10418.7430	10418.7403	+0.0027
C13_carboxyl	B	3900.0642	3900.0634	+0.0008
C13_carboxyl	C	2930.3766	2930.3769	-0.0003
C13_methylene	A	10303.8140	10303.7637	+0.0503
C13_methylene	B	3890.0010	3889.9809	+0.0201
C13_methylene	C	2915.5160	2915.5484	-0.0324
CD2	A	9388.2920	9388.2920	+0.0000
CD2	B	3830.4453	3830.4453	-0.0000
CD2	C	2853.8022	2853.8022	-0.0000
N15	A	10420.2100	10420.2444	-0.0344
N15	B	3792.5870	3792.5436	+0.0434
N15	C	2869.3490	2869.3715	-0.0225

Table S6: Selected primitive structural parameters for the glycine I C_s benchmark with MORPHEUS-generated constraints. Bond lengths are in Å; angles and dihedrals are in degrees.

Parameter	Description	Value	Std. error
R(1,2)	C–N	1.44285	0.00076
R(1,4)	C–C	1.51134	0.00085
R(1,8)	C–H	1.09081	0.00021
R(1,9)	C–H	1.09081	0.00021
R(2,3)	N–H	1.00228	0.00011
R(2,10)	N–H	1.00228	0.00011
R(4,5)	C–O	1.20529	0.00184
R(4,6)	C–O	1.35001	0.00128
R(6,7)	O–H	0.96516	0.00000
A(2,1,4)	N–C–C	115.266	0.062
A(8,1,9)	H–C–H	105.923	0.031
A(3,2,10)	H–N–H	105.634	0.000
A(1,4,5)	C–C–O	125.338	0.095
A(1,4,6)	C–C–O	111.640	0.123
A(5,4,6)	O–C–O	123.022	0.060
A(4,6,7)	C–O–H	106.584	0.000
D(2,1,4,5)	N–C–C–O	180.000	0.000
D(1,4,6,7)	C–C–O–H	0.000	0.000
D(8,1,4,5)	H–C–C–O	-56.781	0.030
D(9,1,4,5)	H–C–C–O	56.781	0.030

S4 Glycine II Functional-Constraint Benchmark

Table S7 reports the Gly-II advisor-constrained benchmark discussed in the main text. The fit uses moments of inertia as observables, but the diagnostic table is reported in MHz. The corrected experimental constants are $B_e = B_0 + \Delta B_{\text{vib}}$ with the Gly-II zero-point corrections. The Cartesian input does not store constraints; the five functional constraints are generated by the MORPHEUS advisor from the QC geometry and the isotopologue coverage.

Table S7: Complete rotational-constant comparison for glycine II. Corrected experimental constants, calculated constants, and differences are in MHz.

Isotopologue	Constant	Corrected exp.	Calculated	Difference
parent	A	10195.5900	10196.2607	-0.6707
parent	B	4105.3670	4105.3251	0.0419

Isotopologue	Constant	Corrected exp.	Calculated	Difference
parent	C	3039.8450	3039.8159	0.0291
N15	A	10194.1150	10194.6721	-0.5571
N15	B	3994.8380	3994.8046	0.0334
N15	C	2978.7993	2978.7971	0.0022
C13_methylene	A	10060.9700	10061.6298	-0.6598
C13_methylene	B	4088.6720	4088.5073	0.1647
C13_methylene	C	3018.8730	3018.8495	0.0235
C13_carboxyl	A	10195.6980	10195.0945	0.6035
C13_carboxyl	B	4093.0000	4093.1241	-0.1241
C13_carboxyl	C	3033.1070	3033.1532	-0.0462
O18_carbonyl	A	10064.2400	10063.7580	0.4820
O18_carbonyl	B	3929.9160	3929.9078	0.0082
O18_carbonyl	C	2931.7130	2931.8037	-0.0907
O18_hydroxyl	A	9580.9720	9580.9886	-0.0166
O18_hydroxyl	B	4090.8498	4090.8965	-0.0467
O18_hydroxyl	C	2975.1370	2975.1377	-0.0007
OD_NH2	A	9792.2000	9792.2302	-0.0302
OD_NH2	B	4096.1490	4096.1104	0.0386
OD_NH2	C	2998.1238	2998.0987	0.0251
CD2	A	9145.3030	9145.3053	-0.0023
CD2	B	4020.3863	4020.3836	0.0027
CD2	C	2947.9923	2947.9913	0.0010
OH_ND2	A	9927.6990	9927.6852	0.0138
OH_ND2	B	3716.3758	3716.3613	0.0145
OH_ND2	C	2841.7341	2841.7204	0.0137
OD_ND2	A	9542.6600	9542.6581	0.0019
OD_ND2	B	3710.5050	3710.5504	-0.0454
OD_ND2	C	2806.0910	2806.1569	-0.0659

S5 Cyclopentadiene Benchmark

Table S8 reports the complete comparison between corrected experimental rotational constants and constants calculated from the final semiexperimental structure. The workflow uses generalized internal coordinates (GICs) or symmetry-adapted Cartesian coordinates generated within MORPHEUS. The data set contains 14 isotopologues and all three principal rotational constants for each isotopologue. The MORPHEUS refinement uses only a Cartesian representation of the molecular structure, whereas the MSR refinement used manual Z-matrix coordinates and constraints.

Table S8: Complete rotational-constant comparison for the cyclopentadiene benchmark derived from MSR data. Corrected experimental and calculated constants are in MHz.

Isotopologue	Component	Corrected exp.	Calculated	Difference
msr_symm_01	A	8489.154993	8489.232117	-0.077124
msr_symm_01	B	8292.100993	8292.093033	0.007960
msr_symm_01	C	4305.664997	4305.658417	0.006580
msr_symm_02	A	8292.098993	8292.093033	0.005960
msr_symm_02	B	8280.278992	8280.319224	-0.040231
msr_symm_02	C	4251.270992	4251.257436	0.013556
msr_symm_03	A	8482.790995	8482.866229	-0.075234
msr_symm_03	B	8105.052002	8105.082639	-0.030637
msr_symm_03	C	4253.082997	4253.084517	-0.001520
msr_symm_04	A	8408.381002	8408.358013	0.022989
msr_symm_04	B	8172.557990	8172.606506	-0.048516
msr_symm_04	C	4252.643011	4252.628488	0.014523
msr_symm_05	A	8408.381002	8408.358024	0.022978
msr_symm_05	B	8172.557990	8172.606496	-0.048506
msr_symm_05	C	4252.643011	4252.628488	0.014523
msr_symm_06	A	8195.611995	8195.593227	0.018768
msr_symm_06	B	7918.397999	7918.289812	0.108187
msr_symm_06	C	4178.352982	4178.398406	-0.045424
msr_symm_07	A	8195.611995	8195.612584	-0.000589
msr_symm_07	B	7918.397999	7918.308530	0.089469
msr_symm_07	C	4178.352982	4178.398587	-0.045604
msr_symm_08	A	8477.175001	8477.228373	-0.053372
msr_symm_08	B	7650.516996	7650.483876	0.033120
msr_symm_08	C	4123.171991	4123.147057	0.024934
msr_symm_09	A	8477.175001	8477.228367	-0.053366
msr_symm_09	B	7650.516996	7650.483883	0.033113
msr_symm_09	C	4123.171991	4123.147057	0.024933
msr_symm_10	A	8371.829003	8371.953330	-0.124328
msr_symm_10	B	7735.107991	7735.111002	-0.003010
msr_symm_10	C	4122.272002	4122.241235	0.030767
msr_symm_11	A	8371.829003	8371.953344	-0.124341
msr_symm_11	B	7735.107991	7735.110990	-0.002998
msr_symm_11	C	4122.272002	4122.241235	0.030767
msr_symm_12	A	7057.547996	7057.656561	-0.108565
msr_symm_12	B	6730.362994	6730.334074	0.028920
msr_symm_12	C	3555.255987	3555.260397	-0.004410
msr_symm_13	A	7057.547996	7057.672705	-0.124709
msr_symm_13	B	6730.362994	6730.349532	0.013462

Isotopologue	Component	Corrected exp.	Calculated	Difference
msr_symm_13	C	3555.255987	3555.260180	-0.004193
msr_symm_14	A	6656.949995	6656.991310	-0.041315
msr_symm_14	B	6653.764996	6653.536122	0.228874
msr_symm_14	C	3469.286997	3469.303412	-0.016415

S6 Nitrobenzene Planar Benchmark

Nitrobenzene is a planar aromatic benchmark. The least-squares fit used the independent AB moment pair, corresponding to (I_a, I_b) , while the C constants were recomputed from the optimized Cartesian geometry and retained as diagnostic quantities. Table S9 reports the complete diagnostic comparison.

Table S9: Complete rotational-constant comparison for the nitrobenzene planar benchmark. Corrected experimental and calculated constants are in MHz. The fit used the independent AB pair; the C component is diagnostic.

Isotopologue	Component	Corrected exp.	Calculated	Difference
parent	A	3995.209000	3995.211821	-0.002821
parent	B	1296.305300	1296.307106	-0.001806
parent	C	979.000500	978.740045	+0.260455
N15	A	3995.193000	3995.211821	-0.018821
N15	B	1287.445000	1287.445036	-0.000036
N15	C	973.938200	973.679678	+0.258522
O18	A	3925.992000	3925.984021	+0.007979
O18	B	1264.689700	1264.689116	+0.000584
O18	C	956.803400	956.552095	+0.251305
6C13	A	3995.172000	3995.211821	-0.039821
6C13	B	1274.812600	1274.812638	-0.000038
6C13	C	966.688800	966.436980	+0.251820
6D	A	3995.241000	3995.211821	+0.029179
6D	B	1253.544800	1253.544803	-0.000003
6D	C	954.416100	954.164457	+0.251643
2C13	A	3949.267000	3949.255685	+0.011315
2C13	B	1295.514600	1295.514106	+0.000494
2C13	C	975.768500	975.508297	+0.260203
2D	A	3856.605000	3856.604007	+0.000993
2D	B	1296.294600	1296.294437	+0.000163
2D	C	970.447500	970.190734	+0.256766
3C13	A	3950.485000	3950.473551	+0.011449
3C13	B	1284.693900	1284.693395	+0.000505

Isotopologue	Component	Corrected exp.	Calculated	Difference
3C13	C	969.688400	969.433703	+0.254697
3D	A	3858.465000	3858.463986	+0.001014
3D	B	1276.881200	1276.881052	+0.000148
3D	C	959.645200	959.390171	+0.255029

The Kraitchman diagnostic for nitrobenzene is reported in Table S10. The comparison uses absolute substitution coordinates because the Kraitchman equations do not determine coordinate signs. The diagnostic was not used in the least-squares objective.

Table S10: Kraitchman diagnostic for the nitrobenzene planar benchmark. Absolute Kraitchman substitution coordinates and absolute fitted coordinates are in Å.

Isotopologue	Atom	Axis	Kraitchman	Fit	Difference
N15	2 N	<i>a</i>	1.64701	1.64723	-0.00022
N15	2 N	<i>b</i>	0.01795	0.00000	+0.01795
N15	2 N	<i>c</i>	0.01379	0.00000	+0.01379
O18	13 O	<i>a</i>	2.22302	2.21369	+0.00933
O18	13 O	<i>b</i>	1.06338	1.08300	-0.01962
O18	13 O	<i>c</i>	0.00896	0.00000	+0.00896
6C13	11 C	<i>a</i>	2.56993	2.56999	-0.00005
6C13	11 C	<i>b</i>	0.03786	0.00000	+0.03786
6C13	11 C	<i>c</i>	0.00000	0.00000	+0.00000
6D	12 H	<i>a</i>	3.65015	3.65027	-0.00012
6D	12 H	<i>b</i>	0.00000	0.00000	+0.00000
6D	12 H	<i>c</i>	0.01902	0.00000	+0.01902
2C13	3 C	<i>a</i>	0.48915	0.48830	+0.00085
2C13	3 C	<i>b</i>	1.21604	1.21671	-0.00066
2C13	3 C	<i>c</i>	0.00000	0.00000	+0.00000
2D	4 H	<i>a</i>	0.05793	0.06125	-0.00332
2D	4 H	<i>b</i>	2.13423	2.13424	-0.00001
2D	4 H	<i>c</i>	0.00000	0.00000	+0.00000
3C13	7 C	<i>a</i>	1.88184	1.87669	+0.00515
3C13	7 C	<i>b</i>	1.19989	1.20775	-0.00786
3C13	7 C	<i>c</i>	0.00000	0.00000	+0.00000
3D	8 H	<i>a</i>	2.43739	2.41687	+0.02052
3D	8 H	<i>b</i>	2.11909	2.14308	-0.02399
3D	8 H	<i>c</i>	0.03090	0.00000	+0.03090

Table S11: p-EBN fit diagnostics from the legacy MSR-import benchmark. The fit used the independent AB moment pair; the C rotational constants are retained as diagnostics.

Diagnostic	Value
Isotopologues	11
Effective rank	6
Effective parameters	6
Condition number	2440.25
Rotational RMS / MHz	0.118975
Max $ \Delta B $ / MHz	0.567403
Weighted RMS	0.141739
Stationary point	minimum

S7 p-EBN Legacy MSR-Import Benchmark

The p-EBN case was recovered from the legacy MSR-import fixtures and rerun with the current MORPHEUS workflow. The current run imports the Z-matrix geometry, isotopologue definitions, rotational constants, vibrational corrections, weights, and frozen Z-matrix parameters. The historical Z-matrix-style equality-constraint block is not part of the current public MORPHEUS constraint API and was therefore not imposed in this compatibility run. Table S11 summarizes the fit, and Tables S12–S14 report the rotational constants, Kraitchman diagnostic, and selected structural parameters.

Table S12: Complete rotational-constant comparison for the p-EBN legacy MSR-import benchmark. Corrected experimental and calculated constants are in MHz. The fit used the independent AB moment pair; C is diagnostic.

Isotopologue	Component	Corrected exp.	Calculated	Difference
parent	A	5690.084365	5690.075486	0.008879
parent	B	711.461697	711.463764	-0.002068
parent	C	632.404642	632.392049	0.012593
$^{13}\text{C}(1)$	A	5689.787883	5690.075486	-0.287603
$^{13}\text{C}(1)$	B	709.591078	709.571546	0.019532
$^{13}\text{C}(1)$	C	630.926783	630.896616	0.030166
$^{13}\text{C}(2)$	A	5598.029883	5598.023971	0.005913
$^{13}\text{C}(2)$	B	711.004858	711.013323	-0.008464
$^{13}\text{C}(2)$	C	630.892369	630.883832	0.008537
$^{13}\text{C}(3)$	A	5598.029883	5598.023627	0.006256
$^{13}\text{C}(3)$	B	711.004858	711.013348	-0.008490
$^{13}\text{C}(3)$	C	630.892369	630.883848	0.008521
$^{13}\text{C}(4)$	A	5598.343855	5598.493203	-0.149349
$^{13}\text{C}(4)$	B	710.960437	710.963815	-0.003379

Isotopologue	Component	Corrected exp.	Calculated	Difference
$^{13}\text{C}(4)$	C	630.861335	630.850813	0.010522
$^{13}\text{C}(5)$	A	5598.343855	5598.496086	-0.152231
$^{13}\text{C}(5)$	B	710.960437	710.963789	-0.003352
$^{13}\text{C}(5)$	C	630.861335	630.850828	0.010506
$^{13}\text{C}(6)$	A	5690.100926	5690.075486	0.025441
$^{13}\text{C}(6)$	B	709.476356	709.476380	-0.000024
$^{13}\text{C}(6)$	C	630.835668	630.821383	0.014285
$^{13}\text{C}(7)$	A	5689.975800	5690.075486	-0.099685
$^{13}\text{C}(7)$	B	703.698492	703.695775	0.002716
$^{13}\text{C}(7)$	C	626.264052	626.247314	0.016738
$^{13}\text{C}(8)$	A	5690.022772	5690.075486	-0.052713
$^{13}\text{C}(8)$	B	703.494496	703.497679	-0.003183
$^{13}\text{C}(8)$	C	626.102657	626.090418	0.012238
$^{13}\text{C}(9)$	A	5690.053400	5690.075486	-0.022086
$^{13}\text{C}(9)$	B	695.483305	695.491125	-0.007820
$^{13}\text{C}(9)$	C	619.748527	619.740932	0.007596
$^{15}\text{N}(10)$	A	5690.642889	5690.075486	0.567403
$^{15}\text{N}(10)$	B	696.225258	696.222764	0.002494
$^{15}\text{N}(10)$	C	620.336151	620.321809	0.014342

Table S13: Kraitichman diagnostic for the p-EBN legacy MSR-import benchmark. Absolute Kraitichman substitution coordinates and absolute fitted coordinates are in Å.

Isotopologue	Atom	Axis	Kraitichman	Fit	Difference
$^{13}\text{C}(1)$	1 C	<i>a</i>	1.37054	1.37943	-0.00889
$^{13}\text{C}(1)$	1 C	<i>b</i>	0.04432	0.00001	0.04431
$^{13}\text{C}(1)$	1 C	<i>c</i>	0.05182	0.00000	0.05182
$^{13}\text{C}(2)$	2 C	<i>a</i>	0.67660	0.67156	0.00505
$^{13}\text{C}(2)$	2 C	<i>b</i>	1.21097	1.21167	-0.00070
$^{13}\text{C}(2)$	2 C	<i>c</i>	0.02620	0.00000	0.02620
$^{13}\text{C}(3)$	3 C	<i>a</i>	0.67660	0.67154	0.00506
$^{13}\text{C}(3)$	3 C	<i>b</i>	1.21097	1.21168	-0.00071
$^{13}\text{C}(3)$	3 C	<i>c</i>	0.02620	0.00000	0.02620
$^{13}\text{C}(4)$	4 C	<i>a</i>	0.70882	0.70753	0.00130
$^{13}\text{C}(4)$	4 C	<i>b</i>	1.20888	1.20858	0.00030
$^{13}\text{C}(4)$	4 C	<i>c</i>	0.02564	0.00000	0.02564
$^{13}\text{C}(5)$	5 C	<i>a</i>	0.70882	0.70755	0.00128
$^{13}\text{C}(5)$	5 C	<i>b</i>	1.20888	1.20856	0.00032
$^{13}\text{C}(5)$	5 C	<i>c</i>	0.02564	0.00000	0.02564
$^{13}\text{C}(6)$	6 C	<i>a</i>	1.41308	1.41379	-0.00071

Isotopologue	Atom	Axis	Kraitchman	Fit	Difference
$^{13}\text{C}(6)$	6 C	<i>b</i>	0.00000	0.00000	0.00000
$^{13}\text{C}(6)$	6 C	<i>c</i>	0.00000	0.00000	0.00000
$^{13}\text{C}(7)$	7 C	<i>a</i>	2.80547	2.80656	-0.00109
$^{13}\text{C}(7)$	7 C	<i>b</i>	0.02098	0.00001	0.02098
$^{13}\text{C}(7)$	7 C	<i>c</i>	0.03553	0.00000	0.03553
$^{13}\text{C}(8)$	8 C	<i>a</i>	2.84256	2.84253	0.00003
$^{13}\text{C}(8)$	8 C	<i>b</i>	0.00000	0.00000	0.00000
$^{13}\text{C}(8)$	8 C	<i>c</i>	0.03179	0.00000	0.03179
$^{13}\text{C}(9)$	9 C	<i>a</i>	4.04886	4.04814	0.00072
$^{13}\text{C}(9)$	9 C	<i>b</i>	0.01219	0.00000	0.01219
$^{13}\text{C}(9)$	9 C	<i>c</i>	0.01835	0.00000	0.01835
$^{15}\text{N}(10)$	10 N	<i>a</i>	3.96474	3.96467	0.00007
$^{15}\text{N}(10)$	10 N	<i>b</i>	0.00000	0.00000	0.00000
$^{15}\text{N}(10)$	10 N	<i>c</i>	0.00000	0.00000	0.00000

Table S14: Selected structural parameters from the p-EBN MORPHEUS fit. Bond lengths are in Å; angles are in degrees.

Parameter	Description	Value	Std. error
R(1,2)	C–C	1.40330	0.01208
R(1,3)	C–C	1.40330	0.01208
R(1,7)	C–C	1.42713	0.04664
R(2,4)	C–C	1.37909	0.03102
R(2,11)	C–H	1.08012	0.00000
R(3,5)	C–C	1.37909	0.03102
R(3,12)	C–H	1.08012	0.00000
R(4,6)	C–C	1.39981	0.00002
R(4,13)	C–H	1.08006	0.00000
R(5,6)	C–C	1.39979	0.00002
R(5,14)	C–H	1.08006	0.00000
R(6,8)	C–C	1.42873	0.03574
R(7,10)	C–N	1.15811	0.03755
R(8,9)	C–C	1.20562	0.03576
R(9,15)	C–H	1.06154	0.00000
A(2,1,3)	C–C–C	119.411	1.873
A(2,1,7)	C–C–C	120.294	0.936
A(3,1,7)	C–C–C	120.295	0.936
A(1,2,4)	C–C–C	120.165	0.983
A(1,2,11)	C–C–H	119.674	0.000
A(4,2,11)	C–C–H	120.161	0.983
A(1,3,5)	C–C–C	120.165	0.983

Parameter	Description	Value	Std. error
A(1,3,12)	C–C–H	119.674	0.000
A(5,3,12)	C–C–H	120.161	0.983
A(2,4,6)	C–C–C	120.429	0.046
A(2,4,13)	C–C–H	120.212	0.046
A(6,4,13)	C–C–H	119.359	0.000
A(3,5,6)	C–C–C	120.430	0.046
A(3,5,14)	C–C–H	120.211	0.046
A(6,5,14)	C–C–H	119.359	0.000
A(4,6,5)	C–C–C	119.398	0.001
A(4,6,8)	C–C–C	120.301	0.000
A(5,6,8)	C–C–C	120.301	0.000
A(1,7,10)	C–C–N	180.000	0.000
A(6,8,9)	C–C–C	180.000	0.000
A(8,9,15)	C–C–H	180.000	0.000

S8 Azulene Comparison with the Reference Refinement

The azulene calculation used fixed C1–H11, C3–H13, and C2–H12 bond lengths; the remaining C–H distances are refined. All internal coordinates were generated algorithmically from Cartesian structures. Because azulene is planar, the fit used the independent AB moment pair and retained the C constants as diagnostics.

Table S15: Complete rotational-constant comparison for the azulene MSR-comparable subset. Corrected experimental and calculated constants are in MHz.

Isotopologue	Component	Corrected exp.	Calculated	Difference
parent	A	2862.097939	2862.075372	+0.022567
parent	B	1262.517835	1262.559239	-0.041404
parent	C	876.138923	876.087228	+0.051695
azulene-4-d1	A	2752.561850	2752.579320	-0.017470
azulene-4-d1	B	1262.294384	1262.218223	+0.076161
azulene-4-d1	C	865.456501	865.387542	+0.068959
azulene-5-d1	A	2794.488840	2794.483227	+0.005613
azulene-5-d1	B	1241.273362	1241.275392	-0.002030
azulene-5-d1	C	859.559161	859.497208	+0.061953
azulene-6-d1	A	2862.003650	2862.075372	-0.071722
azulene-6-d1	B	1223.400793	1223.394588	+0.006205
azulene-6-d1	C	857.105233	857.048897	+0.056336
azulene-4,8-d2	A	2649.532362	2649.523242	+0.009120

Isotopologue	Component	Corrected exp.	Calculated	Difference
azulene-4,8-d2	B	1262.005103	1261.903897	+0.101206
azulene-4,8-d2	C	854.871156	854.788696	+0.082460
azulene-5,7-d2	A	2725.125973	2725.133539	-0.007566
azulene-5,7-d2	B	1221.780703	1221.778074	+0.002629
azulene-5,7-d2	C	843.602156	843.573085	+0.029071
azulene-1-13C	A	2841.331150	2841.459456	-0.128306
azulene-1-13C	B	1251.448996	1251.449765	-0.000769
azulene-1-13C	C	868.849088	868.805922	+0.043166
azulene-2-13C	A	2862.093080	2862.075372	+0.017708
azulene-2-13C	B	1240.233194	1240.296442	-0.063248
azulene-2-13C	C	865.322674	865.309646	+0.013028
azulene-9-13C	A	2853.194180	2852.904191	+0.289989
azulene-9-13C	B	1261.606550	1261.641703	-0.035153
azulene-9-13C	C	874.835186	874.784969	+0.050217
azulene-4-13C	A	2821.753737	2821.770671	-0.016934
azulene-4-13C	B	1261.608919	1261.591491	+0.017428
azulene-4-13C	C	871.875087	871.811446	+0.063641
azulene-5-13C	A	2836.950530	2837.099313	-0.148783
azulene-5-13C	B	1251.104248	1251.070937	+0.033311
azulene-5-13C	C	868.277606	868.215431	+0.062175
azulene-6-13C	A	2862.033150	2862.075372	-0.042222
azulene-6-13C	B	1243.247466	1243.219286	+0.028180
azulene-6-13C	C	866.765213	866.731282	+0.033931

Table S16: Complete structural comparison between MSR and MORPHEUS results for azulene employing analogous constraints. Bond lengths are in Å; angles are in degrees.

Parameter	Unit	MORPHEUS	σ	MSR	Difference
$R(1, 9)$	Å	1.39745	0.00599	1.39900	-0.00155
$R(4, 10)$	Å	1.37923	0.01201	1.37800	+0.00123
$R(4, 5)$	Å	1.40122	0.00653	1.40100	+0.00022
$R(5, 6)$	Å	1.39054	0.00442	1.39200	-0.00146
$R(4, 14)$	Å	1.08356	0.00220	1.08700	-0.00344
$R(5, 15)$	Å	1.08133	0.00230	1.08200	-0.00067
$R(6, 16)$	Å	1.08117	0.00441	1.08200	-0.00083
$R(1, 11)$	Å	1.08116	0.00000	1.08000	+0.00116
$R(2, 12)$	Å	1.08236	0.00000	1.08100	+0.00136
$R(1, 2)$	Å	1.39593	0.00584	1.40400	-0.00807
$R(9, 10)$	Å	1.50860	0.01549	1.50000	+0.00860

Table S17: Summary of the anhydride literature-data tests. “GICs” is the number of final non-redundant coordinates generated by MORPHEUS, “Act.” is the number of active totally symmetric coordinates before projection of hard constraints, and “Eff.” is the final rank/effective number of fitted variables. Residuals are RMS/maximum absolute rotational residuals in MHz.

System	Auxiliary relation	GICs	Act.	Eff.	Cond.	Residuals
Maleic anhydride	none	21	8	8	839	0.0359/0.1453
Phthalic anhydride	weighted r_e^{BO} predicates	39	14	14	4.05	0.5013/0.8482
Succinic anhydride	fixed local H frame	27	9	5	498	0.0353/0.0731

Parameter	Unit	MORPHEUS	σ	MSR	Difference
$A(1, 9, 10)$	degree	106.13001	0.52104	106.50000	-0.36999
$A(5, 6, 7)$	degree	130.09492	0.17695	129.90000	+0.19492
$A(6, 5, 15)$	degree	115.90333	0.30686	115.70000	+0.20333
$A(4, 5, 6)$	degree	128.68339	0.00000	128.68000	+0.00339
$A(5, 4, 10)$	degree	128.79958	0.00000	128.81000	-0.01042
$A(9, 1, 11)$	degree	125.13843	0.00000	125.15000	-0.01157
$A(10, 4, 14)$	degree	115.31629	0.00000	115.31000	+0.00629
$A(1, 2, 12)$	degree	125.06740	0.00000	125.07000	-0.00260
$A(1, 2, 3)$	degree	109.86520	0.00000	109.71000	+0.15520
$A(2, 1, 9)$	degree	108.93739	0.52104	108.66000	+0.27739
$A(4, 10, 9)$	degree	127.46958	0.08847	127.58000	-0.11042
$A(5, 6, 16)$	degree	114.95254	0.08847	115.07000	-0.11746

S9 Anhydride Literature-Data Series

The three anhydrides form a homologous series used to test how MORPHEUS handles progressively less complete experimental information. In all cases the structural refinement starts from a Cartesian parent geometry and tabulated rotational constants. When B_0 and ΔB_{vib} are available, the corrected semiexperimental observables are defined directly as $B_e = B_0 + \Delta B_{\text{vib}}$, with any electronic correction added as a separate term when reported. No line list, SPFIT input, or molecule-specific coordinate model is required by the refinement step.

Maleic anhydride is the complete-data limiting case: the isotopic set is rich enough that no auxiliary structural relation is needed. Phthalic anhydride uses weighted primitive-coordinate predicates because the literature analysis itself treats the r_e^{BO} distances and angles as mixed-estimation information. Succinic anhydride has only heavy-atom substitutions; the hydrogen frame is therefore unsupported by direct isotope information. The production comparison in Table S17 uses fixed local hydrogen parameters. Runs with only soft C–H predicates reproduce the intended local predicate values but can enter a nonphysical hydrogen-frame valley before final topology reporting, which is the expected distinction between soft observations and hard removal of unsupported directions.

S9.1 Maleic Anhydride Literature-Data Reproduction

Maleic anhydride was added as a reproducibility test based entirely on the published semiexperimental dataset of Vogt, Demaison, and Rudolph. The starting geometry was reconstructed from their reported r_e^{SE} parameters, and the ten isotopologue records use the published B_0 constants and rovibrational corrections. The small electronic correction was included by using the tabulated B_e values, equivalently as $\Delta B_{\text{el}} = B_e - B_0 - \Delta B_{\text{vib}}$. No Z-matrix, manual coordinate model, constraints, predicates, or parameter classes were used.

S9.2 Input and Manifest Excerpt

The complete example distributed with the code consists only of the parent Cartesian geometry and the isotopologue records. The command used for the calculation was

```
python -m merlino semiexp \  
  --xyz examples/semiexp/maleic_anhydride/parent.xyz \  
  --observations examples/semiexp/maleic_anhydride/isotopologues.toml \  
  --outdir working/semiexp/maleic_anhydride \  
  --coordinate-model gic \  
  --observable moments \  
  --components AB \  
  --robust-loss huber
```

The starting geometry was the following Cartesian file, reconstructed from the published semiexperimental structure and used only as an initial point:

```
9  
maleic anhydride rse start reconstructed from Vogt Demaison Rudolph Struct Chem  
  2011 Table 1  
O 0.0000000000 0.0000000000 0.0000000000  
C 1.1240088423 -0.8088727727 0.0000000000  
C 0.6662000000 -2.2214374877 0.0000000000  
C -0.6662000000 -2.2214374877 0.0000000000  
C -1.1240088423 -0.8088727727 0.0000000000  
O 2.2290423505 -0.3689031082 0.0000000000  
O -2.2290423505 -0.3689031082 0.0000000000  
H 1.3567205284 -3.0472907736 0.0000000000  
H -1.3567205284 -3.0472907736 0.0000000000
```

Each isotopologue record contains the observed B_0 constants and the corrections needed to form the effective equilibrium constants. For example, the parent species is represented as

```
[[isotopologues]]  
label = "Main"  
substitutions = ""  
[isotopologues.constants]  
A_MHz = 6842.489000  
B_MHz = 2467.614000  
C_MHz = 1813.628000
```



```
[isotopologues.vibrational_correction]
delta_A_MHz = 49.780000
delta_B_MHz = 14.180000
delta_C_MHz = 11.160000
convention = "add"
[isotopologues.electronic_correction]
delta_A_MHz = 0.421000
delta_B_MHz = 0.066000
delta_C_MHz = 0.012000
convention = "add"
```

and substituted species are specified by isotope substitutions rather than by new structural models, for example

```
[[isotopologues]]
label = "13C2d2"
substitutions = "2:13;8:2;9:2"
```

The full output manifest is archived with the example. The excerpt below shows the information needed to audit model construction and the absence of manual constraints:

```
[method]
program = MORPHEUS
method = semiexperimental equilibrium-geometry least squares
coordinate_model = gic
coordinate_basis = MORPHEUS non-redundant symmetry-adapted GICs; active subspace
    is totally symmetric
observable = moments
components = Ia,Ib
isotopologues = Main, d2, 13C2, 13C3, 1801, 1806, 13C2d2, 13C3d2, 1801d2, 1806d2

[constraints]
fixed_parameter = none
parameter_class = none
qm_predicate = none

[constraint_diagnostics]
input_records = 0
input_expression_constraints = 0
input_coordinate_definitions = 0
symmetry_expanded_fixed_primitives = 0
parameter_classes = 0
```

The final diagnostic excerpt reports the rank-consistent fitted model:

```
[fit_statistics]
convergence = step_tolerance
iterations = 71
n_optimized_parameters = 8
rank = 8
```

Table S18: Maleic anhydride refinement diagnostics. The GIC fit used moments of inertia for the independent AB planar pair; the omitted C constants are reported as diagnostics.

Quantity	Value
Generated non-redundant GICs	21
Totally symmetric A_1 coordinates	8
Independent fitted variables	8
Rank	8
Condition number	838.856
Rotational RMS residual / MHz	0.03593
Maximum absolute residual / MHz	0.14526
Iterations	71

```

condition_number = 838.856024878
robust_loss = huber
coordinate_builder_calls = 1

[rotational_residual_statistics]
n_rotational_constants = 30
rotational_rms_MHz = 0.0359332113823
rotational_max_abs_MHz = 0.145260439303

[working_coordinates]
coordinate_count = 21
index active class value sigma label
1 yes - 3.22390949563 0.000251446997267 GIC001 MORPHEUS A1Str0001 irrep=A1
...
8 yes - -0.00433581007571 0.000311593700452 GIC008 MORPHEUS A1Ang0003 irrep=A1
9 no - -2.02080870849 0 GIC009 MORPHEUS A2Tor0001 irrep=A2
...
21 no - 6.53687463342e-16 0 GIC021 MORPHEUS B2Ang0003 irrep=B2

```

Table S19: Complete rotational-constant comparison for maleic anhydride. Corrected experimental and calculated constants are in MHz. The corrected experimental constants reproduce the published B_e values.

Isotopologue	Constant	Corrected exp.	Calculated	Difference
Main	A	6892.6900	6892.7047	-0.0147
Main	B	2481.8600	2481.8511	+0.0089
Main	C	1824.8000	1824.7976	+0.0024
d ₂	A	6148.0000	6148.0220	-0.0220
d ₂	B	2437.5100	2437.5096	+0.0004
d ₂	C	1745.4800	1745.4787	+0.0013

Table S20: Selected maleic anhydride structural parameters from the MORPHEUS reproduction compared with the published r_e^{SE} structure. Bond lengths are in Å; angles are in degrees.

Parameter	Atoms	MORPHEUS	Published	Difference
O–C	1–2	1.38489	1.3848	+0.00009
C–C	2–3	1.48479	1.4849	-0.00011
C=C	3–4	1.33241	1.3324	+0.00001
C=O	2–6	1.18945	1.1894	+0.00005
C–H	3–8	1.07652	1.0765	+0.00002
C–O–C	2–1–5	108.513	108.52	-0.007
O–C–C	1–2–3	107.784	107.78	+0.004
O–C–O	1–2–6	122.537	122.55	-0.013
C–C–O	3–2–6	129.679	129.67	+0.009
C–C–H	4–3–8	129.896	129.90	-0.004

Isotopologue	Constant	Corrected exp.	Calculated	Difference
$^{13}\text{C}2$	A	6891.4300	6891.4558	-0.0258
$^{13}\text{C}2$	B	2466.6500	2466.6492	+0.0008
$^{13}\text{C}2$	C	1816.4800	1816.4793	+0.0007
$^{13}\text{C}3$	A	6739.8700	6739.8341	+0.0359
$^{13}\text{C}3$	B	2476.4200	2476.4210	-0.0010
$^{13}\text{C}3$	C	1811.0000	1811.0031	-0.0031
$^{18}\text{O}1$	A	6738.2200	6738.1958	+0.0242
$^{18}\text{O}1$	B	2481.8500	2481.8511	-0.0011
$^{18}\text{O}1$	C	1813.7800	1813.7867	-0.0067
$^{18}\text{O}6$	A	6839.8800	6839.9516	-0.0716
$^{18}\text{O}6$	B	2367.8700	2367.8784	-0.0084
$^{18}\text{O}6$	C	1758.9400	1758.9566	-0.0166
$^{13}\text{C}2\text{d}_2$	A	6146.3100	6146.1647	+0.1453
$^{13}\text{C}2\text{d}_2$	B	2422.8100	2422.8401	-0.0301
$^{13}\text{C}2\text{d}_2$	C	1737.7900	1737.7951	-0.0051
$^{13}\text{C}3\text{d}_2$	A	6033.8400	6033.8538	-0.0138
$^{13}\text{C}3\text{d}_2$	B	2432.2700	2432.2723	-0.0023
$^{13}\text{C}3\text{d}_2$	C	1733.4900	1733.4936	-0.0036
$^{18}\text{O}1\text{d}_2$	A	6013.3300	6013.3614	-0.0314
$^{18}\text{O}1\text{d}_2$	B	2437.5200	2437.5096	+0.0104
$^{18}\text{O}1\text{d}_2$	C	1734.4500	1734.4516	-0.0016
$^{18}\text{O}6\text{d}_2$	A	6099.5800	6099.4994	+0.0806
$^{18}\text{O}6\text{d}_2$	B	2327.3500	2327.3468	+0.0032
$^{18}\text{O}6\text{d}_2$	C	1684.5800	1684.5745	+0.0055

Table S21: Testosterone T1-H2⁻-gp rotational constants used for the large-molecule scale check. Constants and residuals are in MHz. The QC column is the unrefined rotational constant calculated from the input geometry; the class-refined column is the parent-only MORPHEUS result with a single C–C skeletal stretch class.

Constant	B_0	ΔB_{vib}	B_e^{SE}	QC	Class-refined	Final diff.
<i>A</i>	801.5	8.1	793.4	788.762	793.427	-0.027
<i>B</i>	171.9	1.6	170.3	169.626	169.667	+0.633
<i>C</i>	156.6	1.4	155.2	154.637	154.909	+0.291

S10 Testosterone Large-Molecule Scale Check

The testosterone calculation in the main text is not an isotope-rich semiexperimental structure determination. It is a scale and automation check using the published data of Uribe et al. for the detected T1-H2⁻-gp conformer. Its purpose is to document the full large-molecule chain: the QC protocol provides an accurate equilibrium reference geometry, the accelerated ΔB_{vib} workflow makes vibrational corrections available at this size, and MORPHEUS adds the algorithmic refinement-model layer. The only experimental input is the parent ground-state rotational constant set. The semiexperimental target is obtained by subtracting the B3 vibrational corrections reported in the same table:

$$B_e^{\text{SE}} = B_0 - \Delta B_{\text{vib}}.$$

Thus the data used by MORPHEUS are $B_0 = (801.5, 171.9, 156.6)$ MHz and $\Delta B_{\text{vib}} = (8.1, 1.6, 1.4)$ MHz for (*A*, *B*, *C*), giving $B_e^{\text{SE}} = (793.4, 170.3, 155.2)$ MHz.

The calculation starts from the QC Cartesian geometry and does not require a Z-matrix, hand-selected internal coordinates, or a stored refinement model. Topology perception identifies the C₁₉H₂₈O₂ molecular graph, MORPHEUS builds the 141-coordinate non-redundant GIC space, and the MORPHEUS class layer applies the reduced-dimensionality model after symmetry and rank reduction. For this parent-only diagnostic, all angular, torsional, and out-of-plane GIC families are fixed. Stretch combinations involving C–H, O–H, and C–O primitives are also fixed. The retained C–C skeletal stretch block is then assigned to one shared parameter class, so the fit contains one effective variable. This is a hard class constraint: it does not add an observation or a statistical prior, but replaces the selected active coordinates by a single shared displacement direction before the least-squares normal equations are formed.

```
parameter_class = CC_skeleton; mode=shared;
patterns = R(1,2)|R(1,6)|R(2,3)|R(2,45)|R(3,4)|R(3,10)|...

parameter_class = non_CC_stretches; mode=fixed;
patterns = all retained stretch combinations involving C-H, O-H, or C-O

parameter_class = Angles_fixed; mode=fixed; patterns=AAng
parameter_class = Torsions_fixed; mode=fixed; patterns=ATor
parameter_class = OOP_fixed; mode=fixed; patterns=AOop
```

Table S22: Testosterone model-construction diagnostics. The class-refined calculation used rotational constants as observables and all three parent components.

Quantity	Value
Atoms	49
Non-redundant GICs	141
Retained active C–C stretch combinations	5
Effective fitted variables	1
Rank	1
Condition number	1.0
Unrefined QC RMS residual / MHz	2.725
Class-refined RMS residual / MHz	0.402
Iterations	5

The important point is that the class is applied in the final working basis, not as a molecule-specific Z-matrix instruction. The user supplies Cartesian coordinates and the literature rotational data; the topology labels determine which retained GICs belong to the C–C skeletal class.

The exact repository command used for this diagnostic is archived in the testosterone example directory as `run_testosterone_cc_class.sh`. It can be run from the repository root and reads the QC Cartesian geometry, the published parent rotational constants, and the published vibrational corrections:

```
sh examples/semiexp/testosterone/run_testosterone_cc_class.sh
```

S11 Characteristics of the Test Cases

Table S23: Constraints and coordinate choices (symmetry-adapted internal or Cartesian coordinates, SAI and SAC, respectively) used for the benchmark calculations.

Case	Constraints	Coordinates
Glycolaldehyde	No fixed primitive constraints	SAI, ABC
Cyclopentadiene	No fixed primitive constraints	SAI and SAC, ABC
Nitrobenzene	No fixed primitive constraints	SAI, AB
p-EBN	Legacy frozen Z-matrix parameters	SAI, AB
Azulene	C–H bond lengths and selected angles	SAI, AB
Norcamphor	H-atom constraints	SAI and SAC, ABC
Glycine I	Four generated QC-reference constraints	SAI, ABC
Glycine II	Five generated QC constraints	SAI, ABC
Maleic anhydride	No fixed primitive constraints	SAI and SAC, AB
Phthalic anhydride	Weighted r_e^{BO} predicates	SAI, AC
Succinic anhydride	Fixed local H-frame parameters	SAI, AC
Testosterone	One shared C–C skeletal stretch class	SAI, ABC