47

EE22BTECH11059

47.2023 Let (X, Y) have joint probability mass function

$$p(x, y) = \begin{cases} \frac{c}{2^{x+y+2}} & if\, x = 0, 1, 2, \dots y = 0, 1, 2, \dots; x \neq y \\ 0 & otherwise \end{cases}$$

(1)

Then which of the following is true?

1) $c = \frac{1}{2}$
2) $c = \frac{1}{4}$
3) $c > 1$
4) X and Y are independent

Solution: For p(x, y) to be joint probability mass function

$$\sum_{y=-\infty}^{\infty} \sum_{x=-\infty}^{\infty} p(x, y) = 1 \,|x \neq y \qquad (2)$$

$$\sum_{y=0}^{\infty} \sum_{x=0}^{\infty} \frac{c}{2^{x+y+2}} - \sum_{x=y} \frac{c}{2^{x+y+2}} = 1 \qquad (3)$$

$$\sum_{y=0}^{\infty} \frac{c}{2^{y+2}} \sum_{x=0}^{\infty} 2^{-x} - \frac{c}{4} \sum_{x=0}^{\infty} \frac{1}{4^x} = 1 \qquad (4)$$

$$\sum_{y=0}^{\infty} \frac{2c}{2^{y+2}} - \frac{c}{3} = 1 \qquad (5)$$

$$\frac{2c}{4} \sum_{y=0}^{\infty} 2^{-y} - \frac{c}{3} = 1 \qquad (6)$$

$$c - \frac{c}{3} = 1 \qquad (7)$$

$$c = \frac{3}{2} \qquad (8)$$

1) Marginal probability mass function of X

$$p_X(x) = \sum_{y=0}^{\infty} p(x, y) \qquad (9)$$

$$= \sum_{y=0}^{\infty} \frac{3}{2^{x+y+3}} \qquad (10)$$

$$= \frac{3}{2^{x+3}} \sum_{y=0}^{\infty} 2^{-y} \qquad (11)$$

$$= \frac{3}{2^{x+2}} \qquad (12)$$

2) Marginal probability mass function of Y

$$p_Y(y) = \sum_{x=0}^{\infty} p(x, y) \qquad (13)$$

$$= \sum_{x=0}^{\infty} \frac{3}{2^{x+y+3}} \qquad (14)$$

$$= \frac{3}{2^{y+3}} \sum_{x=0}^{\infty} 2^{-x} \qquad (15)$$

$$= \frac{3}{2^{y+2}} \qquad (16)$$

For X and Y to be independent,

$$p(x, y) = p_X(x)\, p_Y(y) \qquad (17)$$

$$p_X(x)\, p_Y(y) = \frac{3}{2^{x+2}} \frac{3}{2^{y+2}} \qquad (18)$$

$$= \frac{9}{2^{x+y+4}} \qquad (19)$$

$$p(x, y) \neq p_X(x)\, p_Y(y) \qquad (20)$$

Option (4) is incorrect.
$\therefore$ Only option (3) is correct.
Joint cumulative distribution function F(x, y)

$$= P(X \leq x, Y \leq y) \qquad (21)$$

$$= \sum_i \left(\sum_j p(i, j) \right) |x \neq y \qquad (22)$$

1) x<y

$$F(x, y) = \sum_{i=0}^{x} \sum_{j=1}^{i} p(i, j) \tag{23}$$

$$= \sum_{i=0}^{x} \sum_{j=1}^{i} \frac{3}{2^{i+j+3}} \tag{24}$$

$$= \frac{3}{8} \sum_{i=0}^{x} \frac{1}{2^i} \sum_{j=1}^{i} 2^{-j} \tag{25}$$

$$= \frac{3}{8} \sum_{i=0}^{x} \frac{1}{2^i} \frac{\left(1 - \frac{1}{2^i}\right)}{\frac{1}{2}} \tag{26}$$

$$= \frac{3}{4} \left(\sum_{i=0}^{x} \frac{1}{2^i} - \frac{1}{4^i} \right) \tag{27}$$

$$= \frac{3}{4} \left(\left(\frac{1 - \frac{1}{2^{x+1}}}{\frac{1}{2}} \right) - \left(\frac{1 - \frac{1}{4^{x+1}}}{\frac{3}{4}} \right) \right) \tag{28}$$

$$= \frac{3}{2} - \frac{3}{2^{x+2}} - 1 + \frac{1}{2^{2x+2}} \tag{29}$$

$$= \frac{1}{2} + \frac{1}{2^{2x+2}} - \frac{3}{2^{x+2}} \tag{30}$$

2) x>y

$$F(x, y) = \sum_{j=0}^{y} \sum_{i=1}^{j} p(i, j) \tag{31}$$

$$= \sum_{j=0}^{y} \sum_{i=1}^{j} \frac{3}{2^{i+j+3}} \tag{32}$$

$$= \frac{3}{8} \sum_{j=0}^{y} \frac{1}{2^j} \sum_{i=1}^{j} 2^{-i} \tag{33}$$

$$= \frac{3}{8} \sum_{j=0}^{y} \frac{1}{2^j} \frac{\left(1 - \frac{1}{2^j}\right)}{\frac{1}{2}} \tag{34}$$

$$= \frac{3}{4} \left(\sum_{j=0}^{y} \frac{1}{2^j} - \frac{1}{4^j} \right) \tag{35}$$

$$= \frac{3}{4} \left(\left(\frac{1 - \frac{1}{2^{y+1}}}{\frac{1}{2}} \right) - \left(\frac{1 - \frac{1}{4^{y+1}}}{\frac{3}{4}} \right) \right) \tag{36}$$

$$= \frac{3}{2} - \frac{3}{2^{y+2}} - 1 + \frac{1}{2^{2y+2}} \tag{37}$$

$$= \frac{1}{2} + \frac{1}{2^{2y+2}} - \frac{3}{2^{y+2}} \tag{38}$$