

Twelve Unit Cubes in a Cube of Side 2.9315, Found by Hardening Balls into Cubes

Yohei Nakajima*

8 October 2026

Abstract

We give a packing of 12 unit cubes in a cube of side 2.9315185, with every pair and every wall at least 10^{-6} apart, improving the previous best known side 2.9327717687 (H. Lin, July 2026 [8]). The packing was found by a soft-to-rigid homotopy: unit-diameter balls are compressed in a shrinking box and then deformed through rounded cubes into rigid unit cubes while inward pressure on the box continues. The final configuration is tightened with a constrained solver that keeps every corner inside the box and a separating plane between every nearby pair, and validated by an exact rational certificate. On unit squares in a square, the same method recovers the best known packings for $n = 11$ (now proved optimal), 17, 18, 19, 26 and 27, and outperforms a rigid-start control on the same seeds. We report where the method fails ($n = 28, 29$ squares; $n = 10, 11$ cubes so far), an ablation showing that an oriented intermediate shape does not help, and code to reproduce every result.

1 Introduction

Let $s_d(n)$ be the side of the smallest d -dimensional cube that contains n non-overlapping unit cubes, free to rotate. For squares ($d = 2$) the problem has a long record history [1, 7, 3, 9]; the smallest unsettled case, $n = 11$, was recently closed [9]. For cubes ($d = 3$) far fewer results exist. Friedman’s catalogue [2] lists non-trivial packings for $n = 9$ –14 and 28–33; the entries for $n = 11$ and 12 were improved in 2026 (T. Berthold et al. in March for $n = 11$; H. Lin in July for $n = 11$ and 12, with coordinates published in [8]), and for every other $n < 34$ the trivial packing is the best known.

Contribution.

1. A packing of 12 unit cubes in a cube of side 2.9315185094797 with every pair and every wall separated by at least 10^{-6} (§3), 0.0012533 below the previous record 2.9327717687048653. It is certified in exact rational arithmetic.
2. A simple search method, soft-to-rigid homotopy with inward pressure (§2), and an exact tightening step that turns its output into a locally optimal packing to 10^{-12} .
3. An empirical record of what the method does and does not reach: in 2D it recovers six published records from random starts and beats a rigid-start control; it fails on $n = 28, 29$ squares; an oriented intermediate shape does not raise its hit rate (§4, §5).

This is a search result, not an optimality proof.

*yohei@untapped.vc. Experiments, code and verification were developed with the assistance of Claude (Anthropic); see the acknowledgements.

Related methods. Gensane and Ryckelynck’s perturbation–compression algorithm found dense square packings, including $n = 11$, from random starts [4]. Recent square records ($n = 28, 29, 39, 41, 50, 51, 55$) come from rigid simulated annealing and exact refinement [9]. Our method differs in starting from balls and changing the shapes, not just their positions, during the search.

2 Method

Shape family. A *rounded cube* of softness $r \in [0, \frac{1}{2}]$ is a cube of half-side $\frac{1}{2} - r$ dilated by a ball of radius r (the Minkowski sum). At $r = \frac{1}{2}$ it is a unit-diameter ball, at $r = 0$ a unit cube, and every member lies inside the unit cube with the same centre and orientation. Hardening ($r \downarrow 0$) therefore only grows the shapes, and the container must expand to accommodate them. The square version uses rounded squares.

Energy and pressure. For centres c_i , orientations q_i (quaternions in 3D, angles in 2D) and container side L ,

$$E = \sum_{i < j} (2r - \text{sd}(i, j))_+^2 + \sum_i \sum_{\text{walls}} (\text{violation})_+^2 + \mu L,$$

where $\text{sd}(i, j)$ is the signed distance between the cores (cube of half-side $\frac{1}{2} - r$): exact Euclidean distance when separated (vertex–face and edge–edge features), and minus the separating-axis penetration depth when overlapping (15 axes for cubes [5]). The term μL is inward pressure; the box resizes symmetrically so all walls push. Positions, orientations and L follow momentum gradient descent with analytic gradients (checked against finite differences on 72,000 random cube-pair gradient components and 120,000 square-pair components).

Schedule. Compress balls under pressure 3μ (5,000 steps); morph $r : \frac{1}{2} \rightarrow 0$ linearly at pressure $\mu = 0.02$ (45,000 steps); settle rigid cubes while μ decays to $2 \cdot 10^{-6}$ (3,000 steps). Optional *annealing* adds Gaussian kicks to positions and angles that decay to zero during compression and morphing. The *rigid control* runs the identical schedule with $r = 0$ throughout.

Legalization. Penalty pressure tolerates small overlaps. Each run is legalized by scaling centres apart about their mean (bisection to the smallest factor at which no pair overlaps by more than 10^{-12} under the separating-axis test) and measuring the tight bounding cube. All sides reported below are legal packings.

Exact tightening. Runs within a threshold of the catalogue are passed to a constrained solver: minimize L over centres, rotations and per-pair separating planes $u_{ij} \cdot p = c_{ij}$, subject to every corner of every cube lying in $[0, L]^3$ and the corners of i and j lying on opposite sides of their plane. We use SLSQP [6] in outer iterations with a trust region, rebuilding the pair list and re-basing rotations each iteration, and accept a step only if the independently legalized side decreases. From near-miss runs the solver converges to Trump’s $s_2(11)$ [10] and Bidwell’s $s_2(17)$ [3] to 10^{-12} , and from a perturbed copy to Friedman’s $s_3(9) = 2 + 1/\sqrt{2}$.

Exact certificate. For a reported 3D packing we expand centres by a factor $1 + 10^{-6}$ about their mean and inflate each half-side to $\frac{1}{2}(1 + 10^{-12})$, which covers floating-point non-orthogonality of the axes. We convert every coordinate to an exact rational, compute all corners exactly, and check (i) every corner lies in $[0, S]^3$ for a rational S and (ii) every pair is strictly separated by an explicit rational plane. A linear program proposes each plane; the check itself uses only exact arithmetic.




Figure 1: Twelve unit cubes in a cube of side 2.931515. Blue: tilt below 1.5° ; red: tilted.

3 Twelve cubes

Theorem 1. *Twelve unit cubes fit in a cube of side 2.9315185094797 with pairwise and wall clearance at least 10^{-6} . Hence $s_3(12) \leq 2.9315185094797 < 2.9327717687048653$.*

Certificate. The claim file (`claims/cubincub_n12/cubincub_n12.json`, Table 3) lists one pose $[x, y, z, q_w, q_x, q_y, q_z]$ per cube, the format of the previous record file [8]. It was obtained from the tightened packing (side 2.931514577965, with touching contacts) by spreading centres by a factor $1 + 1.0 \cdot 10^{-6}$ until every pair is separated by 10^{-6} , adding a 10^{-6} wall gap and rounding the side up. Two checks pass. (i) A floating-point 15-axis separating-axis test (`verify.py`, 30 lines): minimum wall gap $1.000000000 \cdot 10^{-6}$, minimum pair gap $1.000000001 \cdot 10^{-6}$. (ii) An exact check (`certify_exact.py`): every number is converted to a rational, rotations are the exactly orthogonal $H(q)/|q|^2$, every corner lies in $[0, s]^3$ and every pair is strictly separated by an explicit rational plane.

Four cubes are exactly axis-aligned, two are tilted by less than 1° , and six are tilted by 9° or more (Figure 1). Several quaternions have repeated components (e.g. cube 1: $(0.58389, -0.58389, -0.39884, -0.39884)$), suggesting a closed form that we have not derived. We have not derived it.

How it was found. Seed 52 of the first 64 ball $\rightarrow$ cube starts with annealing ended at legalized side 2.931821; exact tightening lowered it to 2.931514577965. A second start (seed 83) ended at 2.931910 and tightens to 2.931619541: a distinct local optimum, also below the catalogue. Two of 192 starts land below 2.93277. The rigid control’s best of 32 starts is 2.99257.

Other cube counts. Table 1 summarizes the first 3D screen. The method matches Friedman’s $n = 9$ packing often, settles in worse arrangements at $n = 10$ and 11, and is the only setup that comes near the catalogue at $n = 12$.

4 Validation on squares

We developed the method on unit squares, where records are dense and recently audited [9, 3].

n	catalogue	starts (ball / rigid)	best ball→cube	best rigid	matches (ball / rigid)
9	2.70711	64 / 32	2.707107	2.707107	16 / 5
10	2.70711	64 / 32	2.767767	2.894428	0 / 0
11	2.88295	64 / 32	2.894427	2.918992	0 / 0
12	2.93277	192 / 32	2.931515	2.992573	—
13	2.956	48 / —	2.997184	—	0 / —
14	2.98995	42 / —	3.000000	—	0 / —

Table 1: First 3D screen (annealed ball→cube vs. rigid control, 53,000 steps per start; best after exact tightening). The best $n = 10$ run is within $2 \cdot 10^{-7}$ of $1 + 5\sqrt{2}/4$; the tightened best $n = 11$ run equals $2 + 2/\sqrt{5} \approx 2.894427$ to 10^{-9} , below the March 2026 value but above Lin’s 2.8829529.

$n = 11$. With annealing and the long schedule, 6 of 160 random starts reach Trump’s packing (tilt 40.18°); 0 of 120 rigid-control starts do (best 3.8877, median 4.000). Trump’s packing has since been proved optimal [9], so this is a sanity check.

$n = 2$ –30 **at a fixed schedule**. Among cases whose best known packing has tilted squares, the method matches $n = 5, 10, 11$ and none from 17 upward at the standard step budget; the rigid control matches 5 and 10. Grid cases split by slack: when the grid has empty cells ($n = 6, 7, 12, 13, 20, 21$) rigid starts fall into it more often (46–100% vs. 2–19%), while for full or nearly full grids ($n = 9, 15, 16, 22$ –25, 30) rigid starts jam at tilts they cannot rotate out of (e.g. $n = 25$: rigid 0/24, balls 7/48).

n	best known (finder)	recovered to	hit rate
17	4.675530 (Bidwell 1998)	10^{-12}	10/960 blob starts; 0/80 rigid
18	4.822876 (Hämäläinen 1980)	10^{-5}	2/24
19	4.885618 (Wainwright 1979)	10^{-12}	2/960
26	5.621320 (Friedman 1997)	10^{-12}	2.5% / 6.3% / 10.4% at $3\times/5\times/10\times$ steps
27	5.707107 (Göbel 1979)	10^{-12}	1/416
28	5.824445 (Ellsworth 2025)	—	best 5.844498
29	5.933833 (Schadt 2025)	—	best 5.938996

Table 2: Squares: best known packings reached from random starts. No run went below a catalogue value in about 5,000 starts at $n \geq 17$.

Old records recovered with more steps and exact tightening. Exact tightening matters: raw runs stop about 10^{-4} above the record because of penalty slack, and the solver also pulls runs from up to 0.022 away into the record basin ($n = 27$). After tightening, every arrangement within 0.001 of the catalogue at $n = 17$ and 19 was the catalogue packing itself.

5 Ablations and negative results

Oriented intermediate shape. Disks carry no orientation, and on the $n = 11$ hit the angles lock in only once $r < 0.06$. We tested morphing through regular octagons cut back to squares, which exert torque from the start. On the same seeds and steps ($n = 17, 19$, seeds 1–160), octagons hit 0/320 vs. 4/320 for annealed disks (one-sided $p \approx 0.06$), with no better medians and $3\times$ the cost. Late orientation commitment appears to help rather than hurt.

Penalty polish. A fixed-box squeeze-and-shake stage changed sides by at most 0.009 and never changed the arrangement; it is superseded by exact tightening.

Scaling. Hit rates fall with n at a fixed budget. At $n = 26$ they rise with hardening time; at $n = 27$ – 29 they do not.

6 Discussion

The method’s value is the basin it lands in: starting from balls, configurations are dense before orientation matters, and rotation is decided only as corners appear. For 3D, where the catalogue is sparse, this was enough to improve $n = 12$: 2 of 192 starts land below the record. At $n = 13$ and 14 (48 and 42 starts, no rigid control) no run came within 0.04 and 0.01 of the catalogue; the best $n = 14$ run is the trivial side 3.

Limitations. (i) No optimality claims. (ii) The catalogue values for cubes were read from Friedman’s page images, and Lin’s packings are not described in text; 2.93277 is truncated (“2.93277+”), which does not affect a 0.00126 margin. (iii) Hit rates are from modest sample sizes and are reported with counts.

Reproducibility. All code, seeds, run logs and certificates are at <https://github.com/yoheinakajima/soft-to-rigid-packing>. `scripts/reproduce_n12.sh` regenerates the $n = 12$ packing and its certificate in a few minutes on a laptop.

Acknowledgements. Code, experiments and verification were developed with Claude (Anthropic), an AI system, under the author’s direction. The author takes responsibility for the content.

References

- [1] Paul Erdős and Ronald L. Graham. On packing squares with equal squares. *Journal of Combinatorial Theory, Series A*, 19:119–123, 1975.
- [2] Erich Friedman. Cubes in cubes. Erich’s Packing Center, <https://erich-friedman.github.io/packing/cubincub/>, . Updated 3 July 2026; accessed 8 October 2026.
- [3] Erich Friedman. Packing unit squares in squares: a survey and new results. *Electronic Journal of Combinatorics*, Dynamic Survey DS7, . Squares-in-squares catalogue now maintained by D. Ellsworth, https://kingbird.myphotos.cc/packing/squares_in_squares.html.
- [4] Thierry Gensane and Philippe Ryckelynck. Improved dense packings of congruent squares in a square. *Discrete & Computational Geometry*, 34:97–109, 2005.
- [5] Stefan Gottschalk, Ming C. Lin, and Dinesh Manocha. OBBTree: a hierarchical structure for rapid interference detection. In *Proceedings of SIGGRAPH ’96*, pages 171–180, 1996.
- [6] Dieter Kraft. A software package for sequential quadratic programming. Technical Report DFVLR-FB 88-28, DFVLR, 1988.
- [7] Walter Stromquist. Packing 10 or 11 unit squares in a square. *Electronic Journal of Combinatorics*, 10:R8, 2003.

- [8] Tencent Hunyuan. Hyra-results: packing records. https://github.com/Tencent-Hunyuan/Hyra-results/tree/main/AI4Science/packing_records. File `cubincub_n12.json`, side 2.9327717687048653; accessed 8 October 2026.
- [9] The Squares Project. An atlas of the square packing problem. <https://jlevy.github.io/squares/>. Accessed 8 October 2026.
- [10] Walter Trump. Packing 11 unit squares in a square. <https://trump.de/square-packing/>, 1979.

A Coordinates for $n = 12$

Centres and unit quaternions (w, x, y, z) ; cube i occupies $c_i + R(q_i)[- \frac{1}{2}, \frac{1}{2}]^3$ in the box $[0, 2.9315185094797]^3$. Full-precision values: `claims/cubincub_n12/cubincub_n12.json`.

Table 3: Coordinates of the $n = 12$ packing.

i	x	y	z	w	q_x	q_y	q_z
1	1.5243900087	2.4315173049	1.3320181026	0.5838886869	-0.5838886869	-0.3988408220	-0.3988408220
2	2.4315175095	2.4315173049	0.5000010000	0.7071067812	0.7071067812	-0.0000000000	0.0000000000
3	0.5000010000	1.4369556928	1.4377218077	0.5395289954	0.4570650535	0.4570650535	-0.5395289954
4	0.5098745912	0.5769319953	2.3545775829	0.4560511351	-0.5403863020	-0.4560472394	0.5403896022
5	0.6086104699	2.2929309153	2.3887769910	0.0258332833	0.7609821302	-0.6482073268	-0.0081302199
6	2.4315175095	1.4315163049	0.5000010000	0.0000000000	-0.0000000000	-0.0000000000	-1.0000000000
7	2.4298431049	2.4215276465	2.2620542416	0.0000269579	0.0016621483	-0.9999712786	0.0073944948
8	1.5800409188	1.4806927085	2.4315175095	0.0000000000	-0.9765112947	-0.2154662186	-0.0000000000
9	1.6870691741	0.7046058868	1.3931640729	0.1630483719	-0.8266039691	-0.3023204668	0.4458065074
10	0.6714203714	0.5125513149	0.5042794969	0.0019279356	0.0017767559	-0.7100527881	-0.7041435680
11	2.4315175095	0.5648571561	2.4315175095	0.5000000000	-0.5000000000	-0.5000000000	0.5000000000
12	0.5000010000	2.3777515563	0.5000010000	0.7071067812	-0.7071067812	0.0000000000	0.0000000000