PH3244 — PHYSICS LABORATORY - IV

Yojit Kumar

Reg. no.: 20231289

Batch: (Friday)

Lab Partner: Nachiketa Kulkarni

Index of Experiments

#	Name of Experiment	Page	Date	
			Conducted	Submitted

Experiment No. 1

Quincke's Method

Date: 23 January, 2026

1.1 Aim

The aim of the experiment is to measure the magnetic susceptibility of the salt solution based on the principle that in a non-uniform magnetic field, magnetic dipoles within the liquid align with the field. This creates a force that pulls the liquid into the region of the stronger field (between the magnet poles), causing the liquid level to rise against gravity.

1.2 Theory

When an object is placed in a magnetic field, a magnetic moment is induced in it. The magnetic moment can then be measured by measuring the force exerted by the sample in an inhomogeneous magnetic field. The force on the sample is negative of the gradient of the change in energy density when the sample is placed. And a simplified expression can be written as:

$$\chi = \frac{2\rho g h}{\mu_0 H^2}$$

1.3 Experimental Observation

1.3.1 Preparation of the Solution

Solution - I

Weight of salt($MnSO_4.H_2O$) = 5.0 g
Volume of water added = 30 mL
Molarity of the salt =

$$\left(\frac{5.0}{169.02}\frac{1000}{30}\right) = 0.99\,\mathrm{M}$$

Solution - II

Weight of salt($MnSO_4.H_2O$) = 7.5 g
Volume of water added = 15 mL
Molarity of the salt =

$$\left(\frac{7.5}{169.02}\frac{1000}{15}\right) = 2.96\,\mathrm{M}$$

1.3.2 Calibration Data

Table 1.1: Calibration Data (Increasing Current)

#	Current I (A)	Magnetic Field B (10G)	B^2 (G^2)
1	0.00	40	160000
2	0.10	68	462400
3	0.40	159	2528100
4	0.70	266	7075600
5	1.00	374	13987600
6	1.30	478	22848400
7	1.60	575	33062500
8	1.90	670	44890000
9	2.20	762	58064400
10	2.50	848	71910400
11	2.80	931	86676100

Table 1.2: Calibration Data (Decreasing Current)

#	Current I (A)	Magnetic Field B (10G)	B^2 (G^2)
1	2.80	931	86676100
2	2.50	866	74995600
3	2.20	790	62410000
4	1.90	707	49984900
5	1.60	619	38316100
6	1.30	525	27562500
7	1.00	423	17892900
8	0.70	315	9922500
9	0.40	198	3920400
10	0.10	85	722500
11	0.00	46	211600

1.3.3 Microscope Measurment

Table 1.3: Microscope Readings for SolOne Inc

#	Current (A)	Field (G)	Readings (in mm)	Height Difference (in mm)
1	0.00	300.68	17.63	0.28
2	1.90	6657.42	17.35	0.00
3	2.20	7580.85	19.34	1.99
4	2.35	8034.36	19.45	2.10
5	2.50	8482.39	19.54	2.19
6	2.65	8924.95	19.63	2.28
7	2.80	9362.04	19.72	2.37

Table 1.4: Microscope Readings for SolOne Dec

#	Current (A)	Field (G)	Readings (in mm)	Height Difference (in mm)
1	2.80	9337.36	19.72	2.43
2	2.65	8997.67	19.66	2.37
3	2.50	8642.40	19.60	2.31
4	2.35	8271.54	19.53	2.24
5	2.20	7885.11	19.43	2.14
6	2.00	7345.64	19.30	2.01
7	1.90	7065.52	19.24	1.95
8	1.75	6632.35	19.15	1.86
9	1.60	6183.61	19.03	1.74
10	1.45	5719.28	18.87	1.58
11	1.30	5239.38	18.68	1.39
12	1.15	4743.90	18.64	1.35
13	1.00	4232.84	18.40	1.11
14	0.85	3706.20	18.05	0.76
15	0.70	3163.98	17.83	0.54
16	0.54	2568.45	17.59	0.30
17	0.40	2032.81	17.43	0.14
18	0.25	1443.86	17.37	0.08
19	0.00	427.65	17.29	0.00

Table 1.5: Microscope Readings for SolTwo Inc

#	Current (A)	Field (G)	Readings (in mm)	Height Difference (in mm)
1	0.00	300.68	16.82	0.00
2	0.15	834.45	16.93	0.11
3	0.30	1362.75	17.08	0.26
4	0.45	1885.58	17.29	0.47
5	0.60	2402.93	17.65	0.83
6	0.75	2914.81	18.09	1.27
7	0.90	3421.22	18.60	1.78
8	1.05	3922.16	19.17	2.35
9	1.20	4417.62	19.82	3.00
10	1.35	4907.61	20.40	3.58
11	1.50	5392.13	20.71	3.89
12	1.65	5871.17	21.16	4.34
13	1.80	6344.75	21.75	4.93
14	2.00	6967.66	22.07	5.25
15	2.20	7580.85	22.32	5.50
16	2.40	8184.31	23.55	6.73
17	2.60	8778.04	24.65	7.83

Table 1.6: Microscope Readings for SolTwo Dec

#	Current (A)	Field (G)	Readings (in mm)	Height Difference (in mm)
1	2.50	8642.40	25.19	8.06
2	2.30	8144.47	25.05	7.92
3	2.10	7618.84	24.70	7.57
4	1.90	7065.52	23.84	6.71
5	1.70	6484.50	22.82	5.69
6	1.50	5875.79	21.81	4.68
7	1.30	5239.38	19.95	2.82
8	1.00	4232.84	19.48	2.35
9	0.70	3163.98	18.58	1.45
10	0.40	2032.81	17.63	0.50
11	0.20	1244.08	17.28	0.15
12	0.00	427.65	17.13	0.00

1.3.4 Other Observations

1. The solution rises on applying an external magnetic field. → **its paramagnetic**.

2. You need a strong concentration for even $MnSO_4$, as we shown that there was very less height difference for our **solution 1**

3. We find it easier to measure readings going down compared to going up, as we believe that was solution passing through already wetted glass. Whilst for the first solution we were unable to measure a consistent trend, as we believe that was

because of the solution sticking to the tube. This has introduced a random error in our experiment.

4. We observed the meniscus changing its shape towards the end of the our second solution readings, when the meniscus was reaching the borders of the pole of the electromagnet. Suggesting, non-uniform fields at the boundaries.

5. We calibrated our setup for both increasing and decreasing current to check for hysterisis. We observed it to be very insignificant from our readings, although very clear to see.

1.4 Analysis

The first part in our analysis is to create the calibration plot for the Magnetic Field v/s Current curve using our measurments. We have calibrated the setup for both increasing and decreasing trend of current, wo we can utilise both sets of our measurments and also study the hysterisis in the coil.

Residual magnetic field strength before we started any measurments was 400 G and the same after we concluded our calibration was 460 G. So we saw a definite difference between field strength if we are increasing or decreasing, so we can confirm hysterisis. But the difference is pretty negligible.

We created simple python plots and curve fitted the calibration data onto a quadratic polynomial. The corresponding equation that we get for increasing and decreasing trend are

$$B = (-1.2162 \times 10^2) \times I^2 + (3.5767 \times 10^3) \times I + (3.0068 \times 10^2)$$
$$B = (-3.4619 \times 10^2) \times I^2 + (4.1514 \times 10^3) \times I + (4.2765 \times 10^2)$$

respectively.



Figure 1.1: Calibration Curves for increasing & decreasing trend

Next, we move on to plot the Height Difference v/s B^2 curve, which we will do by finding the corresponding values for magnetic field from the calibration curve fit. **Note:** This has been linearly plotted after converting all the units in SI.



Figure 1.2: curve fit for first solution with less concentration



Figure 1.3: curve fit for second solution with higher concentration

Before moving on two calculate χ value, we need to find the specific density of the solution. We didn't had access to the specific gravity bottle as said in the lab manual, So wi will be taking some assumptions and calculating the values in a ideal sense.

- density of water $(\rho_w) = 0.9970\,\mathrm{g\,cm^{-3}}$ (at $25\,^\circ\mathrm{C}$)
- density of $MnSO_4.H_2O$ $(\rho_{salt}) = 2.95\,\mathrm{g\,cm^{-3}}$

Solution - I

- weight of water $= 30 \times 0.9970 = 29.91\,\mathrm{g}$
- volume of salt $= 5.0 \times \frac{1}{2.95} = 1.69\,\mathrm{mL}$
- Total weight of solution $= 5.0 + 29.91 = 34.91\,\mathrm{g}$
- Total volume of solution $= 30 + 1.69 = 31.69\,\mathrm{mL}$
- density of solution $(\rho_{sol}) = \frac{34.91}{31.69} = \underline{1.10\,\mathrm{g\,cm^{-3}}}$

Solution - II

- weight of water $= 15 \times 0.9970 = 14.96\,\mathrm{g}$
- volume of salt $= 7.5 \times \frac{1}{2.95} = 2.54\,\mathrm{mL}$
- Total weight of solution $= 7.5 + 14.96 = 22.46\,\mathrm{g}$
- Total volume of solution $= 15 + 2.54 = 17.54\,\mathrm{mL}$
- density of solution $(\rho_{sol}) = \frac{22.46}{17.54} = \underline{1.28\,\mathrm{g\,cm^{-3}}}$

Using the density and the slope and putting it in the standard formulae, we can find the value of χ_{sol} for us.

$$\chi_{SI} = \frac{2\mu_0 \rho g h}{B^2}$$

For Solution I (increasing),

$$\chi_1 = 2 \times 4\pi \times 10^{-7} \times 1100 \times 2.8711 \times 10^{-3}$$
$$\chi_1 = 9.0514 \times 10^{-5}$$

For Solution I (decreasing),

$$\chi_1 = 2 \times 4\pi \times 10^{-7} \times 1100 \times 2.8277 \times 10^{-3}$$
$$\chi_1 = 8.9147 \times 10^{-5}$$

For Solution II (increasing),

$$\chi_2 = 2 \times 4\pi \times 10^{-7} \times 1280 \times 9.8632 \times 10^{-3}$$
$$\chi_2 = 3.1095 \times 10^{-4}$$

For Solution II (decreasing),

$$\chi_2 = 2 \times 4\pi \times 10^{-7} \times 1280 \times 1.1890 \times 10^{-2}$$
$$\chi_2 = 3.7486 \times 10^{-4}$$

We will use a simple average to continue on from here;

$$\chi_1 = 8.9830 \times 10^{-5}$$
$$\chi_2 = 3.4390 \times 10^{-4}$$

This quantity χ can be referred to as Volume Magnetic susceptibility, to calculate mass susceptibility and molar susceptibility. We need to use the following formulae:

$$\chi'_{mass} = \chi/\rho$$
$$\chi''_{molar} = M\chi'$$

Water is a diamagnetic and we can remove its contribution from the volume susceptibility to calculate the mass susceptibility for just the salt.

$$\chi_{sol} = \chi_{salt} + \chi_{water}$$

For Solution - II,

$$3.7486 \times 10^{-4} = \chi_{salt} - (9.04 \times 10^{-6})$$
$$\chi_{salt} = 3.8390 \times 10^{-4}$$

Similar relation can be written about the mass susceptibility of the solution:

$$\chi'_{sol} = \frac{m_{salt}}{m_{sol}}\chi'_{salt} + \frac{m_{water}}{m_{sol}}\chi'_{water}$$

$$\chi'_{sol} = \frac{7.5}{22.46}\chi'_{salt} + \frac{14.96}{22.46}(-9.04 \times 10^{-9})$$
$$\frac{3.7486 \times 10^{-4}}{1280} = 0.3339\chi'_{salt} - (6.02 \times 10^{-9})$$
$$\chi'_{salt} = 8.9511 \times 10^{-7}$$

Now, if we were to calculate the Molar susceptibility we get:

$$\chi_{molar} = 169.02 \times 10^{-3} \times 8.9511 \times 10^{-7}$$
$$\chi_{molar} = 1.5127 \times 10^{-7}$$

Table 1.7: Experimental Results for the given salt

#	Parameter	Value
1	Volume Susceptibility (χ)	3.7486×10^{-4}
2	Mass Susceptibility (χ')	$8.9511 \times 10^{-7}\,\mathrm{m^3\,kg^{-1}}$
3	Molar Susceptibility (χ'')	$1.5127 \times 10^{-7}\,\mathrm{m^3\,mol^{-1}}$

Theoritical Molar Susceptibility for Manganese(II) Sulfate Monohydrate ($\boldsymbol{MnSO_4.H_2O}$) $= 1.7844 \times 10^{-7}\,\mathrm{m^3\,mol^{-1}}$

Error Analysis

$$\text{Error in Molar Susceptibility result} = \frac{1.7844 \times 10^{-7} - 1.5127 \times 10^{-7}}{1.7844 \times 10^{-7}} \times 100\%$$
$$= 15.22\%$$

The propagation of uncertainty while taking measurement can be written as:

$$
\begin{aligned}
\frac{\sigma_{\chi''}}{\chi''} &= \frac{\sigma_{\chi'}}{\chi'} \\
&= \sqrt{\left(\frac{0.1}{7.5}\right)^2 + \left(\frac{0.1}{22.46}\right)^2 + \left(\frac{\sigma_{\chi'_{sol}}}{\chi'_{sol}}\right)^2} \\
&= \sqrt{\left(\frac{0.1}{7.5}\right)^2 + \left(\frac{0.1}{22.46}\right)^2 + \left(\frac{\sigma_{\chi}}{\chi}\right)^2} \\
&= \sqrt{\left(\frac{0.1}{7.5}\right)^2 + \left(\frac{0.1}{22.46}\right)^2 + \left(\frac{\sigma_{slope}}{\text{slope}}\right)^2} \\
\sigma_{\chi''} &= 1.5127 \times 10^{-7} \sqrt{\left(\frac{0.1}{7.5}\right)^2 + \left(\frac{0.1}{22.46}\right)^2 + \left(\frac{5.8082 \times 10^{-4}}{1.1890 \times 10^{-2}}\right)^2} \\
\sigma_{\chi''} &= 7.68 \times 10^{-9}
\end{aligned}
$$

This uncertainty due to measurment comes around to be about 5%. Using this we can safely say that our experiment was able to measure the magnetic susceptibilty of the salt under certain standard deviation limits.

Why is our result slightly lower than theoritcal?
We attribute this deviation to be mostly arising from a improper setup. Before our second set of readings, we tried to make sure our tube was clean, and used IPA to help us aid in that process. We waited for long enough time for the IPA to dry off, but as we started taking readings for our second solution we see that some bubbles are rising in the tube, especially at the thinner end. This was obvious to see through the microscope, and minorly affected our ability to take the microscope reading. In hindsight, we realised that this must be the IPA rising as it is lighter compared to our solution. This introduced a IPA which slightly diluted our solution and affected the density value. Magnetically this doesn't had a significant as it is dimagnetic like water, but has even lesser magnetic moment. In an extreme case, oxidation of manganese ions in the solution could also have played a role in the decreased value. Also the theoritcal value is temperature dependent and measured at 293 K which is definitly lower than the temperature setting for our experiment. And a higher temperature would lead to a lower value as the magnetization is related to temperature as inverse dependency as stated by Curie's Law.

1.5 Conclusion

We successfully completed the experiment by measuring the magnetic susceptibility of the salt, manganese sufate monohydrate. Our measured value for the molar magnetic susceptibility (1.5217×10^{-7}) was about **15%** deviation form the theoritical value.

Our value being lower than the theoritical one can be easily explained because of the uncertainty in the solution density and temperature conditions.

Experiment No. 2

Bipolar Junction Transistor

Date: 30 January, 2026

2.1 Aim

The aim of the experiment is to study the input and output characterstic of an n-p-n Bipolar Junction Transistor that is setup in common emitter bias mode. We will also construct an amplifier circuit and test the current and voltage gains for the circuit. We will also create a phase-shift oscillator circuit and test its function.

2.2 Theory

2.3 Experimental Observation

Table 2.1: Base Characterstic

#	V_{BE} (mV) 0	I_B (mA) 0	V_{BE} (mV) 1	I_B (mA) 1	V_{BE} (mV) 2	I_B (mA) 2
1	3.60	0.00	3.60	0.00	269	0.00
2	575.00	4.23	241.00	0.00	435	0.03
3	589.00	6.57	458.00	0.08	547	1.19
4	455.00	0.07	560.00	2.54	570	2.57
5	533.00	1.02	586.00	6.04	588	4.50
6	558.00	2.33	604.00	10.25	604	7.01
7	598.00	8.61	615.00	14.50	614	9.96
8	604.00	10.16	625.00	19.45	623	13.49
9	614.00	13.95	632.00	23.65	633	19.34
10	621.00	17.05	644.00	33.55	637	22.40

Table 2.2: Collector Characterstic

#	V_{CE} (V) 0	I_C (mA) 0	V_{CE} (V) 1	I_C (mA) 1	V_{CE} (V) 2	I_C (mA) 2
1	0.00	0.00	0.01	0.00	0.00	0.00
2	0.01	0.02	0.06	1.24	0.06	2.05
3	0.07	0.46	0.13	7.19	0.11	7.13
4	0.12	1.64	0.21	10.25	0.18	13.84
5	0.24	2.88	0.41	11.04	0.33	17.01
6	0.34	2.97	0.61	12.29	0.53	18.73
7	0.54	2.97	0.84	12.38	0.82	20.32
8	0.80	2.97	1.13	12.44	1.19	20.95
9	1.21	2.98	2.23	12.67	1.96	21.80
10	1.92	2.99	3.26	12.94	3.21	23.09
11	2.83	3.01	4.80	13.54	5.66	25.96
12	4.91	3.07	7.32	14.05	9.68	29.76
13	9.11	3.16	NaN	NaN	NaN	NaN
14	NaN	NaN	NaN	NaN	NaN	NaN

2.4 Analysis

2.5 Conclusion