

NDA: Time Series Analysis - Part 2

Lionel Tabourier

LIP6 – CNRS and Sorbonne University

`first_name.last_name@lip6.fr`

January 2021

Outline

- 1 What to do with the residuals?
- 2 Autoregressive models
- 3 Moving Average models
- 4 ARMA models

Outline

- 1 What to do with the residuals?
- 2 Autoregressive models
- 3 Moving Average models
- 4 ARMA models

How to analyze a time series? (3)

Third step:

- fit the residuals

If the residuals is an **IID time series**, nothing else to model. . .

Simple method to test if a time series is IID

General idea: suppose IID random variables
What should we observe? Is it the case?

Simple method to test if a time series is IID

General idea: suppose IID random variables
What should we observe? Is it the case?

Sample ACF criterion

Because of the Central Limit Theorem:

- suppose x_t IID with mean 0 and variance 1 (white noise)
- if n large enough, $\hat{\rho}_x(h)$ is approx. distributed as $\mathcal{N}(0, \frac{1}{\sqrt{n}})$

In practice, consider the 95% confidence interval:
How many values fall out of $\left[\frac{-1.96}{\sqrt{n}}, \frac{+1.96}{\sqrt{n}} \right]$? By how much?


Simple method to test if a time series is IID

General idea: suppose IID random variables
What should we observe? Is it the case?

Sample ACF criterion

Because of the Central Limit Theorem:

- suppose x_t IID with mean 0 and variance 1 (white noise)
- if n large enough, $\hat{\rho}_x(h)$ is approx. distributed as $\mathcal{N}(0, \frac{1}{\sqrt{n}})$




Simple method to test if a time series is IID

General idea: suppose IID random variables
What should we observe? Is it the case?

Sample ACF criterion

Because of the Central Limit Theorem:

- suppose x_t IID with mean 0 and variance 1 (white noise)
- if n large enough, $\hat{\rho}_x(h)$ is approx. distributed as $\mathcal{N}(0, \frac{1}{\sqrt{n}})$



not IID noise

How to analyze a time series? (3)

Third step:

- fit the residuals

If the residuals is an IID time series, nothing else to model. . .

Otherwise, we use [ARMA models](#)

Outline

- 1 What to do with the residuals?
- 2 Autoregressive models**
- 3 Moving Average models
- 4 ARMA models

AR(1) process

What is autoregression?

auto means self \Rightarrow regression from itself

The most basic AR model: 1st order regression or AR(1)

$\{X_t\}$ is a series satisfying:

$$X_t = \phi X_{t-1} + W_t, \quad |\phi| < 1$$


where W_t is a white noise (mean 0, variance σ^2)

Reminder: stationary

- $\mathbb{E}[X_t] = 0$
- $\gamma_X(h) = \phi^{|h|} \gamma_X(0) = \phi^{|h|} \frac{\sigma^2}{1-\phi^2}$

AR(1) process

Consider the following time series r_t :




Suppose AR(1) model for a time series r_t , how to compute ϕ ?

- plot r_t as a function of r_{t-1} (*lag-1 plot*)
- linear fit, slope is an estimation of ϕ

AR(1) process

Suppose AR(1) model for a time series r_t , how to compute ϕ ?

- plot r_t as a function of r_{t-1} (*lag-1 plot*)
- linear fit, slope is an estimation of ϕ




on the example $\phi \simeq -0.659$

AR(1) process

Now, we compute the residuals of the TS – AR(1) model:

$$r_t - \phi r_{t-1}$$




Is it closer to an IID noise?

AR(1) process

Now, we compute the residuals of the TS – AR(1) model:

$$r_t - \phi r_{t-1}$$




Is it closer to an IID noise?

AR(1) process

Now, we compute the residuals of the TS – AR(1) model:

$$r_t - \phi r_{t-1}$$

Is it closer to an IID noise?




before AR(1): not IID

AR(1) process

Now, we compute the residuals of the TS – AR(1) model:

$$r_t - \phi r_{t-1}$$

Is it closer to an IID noise?



after AR(1): **closer to IID** → other tests?

Generalized AR models

AR(1) model

W_t is white noise

signal = noise and (weighted) influence of the signal at $t - 1$

- $X_t = \phi X_{t-1} + W_t$
- $X_t = \phi_1 X_{t-1} + \phi_2 X_{t-2} + \dots + \phi_p X_{t-p} + W_t$

Generalized AR models

AR(p) model

W_t is white noise

signal = noise and influence of the signal at p previous steps


- $X_t = \phi X_{t-1} + W_t$
- $X_t = \phi_1 X_{t-1} + \phi_2 X_{t-2} + \dots + \phi_p X_{t-p} + W_t$

Generalized AR models

$$X_t - \phi_1 X_{t-1} - \phi_2 X_{t-2} - \dots - \phi_p X_{t-p} = W_t$$

Some characteristics

- **Stationary** process? see characteristic polynomial
 $P(x) = 1 - \phi_1 x - \phi_2 x^2 - \dots - \phi_p x^p \rightarrow$ if no unit root AR(p) stationary
- typical shape: smooth decay, no cut-off




Generalized AR models

$$X_t - \phi_1 X_{t-1} - \phi_2 X_{t-2} - \dots - \phi_p X_{t-p} = W_t$$

Some characteristics

- **Stationary** process? see characteristic polynomial
 $P(x) = 1 - \phi_1 x - \phi_2 x^2 - \dots - \phi_p x^p \rightarrow$ if no unit root AR(p) stationary
- typical shape: **smooth decay, no cut-off**



How to find AR(p) ACF coefficients?

$$X_t = \phi_1 X_{t-1} + \phi_2 X_{t-2} + \dots + W_t$$

Yule-Walker equations

We have seen that for AR(1):

$$\gamma(h) = \phi^{|h|} \gamma(0) = \phi^{|h|} \frac{\sigma^2}{1 - \phi^2}$$

What about the general case?

How to find AR(p) ACF coefficients?

$$X_t = \phi_1 X_{t-1} + \phi_2 X_{t-2} + \dots + W_t$$

Yule-Walker equations

We have seen that for AR(1):

$$\gamma(h) = \phi^{|h|} \gamma(0) = \phi^{|h|} \frac{\sigma^2}{1 - \phi^2}$$

What about the general case?

- $\gamma(h) = \phi_1 \gamma(h-1) + \phi_2 \gamma(h-2) + \dots$
- $\rho(h) = \phi_1 \rho(h-1) + \phi_2 \rho(h-2) + \dots$

these are called the Yule-Walker equations

How to find AR(p) ACF coefficients?

$$X_t = \phi_1 X_{t-1} + \phi_2 X_{t-2} + \dots + W_t$$

Yule-Walker equations

We have seen that for AR(1):

$$\gamma(h) = \phi^{|h|} \gamma(0) = \phi^{|h|} \frac{\sigma^2}{1 - \phi^2}$$

What about the general case?

- $\gamma(h) = \phi_1 \gamma(h-1) + \phi_2 \gamma(h-2) + \dots$
- $\rho(h) = \phi_1 \rho(h-1) + \phi_2 \rho(h-2) + \dots$

these are called the Yule-Walker equations

How to find AR(p) ACF coefficients?

$$X_t = \phi_1 X_{t-1} + \phi_2 X_{t-2} + \dots + W_t$$

Yule-Walker equations

We have seen that for AR(1):

$$\gamma(h) = \phi^{|h|} \gamma(0) = \phi^{|h|} \frac{\sigma^2}{1 - \phi^2}$$

What about the general case?


- $\gamma(h) = \phi_1 \gamma(h-1) + \phi_2 \gamma(h-2) + \dots$
- $\rho(h) = \phi_1 \rho(h-1) + \phi_2 \rho(h-2) + \dots$

these are called the **Yule-Walker equations**

How to find AR(p) ACF coefficients?

Illustration on a practical case

$$X_t = \frac{1}{3}X_{t-1} + \frac{1}{2}X_{t-2} + W_t$$




Sample ACF (blue) vs Yule-Walker coefficients (orange)

Outline

- 1 What to do with the residuals?
- 2 Autoregressive models
- 3 Moving Average models**
- 4 ARMA models

MA(q) processes

Moving average to smooth a signal:



Original signal $f(x)$ smoothed over a window size w , typically:

ex1: $g_1(x) = \frac{f(x)+f(x-1)+f(x+1)}{3}$ (no weight, two-sided, $w = 3$)

ex2: $g_2(x) = \frac{f(x)+f(x-1)+f(x-2)}{3}$ (no weight, one-sided, $w = 3$)

ex3: $g_3(x) = \frac{f(x)+a.f(x-1)+b.f(x-2)+c.f(x-3)}{(1+a+b+c)}$ (weighted, one-sided, $w = 4$)

MA(q) processes

MA(1) model

W_t is white noise

signal = weighted average of noise at t and of noise at $t - 1$

- $X_t = \beta_0 W_t + \beta_1 W_{t-1}$
- $X_t = \beta_0 W_t + \beta_1 W_{t-1} + \dots + \beta_q W_{t-q}$

MA(q) processes

MA(q) model

W_t is white noise

signal = weighted average of noise at t and q previous steps

- $X_t = \beta_0 W_t + \beta_1 W_{t-1}$
- $X_t = \beta_0 W_t + \beta_1 W_{t-1} + \dots + \beta_q W_{t-q}$

MA(q) processes

$$X_t = \beta_0 W_t + \beta_1 W_{t-1} + \dots + \beta_q W_{t-q}$$


Some characteristics

- **Stationary** process

mean and autocovariance at lag h do not depend on time

Ex: prove if $h \leq q$, $\text{Cov}(X_t, X_{t+h}) = \sigma^2 \sum_{i=0}^{q-h} \beta_i \beta_{i+h}$

- ACF cuts off at lag q



MA(q) processes

$$X_t = \beta_0 W_t + \beta_1 W_{t-1} + \dots + \beta_q W_{t-q}$$


Some characteristics

- **Stationary** process

mean and autocovariance at lag h do not depend on time

Ex: prove if $h \leq q$, $\text{Cov}(X_t, X_{t+h}) = \sigma^2 \sum_{i=0}^{q-h} \beta_i \beta_{i+h}$

- **ACF cuts off at lag q**



Connection between MA and AR

From MA(1) to AR

Considering the MA(1) process:

$$\begin{aligned}X_t &= W_t + \beta W_{t-1} \\ \Rightarrow W_t &= X_t - \beta W_{t-1} \\ \Rightarrow W_t &= X_t - \beta(X_{t-1} - \beta W_{t-2}) \\ \Rightarrow W_t &= X_t - \beta X_{t-1} + \beta^2 X_{t-2} - \beta^3 X_{t-3} + \dots \\ \Rightarrow X_t &= W_t + \beta X_{t-1} - \beta^2 X_{t-2} + \beta^3 X_{t-3} - \dots\end{aligned}$$

In other words, MA(1) is an AR(∞) process

More generally, MA(q) can (often) be seen as AR(∞) process

Connection between MA and AR

From MA(1) to AR

Considering the MA(1) process:

$$\begin{aligned}X_t &= W_t + \beta W_{t-1} \\ \Rightarrow W_t &= X_t - \beta W_{t-1} \\ \Rightarrow W_t &= X_t - \beta(X_{t-1} - \beta W_{t-2}) \\ \Rightarrow W_t &= X_t - \beta X_{t-1} + \beta^2 X_{t-2} - \beta^3 X_{t-3} + \dots \\ \Rightarrow X_t &= W_t + \beta X_{t-1} - \beta^2 X_{t-2} + \beta^3 X_{t-3} - \dots\end{aligned}$$

In other words, MA(1) is an AR(∞) process

More generally, MA(q) can (often) be seen as AR(∞) process

Connection between MA and AR

From AR(1) to MA

$$\begin{aligned}X_t &= W_t + \phi X_{t-1} \\ \Rightarrow X_t &= W_t + \phi(W_{t-1} + \phi X_{t-2}) \\ \Rightarrow X_t &= \dots \\ \Rightarrow X_t &= W_t + \phi W_{t-1} + \phi^2 W_{t-2} + \phi^3 W_{t-3} + \dots\end{aligned}$$

In other words, AR(1) is an MA(∞) process

More generally, AR(p) can (often) be seen as MA(∞) process

Connection between MA and AR

From AR(1) to MA

$$\begin{aligned}
 X_t &= W_t + \phi X_{t-1} \\
 \Rightarrow X_t &= W_t + \phi(W_{t-1} + \phi X_{t-2}) \\
 \Rightarrow X_t &= \dots \\
 \Rightarrow X_t &= W_t + \phi W_{t-1} + \phi^2 W_{t-2} + \phi^3 W_{t-3} + \dots
 \end{aligned}$$

In other words, AR(1) is an MA(∞) process

More generally, AR(p) can (often) be seen as MA(∞) process

Outline

- 1 What to do with the residuals?
- 2 Autoregressive models
- 3 Moving Average models
- 4 ARMA models**

ARMA models

ARMA(p,q) model

ARMA(p,q) model is a combination of AR(p) and MA(q) model:

$$X_t = \phi_1 X_{t-1} + \dots + \phi_p X_{t-p} + W_t + \beta_1 W_{t-1} + \dots + \beta_q W_{t-q}$$

As for AR(p) and MA(q) parameters can be found from the ACF

In practice

- fit the residuals with several (low) values of p and q
- select what is the *best* model
- \Rightarrow complete model:
trend + seasonality + ARMA(p,q) residuals

ARMA models

ARMA(p,q) model

ARMA(p,q) model is a combination of AR(p) and MA(q) model:

$$X_t = \phi_1 X_{t-1} + \dots + \phi_p X_{t-p} + W_t + \beta_1 W_{t-1} + \dots + \beta_q W_{t-q}$$

As for AR(p) and MA(q) parameters can be found from the ACF

In practice

- fit the residuals with several (low) values of p and q
- select what is the *best* model
- \Rightarrow complete model:
trend + seasonality + ARMA(p,q) residuals

A few questions that we have left aside . . .

- How to choose p and q of an ARMA process?
→ *useful to define **partial ACF***
- How to find the coefficients of an ARMA process?
→ *transform it in $AR(\infty)$ (or $MA(\infty)$) process*
- How do we select the best model?
→ ***complexity criteria** and the overfitting problem (AIC, BIC)*
- How to deal more systematically with the trend?
→ ***differentiation method***

A few questions that we have left aside . . .

- How to choose p and q of an ARMA process?
→ *useful to define **partial ACF***
- How to find the coefficients of an ARMA process?
→ *transform it in $AR(\infty)$ (or $MA(\infty)$) process*
- How do we select the best model?
→ ***complexity criteria** and the overfitting problem (AIC, BIC)*
- How to deal more systematically with the trend?
→ ***differentiation method***

A few questions that we have left aside . . .

- How to choose p and q of an ARMA process?
→ *useful to define **partial ACF***
- How to find the coefficients of an ARMA process?
→ *transform it in $AR(\infty)$ (or $MA(\infty)$) process*
- How do we select the best model?
→ ***complexity criteria** and the overfitting problem (AIC, BIC)*
- How to deal more systematically with the trend?
→ ***differentiation method***

A few questions that we have left aside . . .

- How to choose p and q of an ARMA process?
→ *useful to define **partial ACF***
- How to find the coefficients of an ARMA process?
→ *transform it in $AR(\infty)$ (or $MA(\infty)$) process*
- How do we select the best model?
→ ***complexity criteria** and the overfitting problem (AIC, BIC)*
- How to deal more systematically with the trend?
→ ***differentiation method***

Studying time series in python

Among several options, `pandas` library

A few useful functions:

- Load data as dataframe:
`read_csv` from `pandas` library
- Moving average:
`rolling` from `pandas` library
- Fitting:
`curve_fit` in `scipy.optimize` library
- Autocorrelation function:
`plot_ACF` in `statsmodels` library
- ARMA model fit:
`ARMA.fit` in `statsmodels` library