

Anticoagulants

Dr.Nathasha Luke
Department of
Pharmacology

Anticoagulant drugs

- used for the prevention and treatment of arterial and venous thromboembolism
- Thrombosis –thrombus formation within the circulation

Types of anticoagulants

- heparinoids
 - unfractionated heparin (UFH)
 - low-molecular- weight heparin (LMWH)
 - Fondaparinux
- vitamin K antagonists (VKA) — warfarin,
- new oral agents — dabigatran, rivaroxaban, apixaban

Heparin

Mechanism of action

Unfractionated Heparin (UFH)

- Inhibits Thrombin (IIa), IXa, Xa, XIa and XIIa
- Therapeutic action: Xa, thrombin



Potentiates the action of antithrombin 111

Low Molecular Weight Heparin (LMWH)


- Predominantly inhibit factor Xa

Fondaparinux

- Purely inhibits factor Xa



Mechanism of action



Unfractionated heparin

- 1st anticoagulant discovered
- consists of a series of glycosaminoglycan chains and is processed from animal intestinal mucosa
- anticoagulant effect mainly by potentiating the action of the physiological anticoagulant antithrombin ten thousand-fold
- unpredictable pharmacokinetics and a short elimination half-life

Unfractionated heparin

- Adverse effects
 - bleeding
 - allergy
 - heparin-induced thrombocytopenia (HIT)
 - osteopenia (long term use)
- monitoring of heparin -using the activated partial thromboplastin time (target -2-3 times of normal APTT)
- Dosing of UFH requires a bolus dose for initiation of an UFH infusion.

Low molecular weight heparin

- produced by depolymerization of UFH chains to produce a shorter glycosaminoglycan chain.
- more predictable pharmacokinetics, a longer half-life
- monitoring is required only in patients with renal failure and those at extremes of body weight, and is performed using anti-Xa concentration

UFH vs LMWH.

UFH	LMWH
intravenous (IV) infusion or subcutaneous (SC) injection	easier mode of administration (SC injection) than UFH.
Unpredictable pharmacokinetics	more predictable pharmacokinetics
short elimination half-life	longer half-life
Needs infusion followed by bolus	Bolus doses
potentiating the action of the physiological anticoagulant antithrombin	binds to antithrombin but has a greater inhibitory effect on factor Xa.
SE- heparin-induced thrombocytopenia (HIT) and osteopenia	Less compared to UFH
No dose adjustment in RF	Reduce dose in RF
Need monitoring in all	Monitoring only special circumstances
Monitoring - aPTT	Factor Xa levels
Reverse anticoagulant effect - protamine sulphate	Only partially reversed by protamine

fondaparinux

- is a synthetic form of the heparin pentasaccharide molecule
- Anti Xa activity
- Indications similar to other types
- However more benefit in MI
- Costly
- Long half life- single daily dose (LMWH is given bd)

Indications of heparin

- Prevention of thrombosis of peripheral catheters
- Prevention and treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE),
- Treatment of cardiac ischaemia and critical ischaemia of the lower limb


Types of anticoagulants

- heparinoids
 - unfractionated heparin (UFH)
 - low-molecular- weight heparin (LMWH)
 - Fondaparinux
- vitamin K antagonists (VKA) — warfarin,
- new oral agents — dabigatran, rivaroxaban, apixaban

Warfarin

- **Mode of action**

warfarin prevents gamma carboxylation of vitamin K-dependent clotting factors (II, VII, IX and X), producing an anticoagulant effect takes several days to have full anticoagulant effect.



Warfarin

- Monitoring –Prothombin time
- Contraindicated in T1 of pregnancy (Is continued on heparin)- warfarin embrayopathy/fetal warfarin syndrome
- Dosing and initiation –patient dependant
- Acute thrombosis -----initiate warfarin with heparin
- Why?

the initial reduction proteins C & S by warfarin may lead to an enhanced prothrombotic effect.

- Initiating high doses –not beneficial

INR targets with warfarin and indications

Indication	Target INR
Pulmonary embolism	2.5
Proximal deep vein thrombosis	
Calf vein thrombosis	
Non-rheumatic atrial fibrillation (CHADS ₂ score >1)	
Mural thrombus	
Cardiomyopathy	
Mechanical prosthetic aortic heart valve	
Bioprosthetic valve	3.5
Mechanical prosthetic mitral heart valve	
Recurrent venous thrombosis on warfarin therapy	

Adverse events

- Hemorrhage
- Necrosis of skin
- Systemic atheroemboli and cholesterol microemboli
- Vascular disorders: vasculitis
- Hepatobiliary disorders
- Gastrointestinal disorders: nausea, vomiting, diarrhea, taste perversion, abdominal pain, flatulence, bloating

Skin necrosis



Contraindications

- Hemorrhagic tendencies
- Recent major surgery
- Threatened abortion, eclampsia, and preeclampsia
- Unsupervised patients with conditions associated with potential high level of non-compliance
- Hypersensitivity
- Malignant hypertension

Drug interactions

- CYP450 isozymes involved in the metabolism of Warfarin
- Drugs that inhibit these enzymes increase risk of bleeding
- Drugs that activate these enzymes reduce the anticoagulant effect
- What are the drugs interacting with warfarin?

Some common drug interactions

Medscape®

www.medscape.com

Pharmacodynamic interactions

Mechanism

ASA/NSAIDs

Antiplatelet, gastrointestinal injury

Clopidogrel/Ticlopidine

Antiplatelet

Tramadol

INR elevation (unclear mechanism)³³

Levothyroxine

Increased catabolism of clotting factors

Vitamin K-containing food/supplements

INR depression (circumvent warfarin mechanism of action)

Clinical case

- A 56 year old female on warfarin for non valvular AF presents with lethargy and found to be pale. She has a history of tarry stools for five days . Her INR is 3.5
- What is the next step in management ?

Reversal of anticoagulation

- Life threatening bleeds - dried prothrombin complex (prothrombin complex concentrate; PCC) contains factors II, VII, IX and X in combination with IV. Vitamin K
FFP if PCC not available
- non-major bleeding -discontinuation of warfarin or a decrease in dosage and/or administration of vitamin K will usually suffice

