

Alcohol & it's effects on the body

PROF. ANURUDDHI EDIRISINGHE

MBBS, MD,DLM,DMJ(LOND), MFFLM(UK)

DEPARTMENT OF FORENSIC MEDICINE



To produce a graduate who will be able to examine victims and suspects involved in criminal/civil or traffic offences where an allegation of alcohol consumption or intoxication is present and provide an opinion to the court

Examine individuals who are dead due to alcohol or alcohol related illnesses and assist courts

Knowledge, skills and attitude

How it is achieved

- Knowledge on the effect of alcohol in the body (short & long term)
- Knowledge on how a diagnosis of intoxication is arrived by History, examination , & investigation
- Knowledge on the effect of alcohol in the body in relation to driving ability
- Develop skills on examination and forming opinions based on findings

Alcohol

Stages of intoxication -A

Clinical examination for drunkenness -A

Other methods available to establish drunkenness - A

Effects of alcohol on the ability to drive -A

Laws related to drunkenness -A

Differential diagnosis of alcohol intoxication -A

Sudden death due to alcohol abuse -B

(A) Comprehensive knowledge of specified area and competency in specified medico-legal work

(B) Knowledge of the basic principles with ability to identify where referral is needed

(C) Nice to know



Alcohol & Forensic

Association of Crime statistics and heavy drinking and criminal behaviors

Relationship of alcohol to violent(24 to 85%) vs non violent crimes 12–38%

Alcohol and ability to drive


Alcohol and accidents

Alcohol related natural deaths

Alcohol and fitness for detention in custody

Liquor- beverage
containing alcohol

Alcohol VS liquor



In chemistry, an alcohol is any organic compound where a hydroxyl functional group (-OH) is bound to a carbon or hydrogen atoms

Ethanol produced by the fermentation of sugar by yeast

Ethanol ($\text{C}_2\text{H}_5\text{OH}$) is the alcohol found in alcoholic beverages



Alcohol is ethanol

Content of alcohol in various liquors

Fortified Wine GLASS: 60ml
ALCOHOL CONTENT: 10.ml - 17.5%

Beer GLASS: 285ml
ALCOHOL CONTENT: 11.4ml - 4%

Wine GLASS: 100ML
ALCOHOL CONTENT: 11.5ml-11.5%

Spirit GLASS: 200ML
ALCOHOL CONTENT: 11.4ml
- contains one 30ml nip of spirits (38%)



Fruit Juice < 0.1%
Pilsner 3–6%
ESB (Bitter) 3-6%
Lager 4-5%
Porter 4-5%
Alcopops/Breezers/Coolers 4–7%
PA (India Pale Ale) 6-7%
Cider 4–8%
Sparks 6-7%
Stout 5-10%
Sparkling Wine 8 – 12%
Table Wine 8 – 14%
Retsina 10–11%
Barley Wine 10–15%
Wine (general) 10–15%
Port Wine 20%
Fortified Wine 17 – 22%
Liqueur 15–55%
Light Liquors 20%
Liquor/Spirits (general) 40%
Cask Strength Whisky/Rum 60%
Absinthe 55–89.5%
Neutral Grain Spirit 95%
Rectified Spirit 96%
Absolute Alcohol 99-100%

Contents of alcohol & in liquors used in Sri Lanka

Arrack 30-50%

Toddy 4%- 6%

Pot Arrack- 25-30%

Kassipu- 20%

Beer- 4%-6%



Effects of alcohol in the body

Entry

- GIT
- ? RS,SKIN v(50 mg/100 mL after breathing alcohol/air mixtures several hours),

Absorption


- Mouth & oesophagus (small)
- Stomach
- Intestine (duodenum highest)

Blood

- Carried in water
- Circulation and effects in the systems


Excretion

- 5-10% excreted unchanged in breath, urine, and sweat
- 90–95% of oxidized in the liver by alcohol dehydrogenase – acetaldehyde & acetate (acetic acid)



Blood alcohol curve

BLOOD ALCOHOL CURVE



IMPORTANT

Allows to measure the alcohol content in the body

Curves varies from person to person and within the same person over a time

Alcohol is distributed in body water

Alcohol Dehydrogenase is saturated at low alcohol concentrations , therefore elimination is constant

Blood alcohol level may vary

Sex & weight

- Male vs females i.e. females become more drunk than male due to body fat
- body water (weight i.e. smaller body size higher peak)

Duration of drinking

- *Volume of alcohol longer period- elimination starts & low levels of blood alcohol is reached*
- *Sipping of a drink vs gulping of drink*

Nature of the Drink Consumed

- *Beer, Wine, Sprits*
- *>20% alcohol in a drink slows gastric emptying by irritation of gastric mucosa*



Food in the Stomach

- full meal before drinking can reduce the peak alcohol level by an average of 9–23%
- *Fat containing food slower more*
- *Carbonated water increases absorption more*



Physiological Factors

- *Stomach wall permeability, blood supply to intestine*
- *Gastric surgery*



Genetic Variation

- *Chinese races low alcohol dehydrogenase*

Drugs that affects the rate of stomach emptying and alcohol absorption- atropine; chlorpromazine; tricyclic antidepressants, amphetamines, antidiarrheal medicines; codeine and dihydrocodeine; diamorphine (heroin); methadone; dextropropoxyphene (in co-proxamol) metoclopramide; cisapride; erythromycin

Rate of Elimination

- *Chronic alcoholics AD enzyme stimulated*



Social
drinker



Health
drinker



Types of
alcohol
drinkers

Binge
drinker



Alcoholic



Problem
drinker



Underage
drinker



Effects of Alcohol

Alcohol acts as a CNS depressant

- small doses interferes with cortical function,
- larger doses may depress medullary processes.

The apparent stimulatory effects : acts higher centers of the brain that govern inhibition


Effects are more pronounced when blood alcohol levels are rising than when falling.
(Mellan by effect)

Sequence of Central Nervous Depressant Effects of Alcohol


Stage of influence	Blood alcohol concentration mg/100 mL	Clinical effect
Sobriety	10–50	<ul style="list-style-type: none">• Often no obvious effect; may feel “relaxed”
Euphoria	30–120	<ul style="list-style-type: none">• Mild euphoria with increased talkativeness• Decreased inhibitions• Increased self-confidence• Impaired fine motor skills
Excitement	90–200	<ul style="list-style-type: none">• Emotional instability• Poor sensory perception• Impaired memory and comprehension• Incoordination and loss of balance
Drunkenness	150–300	<ul style="list-style-type: none">• Disorientation, mental confusion• Disturbances of vision (e.g., diplopia)• Decreased pain sense• Increased incoordination with staggering gait• Slurred speech
Stupor	250–400	<ul style="list-style-type: none">• General inertia approaching paralysis• Marked lack of response to stimuli• Inability to stand or walk• Vomiting, incontinence of urine and feces
Coma	350–500	<ul style="list-style-type: none">• Coma and anesthesia• Depressed or absent reflexes• Cardiovascular and respiratory depression• Possible death
Death	Over 450	<ul style="list-style-type: none">• Death from respiratory depression

Stages of drunkenness & sobering up

Clever
Attractive
Rich
Invincible
Invisible
Stupid
Ugly
Poor
Fragile
Conspicuous



Sobriety & Euphoria



Excitement



Drunkenness



Stupor

Effects of Alcohol- short term & long term Signs & symptoms

CNS- orientation, Memory, Gait, Coordination, special séances-

- *Speech,*
- *Eye changes – Nystagmus, Pupillary Changes, double vision, tunnel vision*
- *Wernike's encephalopathy, Kosocoff syndrome*

Cardiovascular Effects- pulse, BP, heart, alcoholic cardiac myopathy

Metabolic Effects- hypoglycemia

GIT- Gastritis, GI Bleeding, alcoholic cirrhosis

Nystagmus

eye is easiest to examine to detect the effects of alcohol



Alcohol can cause nystagmus through at least two mechanisms.

- on the vestibular system causing positional alcohol nystagmus (PAN) i.e. when patient is lying supine with the head turned to either the left or right (Behrens, 1978).
- on ocular movements via neural mechanisms by inhibiting smooth pursuits causing impaired ability to maintain eccentric gaze-i.e. Horizontal gaze nystagmus (HGN)

Positional Alcohol Nystagmus

Positional alcohol nystagmus occurs in two stages

PAN I

- is associated with acute elevation of blood alcohol, tending to occur approx 30 minutes after alcohol ingestion.
- nystagmus is in the direction toward which the head is turned.

PAN II

- occurs at approx 5–6 hours after drinking
- Nystagmus in the opposite direction to that seen in PAN I.



Horizontal gaze nystagmus



HGN is a jerky eye movement noted when gaze is directed to one side.

HGN is in the direction of gaze, and intensified at a more eccentric gaze position

HGN can be seen in normal individuals at extreme lateral gaze

HGN when detected at lesser deviations, it is considered pathological.

An angle of onset of 40° or less from the midline is a sensitive indicator of a blood alcohol level in excess of 100 mg/100 mL (Belton H 1987)

Eye changes

PUPILLARY CHANGES

Early stages pupils are dilated



Late stage (coma) pupils are pinpoint



VISION


Double vision

- (BAL 200mg/100mL)

Tunnel vision

Night vision impaired (pupils to adapting)

Contrast sensitivity (reduced)



Speech

Speech production is a complex motor activity.

Requires a high degree of coordination

A sensitive index of alcohol intoxication

Early stages of alcohol intoxication results talkativeness

Slurring of speech : 100 mg/100 mL.

Cardio vascular effects

Moderate doses leads slight increase in blood pressure and pulse rate (rapid bounding pulse due to increase catecholamine)

higher doses of alcohol cause depression

Due to combination of central effects and direct depression of the myocardium

Metabolic effects

Severe Hypoglycemia because of inhibition of gluconeogenesis.

Alcohol induced hypoglycemia within 6–36 hours of heavy drinking

typically seen : undernourished individuals or one who has not eaten for the previous 24 hours.

Death From Alcohol Poisoning

Alcohol intoxication may result in death

- respiratory failure

- circulatory failure

- result of aspiration of stomach contents in the absence of a gag reflex.

blood alcohol above 500 mg/100 mL are “probably fatal”

Lower level of blood alcohol may result death when there are other therapeutic and non therapeutic drugs



Diagnosis of Intoxication

Alcohol intoxication and drunkenness are often used interchangeably.

behavioral changes associated with drunkenness are seen when they believe they have consumed alcohol but actually have not

Therefore drunkenness means the behavioral aspect that is resulted after consumption of alcohol

Intoxication with alcohol

behavioural and physical abnormalities seen when the quantity of alcohol the person consumes exceeds the individual's tolerance for alcohol.
(drinking too much too fast)



Diagnostic Criteria for Alcohol Intoxication—DSM-IV

- Recent ingestion of alcohol
- Clinically significant maladaptive behavioral or psychological changes (e.g. inappropriate sexual or aggressive behavior, mood lability, impaired judgment, impaired social or occupational functioning) that developed during, or shortly after, alcohol ingestion.
- One (or more) of the following signs, developing during, or shortly after, alcohol use:
 - slurred speech
 - unsteady gait
 - impairment in attention or memory
 - incoordination
 - nystagmus
 - stupor or coma
- The symptoms are not due to a general medical condition and are not better accounted for by another mental disorder.

DSM-IV, *Diagnostic and Statistical Manual of Mental Disorders*, 4th ed.

Pathological States Simulating Alcohol Intoxication

Severe head injuries

Metabolic disorders (hypoglycemia, hyperglycemia, uremia, hyperthyroidism)

Neurological conditions associated with dysarthria, ataxia, tremor, drowsiness (Multiple sclerosis, intracranial tumors, Parkinson's disease, epilepsy, acute vertigo)

The effects of drugs, either prescribed or illicit (insulin, barbiturates, benzodiazepines, cocaine)

Psychiatric disorders (e.g., hypomania, general paresis)

High fever

Carbon monoxide

Alcohol Dependence & Withdrawal

Diagnosis of alcohol dependence (criminality is associated)

Uncomplicated Alcohol Withdrawal

- Nausea and vomiting.
- Malaise and weakness.
- Autonomic hyperactivity (raised blood pressure and tachycardia).
- Anxiety, depressed mood, and irritability.
- Transient hallucinations and illusions.
- Headache and insomnia.

Alcohol Withdrawal Delirium (72-96hrs after drink)

- impaired attention and memory, disorganized thinking, disorientation,
- reduced level of consciousness,
- perceptual disturbances, visual hallucinations, tactile or auditory hallucination, and agitation.
- Withdrawal seizures
- Cardiac arrhythmia
- Metabolic disorders - Wernicke's encephalopathy & Korsakoff's psychosis

Medical opinion of a person referred for medico-legal purpose with a history of consumption of alcohol /drunkenness

Whether a person has consumed alcohol/ under the influence of alcohol /intoxicated with alcohol

Whether there are other associated conditions including injuries

Whether the person taken to custody is fit to detain (there is existing threat to life, eg withdrawal status, or other conditions)

Summary

Alcohol is associated with criminal behaviors, accidents and death.

Liquors contain alcohol i.e. ethanol ($\text{C}_2\text{H}_5\text{OH}$)

Different liquors/ beverage contain different % of alcohol varying from 4%-60%

Rate of alcohol absorption is $>$ elimination.

5-10% excreted unchanged in breath, urine, and sweat.

Increasing blood alcohol levels in the body produce different in the behavior stages

- euphoria, excitement, drunkenness, stupor, coma and death

Effects of alcohol in body is important to diagnose intoxication especially those in CNS & eye.

Diagnosis of alcohol intoxication include recent ingestion of alcohol, maladaptive behavior, one or more of CNS depressant sings and exclusion of illnesses that mimic those signs

Death from alcohol poisoning occur above 300mg/100mL of blood

Knowledge on alcohol effect and withdrawal is important in recommendation of fitness for detention

Recommended reading

Clinical forensic Medicine – A physician's guide 2nd Edition by Margaret M Stark

Simpson's Forensic Medicine

Lecture notes in Forensic Medicine- Dr. L.B.L de Alwis

Clinical Forensic Medicine eds W.D.S. MacLay

A close-up photograph of several purple orchids with white centers. The flowers are in various stages of bloom, with some fully open and others as buds. The background is a soft, out-of-focus purple. The text "THANK YOU" is overlaid in the center in a white, sans-serif font. A thin horizontal line is visible on the right side of the image, extending from the text area towards the right edge.

THANK YOU