


The background of the slide is a microscopic image of lung tissue, likely stained with hematoxylin and eosin (H&E). It shows a network of alveolar spaces (appearing as white or light pink areas) separated by thin, pink-stained alveolar septa. Some areas show more dense staining, possibly indicating inflammation or consolidation.

Pneumonia

Dr Roshitha de Silva
Department of Pathology
Faculty of Medicine
University of Kelaniya

Pneumonia

- Acute inflammation of lung parenchyma distal to the terminal bronchioles
- Consist of
 - Respiratory bronchiole
 - Alveolar ducts
 - Alveolar sacs
 - Alveoli



Pneumonia Pathophysiology

Microbial pathogens enter the lung by:


- Inhalation of infectious aerosols
- Hematogenous dissemination
- Aspiration of organisms from oropharynx
- Tracheal intubation

Classification

- Aetiology
 - Bacteria
 - Virus
 - Fungal
- Patient setting
 - Community acquired
 - Hospital acquired
- Immune status
 - Healthy
 - Immuno-compromised
- Site
 - Lobar pneumonia
 - Bronchopneumonia
 - Interstitial pneumonia


Lobar pneumonia


- Acute bacterial infection of an entire lobe or a part of the lobe
- 95 % of cases by *Streptococcus pneumoniae*
- Lobar pneumonia evolves in four stages.
 - Congestion
 - Red hepatisation
 - Grey hepatisation
 - Resolution



Congestion (1 to 2 days)


- Micro
 - Dilatation and congestion of capillaries in the alveolar walls
 - Pale eosinophilic oedema fluid in the air spaces
 - Numerous bacteria in the alveolar fluid
 - Few rbc and neutrophils in the intra-alveolar fluid






Congestion


- Early acute inflammatory response to bacterial infection.
- Macro
 - Enlarged
 - Heavy
 - Dark red
 - Congested



Red Hepatisation (2-4 days)


- Micro
 - Oedema fluid is replaced by strands of fibrin
 - Marked cellular exudate of neutrophils and extravasated rbc





Red Hepatisation

- Liver-like consistency
- Macro
 - Red
 - Firm
 - Consolidated (airless)
 - Cut surface
 - Airless
 - Dry




Pleuritis


- A pleural reaction is also associated if the consolidation extends to the surface
- Inflammation of the pleura
- Fibrinopurulent exudate → pleural effusion
- Exudate
 - Resorbed
 - Organized by fibroblasts
 - Leading to fibrous thickening
 - Adhesions → fibrinous tags



Grey Hepatisation (4-8 days)

- Micro
 - Fibrin strands are dense and more numerous
 - Cellular exudate of neutrophils is reduced
 - rbc are sparse
 - Macrophages begin to appear
 - Cellular exudate is often separated from the septal walls by a thin clear space
 - Organisms are less





Grey Hepatisation

- Macro
 - Firm
 - Heavy
 - Cut surface
 - dry
 - Granular
 - Grey



Resolution

(Begins by 8th day completed in 1-3 weeks)

- Macs are the predominant cells in alveolar spaces
- Fragments of fibrin seen in alveolar spaces
- Alveolar capillaries are engorged
- Collected fluid removed
 - Lymphatics
 - Expectoration
- Restoration of normal lung parenchyma
- Pleuritis may resolve or undergo organisation → fibrous thickening or permanent adhesions

Resolution

- Macro
 - Solid fibrinous material is liquefied by enzymatic action
 - This restores the normal aeration
 - Cut surface
 - Dirty brown
 - Pleural reaction show resolution

Bronchopneumonia

- Infection of the terminal bronchioles
- First cause bronchiolitis → spread through resp passage → reach alveoli
- Results in patchy consolidation
- Common in
 - Extremes of life
 - 2^{ry} infection following viral infection
 - Chronic debilitating illness



Bronchopneumonia

- Common causative organisms
 - Staphylococci
 - Streptococci
 - Pneumococci
 - *Klebsiella pneumoniae*

Bronchopneumonia morphology

Macroscopy


- Frequently bilateral
- Patchy areas of red/grey consolidation
- More often involves the lung bases
- Cut surface
 - 3-4 cm in diameter
 - Poorly demarcated margins
 - Elevated, dry, granular, firm
 - Centered around a bronchiole
- Unaffected areas appear normal




Bronchopneumonia morphology

Microscopy

- Acute bronchiolitis
- Suppurative exudate in bronchioles
- Inflammation in surrounding alveoli
- Less involved alveoli contain oedema fluid





Complications

- Organisation
 - Resolution does not occur
 - Fibrosed, tough, airless leathery lung tissue
 - Results in carnification

Complications


- Pleural effusion
- Empyema
- Lung abscess
- Metastatic infection
 - Infection may extend into the pericardium and the heart → purulent pericarditis, bacterial endocarditis, myocarditis

Complications

- Bronchopneumonia
 - Complete resolution is uncommon
 - There is some degree of destruction to the bronchiole
 - This results in foci of bronchiolar fibrosis and end up with bronchiectasis

Atypical pneumonia

- Caused by atypical bacteria and viruses
- ‘Walking pneumonia’
- Macroscopy
 - Patchy or lobar areas of congestion without consolidation
- Microscopy
 - Interstitial inflammation
 - Alveoli are unaffected



Lung abscess



- Localised area of necrosis of lung tissue with suppuration


Lung abscess

- Macro
 - Single/multiple
 - Basal/scattered
 - Poorly defined ragged wall
 - Abscess cavity is filled with exudate
 - Advance abscess- fibroblastic proliferation
→ fibrous wall



Lung abscess

- Micro
 - Destruction of the lung parenchyma
 - Suppurative exudate
 - Cavity initially surrounded by Acute infla cells then by macs, lymphocytes



Thank You