Emerging & Re-emerging Diseases

Dr.B.D.W.Jayamanne Department of Public Health


Outline

- ► Introduction
- ► Causes
- Epidemiology
- Examples
- Control Measures
- ▶ Challenges



Introduction


- Disease Infections, Genetic, Toxines
 - "conditions that impair normal tissue function"
 - ► Eg: cystic fibrosis, atherosclerosis, measles, etc
- Infectious disease
 - A disease that is caused by the invasion of a host by agents whose activities harm the host's tissues
 - Can be transmitted to other individuals -infectious
 - "harm the host's tissues" -disease
 - ► Infectious diseases have for centuries ranked with wars and famine as major challenges to human progress and survival (Morens et al.2014,Nature)



Infectious diseases	Annual deaths (millions)
Respiratory infections	3.96
HIV/AIDS	2.77
Diamhoeal diseases	1.80
Tuberculosis	1.56
Vaccine-preventable childhood diseases	1.12
Malaria	1.27
STDs (other than HIV)	0.18
Meningitis	0.17
Hepatitis B and C	0.16
Tropical parasitic diseases	0.13
Denique	0.02
Other infectious diseases	1.76

- Infectious diseases are the 2nd leading cause of mortality in the world
- ▶ 15% of all deaths worldwide
- Emergence of new infections -continuous process

NUMBER OF EMERGING INFECTIOUS DISEASE EVENTS



- What are Neglected Tropical Diseases?
 - ► A diverse group of communicable diseases that prevail in tropical and subtropical conditions in 149 countries
 - ► Affect more than one billion people and cost developing economies billions of dollars every year
 - ► Populations living in poverty, without adequate sanitation and in close contact with infectious vectors and domestic animals and livestock are those worst affected.




- Who defines an emerging or re-emerging infectious disease?
 - National Institute of Allergy and Infectious Disease (NIAID), Bethesda, U.S.A. (Established in 1887)
- Emerging Infectious Disease (EID)
 - A completely new infection or has recently increased in incidence or impact and severity, affected newer geographical locations, or is an existing disease that has recently developed new clinical pattern, or developed resistance to existing therapy
 - (Daszak at. el. Anthropogenic change, biodiversity loss, and a new agenda for emerging diseases. J Parasitol Suppl. 2003;89:37–41)
 - During past 2 decades



- Emerging Infectious Disease (EID)
 - Previously unknown or known infectious diseases whose incidence in humans has significantly increased in the past 2 decades
 - ► Eg: SARS ,Novel Influenza(H1N1) ,Avian Influenza,MERS
- Re-emerging Infectious Diseases
 - ► Known infectious diseases that have reappeared after a significant decline in incidence
 - ► Eg: Multidrug resistant TB(MDR-TB), MRSA, VRE, PRSP




335 emerging infectious disease events during 1940 to 2004



Jones KE et al, Nature, 2008




Causes



Causes...

- Antibiotic resistance and bacterial evolution
 - Eg TB 1940 Streptomycin -> 1980 resistance -> MD -TB
 - o 1970 -Hope to eradicate TB ???
 - 1993 -TB -Global Health emergency by WHO
 - MRSA, "pan resistant" Klebsiella pneumoniae
- Insecticide resistance
 - 1960s -Sri Lanka eliminated Malaria with DDT
 - 1970 Resistant mosquitoes to DDT
- Vaccine driven evolution
 - Vaccine induced immunity -Hep.B,Strep. pneumoniae
- Vaccine refusal
 - MMR,DPT
 - Due to side effects
 - Cause outbreaks

Contributing Factors



HOST

- Human demographic change (inhabiting new areas)
- Human behaviour (sexual activity & drug use)
- Human susceptibility to infection (Immunosuppression)
- Poverty & social inequality
- Increased international travel



Contributing Factors...

ENVIRONMENT

- Climate & changing ecosystems (El Nino, deforestation , Global warming, etc)
- ► Economic development & Land use (urbanization, deforestation)
- Technology & industry (food processing, food trade & handling)
- International travel & commerce
- Breakdown of public health measure (war, unrest, overcrowding)
- Deterioration in surveillance systems (lack of political will)




Contributing Factors...

AGENT

- Evolution of pathogenic infectious agents (microbial adaptation & change, virulence)
 - High replication rate -new generations
- Development of resistance to drugs
- Resistance of vectors to pesticides- behavioural change-Malaria- bed nets

ANTIMICROBIAL RESISTANCE



- Pathogens -
 - Group I Pathogens newly recognized during past 2 decades
 - Group II Re-emerging pathogens
 - Group III Agents with Bioterrorism potential (NIAID classification)
 - NIAID Category A Priority (person to person transmitted agents + Viral Haemorrhagic Fevers)
 - NIAID Category B Priority(food and water borne + Viral Encephalitis)
 - NIAID Category C Priority(Emerging infectious diseases)



Group I pathogens

- Acanthamebiasis
- Australian bat lyssavirus
- Babeasia (Atypical)
- Bartonella henselae
- Ehrlichiosis
- Hepatitis C (new in FY14)
- Hepatitis E (new in FY14)
- Human herpesvirus 6
- Human herpesvirus 8
- Parvovirus(B19)
- Borrelia mayonii (new in FY18)
- Borrelia miyamotoi (new in FY14)
- Helicobacter pylori
- etc.

Group II pathogens

- Enterovirus 71
- Clostridium difficile
- Mumps virus
- Streptococcus Group A
- Staphylococcus aureus



Group III- Cat.A

- Can be easily disseminated or transmitted from person to person
- Result in high mortality rates and have the potential for major public health impact
- Might cause public panic and social disruption
- Require special action for public health preparedness

Group III- Cat. B

- Are moderately easy to disseminate
- Result in moderate morbidity rates and low mortality rates
- Require specific enhancements for diagnostic capacity and enhanced disease surveillance

GroupIII- Cat. C

- Emerging pathogens that could be engineered for mass dissemination in the future because of
 - Availability
 - ► Ease of production and dissemination
 - Potential for high morbidity and mortality rates and major health impact Category A Priority Pathogens



Group III- Cat.A

- ► Bacillus anthracis (anthrax)
- Clostridium botulinum toxin (botulism)
- Yersinia pestis (plague)
- Variola major (smallpox) and other related pox viruses
- Francisella tularensis (tularemia)
- Viral hemorrhagic fevers
 - Arenaviruses
 Junin, Machupo, Guanarito, Chapare (new in fiscal year (FY14), Lassa, Lujo (new in FY14)
 - Bunyaviruses
 Hantaviruses causing Hanta Pulmonary syndrome, Rift Valley Fever, Crimean
 Congo Hemorrhagic Fever
 - ► Flaviviruses
 - Dengue
 - Filoviruses
 - Ebola
 - Marburg



Group III- Cat. B

- ► Burkholderia pseudomallei (melioidosis),Coxiella burnetii (Q fever)
- Brucella species (brucellosis), Burkholderia mallei (glanders)
- Chlamydia psittaci (Psittacosis)
- Ricin toxin (Ricinus communis), Epsilon toxin (Clostridium perfringens),
 Staphylococcus enterotoxin B
- Typhus fever (Rickettsia prowazekii)
- Food- and waterborne pathogens
 - -Bacteria, Diarrheagenic E.coli , Pathogenic Vibrios, Shigella species, Salmonella, Listeria monocytogenes, Campylobacter jejuni, Yersinia enterocolitica
 - -Viruses, Caliciviruses, Hepatitis A, Protozoa, Cryptosporidium parvum, Cyclospora cayatanensis
 - -Giardia lamblia,Entamoeba histolytica,Toxoplasma gondii,Naegleria fowleri (new in FY14),Balamuthia mandrillaris (new in FY14)
 - Fungi Microsporidia
 - -Mosquito-borne viruses West Nile virus (WNV),LaCrosse encephalitis (LACV), California encephalitis,Venezuelan equine encephalitis (VEE),Eastern equine encephalitis (EEE),Western equine encephalitis (WEE),Japanese encephalitis virus
- (JE),St. Louis encephalitis virus (SLEV),Yellow fever virus (YFV),Chikungunya virus Zika virus www.niaid.nih.gov/research/emerging-infectious-diseases-pathogens

GroupIII- Cat. C

- ► Nipah and Hendra viruses
- Additional hantaviruses
- Tick Borne hemorrhagic fever viruses
 - Bunyaviruses -Severe Fever with Thrombocytopenia Syndrome virus (SFTSV),
 Heartland virus
 - Flaviviruses Omsk Hemorrhagic Fever virus, Alkhurma virus, Kyasanur Forest virus
- Tickborne encephalitis complex flaviviruses
- Tickborne encephalitis viruses European subtype,Far Eastern subtype,Siberian subtype,Powassan/Deer Tick virus
- ► Tuberculosis, including drug-resistant TB
- Influenza virus
- Other Rickettsias
- Rabies virus
- Prions
- Coccidioides spp.
- Severe acute respiratory syndrome associated coronavirus (SARS-CoV),
- ► MERS-CoV, and other highly pathogenic human coronaviruses (new in FY14)

NTD's


- 1.Buruli ulcer
- 2. Chagas disease
- 3. Dengue and Chikungunya
- 4. Dracunculiasis (guinea-worm disease)
- 6. Echinococcosis
- 7. Foodborne trematodiases
- 8. Human African trypanosomiasis (sleeping sickness)
- 9.Leishmaniasis
- 10.Leprosy(Hansen's disease)
- 11.Lymphatic filariasis
- 12. Mycetoma, chromoblastomycosis and other deep mycoses
- 13.Onchocerciasis (river blindness) 14.Rabies
- 15. Scabies and other ectoparasites 16. Schistosomiasis
- 17. Soil-transmitted helminthiases 18. Snakebite envenoming
- 19.Taeniasis/Cysticercosis 20.Trachoma
- 21. Yaws (Endemic treponematoses)

Epidemiology...


- ▶ 61% of human infections derived from wildlife(Zoonotic diseases)
 - ► Eg.Bubonic plague (14th century) 1/3 of human population died -transmitted through rat flea
 - Smallpox-?rodents -10,000 years ago
 - Influenza (1918) 100 million deaths Birds-> humans & pigs
 - ► HIV 1 Million deaths per year-Chimpanzee
 - Dengue -primates (by Aedes mosquito)
- Many of them RNA viruses
 - High mutation rate -enzymes that replicate RNA viruses are more prone to error
 - More agile at change- when they cross barriers- adapting to new hosts



Epidemiology



Emerging and re-emerging infectious diseases Many are caused by RNA viruses



Adapted from Figure 1 of Morens et al, Nature, 2004



Epidemiology...

- Can be
 - Region specific
 - Country specific
 - ▶ Local area specific
 - Time/Period specific
 - Regions
 - Latin America
 - South East Asia
 - ► China
 - ▶ Africa
- But can be a global problem



Emergent zoonotic virus diseases

Table 1. Examples of novel, emergent zoonotic virus diseases

Year of isolation	Place of isolation	Virus	Reservoir/spillover host
1991	Venezuela	Guanarito virus ²⁰	Rodents
1992	Slovenia	Dobrava virus ²¹	Rodents
1993	United States	Sin Nombre virus ²²	Rodents (Peromyscus maniculatus)
1994	Brisbane, Australia	Hendra virus ²³	Fruit bats (Pteropus sp.)/horses*
	Sao Paolo, Brazil	Sabia virus ²⁴	Rodents
1995	Florida, USA	Black Creek Canal virus ²⁵	Rodents
1996	Ballina, Australia	Australian bat lyssavirus ²⁶	Fruit and insectivorous bats
	Argentina	Andes virus ²⁷	Rodents
1997	Hong Kong (China)	Influenza H5N1 ²⁸	Wild birds/domestic poultry*
	Menangle, Australia	Menangle virus ²⁹	Fruit bats
	Saudi Arabia	Alkhurma virus ^{30,31}	Camels and sheep [†]
1999	Peninsular Malaysia	Nipah virus ^{32,33}	Fruit bats/pigs*
2000	Peninsular Malaysia	Tioman virus ³⁴	Fruit bats
2002-2003	China, Hong Kong (China)	SARS coronovirus35-38	Bats/civets?*
2003-2004	Viet Nam, China	Influenza H5N139,40	Wild birds/domestic poultry*
2007	Melbourne, Australia	Dandenong arenavirus ⁴¹	Rodents?
	Peninsular Malaysia	Melaka virus ⁴²	Fruit bats?
	Uganda	Bundibugyo ebolavirus ⁴³	Fruit bats?/various animals (bush meat)
2008	Lukasa, Zambia	Lujo virus ⁴⁴	Unidentified rodents
	Perak, Malaysia	Kampar virus ⁴⁵	Fruit bats?

^{*} Spillover host; † Tick-borne



Challenges

- Preparedness
- Early & accurate surveillance
- Rapid response
- Prevention and control measures
- Early identification of pathogen
- Multi-sectoral cooperation
- Political commitment to invest of the uncertainty
- Cross-border issues



Sri Lanka -CDC

Podcasts

Announcements

More Content

FIDeas for the Public

EMERGING INFECTIOUS DISEASES®

CDC **EID** journal September 2018 f 💆 🛨 Manuscript Submission << Return to Article Map About the Journal Article Information - Sri Lanka Past Issues All Issues Subscribe Ahead of Print / In Press Click column headers to sort. Advanced Article Search Show 50 ▼ entries Search: Articles by Country Search Title Volume -Issue 🔻 Article Type Year -Month Author Resource Center Dengue Virus Type 2 in Travelers Returning to Japan from Sri Lanka, 2017 11 Research Letter 2017 11 Medscape CME Lyssavirus in Indian Flying Foxes, Sri Lanka Dispatch 2016 8

Reemergence and Autochthonous Transmission of Dengue Virus, Eastern

Molecular Epidemiology of Influenza A(H1N1)pdm09 Virus among

Donmie Virus Transmission by Blood Stem Cell Donor after Travel to Sri

Humans and Swine, Sri Lanka

ISSN: 1080-6059

2015

2014

9

12

21

20

9

12

Dispatch

Dispatch

Pandemic

- An epidemic occurring worldwide, or over a very wide area, crossing international boundaries and usually affecting large number of people
- WHO requirements
 - Ability to infect humans
 - Ability to cause disease in humans
 - Spread from human to human
- Pandemic phase groupings
- Predominantly animal infections; few human infections (phases 1-3)
- 2. Sustained human-to-human transmission (phase 4)
- 3. Widespread human infection (phases 5-6/pandemic)
- 4. Possibility of recurrent events (post peak)
- 5. Disease activity at seasonal levels (post pandemic)

Emerging diseases - in 21st century

- SARS (1st to emerge in 21st century)
- Highly Pathogenic Avian Influenza (H5N1)
- Novel Swine origin Influenza A (H1N1)
- Ebola virus disease
- Middle East Respiratory Syndrome (MERS)
- HIV
- H.Pylori
- Listeriosis
- Zika
- Nipah Virus



Re-emerging Diseases

- Diphtheria Early 1990s epidemic in Eastern Europe(1980- 1% cases; 1994- 90% cases)
- Cholera in 1998 (new strain eltor, 0139)
- Human Plague India (1994) after 15-30 years absence
- Dengue/ DHF- Over past 40 years, 20-fold increase to nearly 0.5 million
- Polio -
- ► TB
- Malaria -Drug resistant
- Leptospirosis




SARS: The First Emerging Infectious Disease Of The 21st Century


- Severe Acute Respiratory Syndrome (SARS) emerged in 2002/2003.
- First case was recognized in Hanoi, Vietnam in February 2003 in a businessman who had travelled from Guandong Province, China through Hong Kong.
- Cases reported from November 2002 to July 2003
- Origin Guandong Province, China



SARS- Spread



SARS...



Source: www.who.int.csr/sars

SARS - Prevention

- Hand hygiene in both healthcare settings and the community
- Cough etiquette
- Protective barriers
- Good waste disposal

Dr Carlo Urbani (1956-2003)



- Italian microbiologist and expert on communicable diseases
- Worked for WHO in Vietnam was the first to identify SARS as an emerging infectious disease
- His work led to strengthening of global surveillance and preventive activities worldwide and ultimately to the control of the epidemic
- He died of SARS on 29th March 2003

Lesson learnt from SARS

- An infectious disease in one country is a threat to all
- Important role of air travel in international spread
- Tremendous negative economic impact on trade, travel and tourism, estimated loss of \$ 30 to \$150 billion
- High level commitment is crucial for rapid containment
- WHO can play a critical role in catalyzing international cooperation and support
- Global partnerships & rapid sharing of data/information enhances preparedness and response



Highly Pathogenic Avian Influenza (H5N1)

- First infected humans in 1997 in China
- Re-emerged in 2003-2004
- Since Nov 2003, avian influenza H5N1 in birds affected 60 countries across Asia, Europe, Middle-East & Africa
- >220 million birds killed by AI virus or culled to prevent further spread
- Majority of human H5N1 infection due to direct contact with birds infected with virus



Novel Swine origin Influenza A (H1N1)

- Swine flu causes respiratory disease in pigs high level of illness, low death rates
- Pigs can get infected by human, avian and swine influenza virus
- Occasional human swine infection reported
- In US from December 2005 to February 2009, 12 cases of human infection with swine flu reported
- Outbreak occurred in 2009 in Mexico & USA
- Spread across states in USA
- Cases also recorded in Canada, Germany, Spain and UK



Novel Swine origin Influenza A (H1N1)...


- April 15th CDC identifies H1N1 (swine flu)
- April 25th WHO declares public health emergency
- April 27th Pandemic alert raised to phase 4
- April 29th Pandemic alert raised to phase 5
- By May 5th more than 1000 cases confirmed in 21 countries
- Screening at airports for flu like symptoms (especially passengers coming from affected area)
- Schools closed in many states in USA
- May 16th India reports first confirmed case
- Stockpiling of antiviral drugs and preparations to make a new effective vaccine



Novel Swine origin Influenza A (H1N1)...

New Influenza A (H1N1), Number of laboratory confirmed cases and deaths as reported to WHO

Status as of 27 May 2009 06:00 GMT



The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

Data Source: World Health Organization
Map Production: Public Health Information
and Geographic Information Systems (GIS)
World Health Organization



© WHO 2009, All rights reserved



Novel Swine origin Influenza A (H1N1)...

Worldwide- 162,380 cases 1,154 deaths

India- 558 cases 1 death



Ebola Virus Disease (EVD)

- Acute serious illness with high fatality
- Two simultaneous outbreaks in 1976 Sudan and Congo
- Emerged close to Ebola river in Congo
- Case fatality rate is around 50%. (varied from 25% to 90% in past outbreaks)
- Declared a 'Public Health Emergency of International Concern' in 2005
- First case notified in West Africa in March 2014
- Spread across the continent through air & land travel
- Spread to Guinea, Sierra Leone, Liberia, Nigeria, USA, Senegal & Mali



EVD - Recent outbreaks



About us v

Health topics ∨


News v

Countries >


Table: Chronology of previous Ebola virus disease outbreaks

Year	Country	Ebolavirus species	Cases	Deaths	Case fatality
2015	Italy	Zaire	1	0	0%
2014	DRC	Zaire	66	49	74%
2014	Spain	Zaire	1	0	0%
2014	UK	Zaire	1	0	0%
2014	USA	Zaire	4	1	25%
2014	Senegal	Zaire	1	0	0%
2014	Mali	Zaire	8	6	75%
2014	Nigeria	Zaire	20	8	40%
2014-2016	Sierra Leone	Zaire	14124*	3956*	28%
2014-2016	Liberia	Zaire	10675*	4809*	45%
2014-2016	Guinea	Zaire	3811*	2543*	67%
2012	Democratic Republic of Congo	Bundibugyo	57	29	51%
2012	Uganda	Sudan	7	4	57%
2012	Uganda	Sudan	24	17	71%
2011	Uganda	Sudan	1	1	100%
2008	Democratic Republic of	7aire	32	14	44%

EVD -Spread



EVD -Transmission



- Natural hosts are fruit bats
- Human to human transmission through direct contact with blood, secretions, organs or other bodily fluids of infected people
- Contamination through surfaces and material (eg: bedding, clothing)
- No formal evidence of sexual transmission



EVD -Prevention

- Personal hygiene hand washing following contact with body fluids
- Avoid handling items with possible contact with infected person
- Avoid funeral or burial rituals that require handling infected dead persons
- Avoid contact with bats or non-human primates or body fluids
- Avoid facilities where patients are treated
- Prevent sexual contact with a person recovered from EVD



Others

- Hanta viral disease
 - Hantavirus pulmonary syndrome(HPS)
 - Haemorrhagic fever with renal syndrome
- Bovine Spongiform Encephalopathy (Mad Cow Disease) -1986 -UK
- vCJD (Transmissible Spongiform Encephalopathy (TSE))- Novel Prion disease - 1995



Sri Lanka??

Leishmaniasis


- First locally transmitted CL case reported in Ambalantota in 1992
- First VL case reported in 2006 in Anuradhapura
- Reported in clusters around Hambantota & Polonnaruwa district
- 940 suspected cases of CL by end of 2011
 A notifiable Disease

Risk factors


- Socioeconomic conditions poverty, poor housing
- Malnutrition vitamin A, Zinc and protein deficiency
- Population mobility
- Environmental change urbanization
- ▶ Climate change



Key tasks - carried out by whom?



What skills are needed?



Multiple expertise needed!



"Today's world is truly a global village – one can safely predict that infectious diseases will continue to emerge"

"Depending on the adequacy of our response and reactions to this threat, ... the situation could lead to a catastrophic storm of microbial threats"



Thank you