



Antiplatelet drugs and thrombolytics

Dr.S.A.F.Kurukulasuriya



Platelets and haemostasis


- Support in 3 ways

Physical barrier-sticking to exposed collagen


Accelerates activation of coagulation proteins

Vasoconstriction(release of storage granules)


A. VASOCONSTRICTION




B. PRIMARY HEMOSTASIS



C. SECONDARY HEMOSTASIS



D. THROMBUS AND ANTITHROMBOTIC EVENTS






Platelet mechanisms



- Platelet aggregation does not occur
 - As long as the resting conformation of GPIIb/IIIa is maintained.
 - High concentrations of cyclic AMP is maintained within the platelet



Antiplatelet drugs

- Cyclooxygenase inhibitors
 - Aspirin
- ADP receptor inhibitors
 - Clopidogrel, ticlopidine
- Phosphodiesterase inhibitors
 - Cilostazol
- Glycoprotein II b/IIIa inhibitors
 - Abciximab, Tirofiban
- Adenosine reuptake inhibitors
 - Dipyridamole

- 
-
- **Cyclooxygenase (COX)**, officially known as **prostaglandin-endoperoxide synthase (PTGS)**, is an enzyme that is responsible for formation of prostanoids, including prostaglandins such as prostacyclin and thromboxane.





Aspirin

- Acetylates and thus inactivates COX
- Thus can prevent formation of TXA2 and PGI2
- These actions can be separated (abolish only the effect of TXA2) by using a ***low dose (75-100mg)/d***
- The acylation of COX is irreversible and lost for the life time of the platelet
- Even the low dose can cause increased risk of GI bleeds.



Dipyridamole

- Reversibly inhibits platelet phosphodiesterase → increase c.AMP levels in the platelet → platelet reactivity reduced
- T_{1/2} is 12 hrs



Ticlopidine

- Thienopyridine derivative
- Inhibits ADP dependent platelet aggregation
- T_{1/2} is 40 hrs
- More effective than aspirin in reducing strokes in patients with TIAs
- Reduces the risk of the combined outcome of stroke, MI



Ticlopidine

- Neutropenia is the most serious side effect (risk 24%)



Clopidogrel

- Thienopyridine derivative
- More effective than aspirin in prevention of ischaemic stroke, MI or vascular death.
- More expensive than aspirin but cheaper than ticlopidine.



Glycoprotein II b/IIIa antagonists

- Glycoprotein II b/IIIa molecule mediates platelet aggregation via binding of adhesive proteins such as fibrinogen or Von Willibrands factor.
- Block a receptor on the platelet for fibrinogen or Von Willibrands factor.
- E.g.- abciximab, eptifibatide, tirofiban



Abciximab

- Human-murine chimeric monoclonal antibody
- The Fab fragment binds to the GP II b/IIIa complex with high affinity.
- It reduces the rate of death, MI, need for urgent CABG after percutaneous coronary angioplasty.
- Refractory unstable angina prior to percutaneous coronary intervention.



Side effects of abciximab

- Haemorrhage esp. high risk after failed fibrinolytic therapy (for acute MI).
- Platelet transfusion may be necessary.
- Thrombocytopenia may occur.



Other drugs


- Dextrans

Dextran 70 reduces post operative venous thromboembolism



Uses of antiplatelet drugs

- Non fatal MI and non fatal stroke reduced by one third.
- Death from any vascular cause reduced by one sixth.
- Aspirin+/-Dipyridamole reduce the risk of occlusion of vascular grafts.



Thrombolytic (fibrinolytic) drugs

Plasminogen in the blood

Fibrinolytics

Activators of plasminogen in the blood

Antifibrinolytics

Inhibitors of plasminogen & plasmin

Antistreptase

Urokinase

streptokinase

+


-

Tranexaemic acid

Plasminogen binds to fibrin

Plasminogen formed on fibrin which it destroys

Fibrin degradation product





The thrombolytic system

- Removal of products of coagulation is needed for the preservation of the intact vascular system.
- The system depends on formation of fibrinolytic enzyme plasmin from its precursor protein plasminogen.
- Fibrinolytics can remove established thrombi and emboli.



Drugs that promote fibrinolysis

- Important in dissolving thrombi in acute myocardial infarction. It limits necrosis of the ischaemic myocardium. Hence it can improve prognosis.
- The approach is to give a plasminogen activator intravenously so that the fibrinolytic enzyme plasmin is formed in increased amounts.



Fibrinolytics

- Non-fibrin selective-(Not well absorbed by fibrin thrombi)
 1. streptokinase
 2. Anistreplase
 3. urokinase
- Fibrin selective(bind strongly to fibrin,can dissolve aging or resistant thrombi)
 1. Recombinant prourokinase
 2. Altiplase



Streptokinase

- Derived from beta-haemolytic streptococci.
- Converts plasminogen to plasmin.
- Rapid administration may cause a fall in blood pressure.



Anistreplase

- This is the plasminogen-streptokinase complex.
- Converts plasminogen to plasmin.
- Longer action than streptokinase.



Urokinase

- Made from human fetal kidney cells.
- Direct activator of plasminogen



Recombinant prourokinase

- Produced by recombinant DNA technology
- Binds to fibrin and converts to urokinase



Alteplase

- Produced by recombinant DNA technology



Uses of thrombolytic drugs

Coronary artery thrombolysis

- Reduce mortality with an acceptable frequency of adverse effects.
- Streptokinase & t-PA reduce mortality. This effect is better when combined with aspirin.



Adverse effects of coronary thrombolysis

- Bleeding-esp. at a vascular lesion
- Multiple micro emboli
- cardiac arrhythmias (due to reperfusion)
- Allergy (avoid reuse within 5 to 12 months)



Non coronary thrombolysis

- Pulmonary thrombolysis

Beneficial with evidence of
haemodynamic decompression-
Alteplase

- Deep vein thrombosis

Esp. if affected vessels are proximal

- Distal arterial occlusion(Distal to the popliteal)