

Nephrotic syndrome


Prof. Wasantha Karunasekera



Nephrotic syndrome



Introduction

- ▶ Chronic illness
 - ▶ Recurring disease – Relapses and remissions
 - ▶ Usually grow out of the disease after 12 years
 - ▶ Most likely aetiology is minimal change nephrotic syndrome
 - ▶ Risk of end stage renal disease – rare
- 

Definition

- ▶ Oedema
- ▶ Hypoalbuminaemia (s. albumin $<25\text{g/l}$)
- ▶ Gross proteinuria (nephrotic range)
urine protein/creatinine ratio
 $>200\text{mg/mmol}$ or
3+ protein on urine dipstick or
proteinuria $\geq 300\text{mg/dl}$
- ▶ Hyperlipidaemia (Cholesterol $> 200\text{mg/dl}$)

Remission

Urine albumin nil/trace ($<4\text{mg}/\text{m}^2/\text{hr}$) for 3 consecutive early morning specimens

or

Urine protein creatinine ratio $<20\text{mg}/\text{mmol}$



Relapse

- urine albumin $\geq 3+$ (40mg/m²/hr) on three consecutive early morning specimens, having been in remission previously
- or
- Urine protein creatinine ratio ≥ 200 mg/mmol

Frequent relapsing

Two or more relapses during first 6 months after the initial episode or 4 or more relapses in any 12 months

Response to steroids

Steroid responsive

Remission achieved within initial 4 weeks of high dose daily steroid therapy

Steroid dependent

Two consecutive relapses while on alternate day steroids or within 14 days after cessation of steroids

Steroid resistance

Failure to achieve remission within 4 weeks of therapy with daily prednisolone (60mg/m²/d)

Classification

- ▶ **Primary (Idiopathic)** – most common
- ▶ **Secondary** – secondary to systemic disorders, or following drugs and heavy metals
 - (SLE, HSP, Vasculitis, lymphoma, leukemia)
 - (infections – Hepatitis B,C, HIV, leprosy, IMN, syphilis)
 - (captopril, penicillamine, gold, NSAIDS)
- ▶ **Congenital** – manifesting at birth or within first 3 months of life.
 - poor prognosis

Primary (Idiopathic) nephrotic syndrome

- ▶ 90% of children with nephrotic syndrome
- ▶ Cause remains unknown, seen more commonly in atopic families

Histologic types:

Minimal change NS 85% – **>95% SSNS**

Focal segmental glomerular sclerosis

(FSGN) 5%– **20% SSNS**

Diffuse mesangial proliferation 10% –

50% SSNS

Membranous nephropathy

Membranoproliferative glomerulonephritis

Pathophysiology


- ▶ An increase in glomerular capillary wall permeability leading to gross proteinuria & hypoalbuminaemia
- ▶ The cause for the increased permeability is not well understood

Pathophysiology

- ▶ In minimal change disease, possible T cell dysfunction leading to alteration of cytokines has been recognised
- ▶ This leads to loss of negatively charged glycoproteins within the glomerular capillary wall


Mechanisms of oedema

- ▶ Incompletely understood
- ▶ Protein loss leads to **hypoalbuminaemia**. This decreases the plasma oncotic pressure leading to transudation of fluid in to extra cellular space
- ▶ This leads to reduce intravascular volume





- ▶ Reduced intravascular volume decreases renal perfusion pressure. This activates renin angiotensin aldosterone system which stimulates tubular reabsorption of sodium
- ▶ Reduced intravascular volume causes release of ADH enhance reabsorption of water


Mechanisms of elevated serum lipid

- ▶ Hypoalbuminaemia stimulates generalized hepatic protein synthesis including synthesis of lipoproteins
 - ▶ Lipid catabolism is diminished because of reduced plasma lipoprotein lipase due to increased urinary loss
- 


Pathology

- In minimal change disease the glomeruli appear normal or show minimal increase in mesangial cells and matrix
 - 95% responds to corticosteroid therapy
- 


- Mesangial PGN characterized by a diffuse increase in mesangial cells and matrix on light microscopy
 - Immunofluorescent microscopy reveals mesangial deposition of IgM and/or IgA
 - 50% respond to corticosteroid therapy
- 

- ▶ In FSGS glomeruli shows mesangial proliferations and segmental scarring on light microscopy.
 - ▶ Immunofluorescent microscopy shows IgM and C3 staining in the areas of segmental sclerosis
 - ▶ 20% responds to corticosteroid therapy
- 


In clinical practice

- ▶ Those who respond to steroid therapy is labeled as having **steroid sensitive nephrotic syndrome**
 - ▶ Those who respond within 14 days of commencement known as **early responders**.
 - ▶ Response occurs after 14 days known as **late responders**
- 


Clinical Features

- ▶ Incidence is 2–4/100,000
 - ▶ Male: female = 2:1
 - ▶ Onset at 2–6 years
 - ▶ Often a preceding history of URTI
- 


Clinical Features

- ▶ Oedema in periorbital area & in the lower extremities, becomes generalized with development of ascites, pleural effusion and genital oedema
 - ▶ decreased urine output
 - ▶ Lethargy, irritability, decreased appetite, diarrhoea, abdominal pain, hypertension
 - ▶ Hypertension and gross haematuria are uncommon
- 

Atypical features suggest non minimal change NS

- ▶ Age of onset < 1 year or > 8 years
 - ▶ Macroscopic haematuria/ persistent microscopic haematuria
 - ▶ Persistent elevated BP
 - ▶ Evidence of renal impairment
 - ▶ Persistently reduced complement levels
 - ▶ Steroid resistance
 - ▶ Extra renal manifestations (arthritis, rash, anemia, fever)
- 


Differential diagnosis

- ▶ Acute post streptococcal glomerulonephritis
 - ▶ Angioedema
 - ▶ Congestive heart failure
 - ▶ Protein losing enteropathy
 - ▶ Hepatic failure
 - ▶ Chronic glomerulonephritis
 - ▶ Protein malnutrition
- 

Investigations


- ▶ **Urine FR :**
proteinuria $\geq +3$, transient haematuria in 20–30%, hyaline casts
- ▶ **Urine protein to creatinine ratio :**
>200mg/mmol
- ▶ **Serum proteins :** albumin <25g/l
- ▶ **Serum cholesterol and triglyceride**
levels are elevated

Investigations


- ▶ Renal function is usually normal
 - ▶ C3 & C4 levels are normal
 - ▶ Urine culture & ABST
- 

Management

Supportive care and monitoring

- ▶ Height , weight , body surface area (BSA), BP
– on admission
 - ▶ Weight, Input / output, urine ward test – daily
 - ▶ BP – three times a day and more frequently if abnormal
- 

Supportive care and monitoring

- Bed rest – not recommended
activity as tolerated
 - Diet – Normal calories
Normal protein
(no restriction /no need of high protein)
 - No added salt during oedema
 - Restriction of fat will be an advantage
- 

Supportive care and monitoring

▶ Antibiotics –

In case of infection – treat vigorously

Severe oedema and free fluids in the abdomen – might need prophylactic antibiotics to prevent spontaneous bacterial peritonitis

Pneumococci and H. influenzae
(Encapsulated organisms)

Associated sepsis should be treated vigorously

Supportive care and monitoring

Fluids – No restriction needed

Should be closely observed for signs of hypervolemia


Diuretics – should be used in severe oedema , but with caution due to risk of hypovolemia

It should be used only after correction of intravascular volume


Furosemide 1–2 mg/kg in divided doses PO/IV

Should be used with 25% salt free albumin / cryo poor plasma. Albumin should be given as a slow infusion


Supportive care

- ▶ Hyperlipidaemia – Dietary control of saturated fats will be an advantage
 - ▶ Lipid lowering drugs are not needed as hyperlipidaemia is transient
 - ▶ Antihypertensives
Loop diuretics and calcium channel blockers
 - ▶ Advice to mother regarding disease & steroids side effects
- 

Specific management – Steroids




Management of first episode

- ▶ Induction of remission with high dose daily prednisolone
 - ▶ Prednisolone 60 mg/m²/day, daily therapy for 4–6 weeks. 4 weeks for early responders and 6 weeks for late responders
 - ▶ Maximum 60 mg/ day
 - ▶ Preferably given as a single dose in the morning
 - ▶ Should **not** be given on empty stomach
 - ▶ May be given in divided doses to reduce GI side effects and can be combined with a H₂ receptor blocker
- 

Achieving remission

- ▶ 80% of SSNS will do so with in 14 days
- ▶ About 80–90% of children respond to steroid therapy within 3 weeks

Maintenance of remission

- ▶ Prednisolone 40 mg/m²/day EOD (every other day) for 4 weeks
 - ▶ Taper it slowly over 3– 6 months, specially in late responders
 - ▶ Longer tapering regimen will help to maintain the remission
 - ▶ Studies suggested alternative day steroids for 6 months will reduce the relapse by 33%.
- 

Another regime after high dose of therapy (Polish regime)

- ▶ After prednisolone high daily dose i.e. 60 mg/m²/day 4–6 weeks
- ▶ 60 mg/m²/day EOD for 2 weeks then
- ▶ 50 mg/m²/day EOD for 2 weeks, then
- ▶ 40 mg/m²/day EOD for 2 weeks then
- ▶ 30 mg/m²/day EOD for 2 weeks then taper it altogether for a period of 6 months

Prednisolone

- ▶ Wonderful drug if used cautiously
- ▶ Precautions and monitoring
- ▶ **Cushings facies**



Prednisolone

- ▶ **Regular eye checkups**

Posterior sub capsular cataract, glaucoma, secondary ocular infections



Prednisolone

- ▶ **Assessment of growth**

Height – short stature

Weight – obesity (weight gain and increased appetite)

- **Hypocalcaemia and osteopenia**

Regular DEXA scan

Calcium supp/or bisphosphonates for severe osteopenia

Regular calcium monitoring


Prednisolone

- ▶ **Addisonian crisis**


During stress or surgeries

- Might need IV hydrocortisone up to about 1 year after cessation of steroids

On discharge

- ▶ Educate about the illness and compliance
 - ▶ How to check urine for protein at home
 - ▶ Diet & exercise
 - ▶ Vaccination
 - ▶ When to come back
 - ▶ Risk of infections
 - ▶ SE of prednisolone
 - ▶ Maintain a diary mentioned about proteinuria, medications and intercurrent illness
 - ▶ During a GP visit indicate the child's clinic book
 - ▶ Normal activity and school attendance
- 

Immunization – special issues

- ▶ Live vaccines are contraindicated during high dose steroid therapy and 3 months after cessation
 - ▶ All killed vaccines should be offered preferably during EOD therapy
 - ▶ Hepatitis B should be given and antibody levels should be checked
 - ▶ Varicella vaccine (Live vaccine) should be given while off prednisolone for past 3 months
 - ▶ Pneumococcal vaccine
 - ▶ No live vaccines while on cyclophosphamide or within 6 months after cessation of therapy
- 


Urine for protein at home

- ▶ Should be done daily in the morning during a relapse
- ▶ In remission – when there is oedema or infection daily
- ▶ Otherwise 2–3 times per week




Urine for protein at home

Heat test


- ▶ Fill 2/3 of test tube with urine
 - ▶ should be heated at slant and upper half should be heated up to boiling point
 - ▶ If there is turbidity add 10% acetic acid or vinegar to exclude phosphates
 - ▶ Persistent turbidity is read against a black print
- 

Urine for protein at home

- ▶ Dipstick test –



Grading of proteinuria

- ▶ Nil : No turbidity
 - ▶ Trace : Slight turbidity , but no difficulty in reading the black print
 - ▶ + : Clouding of the print, but still readable with difficulty
 - ▶ ++ : Can notice the black but can not read the print
 - ▶ +++ : Can not notice black
 - ▶ ++++: Precipitate
- 

Relapse


- urine albumin $\geq 3+$ (40mg/m²/hr) on three consecutive early morning specimens, having been in remission previously

or

- Urine protein creatinine ratio ≥ 200 mg/mmol


Relapse

- Many children experience at least one relapse(70%)
- Usually precipitated by upper viral respiratory tract or GI infections
- In the past relapse rate is 60 –80% but with longer duration of steroid treatment relapse rate has reduced to 30 –40%
- Look for infections and treat if any
- Can wait for 5–7 days to commence therapy provided the child is free of oedema. Could be an episode of transient proteinuria

- Relapse should be treated with prednisolone 60 mg/m²/day until urine nil of protein for 3 consecutive days (remission)
 - The dose is then changed to 40 mg/m²/day every other day for 28 days. Then the dose of prednisolone may be tapered over 3–6 months
- 

Frequent relapsing

Two or more relapses during first 6 months after the initial episode or 4 or more within a period of one year



Response to steroids

Steroid responsive

Remission achieved with high dose daily steroid therapy alone

Steroid dependent

two consecutive relapses occurring during steroid therapy (EOD) or within 14 das after cessation of steroids

Steroid resistance

Failure to achieve remission within 4 weeks of high dose daily steroid therapy



Treatment of frequent relapses & steroid dependency

- First line of treatment is maintenance therapy with low dose prednisolone 0.1–0.6mg/kg EOD for 6 months followed by slow tapering.
- If a child requires more than 0.6mg/kg of prednisolone on alternate days to remain protein free, and particularly if there are signs of steroid toxicity then alternative therapy should be considered

Alternative therapy

- ▶ Levamisole
- ▶ Cyclophosphamide
- ▶ Cyclosporine
- ▶ Mycophenolate mofetil
- ▶ Azathioprine
- ▶ Chlorambucil
- ▶ Tacrolimus

Levamisole

- ▶ Ant-helminthic drug , steroid sparing effect
- ▶ Dosage – 2.5 mg/kg/day EOD
- ▶ Indications – steroid dependent nephrotic syndrome
- ▶ Not to induce remission but to maintain remission
- ▶ Initially with EOD prednisolone subsequently as a single agent
- ▶ Duration up to 5 years
- ▶ SE – Reversible neutropenia – monitor FBC
- ▶ Be alert in throat infections – might indicate agranulocytosis
- ▶ Drug should be discontinued permanently
- ▶ Itchy skin rash & vasculitis – permanent discontinuation
- ▶ Other SE – Liver toxicity, Convulsions

Cyclophosphamide


- ▶ Indications– to maintain prolong remission and reduces the number of relapses in steroid dependent and frequent relapsing nephrotic syndrome
- ▶ PO/IV
- ▶ PO– 3 mg/kg once a day for 8 weeks / 2 mg/kg once a day 12 weeks
- ▶ IV– 600 mg/m² monthly for 6 months

SE:

- ▶ Are related to total cumulative dose
- ▶ Recommended cumulative dose 168 mg/kg
- ▶ Haemorrhagic cystitis – specially with IV
- ▶ granulocytopenia , disseminated varicella, pigmentation, hair loss, future malignancy, Nausea, Vomiting, amenorrhoea, azoospermia with high cumulative dose

Cyclophosphamide


Precautions :

- ▶ Well hydration of the patient before and after therapy
 - ▶ Increased oral intake of fluids even with oral therapy
 - ▶ Monitor FBC before administration of IV
 - ▶ FBC weekly during oral therapy and monthly during IV therapy
 - ▶ Discontinue in neutropenia Absolute N Count < 1500
- 

Cyclosporin A

- ▶ Recommended as initial therapy for steroid resistant nephrotic syndrome
- ▶ Initial dose 3–5 mg/kg/d in 2 divided doses, 2–3 years


Adverse effects of cyclosporin

- ▶ Hypertrichosis (hirsutism)
 - ▶ Hypertrophy of gums
 - ▶ Hepatic dysfunction
 - ▶ High/ impaired renal functions
 - ▶ Hyperkalemia
 - ▶ Hypertension
- 


Steroid resistant patients

- High dose pulse methylprednisolone intravenously
- Methylprednisolone is usually given as a 30mg/kg/day bolus (max 1g) daily for 3 days
- ACE inhibitors and angiotensin II blockers may be helpful as an adjunct therapy to reduce proteinuria in steroid resistant patients


Indications for renal biopsy

- ▶ Not regularly done
 - ▶ Infants <6 months of age
 - ▶ Steroid resistance
 - ▶ Before treatment with 3rd line drug
 - ▶ Clinical course suggesting a non minimal change disease
- 

Complications

- ▶ Mortality 1–2%
 - ▶ Infections eg. Spontaneous bacterial peritonitis, disseminated varicella
 - ▶ Hypovolaemia
 - ▶ Thrombosis
 - ▶ Steroid side effects
- 


Management of hypovolaemia

- ▶ Rapid protein loss  hypovolaemia
- ▶ **Hypovolaemic crisis:**
Abdominal pain, cold extremities,
prolonged capillary refill time > 2 seconds,
wide central:peripheral temperature gap,
rapid thready pulse & narrow pulse
pressure and hypotension
- ▶ **Treatment** :Rapid IV infusion of cryo-poor
plasma, salt poor albumin, 25% salt free
albumin is the best


Infections

- ▶ Respiratory tract infections & primary peritonitis are frequent infections
- ▶ UTI can precipitate a relapse or may be a cause for poor response to prednisolone
- ▶ Chickenpox & measles could become severe if on prednisolone/cyclophosphamide
- ▶ Increase susceptibility to infection is due to
 - Urinary loss of IgG & properdin factor B
 - Defective cell mediated immunity
 - Immunosuppressive therapy
 - Malnutrition
 - Oedema fluids act as a potential culture medium


Thromboembolic events

- ▶ Increased risk 2–5%
 - ▶ Both arterial & venous thrombosis may be seen including renal vein thrombosis, pulmonary embolus, sagittal sinus thrombosis & thrombosis of catheters
 - ▶ Risk of thrombosis is related to
 - Increased prothrombotic factors (fibrinogen, thrombocytosis, haemoconcentration & relative immobilization)
 - Decreased fibrinolytic factors (urinary loss of antithrombin protein C & S)
- 


Causes of abdominal pain

- Hypovolaemia
 - Peritonitis
 - UTI
 - Gastritis (related to steroids)
 - Cellulitis
 - Renal vein thrombosis
 - Non specific
- 


Prognosis

- ▶ Prognosis – depend on the timely, correct management of the initial episode
 - ▶ Response to steroid therapy is the most important prognostic marker
 - ▶ Children who respond rapidly to steroids and those who have no relapses during the first 6 months after diagnosis are likely to follow an infrequently relapsing course
- 

Prognosis


- ▶ SSNS resolves towards the end of 2nd decade
 - ▶ SRNS most often caused by FSGS, generally have much poorer prognosis
 - ▶ They develop progressive renal insufficiency & lead to end stage renal failure
- 

Congenital NS (Finnish type)

- ▶ Rare, autosomal recessive
 - ▶ C.F. within 1st 3 months with persistent oedema and infections
 - ▶ Additional features – prematurity, large placenta, respiratory distress & skull suture separation.
 - ▶ Treatment: dialysis & renal transplant otherwise death by 5 years.
 - ▶ AN diagnosis : Increased alpha feto protein
- 

Secondary NS

Should be suspected in patients

- ▶ Age > 8 years
 - ▶ Hypertension
 - ▶ Haematuria
 - ▶ Renal dysfunction
 - ▶ Rashes
 - ▶ Arthralgia
 - ▶ Depressed serum complement levels
- 

Causes of secondary NS

- ▶ **Other GN** : membranous, MPGN
- ▶ SLE, Henoch Schonlein purpura
- ▶ **Chronic infections** – P.malaria, Hepatitis B & C, filaria, leprasy & HIV
- ▶ **Tumours** – Hodgkin lymphoma
- ▶ **Drugs** – penicillamine, captopril, gold, NSAID, mercury probenecid, ethosuximide, methimazole, lithium, procainamide, chlorpropamide, phenytoin

Thank
you