This research included forty patients; their age ranged between (12:42) years with a median 17 years and (IQR: 16-24) years, regarding their Age Categories 22 patients (55%) were in 10s, 16 patients (40%) were in 20s, 2 patients (5%) were 40s. According their gender 20 patients were females (50%) and 20 patients were males (50%).

For laterality, 17 cases (43%) involved the left knee, while 23 cases (58%) involved the right knee, and Recurrent patients were 21 (53%).

The clinical symptoms were painful in 16 patients (40%), painful and swelling in 24 patients (60%). Additionally, 21 patients (53%) had History of previous dislocation and 27 patients (68%) had History of trauma.

According to some clinical measures, the insal ratio ranged between (1:2) with a median 1.40 and (IQR: 1.25-1.66), the Caton ranged between (1:1.65) with a median 1.33 and (IQR: 1.16-1.49)and the PTI ranged between (0.1:2.30) with a median 0.42 and (IQR: 0.32-0.50).

Concerning blood pressure, SBP (90:170) mmHg with a median 120 mmHg and (IQR: 110-135) mmHg and DBP ranged between (60:100) mmHg with a median 80 mmHg and (IQR: 70-85) mmHg, finally regarding hypertension status 17 patients (43%) were abnormal and 23 patients (58%) were normal, as shown in Table (1) and Figure 1- 3.

Table (1): Description of Some Parameters Among All Studied Patients.

	N = 40
Recurrence	21 (53%)
Age	
Min-Max	12.0-42.0
Median (Q1,Q3)	17.0 (16.0,24.0)
Gender	
Female	20 (50%)
Male	20 (50%)
Laterality	
LT knee	17 (43%)
RT knee	23 (58%)
Insal	
Min-Max	1.00-2.20
Median (Q1,Q3)	1.40 (1.25,1.66)
Caton	
Min-Max	1.00-1.65
Median (Q1,Q3)	1.33 (1.16,1.49)
PTI	
Min-Max	0.10-2.30
Median (Q1,Q3)	0.42 (0.32,0.50)
SBP	
Min-Max	90-170
Median (Q1,Q3)	120 (110,135)
DBP	
Min-Max	60-100
Median (Q1,Q3)	80 (70,85)
Clinical symptoms	
painful	16 (40%)
painful/swelling	24 (60%)

History of previous dislocation	21 (53%)
History of trauma	27 (68%)
Age categories	
10s	22 (55%)
20s	16 (40%)
40s	2 (5.0%)
Hypertension status	
abnormal	17 (43%)
normal	23 (58%)




Figure (1) Box plot show the distribution of Age among all studied patient according to their gender




Figure (2) Bar Plot show the Categories of age among all studied patient




Figure (3) Box plot show the distribution of Caton among all studied patient

This report describes the selected parameters among recurrent patients only in the study which included 21 patients. Their age ranged between (13:42) years, with a median 23 years (IQR: 16-24) years. In term of Age Distribution 10 patients (48%) were in their teens, 9 patients (43%) were in their twenties, 2 patients (9.5%) were in their forties.

Concerning blood pressure, DBP ranged between (70:100) mmHg with a median 80 mmHg and (IQR: 80-90) mmHg and SBP ranged between (110:170) mmHg with a median 130 mmHg and (IQR: 120-140) mmHg, as insulated in Table (2) and Figure (4) and (5).

Table (2): Distribution of Some Parameters in Recurrent Patients

	N = 21
Age	
Min-Max	13-42
Median (Q1,Q3)	23 (16,24)
Age categories	
10s	10 (48%)
20s	9 (43%)
40s	2 (9.5%)
DBP	
Min-Max	70-100
Median (Q1,Q3)	80 (80,90)
SBP	
Min-Max	110-170
Median (Q1,Q3)	130 (120,140)




Figure (4) Bar Plot show the Categories of Age among reccurent patient




Figure (5) Box plot show the distribution of SBP among recurrent patient