GEOfinder

GSE140053, GSE145525, GSE38901, GSE43579, GSE62309, GSE57398, GSE66448, GSE60984, GSE105028, GSE112379, GSE121182, GSE137558, GSE152144, GSE143591, GSE194364, GSE166456, GSE209667, GSE270671, GSE253873, GSE259305, GSE276302, GSE303857, GSE89382, GSE280307, GSE101731, GSE154744, GSE209675, GSE267931, GSE173536, GSE192370, GSE326102

Experiment Type(s):

- ☐ Expression profiling by RT-PCR
- ☐ Expression profiling by array
- ☐ Expression profiling by high throughput sequencing
- ☐ Genome binding/occupancy profiling by genome tiling array

Species:

- ☐ Arabidopsis thaliana
- ☐ Bos taurus
- ☐ Caenorhabditis elegans
- ☐ Danio rerio
- ☐ Drosophila melanogaster
- ☐ Escherichia coli
- ☐ Gallus gallus

Number of Samples:

- ☑ 1-10
- ☑ 11-50
- ☑ 51-100
- ☑ 101-500
- ☑ 501-1000
- ☑ 1000+

Start Year:

End Year:

Submit

- [x] Genome binding/occupancy profiling by high throughput sequencing
- [ ] Genome variation profiling by SNP array
- [ ] Genome variation profiling by genome tiling array
- [ ] Methylation profiling by array
- [ ] Methylation profiling by genome tiling array
- [ ] Methylation profiling by high throughput sequencing
- [ ] Non-coding RNA profiling by array
- [ ] Non-coding RNA profiling by high throughput sequencing
- [ ] Other

- [x] Homo sapiens
- [ ] Mus musculus
- [ ] Oryza sativa
- [ ] Rattus norvegicus
- [ ] Saccharomyces cerevisiae
- [ ] Schizosaccharomyces pombe
- [ ] Sus scrofa
- [ ] synthetic construct

Top results (up to 100):

GSE ID	Title	Summary
GSE90716	Hijacking of stress response machinery by oncogenes in acute leukaemia [ChIP-seq]	HSF1 orchestrates shock response pat pathway is co-optec and provides critica from oncogenic str has a diverse occup signature dependin type. In this study, v HSF1 ChIP-seq ana the human T-ALL c CUTLL1 and P12. T revealed the HSF1 binding signature ir P12 cells. MYC is a oncogene in T-ALL. oncogene addictior that act downstrear transcription factor anabolic pathways a understood. For thi performed MYC Ch analysis using the h cell line CUTLL1. Th revealed that HSF1 targets are include binding signature.
GSE194098	HSF2 cooperates with HSF1 to drive a transcriptional program critical for the malignant state	Heat shock factor well known for its r heat shock respons where it drives a tra program comprisin protein (HSP) gene tumorigenesis, whe program comprisin many non-canonica genes that support Here, we find that H HSF1 paralog with role in the HSR, phy

GSE ID	Title	Summary
		functionally interac across diverse type HSF1 and HSF2 ha similar chromatin o regulate a common which include both non-canonical trans targets with roles c supporting maligna either HSF1 or HSF dysregulated respo nutrient stresses in reduced tumor pro cancer cell line xen Taken together, the establish HSF2 as a cofactor of HSF1 ir cancer cell transcri program to suppor malignant state.
GSE209687	Transcriptional responses of cancer cells to heat shock-inducing stimuli involve amplification of robust HSF1 binding	This SuperSeries is the SubSeries listed
GSE159724	Heat Shock Factor 1 (HSF1) supports the ESR1 action in breast cancer (ChIP-seq)	Heat shock factor key regulator of tra responses to prote It has been recently signaling of estroge To study the coope HSF1 and ESR1 in transcriptional resp estrogen, we establ estrogen receptor breast cancer cell li reduced HSF1 level specific shRNA or

GSE ID	Title	Summary
		approach. HSF1 de the inhibition of the effect of estrogen i T47D cells. RNA-se revealed that the st effect of E2 on the transcriptome was HSF1-deficient MCI could partially resul higher basal expres dependent genes ir as a consequence c enhanced binding c ESR1 to chromatin, revealed by ChIP-se Thus, we postulate fraction of ESR1 cc released from the ir complex with HSP9 transcriptional com without E2-stimulat
GSE38912	HSF1 drives a transcriptional program distinct from heat shock to support highly malignant human cancers	This SuperSeries is the SubSeries listec
GSE63589	Genome-wide characterization of HSF1 binding in human 451Lu melanoma cancer cells under basal conditions	HSF1 orchestrates transcriptional prog cancer cells. In this assayed for genom localization of HSF1 in the 451Lu melan untreated cells. The revealed a transcrip program enriched f metastasis-related

GSE ID	Title	Summary
GSE274408	Heat Shock Transcription Factor 1 Facilitates Liver Cancer Progression by Driving Super-Enhancer-Mediated Transcription of MYCN	HSF1MYCN axis, a dependent regulatc be a prognostic ind potential therapeut liver cancer.
GSE137560	Heat Shock Factor 1 (HSF1) acquires transcriptional competence under 17b-estradiol in ERa-positive breast cancer cells	This SuperSeries is the SubSeries listec
GSE126151	A systematic analysis of nuclear heat-shock protein 90 identifies a metazoan-specific regulatory module	Molecular chaperor heat-shock proteins in protein folding ar assembly processes in cytosol has been elucidated. Chaperc present in the nucle compartment wher enter after being fu the cytosol, raising question about cha function in this com We have performec analysis of nuclear protein 90 to identi functions of this ch the nucleus. Combi

GSE ID	Title	Summary
		and genetic interac cancer co-expressi allowed us to define functional interacto HSP90 consisting c proteins. Using trar studies we identifie Factor C1 (HCFC1) metazoan-specific t regulator that depe HSP90 for its stabil nucleus. We found required for optima HCFC1 in transcrip study provides the insight into the fun nuclear HSP90.
GSE45852	Tight coordination of protein translation and heat shock factor 1 activation supports the anabolic malignant state [ChIP-Seq]	A unifying characte aggressive cancers profound anabolic s metabolism to enab proliferation and bi expansion. The ribo centrally situated to metabolic states bu impacts systems th cellular survival is u Here, through integ chemical-genetic ar find that a dominan transcriptional effe protein translation i cells is complete ina heat shock factor 1 multifaceted transc regulator of the hea response and many cellular processes e tumorigenesis. Trar through the riboson

GSE ID	Title	Summary
		the transcriptional l and links the funda anabolic processes production and ene metabolism with HS Targeting this link c cancer cells of thei chaperone armame thereby rendering t phenotype unsusta
GSE280712	TSA-sequencing in untreated (UN), sodium arsenite (SA) and heat shock (HS) stressed HeLa cells	Mammalian cells ac defense mechanism response to stresse stress bodies (nSBs transient membran organelles in primat form upon severe s is known on how th contributes to cellu homeostasis or whe have any pathophys roles. We observed novo formation of r accompanies enlarg pericentromeric Sat chromatin loci, pror investigate whether acute alteration imp chromatin accessib expression. To test performed an impr Tyramide Signal Am (TSA) mapping to ic proximal to newly fo
GSE111923	Osmostress induced changes of chromatin architecture and transcription in mammalian cells [ChIP-Seq]	Osmostress causes compaction and sin triggers a massive transcriptional resp eukaryotic cells tha

GSE ID	Title	Summary
		for adaptation and However, thus far it unknown how chan chromatin organiza ongoing DNA trans linked under hyper- conditions. In this s focused on elucidat extent of chromatir reoganization upon osmotic shock, its r and its link with act transcription in hun cells.
GSE198223	HSF1 activates ANXA11-mediated glutamine metabolism by super enhancer long non-coding RNA LINC00857 in colorectal cancer [ChIP-seq]	We shed light on a cooperative machin HSF1 and super en the first time in CR stimulate super enh by the enrichment acetyltransferase P

GSE ID	Title	Summary
		LINC00857 promot contributing to LIN(transcription.
GSE24326	Genome-wide profiling of H3K56 acetylation and transcription factor binding sites in human adipocytes	We use ChIP-seq to genome-wide sites of lysine 56 of the h (H3K56), which is a three histone modif with known roles in insulin resistance, ir adipocytes derived mesenchymal stem Surprisingly, we find large fraction of ge some level of acety H3K56, but the high acetylation are assc genes previously re involved in type 2 d Using computationa we propose that the transcription factor be involved in recru modifying enzymes sites. We confirm th by measuring the b E2F4 using ChIP-se examine the binding proteins using ChIP and C/EBP. HSF-1 i regulator of stress and is a target of th histone modifiers a find a high degree c between HSF-1 bin H3K56 acetylation that are not stresse contrast, C/EBP, wh known to be modifi enzymes, shows mu

GSE ID	Title	Summary
		overlap with the site acetylation. Our res represent the first r the regulatory code adipocytes.
GSE169435	The homeotherm-conserved protein CGGBP1 connects heat stress to chromosomal fusions	We report that heat induces chromoson events in cell cultur describe the finding normal primary fibr [Coriell Cell Reposit GM02639] and sub apply it to a system stable lines called C [described in PMID The stable line CT s Control shRNA and for a knockdown of CGGBP1 using spec [Origene Lentiviral s TL313955V]. Our re suggest that as me frequency of chrom fusions, the CGGBF in KD, when compa imitates the effects stress. Further, ChIl for TP53BP1 show misdirected repairs flanks in KD as com We show that CGGI and heat give rise t chromosomal fusio overlapping mecha involve misdirected flanks of repeats ge
GSE105189	ChIP-seq from GM12878 (ENCSR009MBP)	HSF1 ChIP-seq on h GM12878 | For dat

GSE ID	Title	Summary
		terms and conditio refer to [url] and [u
GSE167552	The stress response regulator HSF1 modulates natural killer cell anti-tumour immunity	Diverse cellular insu on activation of the factor 1 (HSF1), wh the proteotoxic stre to maintain protein homoeostasis. HSF numerous gene pro beyond the proteot response in a cell-ty context-specific ma promote malignanc the role(s) of HSF1 populations of the t microenvironment r elusive. Here, we le vivo model of HSF1 and single-cell trans tumour profiling to augmented HSF1 a natural killer (NK) c cytotoxicity, cytokir and subsequent an immunity. Mechanis directly binds and r expression of key n NK cell effector fun work demonstrates regulates the immu under the stress co the tumour microer These findings have implications for enh efficacy of adoptive therapies and for d combinatorial strate including modulato mediated tumour ki

GSE ID	Title	Summary
		SuperSeries is com SubSeries listed be
GSE139882	Dynamic regulation of stress-responsive non-genomic CTCF complexes [ChIP-seq]	The nuclear protein binding factor (CTC diverse roles in chr architecture and ge regulation. Functior associates with tho genomic sites and i proteins, such as c non-coding RNAs t specific transcriptic programming. In th examined CTCF dur cellular stress respc human primary cell immune-blotting, qu real time-PCR, chro immunoprecipitatio analysis, mass spec RNA immunoprecip sequence analysis, confocal microscop Unexpectedly, we f CTCF is exquisitely diverse forms of st normal patient-deri mammary epithelial (HMECs). In HMECs of CTCF protein for genomic complexes to Serine/arginine-r factor (SC-35)-cont nuclear speckles, e canonical associatic chromatin. Upon st genomic CTCF prot downregulated by c protein stability, res of CTCF from SC-3

GSE ID	Title	Summary
		speckles and chang RNA interactions. C complexes that ass genomic DNA are r stress-induced deg CTCF-DNA binding unchanged. Restora cellular CTCF prote and re-localization t speckles can be acl inhibition of protea mediated degradati Surprisingly, we obs same characteristic stress response du differentiation of hu pluripotent stem ce CTCF forms stress- complexes that loca nuclear speckles du specific stage of ne commitment/develo not in differentiatec speculate that thes canonical CTCF cor a largely non-genor RNA processing, pc maintain cells in a p differentiation state dynamically regulat environmental signa canonical, stress-re activity of CTCF is u persistently stresse epigenetically re-pr variant HMECs and cancer cell lines. Th reveal new insights function in cell diffe and the stress-resp

GSE ID	Title	Summary
		implications for oxi damage-induced ca and neuro-degener diseases.
GSE231486	Genome-wide Transcription Factor binding maps reveal cell-specific changes in the regulatory architecture of human HSPC	This SuperSeries is the SubSeries listec
GSE303801	Pre-occupied ATF4 relies on CEBP co-binding for robust transcription activation during integrated stress response [HeLa_CCCP_CUT&Tag]	Upon encountering need to rapidly and remodel their trans restore homeostasi survival. However, t mechanisms underl induced transcriptic remodeling are not understood. In this employed multi-omi approaches to com profile the changes transcriptome, epig landscapes, and 3D organization during integrated stress re a conserved cellula activated by diverse stimuli. Our findings widespread transcr changes occur 4-6 the induction of ISF with the genome-wi key transcriptional ATF4. Notably, while to most of its targe upon ISR activation occupies the promc

GSE ID	Title	Summary
		hundreds of genes stress conditions, p for enhanced trans activation in respor signals. We further that ATF4-mediatec activation does not traditional transcrip regulatory mechani increased H3K27 a chromatin accessib remodeling of enha promoter loops. Ins data suggest that t localization of ATF4 pronounced genom redistribution of tra factor CEBP-, from sites to a subset of regions, where it cc with ATF4 to elicit t activation. The com occupancy of ATF4 ultimately dictates t transcriptional outc ISR. Interestingly, th binding sites of CEI response to stress determined by the the CEBP- or ATF4 rather by the highe chromatin architect surrounding the bir These findings prov insights into the mc mechanisms that g rapid and efficient t remodeling during I potential implicatior stress responses.

GSE ID	Title	Summary
GSE135807	ChIP-seq assay in human cancer cells	ChIP-seq assay was for the quantitative the binding of TFs a upon acute stimula

GSE ID	Title	Summary
GSE45024	Genome-wide maps of chromatin state for HIRA, UBN1 and ASF1a	The HIRA chaperon comprised of HIRA, CABIN1, collaborate histone-binding pro incorporate histone into chromatin in a replication-indepen To better understar function and mecha integrated HIRA, UB and histone H3.3 C gene expression an HIRA-binding sites with UBN1, ASF1a a active promoters ar weak/poised enhan promoters, binding HIRA/UBN1/ASF1a with the level of ge expression. HIRA is deposition of histor binding sites. There differences in nucle co-regulator compo different classes of regulatory site. Unc this, we report nove interactions betwee complex and transc factors, a chromatii

GSE ID	Title	Summary
		and an ATP-depenc chromatin-remodell Our results map the of the HIRA chaper the chromatin lands point to different in partners at functior regulatory sites.
GSE69304	Genome-wide analysis of ATF4 binding in histidinol-treated cells	Activating Transcrip (ATF4) is a transcri induced by the inte response (ISR). This is a genome-wide o profiling of ATF4 in cells. HAP1 is a nea human cell line that from KBM-7 cells is patient with Chronic Myelogenous Leuke induced ATF4 expre mimicking amino ac with the drug histid identified peaks of using three indeper antibodies.
GSE191243	HSP70 binds RNA and regulates transcription by RNA Polymerase III [ChIP-seq]	We examined the R ability of the 70-kD protein (HSP70) far HSPA1A and identi species bound to H We used a UV-cros immunoprecipitatio approach named Fl (Akta et al., 2017) u 3xFLAG-Histidine-B ectopically expresse HEK293T cells.

GSE ID	Title	Summary
GSE303860	CPSF73 activation and 3 Pol II pausing are lost during read-through transcription after heat shock	This SuperSeries is the SubSeries listed
GSE144786	NELF regulates a promoter-proximal step distinct from RNA Pol II pause-release	RNA polymerase II generally paused at proximal regions in metazoans, and bas studies, this functic attributed to the ne elongation factor (N we show that upon depletion of NELF, I be released into ge stopping instead ar nucleosomal dyad-a region. The transitic pause region is inde positive transcriptic factor P-TEFb. Duri shock response, Pc released from paus shock induced gene genes are paused a transcriptionally do during the heat shc We find that both a heat shock respons intact upon NELF lc that NELF depletior global loss of cap-b complex from chro global reduction of transcript 5 cap sta

GSE ID	Title	Summary
		our studies implicat functioning in early complexes distinct pause-release.
GSE159190	Occupancy of Integrator subunits on DNA after hyperosmotic stress	ChIP-seq was perfo antibodies against I subunits 11 or 3 to occupancy of these DNA after hyperosr
GSE212744	Selective advantage of epigenetically disrupted cancer cells via phenotypic inertia [ATAC-Seq]	Mutations in over o genes, belonging tc classes of epigenet resulted in significa advantage under nu starvation and acidi suggesting that dis epigenetic control, levels of the regula promotes broad res cells to stressful en To uncover the mol mechanisms of suc phenomenon, and t how loss of epigene regulators gives ad during stress, we ex chromatin accessib in nutrient-deprivec using ATAC-seq. AT analysis of melanor expressing 4-7sgRN members of the PR (EED, EZH2, SUZ12 H1B (HIST1H1B), th remodeller SMARCI

GSE ID	Title	Summary
		non-expressed con (TNP2), and grown unperturbed condit under glutamine de (d2).
GSE235444	AFF4 globally affects the release of paused RNA polymerase II in HEL cells [ChIP-seq]	P-TEFb, a heterodin CDK9, and Cyclin T regulator of promo pause release of Pc metazoans. It is cap forming three large complexes, includin elongation complex BRD4/P-TEFb comp 7SK snRNP. In the S BRD4/P-TEFb, P-TE enzymatically active 7SK snRNP, its acti inhibited. The SEC c AFF1 or 4, ENL or A or 3 and EAF1 or 2 P-TEFb, the only su catalytic activity, an noncatalytic subuni found to be able to pause release throu One recent study sl human DLD-1 cells, regulates pause rel shock (HS) genes, v BRD4/P-TEFb comp pause release of th genes. In this study that AFF4 knockdo HEL cells decreased cellular level but als chromatin occupan serine 2 phosphory and notably, increas proximal pause of F

GSE ID	Title	Summary
		several hundred HS thousands of non-H Mechanistically, AFI pause release likely facilitating the bind to Pol II. These resu that the extent of ir AFF4 on pause rele to be context-depe type dependent.
GSE303800	Pre-occupied ATF4 relies on CEBP co-binding for robust transcription activation during integrated stress response [HeLa_CCCP_ATAC-seq]	Upon encountering need to rapidly and remodel their trans restore homeostasi survival. However, t mechanisms underl induced transcriptic remodeling are not understood. In this employed multi-omi approaches to com profile the changes transcriptome, epig landscapes, and 3D organization during integrated stress re a conserved cellular activated by diverse stimuli. Our findings widespread transcr changes occur 4-6 the induction of ISF with the genome-wi key transcriptional ATF4. Notably, while to most of its targe upon ISR activation occupies the promc hundreds of genes stress conditions, p

GSE ID	Title	Summary
		for enhanced trans activation in respor signals. We further that ATF4-mediatec activation does not traditional transcrip regulatory mechani increased H3K27 a chromatin accessib remodeling of enha promoter loops. Ins data suggest that t localization of ATF4 pronounced genom redistribution of tra factor CEBP-, from sites to a subset of regions, where it cc with ATF4 to elicit t activation. The com occupancy of ATF4 ultimately dictates t transcriptional outc ISR. Interestingly, th binding sites of CEI response to stress determined by the s the CEBP- or ATF4 rather by the highe chromatin architect surrounding the bir These findings prov insights into the mc mechanisms that g rapid and efficient t remodeling during I potential implicatior stress responses.

GSE ID	Title	Summary
GSE310047	Hyperosmotic stress induces a large-scale rewiring of 3D chromatin interactions [Cut & Tag]	Cells rapidly adapt hyperosmotic stres coordinated molecu responses. To deter three-dimensional (chromatin structure to this process, we chromatin interacti architectural protei and transcriptional human cells expose induced hyperosmo We performed time C to measure stres remodeling of chro and domains, CUT& quantify CTCF, RAD and H3K27ac occu anchors, and RNA-s capture stress-resp transcriptional prog data reveal global lo existing chromatin concurrent formati novo, transient loop for retained CTCF a as well as transcrip responses that are layered and largely from loop remodeli dataset provides a investigating how n architecture and tra regulation respond hyperosmotic stres
GSE29611	Histone Modifications by ChIP-seq from ENCODE/Broad Institute	This data was gene ENCODE. If you hav about the data, con submitting laborato

GSE ID	Title	Summary
		(mailto:nshoresh@b If you have questio Genome Browser tr associated with this contact ENCODE (mailto:genome@so This track displays chromatin state ge the Broad/MGH EN using ChIP-seq. Che modifications (meth acetylation) to the h proteins present in influence gene expr changing how acce chromatin is to trar The ChIP-seq meth first using formalde cross-link histones DNA-associated pro genomic DNA withi cross-linked chrom subsequently extra mechanically shear immunoprecipitated specific antibodies. reversal of cross-lir immunoprecipitated sequenced and ma human reference g relative enrichment antibody-target (ep the genome is infer density of mapped For data usage terr conditions, please r and [url]

GSE ID	Title	Summary
GSE184139	Chromatin Landscape Analysis of H1 cells with different temperature.	Chromatin Landsca of H1 cells with diff temperature.
GSE202306	Chromatin accessibility changes after BAF complex disruption in neuroblastoma cells by ATAC-Seq	In order to study th regions undergoing accessibility change

GSE ID	Title	Summary
		chromatin remodell is inhibited, ATAC-S BE(2) neuroblastom performed after sil structural subunits ARID1B.
GSE152062	RNA Polymerase II binding profiles of HEK-293T cells exposed to hyperosmotic stress	Anti-Pol II ChIP-seq performed to asses redistribution of RN Polymerase (Pol) II hyperosmotic stres
GSE125220	Histone chaperone CAF-1 is a synthetic lethal vulnerability in the context of ARID1A deficiencies [ATAC-Seq]	Mutations in the BA remodeling comple the most frequent alterations in huma many of which resu function of the resp subunits. Recent ge chemical screens h synthetic lethal inte BAF complex subur other as well as witl chromatin modifier epigenome-focusec library, we identifiec histone chaperone CAF-1 (chromatin a factor 1) as a synth vulnerability of ARI cells that is enhanc androgen receptor testosterone. In line antagonistic roles c and BAF complexes nucleosome deposi

GSE ID	Title	Summary
		remodeling, we sho down of CAF-1 sub in a partial reversal transcriptional and accessibility effects loss of ARID1A. Toc findings identify a r partner of the BAF remodeling comple represent a potenti strategy for targeti deficient cancers.
GSE159803	Heat Shock Factor 1 (HSF1) supports the ESR1 action in breast cancer	This SuperSeries is the SubSeries listec
GSE57399	FBXW7 modulates stress response by post-translational modification of HSF1	This SuperSeries is the SubSeries listec
GSE290558	Mapping breakome of multiple cancer cell-lines	Cancer is character uncontrolled prolife accompanied by th hypertranscription oncogenes, leading transcription stress source of DNA dou breaks (DSBs) that

GSE ID	Title	Summary
		genomic stability. Y transcription stress underexplored. In th utilized maps of DS through in-suspens labeling in situ and (sBLISS), along with stress markers, rev transcription stress coincide with the su regulatory landscap H2AX mapping indi enrichment at trans stress sites while no enriched genes sho H2AX marking, but DSBs tied to transc are distinctly marke Intriguingly, genes v DSBs marked by H2 significantly higher and repair than tho H2AX-low genes, m vulnerability to mut These findings und super-enhancer act determinant of the stress landscape in posing a threat to t stability of oncoger

GSE ID	Title	Summary
GSE33213	Open Chromatin TFBS by ChIP-seq from ENCODE/Open Chrom(UT Austin)	This data was gene ENCODE. If you hav about the data, con submitting laborato (Terry Furey mailto:tsfurey@duk have questions abo Genome Browser tr associated with this contact ENCODE (mailto:genome@so These tracks displa of evidence from di assays as part of th Chromatin track se displays open chror and/or transcriptio binding sites identif multiple cell types b more complementa methodologies, DNa hypersensitivity (HS DNaseI HS), Formal Assisted Isolation o Elements (FAIRE) (U and chromatin immunoprecipitatio select regulatory fa

GSE ID	Title	Summary
		TFBS). Each metho performed on the s using identical grov conditions. (Note: D or all ChIP experime be available for all c Regions that overla methodologies iden regulatory element cross-validated indi confidence regions. multiple lines of evi suggest that regior by a single assay (e only or FAIRE-only) biologically relevant submitted). | DNase DNaseI is an enzym long been used to r chromatin accessib DNaseI hypersensit feature of active cis sequences.The use method has led to t of functional regula elements that inclu enhancers, silencer locus control regior elements. DNaseI hypersensitivity sig chromatin accessib binding of trans-act place of a canonica | FAIRE data: FAIRE (Formaldehyde Ass of Regulatory Elem method to isolate a nucleosome-deplete the genome. FAIRE discovered in yeast

GSE ID	Title	Summary
		subsequently show active regulatory el human cells (Giresi Similar to DNaseI H appears to identify regulatory element promoters, enhanc insulators, locus co and novel element ChIP (Chromatin Immunoprecipitatio method to identify location of protein directly or indirectl genomic DNA. By ic binding location of specific transcriptic general transcriptic components, and c factors, ChIP can h functional annotatic open chromatin reg identified by DNase and FAIRE. | Input c background contro we sequenced the i DNA sample that w ChIP. Crosslinked c sheared and the cr reversed without ca the immunoprecipit This sample is othe processed in a mar to the ChIP sample below. The input tra in revealing potenti arising from the se alignment process number difference reference genome

GSE ID	Title	Summary
		sequenced samples regions of poor sec alignability. | For da terms and condition refer to [url] and [u
GSE144303	Dynamic chromatin accessibility landscape changes following interleukin-1 stimulation	Genome-wide meth examining chromati modification provid information on regu regions of the genc modifications of ch rapid access of the regulatory machine condensed genomi facilitating subsequ expression. Inflamm cytokine stimulation cause rapid gene e changes through di signalling pathway-r transcription factor and regulatory elem
GSE140934	Chromatin Remodeling and Immediate Early Gene Activation by SLFN11 in Response to Replication Stress [ATAC-seq_K562]	Schlafen 11 (SLFN recently arisen as a restriction factor ag replication stress. H that SLFN11 increa chromatin accessib wide, dominantly at response to replica induced by the CHk prexasertib and the topoisomerase I inh camptothecin. Conc SLFN11 selectively cellular stress respo pathways by induci transcription of the Early Genes (IEGs) i

GSE ID	Title	Summary
		FOS, EGR1, NFKB2 Both chromatin ope activation require th helicase activity of whereas extrinsic IE by serum induction independent. SLFN IEG activation is als across 55 non-isog cell lines by compar transcriptome data after camptothecin We conclude that S as a global regulatc chromatin structure intrinsic IEG activat potential to engage immune activation i replicative stress.
GSE45023	Genome-wide maps of chromatin state for H3.3 tagged cells	The HIRA chaperon comprised of HIRA, CABIN1, collaborate histone-binding pro incorporate histone into chromatin in a replication-indepen To better understar function and mecha integrated HIRA, UE and histone H3.3 C gene expression an HIRA-binding sites with UBN1, ASF1a a active promoters ar weak/poised enhan promoters, binding HIRA/UBN1/ASF1a with the level of ger expression. HIRA is deposition of histor

GSE ID	Title	Summary
		binding sites. There differences in nucle co-regulator compc different classes of regulatory site. Unc this, we report nove interactions betwee complex and transc factors, a chromatii and an ATP-depenc chromatin-remodell Our results map the of the HIRA chaper the chromatin lands point to different in partners at functior regulatory sites.
GSE50611	Genomic discovery of potent chromatin insulators for human gene therapy	This data includes r factor profiling usir

GSE ID	Title	Summary
GSE114084	ChIP-seq analysis of CFP1 and related molecules	We report a ChIP-s the CpG binding pr and related protein regulation of chrom transcribed genes i erythroblasts and E transformed B-lymp generated multiple and compared thes available ChIP datas
GSE35583	Histone Modifications by ChIP-seq from ENCODE/University of Washington	This data was gene ENCODE. If you hav about the data, con submitting laborato (Richard Sandstrom mailto:sull@u.washi you have questions Genome Browser tr associated with this contact ENCODE (mailto:genome@so This track was proc

GSE ID	Title	Summary
		of the ENCODE Prc track displays geno maps of histone mc different cell lines (ChIP-seq high-throu sequencing. | For d terms and conditior refer to [url] and [u
GSE91444	ChIP-seq from MCF-7 (ENCSR062HDL)	HSF1 ChIP-seq on I | For data usage te conditions, please r and [url]

GSE ID	Title	Summary
GSE74363	The Common Stress Responsive Transcription Factor ATF3 Binds Active Enhancers and Bookmarks Genomic Sites for Transcriptional Regulation	This SuperSeries is the SubSeries listed
GSE212693	Selective advantage of epigenetically disrupted cancer cells via phenotypic inertia [ChIP-Seq]	Mutations in over o genes, belonging to classes of epigenet resulted in significa advantage under nu starvation and acidi suggesting that dis epigenetic control, levels of the regula promotes broad res cells to stressful en To uncover the mol mechanisms of suc phenomenon, and t how loss of epigene regulators gives ad during stress, we pl examine how the ch landscape changes deprived conditions
GSE243274	Investigating how the SPT2 histone chaperone affects chromatin structure in human osteosarcoma cells.	Histone chaperones essential roles in th of chromatin struct investigated how th histone chaperone chromatin structure accessibility in hum (using U-2 OS osteo cells as model; thes express a H3.3-SNA transgene). To achi performed ATAC-se

GSE ID	Title	Summary
		WT U2OS vs three SPT2 KO clones (KO KO-6
GSE209149	TF ChIP-seq from K562 (ENCSR579XWM)	eGFP-HSF4 ChIP-se K562 | For data usa and conditions, plea [url] and [url]
GSE154445	DNAJC9 Integrates Heat Shock Molecular Chaperones into the Histone Chaperone Network	From biosynthesis t into nucleosomes, h handed through a c histone chaperones histones from non-s interactions. Wheth mechanisms exist t the histone fold dur chaperone handove to release trapped i is unclear. Using str guided and functior proteomics, we iden characterize a histc chaperone function a heat shock co-cha promotes HSP70-m catalysis. We elucid structure of DNAJC histone H3-H4 co-c complex with MCM2 how this dual histor shock co-chaperon histone substrates. that DNAJC9 recru type enzymes via its fold histone H3-H4 upstream in the his chain, during replica transcription-couple nucleosome assembl

GSE ID	Title	Summary
		clean up spurious ir With its dual functic DNAJC9 integrates resourced protein f the histone supply	resolve aberrant int throughout the dyn histones.
GSE140933	Chromatin Remodeling and Immediate Early Gene Activation by SLFN11 in Response to Replication Stress [ATAC-seq_CPTO]	Schlafen 11 (SLFN1 recently arisen as a restriction factor ac replication stress. H that SLFN11 increa chromatin accessib wide, dominantly at response to replica induced by the CHk prexasertib and the topoisomerase I inh camptothecin. Conc SLFN11 selectively cellular stress respc pathways by induci transcription of the Early Genes (IEGs) i FOS, EGR1, NFKB2 Both chromatin ope activation require th helicase activity of whereas extrinsic IE by serum induction independent. SLFN IEG activation is als across 55 non-isog cell lines by compar transcriptome data after camptothecin We conclude that S as a global regulatc chromatin structure

GSE ID	Title	Summary
		intrinsic IEG activat potential to engage immune activation i replicative stress.
GSE135354	Stress-induced RNAchromatin interactions promote endothelial dysfunction (Hi-C)	Chromatin-associat (caRNA) has been p type of epigenomic Here, we test wheth environmental stres cellular dysfunction modulating RNA-ch interactions. We inc endothelial cell (EC) with high glucose a (H+T), that mimic th stress in diabetes n characterize the H+ changes in gene ex single cell (sc)RNA- interactions by Hi-C chromatin interacti iMARGI. H+T induce chromosomal RNA- interactions, particu the super enhancer causal relationship induced RNA-chron interactions and the of EC dysfunction-r we suppress the LIN RNA. This suppress attenuates the expr SERPINE1, a critica inflammatory and p gene. Furthermore, of the co-expressio network between d healthy donor-deriv corroborate the H+ RNA-chromatin inte

GSE ID	Title	Summary
		Taken together, caF dysregulation of ge expression modulat dysfunction, a cruc mechanism underly diseases.
GSE40833	DNaseI/FAIRE/ChIP Synthesis from ENCODE/OpenChrom(Duke/UNC/UTA)	These tracks displa of evidence from di assays as part of th Chromatin track se displays open chro and/or transcriptio binding sites identif multiple cell types b more complementa methodologies, DNa hypersensitivity (HS DNaseI HS), Formal Assisted Isolation o Elements (FAIRE) (U and chromatin immunoprecipitatio select regulatory fa TFBS). Each metho performed on the s using identical grow conditions. (Note: D or all ChIP experime be available for all c Regions that overla methodologies iden regulatory elements cross-validated indi confidence regions. multiple lines of evi suggest that regior by a single assay (e only or FAIRE-only) biologically relevant submitted). | DNase

GSE ID	Title	Summary
		DNaseI is an enzym long been used to r chromatin accessib DNaseI hypersensit feature of active cis sequences.The use method has led to t of functional regula elements that inclu enhancers, silencer locus control regior elements. DNaseI hypersensitivity sig chromatin accessib binding of trans-act place of a canonica | FAIRE data: FAIRE (Formaldehyde Ass of Regulatory Elem method to isolate a nucleosome-deplete the genome. FAIRE discovered in yeast subsequently show active regulatory el human cells (Giresi Similar to DNaseI H appears to identify regulatory element promoters, enhance insulators, locus co and novel elements ChIP (Chromatin Immunoprecipitatio method to identify location of proteins directly or indirectly genomic DNA. By ic binding location of specific transcriptic

GSE ID	Title	Summary
		general transcriptic components, and cl factors, ChIP can h functional annotatic open chromatin reg identified by DNase and FAIRE. | Input c background contro we sequenced the i DNA sample that w ChIP. Crosslinked cl sheared and the cr reversed without ca the immunoprecipit This sample is othe processed in a mar to the ChIP sample below. The input tra in revealing potenti arising from the se alignment process number differences reference genome sequenced samples regions of poor sec alignability. | For da terms and condition refer to [url] and [u
GSE74355	The Common Stress Responsive Transcription Factor ATF3 Binds Active Enhancers and Bookmarks Genomic Sites for Transcriptional Regulation [ChIP-seq]	Comprehensive ana global ATF3-binding the human genome quiescent and stres damage) conditions expressed at a low quiescent ATF3 was bind a large numbe sites that are often with genes required stress responses. W ATF3-binding sites

GSE ID	Title	Summary
		contained motifs fc transcription factor found that ATF3 bc portion of active er characterized by p3 and as a consequer often regulates exp the genes proximal enhancers. While D elicited by camptot dramatically alterec binding profile, mos genes regulated by DNA damage were ATF3 before the str Moreover, we demc ATF3 was co-localiz major stress respor genomic sites, there contributing to indu target gene expres: DNA damage. There results suggest tha bookmarks genomi interacts with other transcriptional regu quiescent and stres conditions.
GSE152063	Hyperosmotic stress induces downstream-of-gene transcription and alters the RNA Polymerase II interactome despite widespread transcriptional repression	Stress-induced reac transcription results synthesis of thousa downstream-of-gen containing transcrip mechanisms underl formation during ce remain unknown. N transcription profile induction in human using TT-TimeLapse revealed that hyper

GSE ID	Title	Summary
		stress induces wide transcriptional repr DoGs are produced of the transcription their upstream gen confirmed that the induced redistributi Polymerase (Pol) II with the transcriptic genes. Stress-induc alterations in the Pc interactome are ob mass spectrometry subunits of the clea polyadenylation ma remained Pol II-assc Integrator complex dissociated from Pc stress conditions. D catalytic subunit of Integrator complex, siRNAs induces hur readthrough transc parental genes part those of stress-indu Our results provide the mechanisms un production and hov activity influences D transcription. | This is composed of the listed below.
GSE140935	Chromatin Remodeling and Immediate Early Gene Activation by SLFN11 in Response to Replication Stress [ChIP-seq_H3K27ac_CPTO]	Schlafen 11 (SLFN recently arisen as a restriction factor ag replication stress. H that SLFN11 increa chromatin accessib wide, dominantly at response to replica

GSE ID	Title	Summary
		induced by the CHk prexasertib and the topoisomerase I inh camptothecin. Conc SLFN11 selectively cellular stress respc pathways by induci transcription of the Early Genes (IEGs) i FOS, EGR1, NFKB2 Both chromatin ope activation require th helicase activity of whereas extrinsic IE by serum induction independent. SLFN IEG activation is als across 55 non-isog cell lines by compar transcriptome data after camptothecin We conclude that S as a global regulatc chromatin structure intrinsic IEG activat potential to engage immune activation i replicative stress.
GSE31755	Histone Modifications by ChIP-seq from ENCODE/Stanford/Yale/Davis/Harvard	This data was gene ENCODE. If you hav about the data, con submitting laboratc (Peggy Farnham mailto:pfarnham@u questions concerni collection and usag Cayting mailto:pcayting@sta for data scoring an inquiries). If you hav

GSE ID	Title	Summary
		about the Genome track associated wi contact ENCODE (mailto:genome@so This track, produce the ENCODE Projec maps of histone mc genome-wide using different cell lines. T method involves firs formaldehyde to cr histones and other associated proteins DNA within cells. Th linked chromatin is extracted, sheared, immunoprecipitatec specific antibodies. reversal of cross-lir immunoprecipitatec sequenced and maj human reference g relative enrichment antibody-target (ep the genome is infer density of mapped Chemical modificati methylation or acet the histone protein chromatin influence expression by chan accessible the chro transcription factor each experiment (d particular antibody particular cell type) enrichment for the modified histone (S with sites that have enrichment (Peaks)

GSE ID	Title	Summary
		included for each c input signal, which r the control conditic antibody targeting ' performed. In gene following chemical r have associated ge phenotypes: | H3K4 H3K9Ac are consid marks of active or p active promoter reg H3K4me1 and H3K considered to be m or potentially active regions. H3K36me3 H3K79me2 are con marks of transcript elongation. H3K27r H3K9me3 are cons marks of inactive re data usage terms a conditions, please r and [url]
GSE286875	Improved spike-in normalization clarifies the relationship between active histone modifications and transcription	Spike-in normalizati quantitative analysi sequencing (ChIP-s Here we introduce a robust dual-spike-ir normalization appro seq (ChIP-wrangler) recent claims that a marks are depende transcription. Conc previous studies im used spike-in norma arrive at their concl used ChIP-wrangler acute depletion of F polymerase II has o impact on the levels

GSE ID	Title	Summary
		H3K4me3 and H3K with other studies, provide proof that maintenance of hist acetylation is not m consequence of on transcription. Our ir ChIP-seq normaliza provides increased guardrails for succe normalization, and a here refines the un of the intricate cros between RNA polyr activity and histone associated with tra
GSE113606	RNA-seq and ATAC-seq analysis of the role of PBRM1 under stress conditions	We used RNA-seq t characterize the co PBRM1 to gene exp NmuMG cells with a oxidative or low-glu conditions. Further, ATAC-seq to charac effects of PBRM1 c accessibility in HK-2 cell lines. Our analy role for PBRM1 in e genes associated w responses and with transcription factor such as the FOXO f
GSE82123	Genome-wide binding of hSTN1 in response to hydroxyurea-induced replication stress	We find that hSTN1 preferentially enrich rich repetitive sequ genome-wide after treatment.
GSE175712	YBX1 mediates translation of oncogenic transcripts to control cell	ChIP-seq for the co protein YBX1 in MC

GSE ID	Title	Summary
	competition in AML [ChIP-seq]	
GSE140932	Chromatin Remodeling and Immediate Early Gene Activation by SLFN11 in Response to Replication Stress [ATAC-seq_CHPTCHK1i]	Schlafen 11 (SLFN1 recently arisen as a restriction factor ac replication stress. H that SLFN11 increa chromatin accessib wide, dominantly at response to replica induced by the CHk prexasertib and the topoisomerase I inh camptothecin. Conc SLFN11 selectively cellular stress respo pathways by induci transcription of the Early Genes (IEGs) i FOS, EGR1, NFKB2 Both chromatin ope activation require th helicase activity of whereas extrinsic IE by serum induction independent. SLFN IEG activation is als across 55 non-isog cell lines by compar transcriptome data after camptothecin We conclude that S as a global regulato chromatin structure intrinsic IEG activat potential to engage immune activation i replicative stress.

GSE ID	Title	Summary
GSE144451	ChIP-seq analyses with eHAP cells	ChIP-seq analyses v cells for TFDP1 anc modifications
GSE43743	ChIP-Seq of transcriptional regulators in MM1S cells upon small molecule treatment.	Discovery of the ge location of proteins Seq has allowed glc of the key transcrip and chromatin regu control gene expres programs in various DNA-associated pro targeted for diseas This study investiga functions of small r therapeutics that ta associated process CDK9 and BRD4.
GSE112608	Targeting Processive Transcription Elongation via SEC Disruption for MYC-Induced Cancer Therapy	The super elongatic (SEC) is required fo productive transcri release of RNA poly (Pol II) with its P-TE and promoting tran processivity with its subunit. Malfunctio contributes to mult diseases including c we identify peptido compounds, KL-1 a structural homolog disrupt the interact the SEC scaffolding AFF4 and P-TEFb, r impaired release of promoter-proximal and a reduced aver processive transcri

GSE ID	Title	Summary
		elongation. SEC is r induction of heat-sh and treating cells w KL-2 attenuates the response from Dros human. SEC inhibiti downregulates MYC dependent transcri programs in mamm and delays tumor p a mouse xenograft MYC-driven cancer, that small-molecule SEC could be used therapy of MYC-ind
GSE133450	BRCA2 ChIP-seq hTert-HME1 cell line	Individuals with a si functional copy of t tumor suppressor h risks for breast, ova other solid tumor m The exact mechanis carcinogenesis due haploinsufficiency r unclear, but one po that at-risk cells are acute periods of de BRCA2 availability a (BRCA2-crisis), whi contribute to diseas establish an in vitro BRCA2-crisis that c chromatin remodeli activation of an NF- pathway in respons BRCA2 depletion. Mechanistically, we BRCA2 chromatin b histone acetylation, associated transcri activity as critical d

GSE ID	Title	Summary
		of the epigenetic re BRCA2-crisis. Thes alterations are refle transcriptional prof malignant tissues fr carriers and, theref reflect natural steps disease. By modelir crisis in vitro, we ha insights into pre-ne molecular alteratior enhance the develo preventative therap
GSE195763	EBF1 and HIF1a ChIPseq in breast cancer cell lines [ChIP-seq]	EBF1 and HIF1a Ch MDA-MB-157 cells normoxia and hypo
GSE209520	Chip sequencing after ER stress induced in HCC cells using tunicamycin	We report the appli CHIP sequencing-b technology in the h throughput analysis modifications in He Endoplasmic reticul related super enhar identified and their target genes were
GSE106252	RNA pol II ChIP-seq LR14 and HFF human fibroblasts	The aim of the expe determine if RNA p occupancy and abu altered human fibro

GSE ID	Title	Summary
		patient that has a n the gene Taf8, whic important role in RI mediated transcript
GSE68052	Nucleotide stress induction of HEXIM1 suppresses melanoma by modulating cancer cell-specific gene transcription [ChIP-Seq]	Cancer metabolism actively studied to g into tumorigenic su mechanisms and su In melanoma, we ide HEXIM1, a transcrip elongation regulato melanoma suppress participates in nucle regulation. HEXIM1 low in melanoma. Its overexpression sup melanoma while its accelerates tumor g HEXIM1 responds t stress. Knockdown rescues neural cres melanoma nucleotic phenotypes in vivo. Mechanistically, unc stress, HEXIM1 is ir form an inhibitory c P-TEFb, the kinase transcription elonga pause transcription tumorigenic genes. alteration in gene e also causes anti-tur transcripts to bind stabilized by HEXIM therefore plays an i in inhibiting cancer gene transcription facilitating anti-can expression. Our stu

GSE ID	Title	Summary
		novel role for HEXI coupling nucleotide with transcriptional melanoma.
GSE272705	PBAF associates with repressed chromatin to enable REST occupancy (ChIP-seq)	Multimeric SWI/SNI remodelers assemb conformations, with functions difficult t Importantly, mutatic SWI/SNF genes are distinct cancers, as specific component melanoma. Through comprehensive epig profiling of SWI/SN and their associate states in melanoma melanocytes, we fo subset of PBAF-exc regions unexpected with PRC2 and repr chromatin. Time-res approaches reveale regions are general sensitive to ATPase remodeling compar sites. Notably, PBAI bound loci are enric REST, a transcriptic represses neuronal turn, absence of AI consequent PBAF l REST ability to binc inactivate its target upregulation of neu synaptic transcripts signature also asso ARID2 mutations in patients. In sum, we a unique role for PI

GSE ID	Title	Summary
		generating accessib silencing transcripti repressed chromati
GSE161684	Role of hNABP2 (hSSB1) in transcriptional regulation	Use Global Run-On seq, chromatin immunoprecipitatio and DNA: RNA immunoprecipitatio to investigate whetl hNABP2 (hSSB1) re transcription.
GSE111924	Osmostress induced changes of chromatin architecture and transcription in mammalian cells	This SuperSeries is the SubSeries listec

GSE ID	Title	Summary
GSE180545	ATAC-seq analysis of HeLa stable cells expressing LAMPS targeting the HS2 enhancer	We describe that lig phase separation at HS2 enhancer incre chromatin accessib
GSE53366	Automated high-throughput chromatin immunoprecipitation	The submitted files seq data for the p3 transcriptional coac GM12878 cells and NRSF transcription GM12878 and Jurk generated using a f automated robotic immunoprecipitatio Cells were fixed usi formaldehyde (NRS 1% formaldehyde a samples).
GSE51065	Proteomic and Genomic Analysis of transcription factor TFII-I reveals insight into the response to cellular stress	We expressed bioti transcription factor in K562 cells and su crosslinked chroma streptavidin mediat We subjected the is to Illumina sequenc identified 18920 TF peaks. Nearest gen revealed 5886 gene with a TFII-I binding proliferating K562 c

GSE ID	Title	Summary
GSE100257	ChIP-seq for histone modifications during MCF10A-ER-Src cell transformation	We performed ChIP histone modificatio chromatin status ar during MCF10A-ER transformation.
GSE202072	EHF and CDX1 co-operate to drive extensive transcriptional reprogramming and chromatin remodelling in CRC cells [ATAC-seq]	EHF and CDX1 co-c drive differentation To identify the char chromatin remodell by these transcripti EHF and CDX1 wer expressed in poorly differentiated HCT1 and ATAC-seq analy performed.
GSE53372	Chromatin immunoprecipitation-sequencing of HEK293 cells (HEK293-HPT-blue) and HEK293 cells stably over-expressing the BAHD1 gene (HEK293-HPT-BAHD1)	We report the genc occupancy of the B protein in HEK293 expressing the BAH (HEK293-HPT-BAHI
GSE109721	ChIP-seq for H3K4me3, H3K27me3, H3K79me2 in MCF10A cells	We performed ChIP histones H3K4me3, and H3K79me2 in M to identify transcrip with bivalent and ps bivalent epigenetic modifications.
GSE79564	Input Strategy for Improving Analysis of ChIP-exo Data and Beyond [RNA-Seq]	Several recently em seq (chromatin immunoprecipitatio sequencing) based perform chemical s bead-bound immun chromatin, posing a for generating simil

GSE ID	Title	Summary
		input controls requ bioinformatics and analyses. Here we p versatile method fo technique-specific i for ChIP-based met utilize additional be processing steps. A this method allowec discovery of a nove binding motif from data.
GSE31417	Genome-wide study of HCFC1 binding sites and its associated transcription factors in cycling Human HeLa cells	Analysis of transcri regulation in humar implicated a large n promoter-specific D proteins that regula transcription via div mechanisms. In son these DNA-sequenc factors associate w intermediaries that interactions with th modifying enzymes intermediary is HCF factor-1; HCFC1). H identified for its inv herpes-simplex viru transcription and su shown to be an imp cycle regulator, has defined role in genc transcriptional regu show here, by chroi immunoprecipitatio high-throughput se analysis (ChIP-seq), a major transcriptic associated factor, v promoter associatic

GSE ID	Title	Summary
		positively with trans activity. Thus, in He 1 is observed on 5⁴ active CpG-island p Examination of the sequences underlyi 1-binding sites reve sequence motifs as the binding of (i) ZN Ronin/THAP11, (ii) (iii) YY1 sequence-s transcription factor Subsequent ChIP-s these four transcrip revealed a large co- of HCF-1 with these transcription factor approximately 90% bound promoters. T suggest that a relat number of transcrip - some (ZNF143 an Ronin/THAP11) in r combinations -- play in HeLa-cell transcr regulation in associ HCF-1.
GSE221869	Antibody-Free and Fast Profiling of Transcription Factor Binding Sites Using Aptamer-Guided Chromatin Cleavage with Sequencing (AptaChC-seq)	A large number of t factors are enriche DNA sites in the nu an important role ir regulation of gene e Chromatin immuno sequencing (ChIP s classic method for of chromatin transc landscape. This me by the complexity c signal to noise ratic number of cells req

GSE ID	Title	Summary
		cost, and heavy reli purity high-perform antibodies. In recen based on MNase ar transposase, CUT& CUT&Tag, ChIL-seq have been develope used, but they still r antibodies and take Here, we develop ar free and fast metho profiling transcripti binding sites, called Guided Chromatin (Sequencing (AptaC AptaChC-seq has tl advantages of high noise ratio, low cost reproducibility.
GSE44221	Identification of chromatin regulatory sites in conditions of basal and increased SF-1 dosage in the H295R adrenocortical tumor cell line	The goal of this stu identify chromatin r sites by FAIRE-seq conditions of basal increased dosage c transcription factor H295R human adre tumor cell line.
GSE163318	Genome-wide ChIP-seq analysis of H2A.Z and H2A.Zac in HeLa Kyoto cells upon knockdown of JAZF1	The goal was to inv genome-wide if JAZ the deposition as w acetylation of the h H2A.Z. ChIP-seq an revealed that deple leads to reduced ac H2A.Z at specific si
GSE30263	CTCF Binding Sites by ChIP-seq from ENCODE/University of Washington	This data was gene ENCODE. If you hav about the data, con

GSE ID	Title	Summary
		submitting laborato (Richard Sandstrom mailto:sull@u.washi you have questions Genome Browser tr associated with this contact ENCODE (mailto:genome@so This track is produc the ENCODE Projec displays maps of ge binding of the CTCI transcription factor cell lines using ChIF throughput sequen data usage terms a conditions, please r and [url]

GSE ID	Title	Summary
GSE83360	Investigation of Rsf1 genomic distribution and effect on gene expression	We report the effec knockdown or delet expression in huma cells. We also repor genomic distributio mouse embryonic s nucleosome positio type and RSF1 kno ESC.
GSE235477	ChIP-seq for Characterization of transcription factor function and patterns fo gene regulation in HepG2 cells	Transcription factor trans-acting proteir cis-regulatory elem in DNA to control g expression. Here, w genomic localizatior 529 sequence-spec 151 cofactors and regulators in the ca HepG2, for a total c broadly-termed DN Proteins (DAPs). We deep collection to r TF's impact on gen including identifying 26 candidate transc repressors. We exa Occupancy Target (the context of three genome organizatic biased motif placen enhancer-promoter involving HOT sites a substantial numbe chromatin regions v DAPs bound and ex properties, finding t MAFF/MAFK TF pa with transcriptional Altogether, these ar

GSE ID	Title	Summary
		provide novel insigh regulatory logic of cell line HepG2 gen demonstrate the us large genomic analy elucidation of indivi functions.

GSE ID	Title	Summary
GSE193713	Genome-wide maps of chromatin state for histone modifications and chromatin remodeling factors in WT and PHF6-KO conditions in Jurkat T-ALL cells.	We have carried ou throughput profilin modifications and t factors in mammali we generated stabl out lines by CRISPF technology in wild t cells. We electropor complex in Jurkat c conatining guide R either non-targetin PHF6 (KO) followed clone isolation and We confirmed abso total protein by WB formation (mutatior at the PHF6 genom Sanger sequencing over six to ten billio sequence from chr immunoprecipitatec generated genome- chromatin-state ma studied histone mo and transcription fa study provides a fr eluciadting the role regulating the chro landscape encompa activating and repr modifications and c remodelers in the c leukemia.
GSE190959	Genome-wide profiling and identification of disordered transcription factors through chemical precipitation mapping [ATAC-seq]	Intrinsically disorde (IDRs) in DNA-asso proteins have recer shown to play impo gene regulation. To understand these d

GSE ID	Title	Summary
		developed an antib independent assay disordered proteins wide by combining mediated precipitat generation sequenc seq, disordered pro precipitation follow sequencing). We fir Ewing sarcoma cell of principle given th like IDR of the oncc protein EWS-FLI1 h shown to play a crit this cancer. We finc seq produces thous strong peaks assoc diverse chromatin s Ewing sarcoma and types tested. These highly enriched for motifs matching EV other transcription prominent IDRs. Mc depletion of EWS-F sarcoma cells leads widespread reorgar DisP-seq signals, in decreases at EWS-I sites and marked ir sites associated wit mesenchymal differ Analysis of these ne signals shows that from redistribution transcription factor initially overlaps EV has an IDR that is r proper localization Our results thus sh

GSE ID	Title	Summary
		antibody-independe precipitation mappi enable genome-wid and identification o transcription factor their roles in compl regulation program
GSE118033	Epigenetic and transcriptional profiling of triple negative breast cancer	The human HCC18 frequently used as model for triple neg cancer (TNBC). Give dysregulated epige mechanisms are inv cancer pathogenes therapeutic strateg chromatin regulato histone deacetylase comprehensive unc the epigenome and profiling in HCC18C the framework for efficacy and molecu mechanisms of epig therapies. Thus, to interplay of transcr chromatin in the HC preclinical model, w nascent transcriptic using Precision Rur to sequencing (PRC Additionally, we ma genome-wide locati polymerase II (Pol II histone variant H2A histone modificatio ChIP-exo. ChIP-exo (ChIP-exo) is a refin ChIP-seq with near precision mapping DNA interactions. Ir

GSE ID	Title	Summary
		Descriptor, we pres information on expe design, data genera control analysis, an validation. We discu data lay the founda future analysis to u the relationship bet nascent transcriptic chromatin.
GSE190960	Genome-wide profiling and identification of disordered transcription factors through chemical precipitation mapping [ChIP-seq]	Intrinsically disorde (IDRs) in DNA-asso proteins have recer shown to play impo gene regulation. To understand these d developed an antib independent assay disordered proteins wide by combining mediated precipitat generation sequenc seq, disordered prc precipitation follow sequencing). We fir Ewing sarcoma cell of principle given th like IDR of the oncc protein EWS-FLI1 h shown to play a crit this cancer. We finc seq produces thous strong peaks assoc diverse chromatin s Ewing sarcoma and types tested. These highly enriched for motifs matching EV other transcription prominent IDRs. Mc

GSE ID	Title	Summary
		depletion of EWS-F sarcoma cells leads widespread reorgar DisP-seq signals, in decreases at EWS-I sites and marked ir sites associated wit mesenchymal differ Analysis of these n signals shows that from redistribution transcription factor initially overlaps EW has an IDR that is r proper localization Our results thus sh antibody-independe precipitation mappi enable genome-wid and identification o transcription factor their roles in compl regulation program
GSE262859	DFF-ChIP: a method to detect and quantify complex interactions between RNA polymerase II, transcription factors, and chromatin	Recently, we introdu chromatin immunop (ChIP) technique ut human DNA Fragm Factor (DFF) to dige prior to immunopre (DFF-ChIP) that pro precise location of complexes and thei with neighboring nu Here we expand the to new targets and useful information purification of DFF, conditions, and the crosslinking. DFF-C was performed indi

GSE ID	Title	Summary
		subunits of Mediat NELF that that do r with DNA directly, b interact with RNA p (Pol II). We found th was associated alm exclusively with pre complexes (PICs) ar and NELF were ass engaged Pol II and, potential intermedia PICs and early initia complexes. DFF-Ch used to analyze the of a tight binding tr factor, CTCF, and a binding factor, gluc receptor (GR), with crosslinking and to these results to thc standard ChIP-Seq sonication and CUT utilizes MNase to fr genomic DNA. The presented indicate ChIP reveals details occupancy that are using other methoc information revealir protein:protein inte
GSE291939	PBAF associates with repressed chromatin to enable REST occupancy [ATAC-Seq]	Multimeric SWI/SNI remodelers assemb conformations, with functions difficult tc Importantly, mutatic SWI/SNF genes are distinct cancers, as specific component melanoma. Through comprehensive epig

GSE ID	Title	Summary
		profiling of SWI/SN and their associate states in melanoma melanocytes, we fo subset of PBAF-exc regions unexpectec with PRC2 and repr chromatin. Time-res approaches reveale regions are general sensitive to ATPase remodeling compar sites. Notably, PBAF bound loci are enric REST, a transcriptic represses neuronal turn, absence of AF consequent PBAF l REST ability to binc inactivate its target upregulation of neu synaptic transcripts signature also asso ARID2 mutations in patients. In sum, we a unique role for PE generating accessik silencing transcripti repressed chromati
GSE67454	A simple method for generating high-resolution maps of genome wide protein binding	Chromatin immuno (ChIP) and its deriv main techniques us determine transcrip binding sites. Howe conventional ChIP v sequencing (ChIP-s problems with poor and newer techniqu significant experime alterations and com

GSE ID	Title	Summary
		bioinformatics. Her upon our high-reso crosslinking ChIP-se seq) method and cc existing methodolo micrococcal nuclea both endo- and exc activity to fragment chromatin and ther precise protein-DN high-resolution X-C achieves single bas resolution of transc binding. A significar of this protocol is t alteration to the co ChIP-seq workflow bioinformatic proce
GSE51117	Genome wide ChIP-Seq profiles of p300 and CBP in control and cells expressing HSP70K71E	There were more e and CBP bindings a promoter regions ir expresing HSP70K compared to the cc
GSE163481	Genome-wide binding of transcription factor IRF3	Genome-wide bindi was analysed in MM using ChIP-seq high approach. MM1.S c fixed in 1% formald Aesar) for 15min at temperature. cells v with hypotonic lysis 15minutes followed lysis buffer for 15m ChIP for IRF3 perfc standard ChIP-seq the genome-wide b data was analysed v revealed peaks enri cell cycle genes TS

GSE ID	Title	Summary
GSE289539	Differential oligomerization regulates PHF13 chromatin function and affinity [ATAC-seq]	PHF13 is a H3K4m reader, that modula chromatin processe transcription, DNA response and chror architecture. PHF13 aberrantly regulate cancers and its mis alters the epigeneti of key transcriptior regulate Epithelial t Mesenchymal trans study we sought to how PHF13s chrom and diverse chroma are intrinsically regu results show that P oligomerize via con ordered regions in terminus increasing chromatin valence a Impressively, a 4-fo expression of PHF1 sufficient to globally chromatin, depende ordered dimerizing oligomerization pot Unexpectedly, we d that PHF13 can sel independent of its domains via intrinsi disordered regions, conversely reduced chromatin affinity. support that there balance between PH ordered and disord that regulates PHF affinity, and sugges can phase transitio

GSE ID	Title	Summary
		polymer-polymer ar phase separation st impact chromatin s function.

About Privacy Notice Cookie Preferences Contact Us