

Your grade: 100%

Your latest: 100% • Your highest: 100% • To pass you need at least 80%. We keep your highest score.

[Next item →](#)

1. Mood's median test uses _____ statistic to make an inference.

1 / 1 point

- ☐ The Students t
- ☐ The Normal
- ☒ The Chi Square
- ☐ No statistic

 **Correct**

Mood's median test uses the Chi Square statistic to make an inference

2. If Levene's test shows significance, it is recommended that we move to the _____

1 / 1 point

- ☐ ANOVA
- ☐ Mood's median test
- ☒ Welch's t-test.
- ☐ Goodness of Fit test

 **Correct**

If Levene's test shows significance, it is recommended that we move to the Welch's t-test

3. The Mann Whitney test is the non-parametric test for the _____

1 / 1 point

- ☐ Goodness of fit test
- ☐ Paired treatment t test
- ☒ Independent t test
- ☐ Pearson correlation test

 **Correct**

The Mann Whitney test is the non-parametric test for the independent t test

4. The assumptions for the Mann Whitney test is (are) _____

1 / 1 point

- ☐ distribution of the differences of the data should be bell shaped
- ☒ simple random samples and each of the samples must have 10 or more values only
- ☐ each of the samples must have 10 or more values
- ☐ simple random samples

 **Correct**

The assumptions for the Mann Whitney test are the simple random samples and each of the samples must have 10 or more values only

5. The Kruskal Wallis test and the Friedman Test use a statistic related to the _____

1 / 1 point

- ☐ F Distribution
- ☐ Students t distribution
- ☐ normal distribution
- ☒ Chi Square distribution

 **Correct**

The Kruskal Wallis test and the Friedman Test use a statistic related to the Chi Square distribution