

MEME16203 Linear Models Marking Guide**Assignment 2****UNIVERSITI TUNKU ABDUL RAHMAN**

Faculty:	FES	Unit Code:	MEME16203
Course:	MAC	Unit Title:	Linear Models
Year:	1,2	Lecturer:	Dr Yong Chin Khian
Session:	May 2023		
Due by:			

- Q1. Show that any two matrices \mathbf{W} and \mathbf{X} have the same column space if there exist matrices \mathbf{F} and \mathbf{G} such that $\mathbf{WG} = \mathbf{X}$ and $\mathbf{XF} = \mathbf{W}$.

Ans.

$$\begin{aligned}
 \mathbf{a} \in \mathcal{C}(\mathbf{X}) &\implies \mathbf{a} = \mathbf{X}\mathbf{b} \text{ for some } \mathbf{b} \\
 &\implies \mathbf{a} = \mathbf{WG}\mathbf{b} \text{ for some } \mathbf{b} \text{ (Because } \mathbf{X} = \mathbf{WG}\text{)} \\
 &\implies \mathbf{a} = \mathbf{W}\mathbf{c} \text{ for some vector } \mathbf{c} \\
 &\implies \mathbf{a} \in \mathcal{C}(\mathbf{W})
 \end{aligned}$$

So, $\mathcal{C}(\mathbf{X}) \subseteq \mathcal{C}(\mathbf{W})$

Then similarly,

$$\begin{aligned}
 \mathbf{g} \in \mathcal{C}(\mathbf{W}) &\implies \mathbf{g} = \mathbf{W}\mathbf{h} \text{ for some } \mathbf{h} \\
 &\implies \mathbf{g} = \mathbf{XF}\mathbf{h} \text{ for some } \mathbf{h} \text{ (Because } \mathbf{W} = \mathbf{XF}\text{)} \\
 &\implies \mathbf{g} = \mathbf{X}\mathbf{k} \text{ for some } \mathbf{k} \\
 &\implies \mathbf{g} \in \mathcal{C}(\mathbf{X})
 \end{aligned}$$

So, $\mathcal{C}(\mathbf{W}) \subseteq \mathcal{C}(\mathbf{X})$

And hence,

$$\mathcal{C}(\mathbf{X}) = \mathcal{C}(\mathbf{W})$$

- Q2. Suppose

$$Y_{ij} = \mu_i + \epsilon_{ij}, i = 1, 2; j = 1, 2, 3.$$

$$\text{Let } \boldsymbol{\epsilon} = \begin{bmatrix} \epsilon_{11} \\ \epsilon_{12} \\ \epsilon_{13} \\ \epsilon_{21} \\ \epsilon_{22} \\ \epsilon_{23} \end{bmatrix} \sim N \left(\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \sigma^2 \begin{bmatrix} 9 & 0 & 0 & 0 & 0 & 0 \\ 0 & 36 & 0 & 0 & 0 & 0 \\ 0 & 0 & 64 & 0 & 0 & 0 \\ 0 & 0 & 0 & 9 & 0 & 0 \\ 0 & 0 & 0 & 0 & 36 & 0 \\ 0 & 0 & 0 & 0 & 0 & 64 \end{bmatrix} \right).$$

What is the BLUE of $3\mu_1 + 5\mu_2$? Explain carefully.

Ans.

$$\begin{bmatrix} Y_{11} \\ Y_{12} \\ Y_{13} \\ Y_{21} \\ Y_{22} \\ Y_{23} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} + \epsilon$$

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

where

$$\mathbf{X} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \mathbf{Y} = \begin{bmatrix} Y_{11} \\ Y_{12} \\ Y_{13} \\ Y_{21} \\ Y_{22} \\ Y_{23} \end{bmatrix}$$

Let $\mathbf{V}^{-1/2} = \text{diag}(\frac{1}{\sqrt{9}}, \frac{1}{\sqrt{36}}, \frac{1}{\sqrt{64}}, \frac{1}{\sqrt{9}}, \frac{1}{\sqrt{36}}, \frac{1}{\sqrt{64}})$ and

$$\mathbf{Z} = \mathbf{V}^{-1/2}\mathbf{Y} = \begin{bmatrix} \frac{1}{\sqrt{9}}y_{11} \\ \frac{1}{\sqrt{36}}y_{12} \\ \frac{1}{\sqrt{64}}y_{13} \\ \frac{1}{\sqrt{9}}y_{21} \\ \frac{1}{\sqrt{36}}y_{22} \\ \frac{1}{\sqrt{64}}y_{23} \end{bmatrix}$$

$$\begin{aligned} E(\mathbf{Z}) &= \mathbf{V}^{-1/2}E(\mathbf{Y}) \\ &= \mathbf{V}^{-1/2}\mathbf{X}\boldsymbol{\beta} \\ &= \mathbf{W}\boldsymbol{\beta} \end{aligned}$$

where

$$\mathbf{W} = \begin{bmatrix} \frac{1}{\sqrt{9}} & 0 \\ \frac{1}{\sqrt{36}} & 0 \\ \frac{1}{\sqrt{64}} & 0 \\ 0 & \frac{1}{\sqrt{9}} \\ 0 & \frac{1}{\sqrt{36}} \\ 0 & \frac{1}{\sqrt{64}} \end{bmatrix}$$

$$\begin{aligned} \text{Var}(\mathbf{Z}) &= \mathbf{V}^{-1/2^T} \text{Var}(\mathbf{Y}) \mathbf{V}^{-1/2} \\ &= \text{diag} \left(\frac{1}{\sqrt{9}}, \frac{1}{\sqrt{36}}, \frac{1}{\sqrt{64}}, \frac{1}{\sqrt{9}}, \frac{1}{\sqrt{36}}, \frac{1}{\sqrt{64}} \right) \times \text{diag} (9\sigma^2, 36\sigma^2, 8\sigma^2, 3\sigma^2, 6\sigma^2, 3\sigma^2) \end{aligned}$$

MEME16203 Linear Models Marking Guide

$$\times \text{diag} \left(\frac{1}{\sqrt{9}}, \frac{1}{\sqrt{36}}, \frac{1}{\sqrt{64}}, \frac{1}{\sqrt{9}}, \frac{1}{\sqrt{36}}, \frac{1}{\sqrt{64}} \right) \\ = \sigma^2 \mathbf{I}$$

Thus, \mathbf{Z} follows a Gauss-Markov model with model matrix \mathbf{W} .

$$(\mathbf{W}^T \mathbf{W}) = \begin{bmatrix} \frac{1}{\sqrt{9}} & \frac{1}{\sqrt{36}} & \frac{1}{\sqrt{64}} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{9}} & \frac{1}{\sqrt{36}} & \frac{1}{\sqrt{64}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{9}} & 0 \\ \frac{1}{\sqrt{36}} & 0 \\ \frac{1}{\sqrt{64}} & 0 \\ 0 & \frac{1}{\sqrt{9}} \\ 0 & \frac{1}{\sqrt{36}} \\ 0 & \frac{1}{\sqrt{64}} \end{bmatrix} = \begin{bmatrix} 0.1545 & 0 \\ 0 & 0.1545 \end{bmatrix}$$

$$(\mathbf{W}^T \mathbf{W})^{-1} = \begin{bmatrix} 6.4725 & 0 \\ 0 & 6.4725 \end{bmatrix}$$

$$\mathbf{W}^T \mathbf{Z} = \begin{bmatrix} \frac{1}{\sqrt{9}} & \frac{1}{\sqrt{36}} & \frac{1}{\sqrt{64}} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{9}} & \frac{1}{\sqrt{36}} & \frac{1}{\sqrt{64}} \end{bmatrix} \begin{bmatrix} \frac{1}{9}y_{11} + \frac{1}{36}y_{12} + \frac{1}{64}y_{13} \\ \frac{1}{9}y_{21} + \frac{1}{36}y_{22} + \frac{1}{64}y_{23} \end{bmatrix}$$

So, the BLUE of $3\mu_1 + 5\mu_2$ is

$$\begin{aligned} & [3 \quad 5][(\mathbf{W}^T \mathbf{W})^{-1} \mathbf{W}^T \mathbf{Z}] \\ &= [3 \quad 5] \begin{bmatrix} 6.4725 & 0 \\ 0 & 6.4725 \end{bmatrix} \begin{bmatrix} \frac{1}{9}y_{11} + \frac{1}{36}y_{12} + \frac{1}{64}y_{13} \\ \frac{1}{9}y_{21} + \frac{1}{36}y_{22} + \frac{1}{64}y_{23} \end{bmatrix} \\ &= 19.4175[\frac{1}{9}y_{11} + \frac{1}{36}y_{12} + \frac{1}{64}y_{13}] = 32.3625[\frac{1}{9}y_{21} + \frac{1}{36}y_{22} + \frac{1}{64}y_{23}] \end{aligned}$$

- Q3. Consider a problem of quartic regression in one variable, X . In particular, suppose that $n = 6$ values of a response y are related to values $x = 0, 1, 2, 3, 4, 5$ by a linear model $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$ for

$$\mathbf{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \\ y_6 \end{bmatrix}, \quad \mathbf{X} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \\ 1 & 4 & 16 & 64 \\ 1 & 5 & 25 & 125 \end{bmatrix}, \quad \boldsymbol{\beta} = \begin{bmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{bmatrix}, \quad \text{and} \quad \boldsymbol{\epsilon} = \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \epsilon_3 \\ \epsilon_4 \\ \epsilon_5 \\ \epsilon_6 \end{bmatrix}$$

Define

$$\mathbf{W} = \begin{bmatrix} 1 & -5 & 5 & -5 \\ 1 & -3 & -1 & 7 \\ 1 & -1 & -4 & 4 \\ 1 & 1 & -4 & -4 \\ 1 & 3 & -1 & -7 \\ 1 & 5 & 5 & 5 \end{bmatrix}$$

- (a) Show that $\mathbf{y} = \mathbf{W}\boldsymbol{\gamma} + \boldsymbol{\epsilon}$ is reparameterization of $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$, where $\boldsymbol{\gamma}^T = [\gamma_1, \gamma_2, \gamma_3, \gamma_4]$.

MEME16203 Linear Models Marking Guide*Ans.* $\mathbf{W} = \mathbf{XF}$ where

$$\mathbf{F} = \begin{bmatrix} 1 & -5 & 5 & -5 \\ 0 & 2 & -7.5 & 137/6 \\ 0 & 0 & 1.5 & -12.5 \\ 0 & 0 & 0 & 5/3 \end{bmatrix}$$

and

 $\mathbf{X} = \mathbf{WG}$ where

$$\mathbf{G} = \begin{bmatrix} 1 & 2.5 & 55/6 & 37.5 \\ 0 & 0.5 & 2.5 & 11.9 \\ 0 & 0 & 2/3 & 5 \\ 0 & 0 & 0 & 0.6 \end{bmatrix}$$

- (b) Notice that $\mathbf{W}^T\mathbf{W}$ is diagonal. Suppose that $\mathbf{y}^T = (-2, 0, 4, 2, 2, 1)$. Find the OLS estimate of $\boldsymbol{\gamma}$ in the model $\mathbf{y} = \mathbf{W}\boldsymbol{\gamma} + \boldsymbol{\epsilon}$ and then OLS estimate of $\boldsymbol{\beta}$ in the original model. (Find numerical values.)

Ans.

$$\mathbf{W}^T\mathbf{W} = \text{diag}(6, 70, 84, 182)$$

$$(\mathbf{W}^T\mathbf{W})^{-1} = \text{diag}(\frac{1}{6}, \frac{1}{70}, \frac{1}{84}, \frac{1}{182})$$

$$\text{Then, } \hat{\boldsymbol{\gamma}}_{OLS} = (\mathbf{W}^T\mathbf{W})^{-1}\mathbf{W}^T\mathbf{y} = \text{diag}(\frac{1}{6}, \frac{1}{70}, \frac{1}{84}, \frac{1}{182}) \begin{bmatrix} 7 \\ 16 \\ -31 \\ 9 \end{bmatrix} = \begin{bmatrix} \frac{7}{6} \\ \frac{8}{35} \\ \frac{-31}{84} \\ \frac{9}{182} \end{bmatrix}$$

Since $\mathbf{W}\hat{\boldsymbol{\gamma}}_{OLS} = \mathbf{XF}\boldsymbol{\gamma}_{OLS}$, we must have $\hat{\boldsymbol{\beta}}_{OLS} = \mathbf{F}\boldsymbol{\gamma}_{OLS}$.

$$\boldsymbol{\beta}_{OLS} = \begin{bmatrix} 1 & -5 & 5 & -5 \\ 0 & 2 & -7.5 & 137/6 \\ 0 & 0 & 1.5 & -12.5 \\ 0 & 0 & 0 & 5/3 \end{bmatrix} \begin{bmatrix} \frac{7}{6} \\ \frac{8}{35} \\ \frac{-31}{84} \\ \frac{9}{182} \end{bmatrix} = \begin{bmatrix} -2.06868132 \\ 4.35412088 \\ -1.17170330 \\ 0.08241758 \end{bmatrix}$$

- Q4. Two varieties of corn (variety A and variety B) were compared in a field trail. In addition to the varieties, three levels of nitrogen were used (0, 30, and 60 pounds per acre (lb/a). Six different fields were used, and the six combinations of varieties and nitrogen levels were randomly assigned to the fields. Let Y_{ij} denote the yield (in bushels per acre) of the i^{th} variety of corn when the j^{th} level of nitrogen is applied. Throughout this question, ϵ_{ij} , $i = 1, 2, j = 1, 2, 3$, denote independent $N(0, \sigma^2)$ random variables where σ^2 is an unknown variance. The following two models were proposed:

MEME16203 Linear Models Marking Guide

$$\begin{array}{lcl}
 \text{Model 1:} & \begin{bmatrix} Y_{11} \\ Y_{12} \\ Y_{13} \\ Y_{21} \\ Y_{22} \\ Y_{23} \end{bmatrix} & = \begin{bmatrix} 1 & 0 & -30 & 900 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 30 & 900 \\ 0 & 1 & -30 & 900 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 30 & 900 \end{bmatrix} \begin{bmatrix} \gamma_1 \\ \gamma_2 \\ \delta_1 \\ \delta_2 \end{bmatrix} + \begin{bmatrix} \epsilon_{11} \\ \epsilon_{12} \\ \epsilon_{13} \\ \epsilon_{21} \\ \epsilon_{22} \\ \epsilon_{23} \end{bmatrix} \\
 \text{Model 2:} & \begin{bmatrix} Y_{11} \\ Y_{12} \\ Y_{13} \\ Y_{21} \\ Y_{22} \\ Y_{23} \end{bmatrix} & = \begin{bmatrix} 1 & 1 & 0 & -1 & 1 \\ 1 & 1 & 0 & 0 & -2 \\ 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & -1 & 1 \\ 1 & 0 & 1 & 0 & -2 \\ 1 & 0 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} \mu \\ \alpha_1 \\ \alpha_2 \\ \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} \epsilon_{11} \\ \epsilon_{12} \\ \epsilon_{13} \\ \epsilon_{21} \\ \epsilon_{22} \\ \epsilon_{23} \end{bmatrix}
 \end{array}$$

- (a) With respect to the effects of varieties and nitrogen levels on corn yields, interpret the parameters γ_1 and δ_1 in Model 1.

Ans.

$E(Y_{12}) = \gamma_1$, thus γ_1 is the mean corn yield for variety A grown with nitrogen applied at 30lb/a.

$E[Y_{13} - Y_{11}] = 60\delta_1$, thus $60\delta_1$ is the difference between mean yields for nitrogen applied at 60lb/a and 0lb/a for either variety A or variety B.

- (b) For Model 1, indicate which of the following quantities are estimable

$$\gamma_1 - \gamma_2; \quad \gamma_1 - 10\delta_1 + 100\delta_2$$

Give a brief explanation, to support your conclusions.

Ans.

Since the model matrix has full column rank, any linear combination of the parameters is estimable. Hence, both quantities above are estimable.

- (c) For Model 2, determine if $\mu + \alpha_1$ is estimable? Give a brief explanation to support your conclusion.

Ans.

$\mu + \alpha_1$ is estimable because $E[(Y_{11} + Y_{12} + Y_{13})/3] = \mu + \alpha_1$

- (d) Expressing Model 2 as $\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$, a solution to the normal equations is $\mathbf{b} = (\mathbf{X}^T\mathbf{X})^{-}\mathbf{X}^T\mathbf{Y}$. Explain how a generalized inverse $(\mathbf{X}^T\mathbf{X})^{-}$ can be computed. (You are not expected to obtain a numerical value for $(\mathbf{X}^T\mathbf{X})^{-}$, just briefly outline a procedure for how it can be computed.)

MEME16203 Linear Models Marking Guide

Ans. Delete the third row and the third column of $(\mathbf{X}^T \mathbf{X})^-$. Invert the remaining 4×4 matrix. Fill in zeros for the third row and third column of the generalized inverse. Alternatively you could use the spectral or singular value decomposition of $(\mathbf{X}^T \mathbf{X})^-$ to obtain a generalized inverse.

- (e) Using $\mathbf{b} = (\mathbf{X}^T \mathbf{X})^- \mathbf{X}^T \mathbf{Y}$ from Part (d), define the estimator

$$\hat{\alpha}_1 - \hat{\alpha}_2 = [0 \ 1 \ -1 \ 0 \ 0] \mathbf{b}$$

What are the properties of this estimator?

Ans.

$E(Y_{21} - Y_{11}) = \alpha_1 - \alpha_2$, hence $\alpha_1 - \alpha_2$ is estimable. Then by Gauss Markov theorem, $[0 \ 1 \ -1 \ 0 \ 0] \mathbf{b}$ is the unique best linear unbiased estimator (BLUE) for $[0 \ 1 \ -1 \ 0 \ 0] \boldsymbol{\beta} = \alpha_1 - \alpha_2$.

- (f) Would the residual sum of squares from fitting models (1) and (2) be the same?

Ans.

Yes, because the columns of the model matrices span the same linear space. Let $\mathbf{W} = [\mathbf{w}_1 | \mathbf{w}_2 | \mathbf{w}_3 | \mathbf{w}_4]$ denote the model matrix for Model 1, and let $\mathbf{X} = [\mathbf{x}_1 | \mathbf{x}_2 | \mathbf{x}_3 | \mathbf{x}_4 | \mathbf{x}_5]$ denote the model matrix for Model 2. Then, $\mathbf{x}_2 = \mathbf{w}_1$, $\mathbf{x}_3 = \mathbf{w}_2$, $\mathbf{x}_1 = \mathbf{w}_1 + \mathbf{w}_2$, $\mathbf{x}_4 = \mathbf{w}_3/30$, and $\mathbf{x}_5 = (3\mathbf{w}_4/900) - 2\mathbf{w}_1 - 2\mathbf{w}_2$. Also, $\mathbf{w}_1 = \mathbf{x}_2$, $\mathbf{w}_2 = \mathbf{x}_3$, $\mathbf{w}_3 = 30\mathbf{x}_4$, and $\mathbf{w}_4 = 900(\mathbf{x}_5 + 2\mathbf{x}_2 + 2\mathbf{x}_3)/3$. Or

$$\mathbf{W} = \mathbf{X}\mathbf{F}, \text{ where } \mathbf{F} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1800/3 \\ 0 & 1 & 0 & 1800/3 \\ 0 & 0 & 30 & 0 \\ 0 & 0 & 0 & 900/3 \end{bmatrix} \text{ and}$$

$$\mathbf{X} = \mathbf{W}\mathbf{G}, \text{ where } \mathbf{G} = \begin{bmatrix} 1 & 1 & 0 & 0 & -2 \\ 1 & 0 & 1 & 0 & -2 \\ 0 & 0 & 0 & 1/30 & 0 \\ 0 & 0 & 0 & 0 & 3/900 \end{bmatrix}$$

- Q5. Two varieties of corn (variety A and variety B) were compared in a field trial. In addition to the varieties, three levels of nitrogen were used (10, 20 and 30 pounds per acre (lb/a)). Six different fields were used, and the six combinations of varieties and nitrogen levels were randomly assigned to the fields. Suppose the data are as follows.

MEME16203 Linear Models Marking Guide

Field	i	j	Amount of Nitrogen (x_{ij})	Bushels per acre (y_{ij})
1	1	1	10	80
2	1	2	20	120
3	1	3	30	140
4	2	1	10	60
5	2	2	20	150
6	2	3	30	170

Consider a Gauss-Markov model

$$y_{ij} = \mu + \alpha_i + \gamma_i X_{ij} + \gamma_3 X_{ij}^2 + \epsilon_{ij}$$

where

- y_{ij} is the yield (in bushels per acre) of the i^{th} variety of corn when the j^{th} level of nitrogen is applied.
- X_{ij} denote the level of nitrogen administered to the corn,
- $\mu, \alpha_1, \alpha_2, \gamma_1, \gamma_2, \gamma_3$ are unknown parameters, and
- ϵ_{ij} denotes a random error with $\epsilon_{ij} \sim NID(0, \sigma^2)$ where $\sigma^2 > 0$.

Let $\boldsymbol{\beta} = (\mu, \alpha_1, \alpha_2, \gamma_1, \gamma_2, \gamma_3)^T$, $\mathbf{y} = [y_{11}, y_{12}, y_{13}, y_{21}, y_{22}, y_{23}]^T$, and $\boldsymbol{\epsilon} = [\epsilon_{11}, \epsilon_{12}, \epsilon_{13}, \epsilon_{21}, \epsilon_{22}, \epsilon_{23}]^T$.

- (a) Determine the design matrix \mathbf{X} so that $E(\mathbf{y}) = \mathbf{X}\boldsymbol{\beta}$.

Ans.

$$\mathbf{X} = \begin{bmatrix} 1 & 1 & 0 & 10 & 0 & 100 \\ 1 & 1 & 0 & 20 & 0 & 400 \\ 1 & 1 & 0 & 30 & 0 & 900 \\ 1 & 0 & 1 & 0 & 10 & 100 \\ 1 & 0 & 1 & 0 & 20 & 400 \\ 1 & 0 & 1 & 0 & 30 & 900 \end{bmatrix}$$

- (b) Determine whether α_2 is estimable. Prove that your answer is correct.

Ans.

let $\mathbf{d}^T = [1 \ -1 \ -1 \ 0 \ 0 \ 0]$, then $\mathbf{X}\mathbf{d} = \mathbf{0}$.

Let $\mathbf{c} = [0 \ 0 \ 1 \ 0 \ 0 \ 0]$, then $\mathbf{c}^T \boldsymbol{\beta} = \alpha_2$, but $\mathbf{c}^T \mathbf{d} = -1 \neq 0$, i.e., there exists a vector \mathbf{d} such that $\mathbf{X}\mathbf{d} = \mathbf{0}$ but $\mathbf{c}_1^T \boldsymbol{\beta} \neq 0$. Thus, α_2 is not estimable.

MEME16203 Linear Models Marking Guide

- (c) Show that
- $\mathcal{C}(\mathbf{X}) = \mathcal{C}(\mathbf{W})$
- , where

$$\mathbf{W} = \begin{bmatrix} 1 & 1 & -1 & 0 & 1 \\ 1 & 1 & 0 & 0 & -2 \\ 1 & 1 & 1 & 0 & 1 \\ 1 & -1 & 0 & -1 & 1 \\ 1 & -1 & 0 & 0 & -2 \\ 1 & -1 & 0 & 1 & 1 \end{bmatrix}.$$

Ans.

$\mathcal{C}(\mathbf{X}) = \mathcal{C}(\mathbf{W})$ means that the space spanned by the columns of \mathbf{X} is a basis for space spanned by the columns of \mathbf{W} and vice versa. Thus there is a matrix \mathbf{F} such that $\mathbf{W} = \mathbf{X}\mathbf{F}$ and a matrix \mathbf{G} such that $\mathbf{X} = \mathbf{W}\mathbf{G}$.

$$\begin{aligned} \mathbf{W} &= \begin{bmatrix} 1 & 1 & -1 & 0 & 1 \\ 1 & 1 & 0 & 0 & -2 \\ 1 & 1 & 1 & 0 & 1 \\ 1 & -1 & 0 & -1 & 1 \\ 1 & -1 & 0 & 0 & -2 \\ 1 & -1 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & 10 & 0 & 100 \\ 1 & 1 & 0 & 20 & 0 & 400 \\ 1 & 1 & 0 & 30 & 0 & 900 \\ 1 & 0 & 1 & 0 & 10 & 100 \\ 1 & 0 & 1 & 0 & 20 & 400 \\ 1 & 0 & 1 & 0 & 30 & 900 \end{bmatrix} \begin{bmatrix} 1 & 0.5 & -1 & -1 & 12.36 \\ 0 & 0.5 & -1 & 1 & 12.36 \\ 0 & -1.5 & 1 & -1 & 12.36 \\ 0 & 0.0 & 1/10 & 0 & -2.43 \\ 0 & 0.0 & 0 & 1/10 & -2.43 \\ 0 & 0.0 & 0 & 0 & 0.05 \end{bmatrix} = \\ \mathbf{X}\mathbf{F} & \\ \mathbf{X} &= \begin{bmatrix} 1 & 1 & 0 & 10 & 0 & 100 \\ 1 & 1 & 0 & 20 & 0 & 400 \\ 1 & 1 & 0 & 30 & 0 & 900 \\ 1 & 0 & 1 & 0 & 10 & 100 \\ 1 & 0 & 1 & 0 & 20 & 400 \\ 1 & 0 & 1 & 0 & 30 & 900 \end{bmatrix} = \begin{bmatrix} 1 & 1 & -1 & 0 & 1 \\ 1 & 1 & 0 & 0 & -2 \\ 1 & 1 & 1 & 0 & 1 \\ 1 & -1 & 0 & -1 & 1 \\ 1 & -1 & 0 & 0 & -2 \\ 1 & -1 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0.5 & 0.5 & 10 & 10 & 436.67 \\ 0 & 0.5 & -0.5 & 10 & -10 & 0 \\ 0 & 0.0 & 0.0 & 10 & 0 & 445.00 \\ 0 & 0.0 & 0.0 & 0 & 10 & 445.00 \\ 0 & 0.0 & 0.0 & 0 & 0 & 18.33 \end{bmatrix} = \\ \mathbf{W}\mathbf{G} & \end{aligned}$$

Thus, $\mathcal{C}(\mathbf{X}) = \mathcal{C}(\mathbf{W})$

- (d) Verify that
- $\tau = \mu + \alpha_1$
- is estimable, then obtained the unique BLUE of
- τ
- .

Ans.

$$\begin{aligned} &E(3Y_{11} - 3Y_{12} + Y_{13}) \\ &= 3(\mu + \alpha_1 + \gamma_1 + \gamma_3) - 3(\mu + \alpha_1 + 2\gamma_1 + 4\gamma_3) + (\mu + \alpha_1 + 3\gamma_1 + 9\gamma_3) \\ &= (3 - 3 + 1)(\mu + \alpha_1) + (3 - 6 + 3)\gamma_1 + (3 - 12 + 9)\gamma_3 \\ &= \mu + \alpha_1. \end{aligned}$$

Thus, $\mu + \alpha_1$ is estimable.

$$\text{Since } \mathbf{c}^T \boldsymbol{\beta} = [1 \ 1 \ 0 \ 0 \ 0 \ 0] \begin{bmatrix} \mu \\ \alpha_1 \\ \alpha_2 \\ \gamma_1 \\ \gamma_2 \\ \gamma_3 \end{bmatrix} = \mu + \alpha_1 \text{ is estimable, thus, by Gauss}$$

MEME16203 Linear Models Marking Guide

Markov theorem, $\mathbf{c}^T \hat{\boldsymbol{\beta}}$ is the unique BLUE of $\mathbf{c}^T \boldsymbol{\beta}$.

Since $\mathcal{C}(\mathbf{X}) = \mathcal{C}(\mathbf{W})$, then $\mathbf{c}^T \hat{\boldsymbol{\beta}} = \mathbf{c}^T \mathbf{F} \hat{\boldsymbol{\gamma}}$, where $\hat{\boldsymbol{\gamma}} = (\mathbf{W}^T \mathbf{W})^{-1} \mathbf{W}^T \mathbf{y}$

$$(\mathbf{W}^T \mathbf{W}) = \begin{bmatrix} 6 & 0 & 0 & 0 & 0 \\ 0 & 6 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 12 \end{bmatrix}; (\mathbf{W}^T \mathbf{W})^{-1} = \begin{bmatrix} \frac{1}{6} & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{6} & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{12} \end{bmatrix} \text{ and } (\mathbf{W}^T \mathbf{y}) =$$

$$\begin{bmatrix} 720 \\ -40 \\ 60 \\ 110 \\ -90 \end{bmatrix}$$

$$\hat{\boldsymbol{\gamma}} = \begin{bmatrix} 120 \\ -6.67 \\ 30 \\ 55 \\ -7.5 \end{bmatrix}$$

$$\mathbf{c}^T \hat{\boldsymbol{\beta}} = [1 \ 1 \ 0 \ 0 \ 0 \ 0] \begin{bmatrix} 1 & 0.5 & -1 & -1 & 12.36 \\ 0 & 0.5 & -1 & 1 & 12.36 \\ 0 & -1.5 & 1 & -1 & 12.36 \\ 0 & 0.0 & 1/10 & 0 & -2.43 \\ 0 & 0.0 & 0 & 1/10 & -2.43 \\ 0 & 0.0 & 0 & 0 & 0.05 \end{bmatrix} \begin{bmatrix} 120 \\ -6.67 \\ 30 \\ 55 \\ -7.5 \end{bmatrix} =$$