

**MEME15203 Statistical Inference****Assignment 2****UNIVERSITI TUNKU ABDUL RAHMAN**

Faculty:	FES	Unit Code:	MEME15203
Course:	MAC	Unit Title:	Statistical Inference
Year:	1,2	Lecturer:	Dr Yong Chin Khian
Session:	January 2023		
Due by:	16/3/2023		

Q1. Consider  $f(x|\theta) = \begin{cases} p, & x = 0 \\ (1-p)\frac{(\ln \theta)^x}{\theta x!}, & x = 1, 2, 3, \dots \\ 0, & \text{otherwise} \end{cases}$

Suppose parameters are  $p \in [0, 1]$  and  $\theta \geq 0$ . Then, for  $X_1, X_2, \dots, X_n$  iid with this, find a method of moments estimator for the parameter vector  $(p, \theta)$  based on the first two sample moments.

(15 marks)

Q2. Let  $X_1, X_2, \dots, X_n$  be a random sample from the probability density function:

$$f(x_i) = \begin{cases} 4\theta x_i^{4\theta-1}, & 0 < x_i < 1, \theta > 0 \\ 0, & \text{otherwise} \end{cases}$$

(a) Find the MLE of  $\theta$ .

(b) Find  $c$  such that  $c\hat{\theta}$  is an unbiased estimator of  $\theta$ , where  $\hat{\theta}$  is the MLE of  $\theta$ .

(15 marks)

Q3. Let  $X \sim POI(\mu)$ . Suppose  $\theta = e^{-\mu}$  and  $\tilde{\theta} = u(x) = \begin{cases} 1, & \text{for } x = 0 \\ 0, & \text{for } x = 1, 2, \dots \end{cases}$ .

Compare the MSEs of  $\hat{\theta}$  and  $\tilde{\theta}$  for estimating  $\theta$  when  $\mu = 5$

(20 marks)

Q4. Let  $X_1, X_2, \dots, X_n$  denote a random sample from the density function given by

$$f(x) = \begin{cases} \frac{4}{\theta} x^3 e^{-x^4/\theta}, & \theta > 0, x > 0, \\ 0, & \text{otherwise} \end{cases}$$

(a) Find the MLE of  $\theta$ .

(b) Find the CRLB of  $\theta$ .

(c) Find the UMVUE for  $\theta$ .

(15 marks)

**MEME15203 Statistical Inference**

Q5. Let  $X_1, X_2, \dots, X_n$  denote a random sample from an exponentially distributed population with mean  $\lambda = \frac{1}{\theta}$ . Let  $\Theta \sim \chi^2(2v)$ .

- (a) Find the Bayes estimator for  $\lambda = \frac{1}{\theta}$  under square error loss.
- (b) Show that it is a biased but consistent estimator for  $\lambda = \frac{1}{\theta}$ .

(20 marks)

Q6. Suppose  $X|\theta \sim U(\theta - \frac{1}{6}, \theta + \frac{5}{6})$  and that a prior distribution of  $\theta$  is  $N(\mu, 1)$ . Find the Bayes estimator of  $\theta$  under squared error loss.

(15 marks)