

Introduction to Statistics - Homework #10

Exercise 8.12 - Crawling babies, Part I

A study recorded data on 12 groups of infants born throughout the year to investigate whether crawling age (x) is associated with environmental temperature (y). For each group, researchers measured the average crawling age (in weeks) and the average temperature (in $^{\circ}\text{F}$) when the babies were six months old. Conduct a hypothesis test to determine if there is a **negative** correlation between temperature and crawling age. (The data come from a bivariate normal distribution.)

(1) State the null and alternative hypotheses. (Use one-sided test)

(2) Find the test statistic and its null distribution.

(3) Find the rejection region.

(4) Compute the correlation coefficient and the observed test statistic using the following summary statistics.

n	s_{xx}	s_{yy}	s_{xy}
12	34.0961	2762.25	-214.735

(5) Using the rejection region, complete the hypothesis test. State the conclusion in the context of the data.

Exercise 8.9 - Speed and height, revised

50 UCLA students were asked to fill out a survey where they were asked about their height, fastest speed they have ever driven, and gender. We are interested in investigating whether height (x) is associated with fastest driving speed (y). Conduct a hypothesis test to determine if there is a correlation between height and fastest speed. (Assume the data come from a bivariate normal distribution.)

(1) State the null and alternative hypotheses. (Use two-sided test)

(2) Find the test statistic and its null distribution.

(3) Compute the correlation coefficient and the observed test statistic using the following summary statistics.

n	s_{xx}	s_{yy}	s_{xy}
50	727.245	18792.32	399.78

(4) Compute the p-value and complete the hypothesis test.