

Optimising NMR Spectroscopy through Method and Software Development

Jonathan Yong

University of Oxford

Contents

Abstract	v
Acknowledgements	vi
Preface	vii
List of figures	xi
List of tables	xiv
List of code listings	xvi
1 NMR theory	1
1.1 Quantum mechanics	2
1.2 The rotating frame	5
1.3 Density operators	8
1.4 Pulse sequences	11
1.4.1 1D pulse-acquire	11
1.4.2 INEPT and product operators	15
1.4.3 2D NMR: general principles	19
1.4.4 The States HSQC experiment	23
1.4.5 The echo-antiecho HSQC: gradients and coherence selection	24
1.5 References	31
2 Pure shift NMR	34
2.1 Theoretical background	35
2.2 Pure shift in practice	39
2.2.1 Acquisition modes	39
2.2.2 Pure shift elements	41
2.2.3 PSYCHE in detail	43

2.3	PSYCHE with a variable number of saltires	47
2.4	Direct optimisation of PSYCHE waveform	50
2.4.1	Techniques for pure shift optimisations	51
2.4.2	Flip angle optimisation	55
2.4.3	Waveform parameterisation and optimisation	57
2.5	Time-reversal method	61
2.6	‘Discrete PSYCHE’	65
2.6.1	Speeding up dPSYCHE simulations	66
2.6.2	Optimisations and experimental evaluation	70
2.7	Ultrafast PSYCHE-iDOSY	76
2.8	References	83
3	POISE	92
3.1	Introduction	93
3.2	Technical overview	95
3.2.1	Routines	95
3.2.2	Optimisation settings	97
3.2.3	Optimisation algorithms	97
3.2.4	Implementation details	103
3.3	What POISE is not	105
3.4	Applications	107
3.4.1	Pulse width calibration	108
3.4.2	Ernst angle optimisation	113
3.4.3	Inversion–recovery	116
3.4.4	NOE mixing time	118
3.4.5	ASAP-HSQC excitation delay	121
3.4.6	Ultrafast NMR	124
3.4.7	HMBC low-pass J-filter	128
3.4.8	PSYCHE pure shift NMR	133
3.4.9	Water suppression	138
3.4.10	Diffusion NMR	143
3.5	POISE for ESR	151
3.6	References	153
4	NOAH	161
4.1	Introduction	163
4.2	Sensitivity analysis of NOAH supersequences	163
4.3	GENESIS: automated pulse programme creation	163

4.4	Discussion of individual modules	164
4.4.1	Sensitivity-enhanced HSQC	164
4.4.2	HSQC-TOCSY	164
4.4.3	HSQC-COSY	164
4.4.4	2DJ and PSYCHE	164
4.4.5	DQF-COSY	164
4.4.6	HMQC	164
4.4.7	HMBC	164
4.4.8	ADEQUATE	165
4.5	Solvent suppression in NOAH	165
4.6	NOAH with short recovery delays (???)	165
4.7	Parallel and generalised NOAH supersequences	165
4.8	References	165
A	Other work	165
A.1	NMR plotting in Python	165
A.2	Citation management	166
A.3	Group website and pulse programming tutorials	166
A.4	References	166

refsection:1

refsection:2

refsection:3

refsection:4

Chapter 4

NOAH

chpt : noah

This final—and long—chapter describes my work on *NOAH* (NMR by Ordered Acquisition using ^1H detection) *supersequences*, pulse sequences which record multiple 2D datasets (*‘modules’*) in the time required for one. This is an attractive NMR technique for several reasons: the time savings are clearly a key factor, but the flexibility of being able to combine almost any set of modules also makes NOAH supersequences applicable to a variety of contexts.

I begin by introducing the concepts underlying NOAH supersequences, as well as a general discussion of the time savings (and sensitivity per unit time) benefits thus realised. I then describe the GENESIS (GENeration of Supersequences In Silico) website, which allows users to generate Bruker pulse programmes for almost every imaginable NOAH supersequence. After this, my work on various aspects of the actual sequences themselves is described, with a special focus on newly developed and/or improved modules. Finally, the design of ‘parallel’ supersequences which use interleaved and/or time-shared modules is discussed.

This work was done in close collaboration with Ěriks Kupče (Bruker UK). However, all results and analysis shown in this chapter are mine, unless explicitly stated. The work in this chapter forms the subject of several publications:

- Yong, J. R. J.; Hansen, A. L.; Kupče, Ě.; Claridge, T. D. W. Increasing sensitivity and versatility in NMR supersequences with new HSQC-based modules. *J. Magn. Reson.* **2021**, 329, 107027, DOI: [10.1016/j.jmr.2021.107027](https://doi.org/10.1016/j.jmr.2021.107027)
- Kupče, Ě.; Yong, J. R. J.; Widmalm, G.; Claridge, T. D. W. Parallel NMR Supersequences: Ten Spectra in a Single Measurement. *JACS Au* **2021**, 1, 1892–1897, DOI: [10.1021/jacsau.1c00423](https://doi.org/10.1021/jacsau.1c00423)
- Yong, J. R. J.; Kupče, Ě.; Claridge, T. D. W. Modular Pulse Program Generation for NMR Supersequences. *Anal. Chem.* **2022**, 94, 2271–2278, DOI: [10.1021/acs.analchem.1c04](https://doi.org/10.1021/acs.analchem.1c04)

964

- Yong, J. R. J.; Kupče, Ě.; Claridge, T. D. W. Uniting Low- and High-Sensitivity Experiments through Generalised NMR Supersequences. 2022, manuscript in preparation

The material in the introductory sections also closely follow two reviews which I have contributed to:

- Kupče, Ě.; Frydman, L.; Webb, A. G.; Yong, J. R. J.; Claridge, T. D. W. Parallel nuclear magnetic resonance spectroscopy. *Nat. Rev. Methods Primers* 2021, 1, No. 27, DOI: [10.1038/s43586-021-00024-3](https://doi.org/10.1038/s43586-021-00024-3)
- Yong, J. R. J.; Kupče, Ě.; Claridge, T. D. W. In *Fast 2D solution-state NMR: concepts and applications*, Giraudeau, P., Dumez, J.-N., Eds., forthcoming, 2022

4.1 Introduction

sec:noah__introduction

The characterisation of small molecules and biomolecules by NMR spectroscopy relies on a suite of standard 2D NMR experiments, which seek to detect heteronuclear scalar couplings (e.g. HSQC and HMBC), homonuclear scalar couplings (e.g. COSY and TOCSY), or through-space interactions (e.g. NOESY and ROESY). Although 2D experiments provide far superior resolution and information content compared to 1D spectra, they also require substantially longer experiment durations, as the indirect dimension must be constructed through the acquisition of many t_1 increments. This problem is further exacerbated by the fact that structural elucidation or verification often necessitates the acquisition of several different 2D experiments.

The acceleration of 2D NMR has thus proven to be a popular area of research, encompassing techniques such as non-uniform sampling (NUS),^{7–10} fast pulsing (i.e. shortening of recovery delays),^{11–14} ultrafast NMR,^{5,15–19} Hadamard encoding,^{20,21} and spectral aliasing.^{22–24} All of these methods seek to directly speed up the acquisition of *individual* 2D spectra. In contrast, *multiple-FID techniques* such as time-shared NMR,^{25,26} multiple-receiver NMR,^{27–29} and—of course—NOAH supersequences,^{5,30,31} instead aim to collecting multiple 2D spectra in (roughly) the time needed to acquire one ‘conventional’ 2D spectrum. This essentially amounts to an increase in *efficiency*, rather than pure speed.

4.2 Sensitivity analysis of NOAH supersequences

sec:noah__snr

General discussion of time savings and sensitivity analysis.

Should be very similar to RSC chapter

4.3 GENESIS: automated pulse programme creation

sec:noah__genesis

I think it makes sense to start with GENESIS;³ that way everything else can be placed in context.

It’s true that this was a full paper, but it included a lot of stuff which wasn’t about the website itself—these will go into later sections.

4.4 Discussion of individual modules

4.4.1 Sensitivity-enhanced HSQC

^{13}C seHSQC

^{15}N seHSQC

4.4.2 HSQC-TOCSY

HSQC + DIPSI + HSQC combos

Extension to HSQC-TOCSY

Cite ASAP work (Luy)

4.4.3 HSQC-COSY

Comparison of several versions of HSQC-COSY (JACS Au SI)

4.4.4 2DJ and PSYCHE

cnst37 scaling

SAPPHIRE

4.4.5 DQF-COSY

Once upon a time, I did some comparisons of States vs EA DQF-COSY. I think this section could probably be left out, though. The differences were *extremely* minor.

4.4.6 HMQC

Suppression of wing artefacts (GENESIS paper)

4.4.7 HMBC

Suppression of $^1J_{\text{CH}}$ artefacts (GENESIS paper)

Investigation of gradient schemes (no difference was really observed, but that's fine)

Also ^{15}N HMBC

4.4.8 ADEQUATE

Recent stuff.

4.5 Solvent suppression in NOAH

GENESIS paper.

4.6 NOAH with short recovery delays (???)

I did like one bit of work on this a while ago. The idea is basically that the spectra are fine but SNR unsurprisingly suffers (in fact SNR/t also decreases).

It may be mildly interesting to compare this against the NORD experiments if I can get their pulse programmes to work... I don't expect the NORD sensitivity to be amazing but it should in fact be better than just doing NOAH with short d1.

It may equally be viable to just leave this entire section out of the thesis as I don't think it adds much...

4.7 Parallel and generalised NOAH supersequences

Blah.

4.8 References

- (1) Yong, J. R. J.; Hansen, A. L.; Kupče, Ě.; Claridge, T. D. W. Increasing sensitivity and versatility in NMR supersequences with new HSQC-based modules. *J. Magn. Reson.* **2021**, 329, 107027, DOI: [10.1016/j.jmr.2021.107027](https://doi.org/10.1016/j.jmr.2021.107027).
- (2) Kupče, Ě.; Yong, J. R. J.; Widmalm, G.; Claridge, T. D. W. Parallel NMR Supersequences: Ten Spectra in a Single Measurement. *JACS Au* **2021**, 1, 1892–1897, DOI: [10.1021/jacsau.1c00423](https://doi.org/10.1021/jacsau.1c00423).
- (3) Yong, J. R. J.; Kupče, Ě.; Claridge, T. D. W. Modular Pulse Program Generation for NMR Supersequences. *Anal. Chem.* **2022**, 94, 2271–2278, DOI: [10.1021/acs.analchem.1c04964](https://doi.org/10.1021/acs.analchem.1c04964).
- (4) Yong, J. R. J.; Kupče, Ě.; Claridge, T. D. W. Uniting Low- and High-Sensitivity Experiments through Generalised NMR Supersequences. **2022**, manuscript in preparation.

- Kupce2021NRMP (5) Kupče, Ě.; Frydman, L.; Webb, A. G.; Yong, J. R. J.; Claridge, T. D. W. Parallel nuclear magnetic resonance spectroscopy. *Nat. Rev. Methods Primers* **2021**, 1, No. 27, DOI: [10.1038/s43586-021-00024-3](https://doi.org/10.1038/s43586-021-00024-3).
- Yong2022RSCBook (6) Yong, J. R. J.; Kupče, Ě.; Claridge, T. D. W. In *Fast 2D solution-state NMR: concepts and applications*, Giraudeau, P., Dumez, J.-N., Eds., forthcoming, 2022.
- Barna1987JMR (7) Barna, J. C. J.; Laue, E. D.; Mayger, M. R.; Skilling, J.; Worrall, S. J. P. Exponential sampling, an alternative method for sampling in two-dimensional NMR experiments. *J. Magn. Reson.* **1987**, 73, 69–77, DOI: [10.1016/0022-2364\(87\)90225-3](https://doi.org/10.1016/0022-2364(87)90225-3).
- Kazimierczuk2010PNMRS (8) Kazimierczuk, K.; Stanek, J.; Zawadzka-Kazimierczuk, A.; Koźmiński, W. Random sampling in multidimensional NMR spectroscopy. *Prog. Nucl. Magn. Reson. Spectrosc.* **2010**, 57, 420–434, DOI: [10.1016/j.pnmrs.2010.07.002](https://doi.org/10.1016/j.pnmrs.2010.07.002).
- Mobli2014PNMRS (9) Mobli, M.; Hoch, J. C. Nonuniform sampling and non-Fourier signal processing methods in multidimensional NMR. *Prog. Nucl. Magn. Reson. Spectrosc.* **2014**, 83, 21–41, DOI: [10.1016/j.pnmrs.2014.09.002](https://doi.org/10.1016/j.pnmrs.2014.09.002).
- Kazimierczuk2015MRC (10) Kazimierczuk, K.; Orekhov, V. Non-uniform sampling: post-Fourier era of NMR data collection and processing. *Magn. Reson. Chem.* **2015**, 53, 921–926, DOI: [10.1002/mrc.4284](https://doi.org/10.1002/mrc.4284).
- Sunninghausen2014JACS (11) Schulze-Sunninghausen, D.; Becker, J.; Luy, B. Rapid Heteronuclear Single Quantum Correlation NMR Spectra at Natural Abundance. *J. Am. Chem. Soc.* **2014**, 136, 1242–1245, DOI: [10.1021/ja411588d](https://doi.org/10.1021/ja411588d).
- Schanda2006JACS (12) Schanda, P.; Van Melckebeke, H.; Brutscher, B. Speeding Up Three-Dimensional Protein NMR Experiments to a Few Minutes. *J. Am. Chem. Soc.* **2006**, 128, 9042–9043, DOI: [10.1021/ja062025p](https://doi.org/10.1021/ja062025p).
- Kupce2007MRC (13) Kupče, Ě.; Freeman, R. Fast multidimensional NMR by polarization sharing. *Magn. Reson. Chem.* **2007**, 45, 2–4, DOI: [10.1002/mrc.1931](https://doi.org/10.1002/mrc.1931).
- Schanda2009PNMRS (14) Schanda, P. Fast-pulsing longitudinal relaxation optimized techniques: Enriching the toolbox of fast biomolecular NMR spectroscopy. *Prog. Nucl. Magn. Reson. Spectrosc.* **2009**, 55, 238–265, DOI: [10.1016/j.pnmrs.2009.05.002](https://doi.org/10.1016/j.pnmrs.2009.05.002).
- Frydman2002PNASUSA (15) Frydman, L.; Scherf, T.; Lupulescu, A. The acquisition of multidimensional NMR spectra within a single scan. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, 15858–15862, DOI: [10.1073/pnas.252644399](https://doi.org/10.1073/pnas.252644399).
- Pelupessy2003JACS (16) Pelupessy, P. Adiabatic Single Scan Two-Dimensional NMR Spectroscopy. *J. Am. Chem. Soc.* **2003**, 125, 12345–12350, DOI: [10.1021/ja034958g](https://doi.org/10.1021/ja034958g).
- Frydman2003JACS (17) Frydman, L.; Lupulescu, A.; Scherf, T. Principles and Features of Single-Scan Two-Dimensional NMR Spectroscopy. *J. Am. Chem. Soc.* **2003**, 125, 9204–9217, DOI: [10.1021/ja030055b](https://doi.org/10.1021/ja030055b).

- Tal2010PNMRS (18) Tal, A.; Frydman, L. Single-scan multidimensional magnetic resonance. *Prog. Nucl. Magn. Reson. Spectrosc.* **2010**, *57*, 241–292, DOI: [10.1016/j.pnmrs.2010.04.001](https://doi.org/10.1016/j.pnmrs.2010.04.001).
- Gouilleux2018ARNMRS (19) Gouilleux, B.; Rouger, L.; Giraudeau, P. Ultrafast 2D NMR: Methods and Applications. *Annu. Rep. NMR Spectrosc.* **2018**, 75–144, DOI: [10.1016/bs.arnmr.2017.08.003](https://doi.org/10.1016/bs.arnmr.2017.08.003).
- Kupce2003JMR (20) Kupče, Ě.; Freeman, R. Two-dimensional Hadamard spectroscopy. *J. Magn. Reson.* **2003**, *162*, 300–310, DOI: [10.1016/s1090-7807\(02\)00196-9](https://doi.org/10.1016/s1090-7807(02)00196-9).
- Kupce2003PNMRS (21) Kupče, E.; Nishida, T.; Freeman, R. Hadamard NMR spectroscopy. *Prog. Nucl. Magn. Reson. Spectrosc.* **2003**, *42*, 95–122, DOI: [10.1016/s0079-6565\(03\)00022-0](https://doi.org/10.1016/s0079-6565(03)00022-0).
- Jeannerat2000MRC (22) Jeannerat, D. High resolution in the indirectly detected dimension exploiting the processing of folded spectra. *Magn. Reson. Chem.* **2000**, *38*, 415–422, DOI: [10.1002/1097-458x\(200006\)38:6<415::aid-mrc665>3.0.co;2-u](https://doi.org/10.1002/1097-458x(200006)38:6<415::aid-mrc665>3.0.co;2-u).
- Bermel2009JACS (23) Bermel, W.; Bertini, I.; Felli, I. C.; Pierattelli, R. Speeding Up ¹³C Direct Detection Biomolecular NMR Spectroscopy. *J. Am. Chem. Soc.* **2009**, *131*, 15339–15345, DOI: [10.1021/ja9058525](https://doi.org/10.1021/ja9058525).
- Njock2010C (24) Njock, G. B. B.; Pegnyemb, D. E.; Bartholomeusz, T. A.; Christen, P.; Vitorge, B.; Nuzillard, J.-M.; Shivapurkar, R.; Foroozandeh, M.; Jeannerat, D. Spectral Aliasing: A Super Zoom for 2D-NMR Spectra. Principles and Applications. *Chimia* **2010**, *64*, 235, DOI: [10.2533/chimia.2010.235](https://doi.org/10.2533/chimia.2010.235).
- Nolis2007ACIE (25) Nolis, P.; Pérez-Trujillo, M.; Parella, T. Multiple FID Acquisition of Complementary HMBC Data. *Angew. Chem. Int. Ed.* **2007**, *46*, 7495–7497, DOI: [10.1002/anie.200702258](https://doi.org/10.1002/anie.200702258).
- Parella2010CMR (26) Parella, T.; Nolis, P. Time-shared NMR experiments. *Concepts Magn. Reson.* **2010**, *36A*, 1–23, DOI: [10.1002/cmr.a.20150](https://doi.org/10.1002/cmr.a.20150).
- Kupce2006JACS (27) Kupče, Ě.; Freeman, R.; John, B. K. Parallel Acquisition of Two-Dimensional NMR Spectra of Several Nuclear Species. *J. Am. Chem. Soc.* **2006**, *128*, 9606–9607, DOI: [10.1021/ja0634876](https://doi.org/10.1021/ja0634876).
- Kupce2008JACS (28) Kupče, Ě.; Freeman, R. Molecular Structure from a Single NMR Experiment. *J. Am. Chem. Soc.* **2008**, *130*, 10788–10792, DOI: [10.1021/ja8036492](https://doi.org/10.1021/ja8036492).
- Kovacs2016MRC (29) Kovacs, H.; Kupče, Ě. Parallel NMR spectroscopy with simultaneous detection of ¹H and ¹⁹F nuclei. *Magn. Reson. Chem.* **2016**, *54*, 544–560, DOI: [10.1002/mrc.4428](https://doi.org/10.1002/mrc.4428).
- Kupce2017ACIE (30) Kupče, Ě.; Claridge, T. D. W. NOAH: NMR Supersequences for Small Molecule Analysis and Structure Elucidation. *Angew. Chem. Int. Ed.* **2017**, *56*, 11779–11783, DOI: [10.1002/anie.201705506](https://doi.org/10.1002/anie.201705506).
- Kupce2021PNMRS (31) Kupče, Ě.; Mote, K. R.; Webb, A.; Madhu, P. K.; Claridge, T. D. W. Multiplexing experiments in NMR and multi-nuclear MRI. *Prog. Nucl. Magn. Reson. Spectrosc.* **2021**, *124–125*, 1–56, DOI: [10.1016/j.pnmrs.2021.03.001](https://doi.org/10.1016/j.pnmrs.2021.03.001).

refsection:5