


MATH7501: Mathematics for Data Science I

Unit 10: Partial Derivatives and Gradient Descent

10.1 Partial Derivatives and Tangent Planes

Consider a surface $f(x, y)$ with a particular point $P = (x_0, y_0)$.



10.1 Partial Derivatives and Tangent Planes

Consider a surface $f(x, y)$ with a particular point $P = (x_0, y_0)$. Then the slope of f at P in the x direction is $\frac{\partial f}{\partial x}$, evaluated at P ; this partial derivative is the derivative of the single-variable function $f(x, y_0)$.

10.1 Partial Derivatives and Tangent Planes

Consider a surface $f(x, y)$ with a particular point $P = (x_0, y_0)$. Then the slope of f at P in the x direction is $\frac{\partial f}{\partial x}$, evaluated at P ; this partial derivative is the derivative of the single-variable function $f(x, y_0)$. We can find the slope of f in the y -direction similarly.



10.1 Partial Derivatives and Tangent Planes

Consider a surface $f(x, y)$ with a particular point $P = (x_0, y_0)$. Then the slope of f at P in the x direction is $\frac{\partial f}{\partial x}$, evaluated at P ; this partial derivative is the derivative of the single-variable function $f(x, y_0)$. We can find the slope of f in the y -direction similarly.

For example, if $f(x, y) = 1 - x^2 - y^2$, and $(x_0, y_0) = (1, -1)$, then $f(x, -1) = 1 - x^2$, so $\frac{\partial f}{\partial x}(P) = -2$.

$$\frac{\partial f}{\partial x} = -2x \quad |_{x=1}$$

10.1 Partial Derivatives and Tangent Planes

Consider a surface $f(x, y)$ with a particular point $P = (x_0, y_0)$. Then the slope of f at P in the x direction is $\frac{\partial f}{\partial x}$, evaluated at P ; this partial derivative is the derivative of the single-variable function $f(x, y_0)$. We can find the slope of f in the y -direction similarly.

For example, if $f(x, y) = 1 - x^2 - y^2$, and $(x_0, y_0) = (1, -1)$, then $f(x, -1) = -x^2$, so $\frac{\partial f}{\partial x}(P) = -2$. Also, $f(1, y) = 1 - y^2$, so $\frac{\partial f}{\partial y}(P) = 2$.

$$\frac{\partial f}{\partial y} = -2y \quad | \quad y = -1$$




Figure 64: The surface $z = f(x, y) = 1 - x^2 - y^2$ and the intersection of the plane $y = -1$ with the surface.




Figure 65: The surface $z = f(x, y) = 1 - x^2 - y^2$ and the intersection of the plane $x = 1$ with the surface.

10.1 Partial Derivatives and Tangent Planes

Question

The volume of a solid with side lengths x, y, z is given by $V(x, y, z) = xyz$. Calculate the partial derivatives of V in all three variables.

10.1 Partial Derivatives and Tangent Planes

Question

The volume of a solid with side lengths x, y, z is given by $V(x, y, z) = xyz$. Calculate the partial derivatives of V in all three variables.

Solution

If y, z are held and x changes, then we can see that

$$\frac{\partial V}{\partial x} = \underline{yz}.$$

$$\frac{\partial (xyz)}{\partial x} = yz \frac{\partial (x)}{\partial x} = yz$$

Similarly, $\frac{\partial V}{\partial y} = \underline{xz}$, $\frac{\partial V}{\partial z} = \underline{xy}$.

10.1 Partial Derivatives and Tangent Planes

For a given function $f(x, y)$, the *second order* partial derivatives are found by differentiating f once with respect to one variable, and then differentiating again with respect to a second variable.

these two variables
do not need to be
the same.

10.1 Partial Derivatives and Tangent Planes

For a given function $f(x, y)$, the *second order* partial derivatives are found by differentiating f once with respect to one variable, and then differentiating again with respect to a second variable. We denote this with

$$\begin{aligned} f_{xx} &= \frac{\partial^2 f}{\partial x^2}, & f_{yy} &= \frac{\partial^2 f}{\partial y^2}, \\ f_{xy} &= \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right), & f_{yx} &= \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right). \end{aligned}$$

10.1 Partial Derivatives and Tangent Planes

For a given function $f(x, y)$, the *second order* partial derivatives are found by differentiating f once with respect to one variable, and then differentiating again with respect to a second variable. We denote this with

$$f_{xx} = \frac{\partial^2 f}{\partial x^2}, \quad f_{yy} = \frac{\partial^2 f}{\partial y^2},$$

$$f_{xy} = \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right), \quad f_{yx} = \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right).$$

$$\begin{aligned} \frac{\partial f}{\partial x} &= \sin(y) - y \sin(x), & \frac{\partial f}{\partial y} &= x \cos(y) + \cos(x) \\ \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) &= \cos(y) - \sin(x), & \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) &= \cos(y) - \sin(x) \end{aligned}$$

Question

Let $f(x, y) = x \sin(y) + y \cos(x)$. Calculate all of the second order partial derivatives, and show that $f_{xy} = f_{yx}$.

— Always true if f is 'sufficiently well-behaved'.

10.1 Partial Derivatives and Tangent Planes

Recall that through linear approximations, we have

$$\Delta f \approx \frac{\partial f}{\partial x} \Delta x + \frac{\partial f}{\partial y} \Delta y.$$

10.1 Partial Derivatives and Tangent Planes

Recall that through linear approximations, we have

$$\Delta f \approx \frac{\partial f}{\partial x} \Delta x + \frac{\partial f}{\partial y} \Delta y.$$


Therefore, if x and y are both changing with time t , and we know how much x and y are changing by, we can approximate how much f is changing over time.

10.1 Partial Derivatives and Tangent Planes

Recall that through linear approximations, we have

top equation

$$\Delta f \approx \frac{\partial f}{\partial x} \Delta x + \frac{\partial f}{\partial y} \Delta y.$$



Therefore, if x and y are both changing with time t , and we know how much x and y are changing by, we can approximate how much f is changing over time.

Theorem (Multi-Variable Chain Rule)

If x, y are differentiable functions of t , and f is a differentiable function of x, y , then

limit

$$\frac{df(x(t), y(t))}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

$\frac{d}{dt} f(x(t), y(t)) \approx \frac{\partial f}{\partial x} x'(t) + \frac{\partial f}{\partial y} y'(t)$
+ (don't count for derivatives).

10.1 Partial Derivatives and Tangent Planes

Question

Suppose the radius of a cylinder decreases at a rate of $r'(t) = -2\text{cm/s}$, and the height is kept constant at $h(t) = 2\text{cm}$. How fast is the volume decreasing when $r(t) = 1\text{cm}$?

10.1 Partial Derivatives and Tangent Planes

Question

Suppose the radius of a cylinder decreases at a rate of $r'(t) = -2\text{cm/s}$, and the height is kept constant at $h(t) = 2\text{cm}$. How fast is the volume decreasing when $r(t) = 1\text{cm}$?

Solution

We have $V = \pi r^2 h$ so

$$V' = \pi 2r h r' + \pi r^2 h' = 2\pi r h r' = -8\pi \text{cm}^3/\text{s}.$$

$$\frac{dv}{dt} = \frac{\partial v}{\partial r} \frac{dr}{dt} + \frac{\partial v}{\partial h} \frac{dh}{dt}$$

10.1 Partial Derivatives and Tangent Planes

Question

Suppose the radius of a cylinder decreases at a rate of $r'(t) = -2\text{cm/s}$, and the height is kept constant at $h(t) = 2\text{cm}$. How fast is the volume decreasing when $r(t) = 1\text{cm}$?

Solution

We have $V = \pi r^2 h$ so

$$V' = \pi 2rhr' + \pi r^2 h' = 2\pi rhr' = -8\pi \text{cm}^3/\text{s}.$$

Question

Suppose now that the height is decreasing at 1cm/s . What is the rate of change of volume?

10.1 Partial Derivatives and Tangent Planes

Question

Suppose the radius of a cylinder decreases at a rate of $r'(t) = -2\text{cm/s}$, and the height is kept constant at $h(t) = 2\text{cm}$. How fast is the volume decreasing when $r(t) = 1\text{cm}$?

Solution

We have $V = \pi r^2 h$ so

$$V' = \pi 2rhr' + \pi r^2 h' = 2\pi rhr' = -8\pi \text{cm}^3/\text{s}.$$

Question

Suppose now that the height is decreasing at 1cm/s . What is the rate of change of volume?

Solution

$$\text{We have } V = \pi r^2 h \text{ so } V' = \pi 2rhr' + \pi r^2 h' = -9\pi \text{cm}^3/\text{s}.$$

10.1 Partial Derivatives and Tangent Planes

Of course, the chain rule extends to higher dimensions as well.

10.1 Partial Derivatives and Tangent Planes

Of course, the chain rule extends to higher dimensions as well.

Question

Suppose a, b, c are the dimensions of a box which are changing over time. From the rate of change of the volume.

10.1 Partial Derivatives and Tangent Planes

Of course, the chain rule extends to higher dimensions as well.

Question

Suppose a, b, c are the dimensions of a box which are changing over time. From the rate of change of the volume.

Solution

The volume is $V = abc$, so

$$\begin{aligned}\frac{dV}{dt} &= V_a \frac{da}{dt} + V_b \frac{db}{dt} + V_c \frac{dc}{dt} \\ &= bc \frac{da}{dt} + ac \frac{db}{dt} + ab \frac{dc}{dt}.\end{aligned}$$

$$\frac{\partial V}{\partial a} = bc, \quad \frac{\partial V}{\partial b} = ac, \quad \frac{\partial V}{\partial c} = ab$$

10.2 Gradient

The partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ tell us the slope of f in the x, y directions, but they can also be used to describe the slope in *any* direction.

Dot product: $(a_1, a_2) \cdot (b_1, b_2)$

$$= a_1 b_1 + a_2 b_2$$

$$= \sqrt{a_1^2 + a_2^2} \times \sqrt{b_1^2 + b_2^2} \times \cos \theta$$

θ is the angle between the \vec{a} and \vec{b} .

10.2 Gradient

The partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ tell us the slope of f in the x, y directions, but they can also be used to describe the slope in *any direction*. Take a point $P \in \mathbb{R}^2$, and say we travel through P with $(x(t), y(t)) = P + \mathbf{u}t$, where $\mathbf{u} = (u_1, u_2)$ is a unit-vector.

$x(t)$ is a scalar, $y(t)$ is a scalar,

$(x(t), y(t))$ is a vector.

10.2 Gradient

The partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ tell us the slope of f in the x, y directions, but they can also be used to describe the slope in *any direction*. Take a point $P \in \mathbb{R}^2$, and say we travel through P with $(x(t), y(t)) = P + \mathbf{u}t$, where $\mathbf{u} = (u_1, u_2)$ is a unit-vector. Then, using the linear approximation,

$$\begin{aligned} f(x(t), y(t)) &\approx f(x(0), y(0)) + \frac{\partial f}{\partial x} u_1 t + \frac{\partial f}{\partial y} u_2 t \\ &= f(P) + t \left(\frac{\partial f}{\partial x}(P), \frac{\partial f}{\partial y}(P) \right) \cdot \mathbf{u}. \end{aligned}$$

$$\begin{aligned} & \underbrace{(a, b) \cdot (c, d)}_{= ac + bd - \text{scalar}} \end{aligned}$$

10.2 Gradient

The partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ tell us the slope of f in the x, y directions, but they can also be used to describe the slope in *any direction*. Take a point $P \in \mathbb{R}^2$, and say we travel through P with $(x(t), y(t)) = P + \mathbf{u}t$, where $\mathbf{u} = (u_1, u_2)$ is a unit-vector. Then, using the linear approximation,

$$\begin{aligned} f(x(t), y(t)) &\approx f(x(0), y(0)) + \frac{\partial f}{\partial x} u_1 t + \frac{\partial f}{\partial y} u_2 t \\ &= f(P) + t \left(\frac{\partial f}{\partial x}(P), \frac{\partial f}{\partial y}(P) \right) \cdot \mathbf{u}. \end{aligned}$$

The expression $\left(\frac{\partial f}{\partial x}(P), \frac{\partial f}{\partial y}(P) \right) \cdot \mathbf{u}$ is therefore referred to as the slope/directional derivative of f at P in the direction of \mathbf{u} .

$$\frac{f(t) - f(0)}{t} \approx \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) \cdot \mathbf{u}$$

10.2 Gradient

The partial derivatives $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ tell us the slope of f in the x, y directions, but they can also be used to describe the slope in *any direction*. Take a point $P \in \mathbb{R}^2$, and say we travel through P with $(x(t), y(t)) = P + \mathbf{u}t$, where $\mathbf{u} = (u_1, u_2)$ is a unit-vector. Then, using the linear approximation,


$$\begin{aligned} f(x(t), y(t)) &\approx f(x(0), y(0)) + \frac{\partial f}{\partial x} u_1 t + \frac{\partial f}{\partial y} u_2 t \\ &= f(P) + t \left(\frac{\partial f}{\partial x}(P), \frac{\partial f}{\partial y}(P) \right) \cdot \mathbf{u}. \end{aligned}$$

The expression $\left(\frac{\partial f}{\partial x}(P), \frac{\partial f}{\partial y}(P) \right) \cdot \mathbf{u}$ is therefore referred to as the *slope/directional derivative of f at P in the direction of \mathbf{u}* . If $\mathbf{u} \neq 0$ is not a unit vector, then the directional derivative is simply $\left(\frac{\partial f}{\partial x}(P), \frac{\partial f}{\partial y}(P) \right) \cdot \frac{\mathbf{u}}{\|\mathbf{u}\|}$.

10.2 Gradient

Question

Find the directional derivative of $f(x, y) = 4 - x^2 - 4y^2$ at the point $(1, 1)$ in the $(1, 1)$ direction.




10.2 Gradient

Question

Find the directional derivative of $f(x, y) = 4 - x^2 - 4y^2$ at the point $(1, 1)$ in the $(1, 1)$ direction.

Solution

The corresponding unit vector is $\mathbf{u} = \frac{1}{\sqrt{2}}\mathbf{i} + \frac{1}{\sqrt{2}}\mathbf{j}$.


$$\begin{aligned}\hat{i} &= (1, 0) \\ \hat{j} &= (0, 1).\end{aligned}$$

10.2 Gradient

Question

Find the directional derivative of $f(x, y) = 4 - \underline{x^2} - \underline{4y^2}$ at the point $(1, 1)$ in the $(1, 1)$ direction.

Solution

The corresponding unit vector is $\underline{\mathbf{u}} = \frac{1}{\sqrt{2}}\mathbf{i} + \frac{1}{\sqrt{2}}\mathbf{j}$. We compute that

$$\underline{\frac{\partial f}{\partial x}} = -2x \text{ and } \underline{\frac{\partial f}{\partial y}} = -8y.$$

10.2 Gradient

Question

Find the directional derivative of $f(x, y) = 4 - x^2 - 4y^2$ at the point $(1, 1)$ in the $(1, 1)$ direction.

Solution

The corresponding unit vector is $\mathbf{u} = \frac{1}{\sqrt{2}}\mathbf{i} + \frac{1}{\sqrt{2}}\mathbf{j}$. We compute that $\frac{\partial f}{\partial x} = -2x$ and $\frac{\partial f}{\partial y} = -8y$. Therefore the slope at $(1, 1)$ in the \mathbf{u} direction is

$$(-2, -8) \cdot \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right) = -\frac{10}{\sqrt{2}} = -5\sqrt{2}.$$

10.2 Gradient

Question

If $f(x, y) = x^2 - 3y^2 + 6y$, find the slope at $(1, 0)$ in the direction of $\mathbf{i} - 4\mathbf{j}$.

$i = (1, 0)$, $j = (0, 1)$.
('x' direction 'y' direction.

10.2 Gradient

Question

If $f(x, y) = x^2 - 3y^2 + 6y$, find the slope at $(1, 0)$ in the direction of $\mathbf{i} - 4\mathbf{j}$.

Solution

The norm of $\mathbf{u} = \mathbf{i} - 4\mathbf{j}$ is $\|\mathbf{u}\| = \sqrt{17}$.

10.2 Gradient

Question

If $f(x, y) = x^2 - 3y^2 + 6y$, find the slope at $(1, 0)$ in the direction of $\mathbf{i} - 4\mathbf{j}$.

Solution

The norm of $\mathbf{u} = \mathbf{i} - 4\mathbf{j}$ is $\|\mathbf{u}\| = \sqrt{17}$. Also, note that $\frac{\partial f}{\partial x} = 2x$ and $\frac{\partial f}{\partial y} = 6 - 6y$.

$\frac{\mathbf{u}}{\sqrt{17}}$ is a unit vector

10.2 Gradient

Question

If $f(x, y) = x^2 - 3y^2 + 6y$, find the slope at $(1, 0)$ in the direction of $\mathbf{i} - 4\mathbf{j}$.

Solution

The norm of $\mathbf{u} = \mathbf{i} - 4\mathbf{j}$ is $\|\mathbf{u}\| = \sqrt{17}$. Also, note that $\frac{\partial f}{\partial x} = 2x$ and $\frac{\partial f}{\partial y} = 6 - 6y$. Therefore

$$f_{\mathbf{u}}(1, 0) = (2, 6) \cdot \frac{(1, -4)}{\sqrt{17}} = -\frac{22}{\sqrt{17}}.$$

$$\left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right)$$

10.2 Gradient

It is useful to give the vector (f_x, f_y) a name, since it comes up a lot when computing directional derivatives.

10.2 Gradient

It is useful to give the vector (f_x, f_y) a name, since it comes up a lot when computing directional derivatives.

Definition (Gradient Vector)

We define $\nabla f = (f_x, f_y)$ to be the *gradient* of f , (or $\nabla f = (f_x, f_y, f_z)$ for functions of three variables). Then the derivative of f in the direction of \mathbf{u} is $\nabla f \cdot \frac{\mathbf{u}}{\|\mathbf{u}\|}$.

10.2 Gradient

It is useful to give the vector (f_x, f_y) a name, since it comes up a lot when computing directional derivatives.

Definition (Gradient Vector)

We define $\nabla f = (f_x, f_y)$ to be the *gradient* of f , (or $\nabla f = (f_x, f_y, f_z)$ for functions of three variables). Then the derivative of f in the direction of \mathbf{u} is $\nabla f \cdot \frac{\mathbf{u}}{\|\mathbf{u}\|}$

(Question)

Find the gradient of $f(x, y) = x^2 - 3(y - 1)^2 + 3$.

10.2 Gradient

It is useful to give the vector (f_x, f_y) a name, since it comes up a lot when computing directional derivatives.

Definition (Gradient Vector)

We define $\nabla f = (f_x, f_y)$ to be the *gradient* of f , (or $\nabla f = (f_x, f_y, f_z)$ for functions of three variables). Then the derivative of f in the direction of \mathbf{u} is $\nabla f \cdot \frac{\mathbf{u}}{\|\mathbf{u}\|}$

(Question)

Find the gradient of $f(x, y) = x^2 - 3(y - 1)^2 + 3$.

Solution

$$\nabla f = (2x, -6(y - 1)) = 2x\mathbf{i} - 6(y - 1)\mathbf{j}.$$

10.2 Gradient

Question

Find the directional derivative of $g(x, y) = e^{x^2} \cos(y)$ at $(1, \pi)$ in the direction of $-3\mathbf{i} + 4\mathbf{j}$.

10.2 Gradient

Question

Find the directional derivative of $g(x, y) = e^{x^2} \cos(y)$ at $(1, \pi)$ in the direction of $-3\mathbf{i} + 4\mathbf{j}$.

Solution

The norm of $\mathbf{u} = -3\mathbf{i} + 4\mathbf{j}$ is $\|\mathbf{u}\| = 5$, and

$$\nabla g = 2xe^{x^2} \cos(y)\mathbf{i} - \sin(y)e^{x^2}\mathbf{j}.$$

$$\frac{\partial g}{\partial x} \quad \left(3^2 + 4^2 = 5^2 \right) \quad \frac{\partial g}{\partial y}$$

10.2 Gradient

Question

Find the directional derivative of $g(x, y) = e^{x^2} \cos(y)$ at $(1, \pi)$ in the direction of $-3\mathbf{i} + 4\mathbf{j}$.


Solution

The norm of $\mathbf{u} = -3\mathbf{i} + 4\mathbf{j}$ is $\|\mathbf{u}\| = 5$, and $\nabla g = 2xe^{x^2} \cos(y)\mathbf{i} - \sin(y)e^{x^2}\mathbf{j}$. Therefore

$$\begin{aligned} g_{\mathbf{u}} &= \nabla g(1, \pi) \cdot \frac{\mathbf{u}}{\|\mathbf{u}\|} \\ &= (2e \cos(\pi), -\sin(\pi)e) \cdot \left(\frac{-3}{5}, \frac{4}{5}\right) = \frac{6e}{5}. \end{aligned}$$

10.2 Gradient


In general, the gradient $\nabla f(a, b)$ points in the direction perpendicular to the contour (line of constant f) passing through (a, b) .



10.2 Gradient

In general, the gradient $\nabla f(a, b)$ points in the direction perpendicular to the contour (line of constant f) passing through (a, b) . In fact, $\nabla f(a, b)$ points in the direction of the shaprest increase of f .

$f(x, y)$
 $= xy$



contours
in blue.
gradient
in purple

<https://www.khanacademy.org/math/multivariable-calculus/multivariable-derivatives/partial-derivative-and-gradient-articles/a/the-gradient>

10.2 Gradient

Question

Consider the function $f(x, y) = \underline{mx} + \underline{ny} + \underline{c}$ for some $m, n, c \in \mathbb{R}$ and $n \neq 0$. Find the gradient of f and compare to the contours.

10.2 Gradient

Question

Consider the function $f(x, y) = mx + ny + c$ for some $m, n, c \in \mathbb{R}$ and $n \neq 0$. Find the gradient of f and compare to the contours.

Solution

The contours are $y = -\frac{mx}{n} + C$. The gradient vector is $\nabla f = (m, n)$, which points in the direction perpendicular to the contours.

$$\nabla f \cdot \left(1, -\frac{m}{n}\right) = (m, n) \cdot \left(1, -\frac{m}{n}\right) = 0.$$

10.2 Gradient

Question

Consider the function $f(x, y) = mx + ny + c$ for some $m, n, c \in \mathbb{R}$ and $n \neq 0$. Find the gradient of f and compare to the contours.

Solution

The contours are $y = -\frac{mx}{n} + C$. The gradient vector is $\nabla f = (m, n)$, which points in the direction perpendicular to the contours.

Question

Consider the function $f(x, y) = x^2 + y^2$. Describe the gradient and the contours.

10.2 Gradient

Question

Consider the function $f(x, y) = mx + ny + c$ for some $m, n, c \in \mathbb{R}$ and $n \neq 0$. Find the gradient of f and compare to the contours.

Solution


The contours are $y = -\frac{mx}{n} + C$. The gradient vector is $\nabla f = (m, n)$, which points in the direction perpendicular to the contours.

Question

Consider the function $f(x, y) = x^2 + y^2$. Describe the gradient and the contours.

Solution

The contours are the concentric circles centered at the origin.



10.2 Gradient

Question

Consider the function $f(x, y) = mx + ny + c$ for some $m, n, c \in \mathbb{R}$ and $n \neq 0$. Find the gradient of f and compare to the contours.

Solution

The contours are $y = -\frac{mx}{n} + C$. The gradient vector is $\nabla f = (m, n)$, which points in the direction perpendicular to the contours.

Question

Consider the function $f(x, y) = x^2 + y^2$. Describe the gradient and the contours.


Solution

The contours are the concentric circles centered at the origin. The gradient is $(2x, 2y)$, which is radial, always points perpendicular to the circles.

10.2 Gradient

Question

The temperature of a metal plate if $T(x, y) = 20 - 4x^2 - y^2$. Find the gradient at the point $(2, -3)$. Hence describe the direction of maximal increase in temperature, and the direction of constant temperature.



10.2 Gradient

Question

The temperature of a metal plate is $T(x, y) = 20 - 4x^2 - y^2$. Find the gradient at the point $(2, -3)$. Hence describe the direction of maximal increase in temperature, and the direction of constant temperature.

Solution

The gradient is $\nabla T = (-8x, -2y)$, which is $(-16, 6)$ at the specified point.

10.2 Gradient

Question

The temperature of a metal plate is $T(x, y) = 20 - 4x^2 - y^2$. Find the gradient at the point $(2, -3)$. Hence describe the direction of maximal increase in temperature, and the direction of constant temperature.

Solution

The gradient is $\nabla T = (-8x, -2y)$, which is $(-16, 6)$ at the specified point. To find the angle this makes with the x-axis, write $-16 = r \cos(\theta)$ and $6 = r \sin(\theta)$, so that $r = 2\sqrt{73}$ and $\theta = \pi - \arctan\left(\frac{6}{16}\right)$.

The direction of constant temperature is perpendicular to ∇T .

10.2 Gradient

Question

The temperature of a metal plate is $T(x, y) = 20 - 4x^2 - y^2$. Find the gradient at the point $(2, -3)$. Hence describe the direction of maximal increase in temperature, and the direction of constant temperature.

Solution

The gradient is $\nabla T = (-8x, -2y)$, which is $(-16, 6)$ at the specified point. To find the angle this makes with the x-axis, write $-16 = r \cos(\theta)$ and $6 = r \sin(\theta)$, so that $r = 2\sqrt{73}$ and $\theta = \pi - \arctan(\frac{6}{16})$. The direction of constant temperature is then $\frac{\pi}{2} - \arctan(\frac{6}{16})$.

10.2 Gradient

Question

A team is mapping the ocean floor. They produce the model


$$D(x, y) = 1700 - 30x^2 - 50 \sin\left(\frac{\pi y}{2}\right) \text{ in } m, \text{ with}$$

$$(x, y) \in [-2, 2] \times [-2, 2] \text{ in } km.$$

10.2 Gradient

Question

A team is mapping the ocean floor. They produce the model $D(x, y) = 1700 - 30x^2 - 50 \sin(\frac{\pi y}{2})$ in m , with $(x, y) \in [-2, 2] \times [-2, 2]$ in km . There is a leaking oil well at $(1, 0.5)$; find the depth at this location, as well as the direction and magnitude of steepest increase here.



10.2 Gradient

Question

A team is mapping the ocean floor. They produce the model $D(x, y) = 1700 - 30x^2 - 50 \sin(\frac{\pi y}{2})$ in m , with $(x, y) \in [-2, 2] \times [-2, 2]$ in km . There is a leaking oil well at $(1, 0.5)$; find the depth at this location, as well as the direction and magnitude of steepest increase here.

Solution

The depth at $(1, 0.5)$ is

$$D(1, 0.5) = 250 - 30 - 50 \sin(\frac{\pi}{4}) \approx 184.6m.$$

$$1700, x=1, y=0.5 \text{ km}$$

10.2 Gradient

Question

A team is mapping the ocean floor. They produce the model $D(x, y) = 1700 - 30x^2 - 50 \sin\left(\frac{\pi y}{2}\right)$ in m , with $(x, y) \in [-2, 2] \times [-2, 2]$ in km . There is a leaking oil well at $(1, 0.5)$; find the depth at this location, as well as the direction and magnitude of steepest increase here.

Solution

The depth at $(1, 0.5)$ is

$D(1, 0.5) = 250 - 30 - 50 \sin\left(\frac{\pi}{4}\right) \approx 184.6m$. The gradient at this point is

$$\nabla D(1, 0.5) = \left(-60, -\frac{50\pi}{2} \cos\left(\frac{\pi}{4}\right)\right) = \left(-60, -\frac{25\pi}{\sqrt{2}}\right).$$

which is measured in m/km .

$$\frac{\partial D}{\partial x} \quad \frac{\partial D}{\partial y}$$

10.2 Gradient

Question

A team is mapping the ocean floor. They produce the model $D(x, y) = 1700 - 30x^2 - 50 \sin(\frac{\pi y}{2})$ in m , with $(x, y) \in [-2, 2] \times [-2, 2]$ in km . There is a leaking oil well at $(1, 0.5)$; find the depth at this location, as well as the direction and magnitude of steepest increase here.

Solution

The depth at $(1, 0.5)$ is

$D(1, 0.5) = 250 - 30 - 50 \sin(\frac{\pi}{4}) \approx 184.6m$. The gradient at this point is

$$\nabla D(1, 0.5) = (-60, -\frac{50\pi}{2} \cos(\frac{\pi}{4})) = (-60, -\frac{25\pi}{\sqrt{2}}),$$

which is measured in m/km . This makes an angle of $\arctan(\frac{25\pi}{60\sqrt{2}})$ with the positive x -axis, and has a magnitude of $\sqrt{3600 + \frac{625\pi^2}{2}}$.