

2 Lie Groups

Definition 2.1 (Lie groups). A **Lie group** is a group that is also a smooth manifold, where the group operations are smooth.

2.1 Generators

Suppose $g \in G$ depends smoothly on a set of continuous parameters; $g(\alpha)$ depends on α_a for $a = 1$ to N . It is useful to set

$$g(\alpha)\Big|_{\alpha=0} = e; \quad D(\alpha)\Big|_{\alpha=0} = 1. \quad (2.1)$$

In some neighbourhood of the identity, the Taylor expansion yields

$$D(d\alpha) = 1 + i d\alpha_a X_a + \cdots. \quad (2.2)$$

Definition 2.2 (generators). For $a = 1$ to N ,

$$X_a \equiv -i \frac{\partial}{\partial \alpha_a} D(\alpha) \Big|_{\alpha=0} \quad (2.3)$$

are called the **generators** of the group.

Remark 2.1. If D is unitary, X_a are hermitian.

Remark 2.2. The generators, X_a 's, form a vector space.

Claim 2.1 (exponential parametrisation).

$$D(\alpha) = \lim_{k \rightarrow \infty} \left(1 + \frac{i\alpha_a X_a}{k} \right)^k = e^{i\alpha_a X_a} \quad (2.4)$$

2.2 Lie Algebras

Claim 2.2.

$$[\alpha_a X_a, \beta_b X_b] \in \text{span}\{X_a\} \quad (2.5)$$

Proof.

$$e^{i\alpha_a X_a} e^{i\beta_b X_b} = e^{i\delta_a X_a}, \quad (2.6)$$

where δ is given by the Baker–Campbell–Hausdorff formula:

$$i\delta_a X_a = i\alpha_a X_a + i\beta_a X_a - \frac{1}{2}[\alpha_a X_a, \beta_b X_b] + \cdots. \quad (2.7)$$

Consider a function

$$Z(t) = \ln(1 + e^{it\alpha_a X_a} e^{it\beta_b X_b} - 1), \quad t \in \mathbb{R}. \quad (2.8)$$

Taylor expansion with respect to t yields

$$Z(t) = t(\alpha_a X_a + \beta_b X_b) + \frac{1}{2}t^2[\alpha_a X_a, \beta_b X_b] + \mathcal{O}(t^3). \quad (2.9)$$

Since

$$e^{i\alpha_a X_a} e^{i\beta_b X_b} = D(\alpha)D(\beta) = D(\alpha\beta), \quad (2.10)$$

$Z(t)$ belongs to $\text{span}\{X_a\}$. Therefore, each coefficient of t^k , including $[\alpha_a X_a, \beta_b X_b]$, in eq. (2.9) must also lie in the vector space. \square

Definition 2.3 (Lie algebras). A **Lie algebra** is a vector space \mathfrak{g} over a field F with a binary operation $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ called the Lie bracket, satisfying bilinearity, alternating property, and the Jacobi identity.

Remark 2.3. The Lie algebra is independent of the choice of representation, unless the representation is unfaithful.³

Definition 2.4 (structure constants). The **structure constants** f_{abc} of the group defines the commutator algebra

$$[X_a, X_b] = if_{abc}X_c, \quad (2.11)$$

where $f_{abc} = -f_{bac}$

Definition 2.5 (Lie-algebra representations). A **representation** of a Lie-algebra \mathfrak{g} is a mapping, D , of the elements of \mathfrak{g} onto a set of linear operators with the following properties:

1. D is linear: $D(aX + bY) = aD(X) + bD(Y)$.
2. D preserves the Lie bracket structure:
 $D([X, Y]) = [D(X), D(Y)] = D(X)D(Y) - D(Y)D(X)$.

Claim 2.3. *If there is any unitary representation of the algebra, then the f_{abc} 's are real.*

Proof.

$$if_{abc}X_c = -if_{bac}X_c = -[X_b, X_a] = -[X_a, X_b]^\dagger = if_{abc}^*X_c \quad (2.12)$$

\square

2.3 The Jacobi Identity

Remark 2.4. The matrix generators satisfy the following identity:

$$[X_a, [X_b, X_c]] + [X_c, [X_a, X_b]] + [X_b, [X_c, X_a]] = 0. \quad (2.13)$$

This is called the Jacobi identity.

Remark 2.5.

$$[X_a, X_b X_c] = [X_a, X_b]X_c + X_b[X_a, X_c]. \quad (2.14)$$

³There is vanishing X_a if and only if the representation is unfaithful; we see only the quotient algebra.

2.4 The Adjoint Representation

Definition 2.6 (adjoint representations). The **adjoint representation** T of an algebra is the one generated by the structure constants:

$$[T_a]_{bc} \equiv -if_{abc}. \quad (2.15)$$

Proof. The Jacobi identity implies

$$f_{bcd}f_{ade} + f_{abd}f_{cde} + f_{cad}f_{bde} = 0, \quad (2.16)$$

which is equivalent to

$$[iT_a, iT_b] = -f_{abc}iT_c, \quad (2.17)$$

since

$$[X_a, [X_b, X_c]] = if_{bcd}[X_a, X_d] = -f_{bcd}f_{abe}X_e. \quad (2.18)$$

□

Remark 2.6. The dimension of the adjoint representation is the number of independent generators, which is the number of real parameters required to describe a group element.

Claim 2.4.

$$\text{tr}(T_a T_b) = k^a \delta_{ab} \quad (\text{no sum}) \quad (2.19)$$

Proof. Consider a linear transformation

$$L : X_a \mapsto X'_a = L_{ab}X_b. \quad (2.20)$$

Then,

$$[X'_a, X'_b] = iL_{ad}L_{be}f_{deg}L_{gc}^{-1}X'_c \quad (2.21)$$

implies

$$L : f_{abc} \mapsto f'_{abc} = L_{ad}L_{be}f_{deg}L_{gc}^{-1}, \quad (2.22)$$

or equivalently,

$$L : T_a \mapsto T'_a = L_{ad}LT_dL^{-1}. \quad (2.23)$$

Since similarity transformations do not change the trace,

$$L : \text{tr}(T_a T_b) \mapsto \text{tr}(T'_a T'_b) = L_{ac}L_{bd} \text{tr}(T_c T_d). \quad (2.24)$$

By choosing an appropriate orthogonal operator L , we get eq. (2.19). □

Remark 2.7. Rescaling L appropriately, we could choose all the non-zero k^a 's to have absolute value 1.

Definition 2.7 (compact Lie algebras). A Lie algebra is said to be **compact** if all k^a 's are positive.

For now, only compact Lie algebras will be considered:

$$\text{tr}(T_a T_b) = \lambda \delta_{ab}, \quad \lambda > 0. \quad (2.25)$$

Claim 2.5.

$$f_{abc} = -i\lambda^{-1} \operatorname{tr}([T_a, T_b]T_c) \quad (2.26)$$

Proof.

$$\operatorname{tr}([T_a, T_b]T_c) = \operatorname{tr}(if_{abd}T_dT_c) = if_{abd} \operatorname{tr}(T_dT_c) = if_{abd}\lambda\delta_{dc} \quad (2.27)$$

□

Claim 2.6. f_{abc} is completely antisymmetric.

Proof.

$$f_{abc} = f_{bca} \quad (2.28)$$

since

$$\begin{aligned} \operatorname{tr}([T_a, T_b]T_c) &= \operatorname{tr}(T_aT_bT_c - T_bT_aT_c) \\ &= \operatorname{tr}(T_bT_cT_a - T_cT_bT_a) = \operatorname{tr}([T_b, T_c]T_a). \end{aligned} \quad (2.29)$$

□

2.5 Simple Algebras and Groups

Definition 2.8 (invariant subalgebras). An **invariant subalgebra** is some set of generators which goes into itself under commutation with any element of the algebra. I.e., if X is any generator in the invariant subalgebra and Y is any generator in the whole algebra, then $[Y, X]$ is a generator in the invariant subalgebra.

Claim 2.7. An invariant subalgebra generates the corresponding invariant subgroup by exponentiation, i.e.,

$$e^{-iY} e^{iX} e^{iY} = e^{iX'}, \quad (2.30)$$

where

$$X' = e^{-iY} X e^{iY}. \quad (2.31)$$

Proof.

$$e^{-iY} (iX)^n e^{iY} = (e^{-iY} iX e^{iY})^n \quad (2.32)$$

□

Definition 2.9 (trivial algebras). An algebra which has no nontrivial invariant subalgebra is called **simple**, where the whole algebra and 0 are trivial invariant subalgebra.

Remark 2.8. A simple algebra generates a simple group.

Claim 2.8. The adjoint representation of a simple Lie algebra is irreducible.⁴

⁴The simplicity of a group does not automatically imply the irreducibility of the adjoint representation.

Proof. Assume the contrary. Then there is an invariant subspace spanned by T_r for $x = 1$ to K , where the rest of the generators T_x for $x = K + 1$ to N . Therefore, since

$$[T_a, T_x] = if_{axr}T_r \quad (2.33)$$

we must have

$$[T_a]_{xr} = -if_{axr} = 0. \quad (2.34)$$

By virtue of the complete antisymmetry of the structure constants, the nonzero structure constants involve either three r 's or three x 's, and thus the algebra falls apart into two nontrivial invariant subalgebras, and is not simple. \square

Definition 2.10 (semisimple algebras). Algebras without Abelian invariant subalgebras are called **semisimple**.

2.7 Fun with Exponentials

Claim 2.9.

$$\frac{\partial}{\partial \alpha_b} e^{i\alpha_a X_a} = \int_0^1 ds \, e^{is\alpha_a X_a} iX_b e^{i(1-s)\alpha_c X_c} \quad (2.35)$$

Proof. We can always define the exponential as a power series:

$$e^{i\alpha_a X_a} = \sum_{n=0}^{\infty} \frac{1}{n!} (i\alpha_a X_a)^n. \quad (2.36)$$

Referring to eq. (2.4),

$$\frac{\partial}{\partial \alpha_b} e^{i\alpha_a X_a} = \frac{\partial}{\partial \alpha_b} \lim_{k \rightarrow \infty} \left(1 + \frac{i\alpha_a X_a}{k} \right)^k \quad (2.37)$$

$$= \lim_{k \rightarrow \infty} \sum_{l=1}^k \left(1 + \frac{i\alpha_a X_a}{k} \right)^{l-1} \left(\frac{iX_b}{k} \right) \left(1 + \frac{i\alpha_a X_a}{k} \right)^{k-l} \quad (2.38)$$

$$= \int_0^1 ds \, e^{is\alpha_a X_a} iX_b e^{i(1-s)\alpha_c X_c}. \quad (2.39)$$

\square