The p-adic height at split anomalous primes: the height matrix modulo p, the digits of the height, and an archimedean threshold

Yoshikazu Hosomura
Independent researcher, Japan · wvbks0@gmail.com

2026. Companion to [A].

Abstract

Let E/Q be an elliptic curve, $p \geq 5$ a prime of good ordinary reduction with $\#E(\mathbb{F}_p) = p$, $p \nmid \prod_\ell c_\ell$, and assume that $\Lambda = E(\mathbb{Q})/\mathrm{tors}$ surjects onto $E(\mathbb{F}_p)$. In the companion paper [A] the author showed that the cyclotomic p-adic height matrix on Λ is p-integral if and only if $E(\mathbb{Q}_p)[p] \neq 0$ (the split case), that in rank one the non-split value of $v_p(\mathrm{Reg}_p)$ is -1 exactly, and that modulo p the height matrix has rank at most two in the split case. The present paper studies the split case further.

First, the whole height matrix modulo p is given in closed form by two local constants and two homomorphisms χ, γ coming from the first two layers of the filtration of $E(\mathbb{Q}_p)$ (Theorem K); the deficit $r - 2$ is exactly what these two layers take away. Second, the second p-adic digit of the height already contains the Fermat quotient of the denominator of the point (Theorem N), so no third layer of the filtration governs $v_p(\mathrm{Reg}_p) \geq 2$. Summed rather than expanded, the digits are the truncations of one identity $n^2 h_p(P) = (1-p)^{-1}[p^{-2}\log(1+p^2\Theta) + 2\log(1+pF)] + p^{-2}\mathcal{L}(z(P''))$ in two Iwasawa logarithms and an even power series $\mathcal{L}$ in the formal logarithm whose coefficients $\mu_2 = -c$, $\mu_4 = -b_4/12$, $\mu_6 = -b_2b_4/360 - b_6/30$ are invariants of E (Theorem N*, Proposition 17); this determines the digit at which each local invariant becomes visible, and shows that $\min(v_p(\mathrm{Reg}_p), k)$ is a function of k layers of the filtration and $k-1$ digits of a Fermat quotient, and of nothing coarser (Theorem $G^{(k)}$). A non-degeneracy needed for these statements, $v_p(\varphi_p') = 1$ at the p-torsion point, is proved for all $p \geq 5$ (Theorem ND). Third, an archimedean input enters: a rational point deep in the formal filtration has large naive height, so if the shortest non-zero point of $\Lambda \cap E_1(\mathbb{Q}_p)$ has canonical height below $4\log p + \beta(E)$ then, in rank two, $v_p(\mathrm{Reg}_p) = 0$ (Propositions 24, 24^c); over several thousand split boxes of rank two this threshold accounts in expectation for the whole observed frequency of non-unit regulators [computation].

The arithmetic steps are formalised in Lean 4 (no sorry, no native_decide, standard axioms only); the analytic and geometric inputs enter as hypotheses. Statements are graded [Lean], [paper], [computation] or [known] throughout.

MSC 2020. 11G50, 11G07, 11S31, 11Y40, 68V20. **Keywords.** p-adic height, p-adic regulator, anomalous prime, Fermat quotient, p-adic sigma function, Lean 4.

1. Introduction

This paper continues [A], to which we refer for the main theorem — p-integrality of the height matrix at an anomalous prime is equivalent to $E(\mathbb{Q}_p)[p] \neq 0$ — and for the criterion through the Fermat quotient λ of the p-th division polynomial. Here we assume throughout that E is split at p, where the height matrix is integral, and ask what determines it modulo p and to higher precision. The answer has three parts: a closed form modulo p in two layers of the filtration of $E(\mathbb{Q}_p)$ (§3.10); the higher digits of the height as truncations of a single identity (§§3.11–3.14); and an archimedean threshold that decides the rank-two case in part (§3.15). §4 contains the proofs, §5 the computations (PARI/GP, curves from Cremona's tables), §6 the boundary between what Lean has checked and what it has not, §7 what is open, §8 what is not claimed and §9 related work.

Numbering. This paper and [A] come from one longer draft [Draft], published by the author under the pen name Kiichi. To keep cross-references stable, statement labels and section numbers of that draft are retained: the statements labelled A–J, S, R, R_s, R′, R′_s, R″, R″_s, R^+, Lemmas 1–5 and Proposition 6 are proved in [A], and §§3.1–3.9 of the draft are replaced by the summary in §3 below. We use in particular Lemma 3 (the decomposition $h_p = 2\Lambda_\sigma + 2A$ of the height), Lemma 4 (split $\iff e = v_p(t(pP')) \geq 2$), Theorem B and Corollary D (the second digit is $-\lambda$, and $\lambda = 0$ in the split case), Theorem H ($\lambda(m\bar{x}) = m^2\lambda(\bar{x})$), Theorem R_s and Proposition R″_s of [A].

Use of AI. The author carried out this research with the assistance of Claude Code (Anthropic). The constructions, proofs, computations and the Lean 4 formalisation were developed with the AI system under the author's direction; independent sessions of the system were used to re-check the Lean development and the computations. The author reviewed the arguments, checked the Lean development and its axiom logs, and takes full responsibility for the content of this paper.

2. Setting and normalisation

The curve. $E/\mathbb{Q}$ is given by a global minimal Weierstrass equation with coefficients $a_1, \ldots, a_6$, and $p \geq 5$ is a prime of good ordinary reduction; $\Lambda := E(\mathbb{Q})/\mathrm{tors}$ has rank r, and c_ℓ is the Tamagawa number at ℓ. Write $t = -x/y$ for the parameter of the formal group $\hat{E}$, $E_n(\mathbb{Q}_p) := \{P : v_p(t(P)) \geq n\}$, and for $P \in E(\mathbb{Q})$

$$P = (\alpha(P)/d(P)^2,\ \beta(P)/d(P)^3),\ \gcd(\alpha, d) = \gcd(\beta, d) = 1,\ d \geq 1.$$

v_p is normalised by $v_p(p) = 1$, $\log_p$ is the Iwasawa logarithm with $\log_p(p) = 0$, and for a p-adic unit u, $q_p(u) := (u^{p-1} - 1)/p \bmod p$ is the Fermat quotient. Then $q_p : \mathbb{Z}_p^\times \to \mathbb{F}_p$ is a surjective homomorphism and $v_p(\log_p u) = 1 + v_p(q_p(u))$, so "the second p-adic digit of $\log_p u$" and "the Fermat quotient of u" are the same datum.

The height. The height is taken in the explicit form of [Har, §2.2], which quotes the definition of [MST]: if P satisfies

> (A1) P reduces to O at p, and (A2) P reduces to a nonsingular point at every bad prime,

then $h_p(P) = 2\log_p(\sigma_p(P)/d(P))$; for a general point $h_p(Q) := h_p(n_{\mathrm{lcm}}Q)/n_{\mathrm{lcm}}^2$ with $n_{\mathrm{lcm}} := \mathrm{lcm}(\#E(\mathbb{F}_p), \mathrm{lcm}_\ell\, c_\ell)$. This is 2p times

the normalisation of [MST], so the two valuations of Reg_p differ by r. Against a computer algebra system the sign is worth recording: with this normalisation **h_p is minus PARI's canonical p-adic height**, so that n^2h_p(P) = $-n^2\cdot$`ellpadicheight(E,p,n,P)*[1,-s_2]~` exactly, in every digit; valuations and the valuation of Reg_p are unaffected.

The letter n. Only n_{lcm} needs the least common multiple; from §3.9 on, **n always denotes the Tamagawa product $\prod_\ell$ c_ℓ**, which is what makes nQ satisfy (A2) and is all the proofs use. The other uses of the letter are kept apart by a subscript: n_{lcm} here, n_Q in §5.1, n_p := (p−1)/2 in §§3.14 and 4.13. In §2 and Theorem C the index of a division polynomial φ_n, ψ_n, ω_n and the order of a cyclic subgroup are bound variables of those statements. The pairing is h_p(P, R) = (h_p(P+R) − h_p(P) − h_p(R))/2 and Reg_p := det(h_p(P_i, P_j)) on a basis of Λ.

Which σ matters. σ_p is the Mazur–Tate sigma function: the unique odd $\sigma(t) = t + c_2t^2 + \cdots$ with **σ_p ∈ tZ_p[[t]]** solving $x(t) + c = -(d/\omega)((1/\sigma)(d\sigma/\omega))$ with $c = (b_2 - E_2(E, \omega))/12$ [Har, (2)], [MT91]; $c_2 = a_1/2$. Another constant in place of E_2 shifts h_p by a multiple of the square of the p-adic elliptic logarithm, and only this choice keeps the coefficients in Z_p. Every statement below is about σ_p.

E_2 modulo p is an invariant of the reduced curve. For p ≥ 5 and E ordinary at p, Katz's congruence E_2^* ≡ E_{p+1}/E_{p−1} (mod p) expresses the p-adic weight-two Eisenstein value through two classical forms, whose values on (E, ω) are isobaric polynomials in $E_4(E, \omega) = c_4$ and $E_6(E, \omega) = -c_6$. Hence

> **E_2(E, ω) mod p — and with it c mod p — is a function of (c_4 mod p, c_6 mod p) alone**,

that is, of the reduced pair ($\bar{E}$, $\bar{\omega}$): the constant of the σ-normalisation is a local quantity at p, and not a global one as its definition through a p-adic modular form makes it look. Two internal checks make the statement independent of the source: E_{p−1} ≡ 1 (mod p) ([Ser73]) and E_{p+1} ≡ E_2 (mod p), the latter from Kummer's congruence B_{p+1}/(p+1) ≡ $B_2/2$ (mod p) together with σ_p(n) ≡ σ_1(n) (mod p). The normalisation $E_6(E, \omega) = -c_6$ is forced by the Tate curve — for E_q : $y^2 + xy = x^3 - 5s_3x - (5s_3 + 7s_5)/12$ one has $c_4 = 1 + 240s_3 = E_4$ and $c_6 = -1 + 504s_5 = -E_6$ — and is calibrated on the Hasse invariant in §5.8. The congruence itself is [known] ([Kat73], p-adic modular forms), and we have not retrieved the original; what §5.8 does is check the consequence against PARI's `ellpadics2` over 315,953 boxes, which makes the *use* of the congruence here safe without making its proof ours. This local description of c is what allows the quantity W of §3.11 to be identified with a datum of the formal group.

Which model matters. h_p(P) and v_p(h_p(P)) are invariants of E/Q, but the two halves b and c of the decomposition of Lemma 3 are not: the substitution (x, y) ↦ (x, y + sx) with s ∈ Z carries a global minimal equation to a global minimal equation, fixes x, α, d, φ_p, ψ_p and λ, and moves only $\beta \mapsto \beta + s\alpha d$, so b and c trade with each other while min(b, c) stands. Since p is odd, s can be chosen with **p | a_1**, and that is the model in which Theorem B and Theorem S are proved; in it b ≥ 2e comes for free, and the whole question is c. Statements about b or c alone are statements about a model, and the invariant form of Theorem S is v_p(h_p(pP′)) = 1.

Anomalous, split, ι. p is *anomalous* when p | #E(F_p), which for p ≥ 7 means #E(F_p) = p (§3.1); E is *split at p* when E(Q_p)[p] ≠ 0; ι says that $\Lambda \to$ E(F_p) is onto.

Division polynomials. φ_n, ψ_n, ω_n are normalised by x(nQ) = φ_n/ψ_n^2, y(nQ) = ω_n/ψ_n^3, with φ_n monic of degree n^2 in x. For odd n and a rational cyclic D of order n, the Vélu isogeny π_D : E → E/D has x-part $\bar{\varphi}/\bar{\psi}^2$ with $\bar{\psi}(x) = \prod$_{{±T}⊂D\{O}}(x −

x(T)) monic of degree (n−1)/2 and $\bar{\varphi}$ monic of degree n. Modulo p, for E ordinary, φ_p(x) = f(x^p) and ψ_p(x) = g(x^p) in F_p[A, B][X] with f monic of degree p in X, g of degree (p−1)/2 with leading coefficient the Hasse invariant, and f/g^2 the x-part of the Verschiebung V; when p is anomalous, ker V = E(F_p).

Grades. [Lean] kernel-checked, no `sorry`, no `native_decide`, axioms within [propext, Classical.choice, Quot.sound]; **[paper]** a complete proof here or cited; **[computation]** verified over a stated range; **[known]** in the literature. Where grades mix, the weakest governs.

3. Main theorems

3.1–3.9 Summary of [A]

Assume #E(F_p) = p, $p \nmid n := \prod_\ell c_\ell$, and ι. Let $\chi : \Lambda \to F_p$ be the discrete logarithm of the reduction with respect to a fixed generator $\bar{x}_0$ of E(F_p), $\Lambda_0 = \ker \chi$. In [A] the following are proved. **Main theorem**: every entry of M_Λ is in Z_p $\iff$ E(Q_p)[p] ≠ 0; in rank one v_p(Reg_p) ≥ 0 $\iff$ split, the non-split value being −1; split $\implies$ v_p(Reg_p) ≥ max(r − 2, 0). **Theorem R** (non-split): p·M_Λ mod p = $2\lambda_1 \cdot \chi\chi^t$. **Theorem R_s** (split): := M_Λ mod p vanishes on ker χ × ker χ, so ρ_1 := rank ≤ 2; and v_p(Reg_p) ≥ r − ρ_1, with v_p(Reg_p) = (r − 2) + v_p(det T_s) when $\rho_1 = 2$ (Corollary R′_s). **Proposition R″_s**: a = h_p(e_1) mod p = $2\lambda^{(2)}/n^2$ with $\lambda^{(2)} := q^{(2)}(\varphi_p(ne_1))/2$ the second Fermat quotient, $q^{(2)}(u) := (u^{p-1} - 1)/p^2$ mod p for q_p(u) = 0. We also use Corollary H″ of [Draft]: in the split case $\lambda^{(2)}(x(mQ)) = m^2\lambda^{(2)}(x(Q))$ for $p \nmid m$, proved exactly as Theorem H of [A] one digit deeper.

One consequence of Proposition R″ can be stated for a whole isogeny class. Let $\varphi : E \to E''$ be an isogeny over Q of degree m prime to p. Then a_p, hence anomalousness, is shared; the reduction $\bar{\varphi} : E(F_p) \to E''(F_p)$ is an isomorphism, its kernel having order dividing m; and E(Q_p)[p] ≅ E″(Q_p)[p], so splitting is shared as well.

Corollary R‴ [paper, conditional; computation.] *If in addition the functoriality $h_p^{E''}(\varphi(P)) = m \cdot h_p^{E}(P)$ of the cyclotomic p-adic height. Then, in the non-split case, $\lambda_{E''}(\bar{\varphi}(\bar{x})) = m \cdot \lambda_E(\bar{x})$ for every $\bar{x} \in E(F_p) \setminus \{O\}$; equivalently, since a change of generator multiplies λ_1 by a square,*

$$(\lambda_1(E)/p) \cdot (\lambda_1(E'')/p) = (m/p) \text{ in } \{\pm 1\}.$$

So the quadratic residue class of λ_1 is constant on the isogeny class when every degree is a square modulo p, and splits it into two otherwise.

The functoriality assumed there is the expected one for the canonical cyclotomic height and we did not verify it in a source, which is why the grade is conditional; the identity itself is [computation] over 2,098 pairs (§5.6). A direct local proof of $\lambda_{E''} \circ \bar{\varphi} = m \cdot \lambda_E$ — one that does not pass through a global point and the height — we do not have.

Two further consequences of Proposition R″_s are quoted throughout §§3.10–3.12 and §4.11, and we state them here. The first is the behaviour of $\lambda^{(2)}$ under a change of lift — Proposition 9 of §3.10 read modulo p — and it is stated separately because what the later sections use is its *slope*.

Proposition R⁗ (the slope) [paper; the arithmetic in Lean.] *Let p be anomalous and split, $\bar{x} \in E(F_p) \setminus \{O\}$, $x_0 \in Z_p$ a lift of $\bar{x}$, and $G := \varphi_p'/p$, integral by Lemma 5. Then for every $k \in Z_p$*

$$\lambda^{(2)}(x_0 + pk) = \lambda^{(2)}(x_0) + k \cdot s(\bar{x}),\ s(\bar{x}) := -G(\bar{x})/(2\varphi_p(\bar{x})) \in F_p,$$

and $\boldsymbol{s(\bar{x}) \neq 0}$*; so* $k \mapsto \lambda\hat{}\{(2)\}(x_0 + pk)$ *is a bijection of* F_p *and* $\lambda\hat{}\{(2)\}$*, unlike* λ*, does depend on the lift. Proof. The congruence is Proposition 9 (§3.10, proved in §4.11) read modulo p: there* $\Theta(x_0 + pk) \equiv \Theta(x_0) + (p-1)kG(x_0)/u - pk^2H(x_0)/(2u) \pmod{p^2}$ *with* $u := \varphi_p(x_0)$ *and* $\Theta \equiv 2\lambda\hat{}\{(2)\} \pmod p$*, whence* $2\lambda\hat{}\{(2)\}(x_0 + pk) \equiv 2\lambda\hat{}\{(2)\}(x_0) - kG(\bar{x})/\varphi_p(\bar{x})$*; the denominator is a unit by Corollary D. The non-vanishing of s is the non-degeneracy (ND) of §3.13 —* $\varphi_p(\bar{x})$ *being a unit,* $s(\bar{x}) \neq 0 \iff G(\bar{x}) \neq 0 \iff v_p(\varphi_p'(x_0)) = 1$ *— and (ND) holds for every* $p \geq 5$ *by Theorem ND (§3.14).* ■

Corollary R′′′′′ (localisation of the off-diagonal block) [paper; the identity in Lean. *] In the situation of Theorem R_s take a basis of* Λ *with* $\chi(e_1) = 1$ *and* $e_2, \ldots, e_r \in \Lambda_0$*, put* $n := \prod_\ell c_\ell$ *and, for* $j \geq 2$*,*

$$\kappa_j := (x(n(e_1 + e_j)) - x(n\ e_1))/p \bmod p \in F_p.$$

Then $b_j = h_p(e_1, e_j) \bmod p$ *satisfies* $\boldsymbol{b_j = s(\bar{x}_0)\cdot\kappa_j/n^2}$ *with* $\bar{x}_0$ *the reduction of* $x(n\ e_1)$*; hence* $\boldsymbol{\rho_1 = 2 \iff \kappa_j \neq 0}$ ***for some j****. Proof. Both* $x(n(e_1 + e_j))$ *and* $x(n\ e_1)$ *are lifts of the same class* $\bar{x}_0$*, since* $\chi(e_j) = 0$*, and they differ by* $p\kappa_j$*; so* κ_j *is well defined. Polarising,* $2b_j \equiv a(e_1 + e_j) - a(e_1) - a(e_j)$ *with* $a(P) := h_p(P) \bmod p$*. Here* $a(e_j) = 0$ *by Lemma 2, and by Proposition R″_s* $a(e_1 + e_j) - a(e_1) = 2(\lambda\hat{}\{(2)\}(x(n(e_1+e_j))) - \lambda\hat{}\{(2)\}(x(n\ e_1)))/n^2$*, which by Proposition R′′′′ with* $k = \kappa_j$ *is* $2s(\bar{x}_0)\kappa_j/n^2$*. The equivalence is the last display of Theorem R_s together with* $s \neq 0$*.* ■

3.10 The height matrix modulo p in closed form, and the second digit

Here E is split at p, #E(F_p) = p, $p \nmid \prod_\ell$ c_ℓ, ι holds, and $n := \prod_\ell$ c_ℓ. Two steps of the filtration of E(Q_p) carry the whole of $\bar{M}$; the *second* p-adic digit of the height then leaves the filtration altogether.

Two homomorphisms. By Lemma 4 the group D := E(Q_p)[p] has order p and D ∩ E_1(Q_p) = 0, while D\{O} → E(F_p)\{O} is a bijection (proof of Corollary D); hence D → E(F_p) is an isomorphism and

$$\mathrm{E(Q_p) = D \oplus E_1(Q_p).}$$

Fix $T_0 \in D$ reducing to the chosen generator G of E(F_p), let χ be the discrete logarithm of the reduction with respect to G, and for $Q \in E(Z_p)$ with $\bar{Q} \neq O$ write $Q = \chi(Q)T_0 + R$ with $R \in E_1(Q_p)$ — well defined, since $pT_0 = O$ — and

$$\gamma(Q) := t(R)/p \bmod p \in F_p,\ t = -x/y \text{ the formal parameter.}$$

Then χ and γ are homomorphisms, $\gamma(T_0) = 0$, $\gamma|_\{E_1(Q_p)\} = t/p \bmod p$, and $\ker\chi \cap \ker\gamma = E_2(Q_p)$.

Lemma 7 (the first digit of the x-coordinate) [paper; the arithmetic step in Lean. *] Let* $P \in E(Z_p)$ *with* $\bar{P} \neq O$ *and let* $R \in E_1(Q_p)$ *with* $t := t(R)$*. Then* $R \mapsto x(P + R)$ *lies in* $Z_p[[t]]$ *and*

$$x(P + R) = x(P) + c_1(P)\cdot t + O(t^2),\ c_1(P) := 2y(P) + a_1x(P) + a_3.$$

Since $\#E(F_p) = p$ *is odd,* $\bar{E}(F_p)[2] = \{O\}$*, so* $\boldsymbol{c_1(P) \in Z_p\hat{}\times}$*.*

With κ_j as in Corollary R′′′′′, Lemma 7 gives $\kappa_j = c_1\cdot\gamma(n\ e_j)$, so $\kappa_j \neq 0 \iff v_p(t(n\ e_j)) = 1$, and Corollary R′′′′′ turns into

$$\rho_1 = \mathbf{2} \iff \gamma|_\{\Lambda_0\} \neq \mathbf{0} \iff \Lambda_0 \not\subset \mathbf{E_2(Q_p)},$$

a condition on the image of Λ in the *second* layer of the filtration, of exactly the same nature as the hypothesis ι in the first.

Proposition 8 (two purely local identities) [paper. *] For every class $\bar{P} \in E(F_p)\backslash\{O\}$, with s as in Proposition R'''', c_1 as in Lemma 7, and $\lambda_T(\bar{P}) := \lambda\hat{}\{(2)\}(x(T))$ for the unique $T \in D\backslash\{O\}$ above $\bar{P}$,*

$$\textit{(L1) } s(\bar{P})\cdot c_1(\bar{P}) = B_0\cdot\chi(\bar{P}),\ \textit{(L2) } \lambda_T(\bar{P}) = \lambda_\{T,1\}\cdot\chi(\bar{P})^2,$$

where $B_0 := s(G)\cdot c_1(G)$ and $\lambda_\{T,1\} := \lambda_T(G)$ depend only on E/Q_p and on the choice of G. No global point and no height enter.

Theorem K (the height matrix modulo p) [paper; the bilinear algebra in Lean. *] Under the hypotheses of Theorem R_s, for all $P, R \in \Lambda$,*

$$h_p(P) \equiv 2\lambda_\{T,1\}\cdot\chi(P)^2 + 2B_0\cdot\chi(P)\gamma(P) \pmod p,$$

$$\bar{M}(P, R) = 2\lambda_\{T,1\}\cdot\chi(P)\chi(R) + B_0\cdot(\chi(P)\gamma(R) + \gamma(P)\chi(R)).$$

The constants $\lambda_\{T,1\}$ and B_0 depend only on E/Q_p; the only global input is the pair of homomorphisms $(\chi, \gamma) : \Lambda \to F_p \times F_p$, and ***the Tamagawa product n cancels****.*

Theorem K contains Theorem R_s (the form is built from two linear functionals, so $\rho_1 \leq 2$), Corollary R'_s, Proposition R''_s and Corollary R'''' in one line. Its consequences are read off:

Corollary K' [paper. *] (i) $\rho_1 = 2 \iff (\chi, \gamma) : \Lambda \to F_p^2$ is onto; (ii) if $\gamma = c\chi$ on Λ then $\bar{M} = (2\lambda_\{T,1\} + 2B_0c)\cdot\chi\otimes\chi$, of rank ≤ 1, and of rank 0 exactly when $c = -\lambda_\{T,1\}/B_0$; (iii) in rank one, $\boldsymbol{v_p(Reg_p) \geq 1 \iff \gamma(e_1)/\chi(e_1) = -\lambda_\{T,1\}/B_0}$, one residue class out of p; (iv) with $\Lambda_1 := \Lambda \cap E_2(Q_p) = \ker\chi \cap \ker\gamma$ of rank $r - 2$ when (χ, γ) is onto, and $\hat{C}$ the Gram matrix of $p\hat{}\{-1\}h_p$ on Λ_1, the Schur complement of Corollary R'_s satisfies $T_s \equiv p\cdot\hat{C} \pmod{p^2}$, so*

$$\boldsymbol{v_p(Reg_p) = r - 2 \iff \det\hat{C} \not\equiv 0 \pmod p}.$$

So the deficit r − 2 is exactly what the two local layers χ and γ take away, and the residual global content of v_p(Reg_p) in rank ≥ 3 is a non-degeneracy of p^{−1}h_p one layer deeper — a statement of Schneider's kind, moved but not removed. The exact formula v_p(Reg_p) = (r − 2) + v_p(det Ĉ) needs det Ĉ to be a unit and fails otherwise (§5.7).

The second digit. Theorem K settles h_p modulo p. The natural continuation would be a third homomorphism $\gamma_2 : \Lambda_1 \to E_2/E_3$, and the natural guess is that v_p(Reg_p) ≥ 2 is a condition on the image of Λ in E(Q_p)/E_3. It is not. Write, for a unit u with q_p(u) = 0,

$$Q_2(u) := (u\hat{}\{p-1\} - 1)/p^2 \in Z_p,\ \Theta(x_0) := Q_2(\varphi_p(x_0)),$$

so that q^{(2)}(u) = Q_2(u) mod p and $\Theta(x_0) \equiv 2\lambda\hat{}\{(2)\}(x_0) \pmod p$. Θ is defined at every lift x_0 of a class of E(F_p)\{O} in the split case, by Corollary D and Lemma 5.

Theorem N (the second digit) [paper. *] In the situation of Theorem R_s let $P \in \Lambda$ with $\chi(P) \neq 0$, put $P' := nP$ and write $x(P') = a/d'^2$ in lowest terms (so $p \nmid d'$). Then*

$$\boldsymbol{n^2\cdot h_p(P) \equiv (1 + p)\cdot\Theta(x(P')) + 2p\cdot q_p(d') \pmod{p^2}},$$

where $q_p(d') = (d'\hat{}\{p-1\} - 1)/p$ mod p is the Fermat quotient of the rational integer d'.

Modulo p this is Proposition R″_s, since $\Theta \equiv 2\lambda\hat{}\{(2)\}$. Modulo p^2 a second term appears, and it is not a function of the image of P in E(Q_p) at all: d′ is the denominator of the point, a global integer, and two points with the same image in E(Q_p)/E_k for every k may have different q_p(d′). Hence:

Remark (no third homomorphism). *The localisation of Theorem K does not continue past the first digit. There is no γ_2 in the answer: v_p(Reg_p) ≥ 2 is not a condition on the position of Λ in the filtration of E(Q_p), and the induction "layer k ↦ a homomorphism to E_k/E_{k+1}" stops at k = 1. What replaces it is the pair (a local p-adic number, the Fermat quotient of a global denominator); see Corollary N′.*

The local half of the second digit is nevertheless completely described, and by a single p-adic number.

Proposition 9 (the second digit under lifting) [paper; the arithmetic step in Lean. *] With G := φ_p′/p (integral by Lemma 5), H := dG/dx and u := φ_p(x₀), for every k ∈ Z_p*

$$\Theta(x_0 + pk) \equiv \Theta(x_0) + (p-1)\cdot k\cdot G(x_0)/u - p\cdot k^2\cdot H(x_0)/(2u) \pmod{p^2},$$

where G(x₀) and u are needed modulo p^2 and H(x₀) modulo p. Modulo p this is Proposition R‴″, with slope $s = -G(\bar{x})/(2u)$.

Proposition 10 (the bad lift) [paper; the arithmetic step in Lean. *] Fix a class $\bar{x} \in E(F_p)\backslash\{O\}$ and $T \in D\backslash\{O\}$ above it. Then $\delta \mapsto \Theta(x(T) + p\delta)$ is a bijective isometry of Z_p; in particular it has a unique zero δ_*(E, p, x̄) ∈ Z_p, a local invariant, and for every lift $x_0 = x(T) + p\delta$*

$$\boldsymbol{v_p(\Theta(x_0)) = v_p(\delta - \delta_*)}.$$

Proposition 10 is the statement that all layers of the local half are governed by the distance to one p-adic number: no new local datum appears at any depth. Combining it with Theorem N gives the criterion at the second layer.

Corollary N′ (rank one, second layer) [paper. *] Let r = 1 and v_p(Reg_p) ≥ 1, that is p | Θ(x(P′)) by Proposition R″_s. Then*

$$\boldsymbol{v_p(Reg_p) \geq 2 \iff \Theta(x(P'))/p + 2\cdot q_p(d') \equiv 0 \pmod{p}},$$

and v_p(Θ(x(P′)))/p is computed from δ_* and δ by Proposition 10. So the second layer is again one residue class out of p, but the class is displaced by a quantity — 2q_p(d′) — that is invisible in E(Q_p).

Finally the bookkeeping of Lemma 3 needs one correction before Theorem N can be proved, and it is the reason the model does not appear in Theorem N.

Lemma 3′ (the formal part, sharpened) [paper. *] In the notation of Lemma 3, b = v_p(Λ_σ) ≥ min(e + v_p(a₁), 2e). In particular the bound b ≥ 2e stated in Lemma 3 for a model with p | a₁ is correct for e = 1 and too strong for e ≥ 2, where the right bound is e + v_p(a₁). It is enough for Proposition R″_s, which needs b ≥ 3; at the next digit the term $a_1\cdot t(pP')$ of 2Λ_σ has to be carried, and it cancels against the term $a_1\varphi\omega\psi$ of the Weierstrass equation (§4.11, Lean `a1_correction_cancels`).*

3.11 The ledger in closed form

Notation as in §3.10: $P \in \Lambda$ with $\chi(P) \neq 0$, $P' := nP$, $P'' := pP'$, $e := v_p(\psi_p(x(P'))) = v_p(d(P'')) \geq 2$, $x(P') = a/d'^2$ in lowest terms. Write

$$\Theta := Q_2(\varphi_p(x(P'))),\ F := (d'^{p-1} - 1)/p \in \mathbb{Z},\ c := (b_2 - E_2(E,\omega))/12,\ W := c/(p^4 \cdot x(P'')),$$

so that $q_p(d') = F \bmod p$, and $1/x(P'') = \psi_p(x(P'))^2/\varphi_p(x(P'))$ has valuation $2e \geq 4$, so that $W \in \mathbb{Z}_p$. Let $z := \log_E(t)$ be the formal logarithm, normalised by $dz = \omega$, so that $v_p(z(Q)) = v_p(t(Q))$ for $Q \in E_1(\mathbb{Q}_p)$. Put $X := x(P'')$, $Y := y(P'')$, $t := t(P'')$, $z := z(P'')$ and

$$R := a_2/X + a_4/X^2 + a_6/X^3 - a_1 Y/X^2 - a_3 Y/X^3,$$

so that $\omega_p(P')^2 = \varphi_p(x(P'))^3 \cdot (1 + R)$ is the Weierstrass equation of E cleared of denominators at P''.

The exact identity of the Remark opening §4 can be summed instead of expanded. Summed, it has no digits in it: what is left is two Iwasawa logarithms of explicit units and one power series in the formal parameter.

Theorem N* (the ledger in closed form) [paper. *] Under the hypotheses of Theorem N, as an identity in $\mathbb{Q}_p$,*

$$\boldsymbol{n^2 \cdot h_p(P) = (1/(1-p)) \cdot [\, p^{-2} \cdot \log(1 + p^2\Theta) + 2 \cdot \log(1 + pF)\,] + p^{-2} \cdot \mathcal{L}(z),}$$

where

$$\boldsymbol{\mathcal{L}(z) := \log(\, x(P'') \cdot \sigma_p(t(P''))^2\,) = 2 \cdot \log(\sigma_p(t)/t) - \log(1 + R),}$$

both logarithms being Iwasawa's. Moreover $v_p(\mathcal{L}(z)) = 2e$ whenever $p \nmid c$.

Proof. The Remark opening §4 gives $p^2 \cdot n^2 h_p(P) = 2\log(\sigma_p(t)/t) + 2\log(\varphi/\omega) - 2p^2 \cdot \log(d')$ with $\varphi := \varphi_p(x(P'))$, $\omega := \omega_p(P')$. From $\omega^2 = \varphi^3(1 + R)$ one has $2\log(\varphi/\omega) = \log(\varphi^2/\omega^2) = -\log \varphi - \log(1 + R)$. For a unit u, $\log u = (p-1)^{-1} \cdot \log(u^{p-1})$, so with $\varphi^{p-1} = 1 + p^2\Theta$ and $d'^{p-1} = 1 + pF$,

$$-\log \varphi - 2p^2 \cdot \log d' = -(p-1)^{-1} \cdot [\, \log(1 + p^2\Theta) + 2p^2 \cdot \log(1 + pF)\,].$$

Dividing by p^2 and using $(1-p)^{-1} = -(p-1)^{-1}$ gives the display, with $\mathcal{L}(z) = 2\log(\sigma_p(t)/t) - \log(1 + R)$. The two expressions for $\mathcal{L}$ agree because

$$t^2 = X^2/Y^2 = (\varphi/\psi^2)^2 \cdot (\psi^3/\omega)^2 = \varphi^2\psi^2/\omega^2 = \psi^2/(\varphi \cdot (1 + R)) = 1/(X \cdot (1 + R)),$$

where $\psi := \psi_p(x(P'))$, using $t = -x/y$, $x = \varphi/\psi^2$, $y = \omega/\psi^3$ at P'' and $\omega^2 = \varphi^3(1 + R)$. Hence $X \cdot \sigma_p(t)^2 = (\sigma_p(t)/t)^2 \cdot Xt^2 = (\sigma_p(t)/t)^2/(1 + R)$, and taking logarithms gives the second form. The valuation is Lemma 1 together with Proposition 17. ∎

The series $\mathcal{L}$ converges at $z = z(P'')$. By Proposition 17 its m-th coefficient has $v_p(\mu_\{2m\}) \geq -v_p((2m+2)!) \geq -(2m+2)/(p-1)$, while $v_p(z^{2m}) = 2me$ with $e \geq 2$ and $p \geq 5$, so the m-th term has valuation $\geq 2me - (2m+2)/(p-1) \to \infty$:

reading $\mathcal{L}$ term by term modulo p^k, as Proposition 18 does, is legitimate. As a series in the formal parameter t the convergence is immediate instead, both $2\log(\sigma_p(t)/t)$ and $\log(1 + R)$ being logarithms of $1 +$ (something of positive valuation).

Nothing in Theorem N* refers to a digit. The two logarithms are functions of the two integers Θ and F alone; the third term is a function of the position of P'' in the formal group, and the next two propositions say what kind of function.

Proposition 17 ($\mathcal{L}$ is an even series with invariant coefficients) [paper; the coefficient identities in Lean. *] As a power series in z,*

$$\mathcal{L}(z) = \mu_2 z^2 + \mu_4 z^4 + \mu_6 z^6 + \cdots, \textbf{ no odd power occurring},$$

and the coefficients are invariants of E over Q:

$$\mu_2 = -\boldsymbol{c},\ \mu_4 = -\boldsymbol{b_4/12},\ \mu_6 = -\boldsymbol{b_2 b_4/360 - b_6/30},$$

*with $\mu_\{2m\} \in \boldsymbol{Z[1/(2m+2)!][b_2, b_4, b_6, E_2]}$ — so $v_p(\mu_\{2m\}) \geq -v_p((2m+2)!)$, which for $p \geq 5$ is $\geq -(2m+2)/(p-1)$ — and $\boldsymbol{E_2}$ **occurring in** μ_2 **alone**. The denominators are **not** bounded independently of m: μ_6 has a 5, μ_8 a 7, and μ_{14} an 11 and a 13. That is the mechanism by which the digit at which a coefficient becomes visible can depend on p (Proposition 18).*

Proof. Mazur and Tate's characterisation of σ_p is the differential equation $(d/dz)^2 \cdot \log \sigma_p(z) = -(x(z) + c)$, and $x(z) + b_2/12 = \wp(z)$ with $\wp(z) = z^{-2} + \Sigma_\{k\geq 1\}\pi_k z^{2k}$ the Weierstrass series of the lattice of E, $\pi_1 = c_4/240$, $\pi_2 = c_6/6048$. Write $a := -b_2/12$. Then

$$(d/dz)^2 \cdot \log(\sigma_p(z)/z) = z^{-2} - \wp(z) - a - c = -(a + c) - \Sigma_\{k\geq 1\}\pi_k z^{2k},$$

whose right-hand side is even; σ_p is odd and $\sigma_p(z)/z \in 1 + z\mathbb{Z}_p[[z]]$, so $\log(\sigma_p(z)/z)$ is even with no constant term, and integrating twice gives

$$2 \cdot \log(\sigma_p(z)/z) = -(a + c)z^2 - 2 \cdot \Sigma_\{k\geq 1\}\pi_k z^{2k+2}/((2k + 1)(2k + 2)).$$

Also $x(z)z^2 = z^2\wp(z) + az^2 = 1 + az^2 + \pi_1 z^4 + \pi_2 z^6 + \cdots$ is a series in z^2, so $\log(x(z)z^2)$ is even, and $\mathcal{L} = \log(x(z)z^2) + 2\log(\sigma_p(z)/z)$ is even: that is the first claim. For the coefficients, expand $\log(1 + az^2 + \pi_1 z^4 + \pi_2 z^6 + \cdots)$:

- z^2: a from the first logarithm and $-(a + c)$ from the second, so $\mu_2 = -\mathbf{c}$; explicitly $-b_2/12 + E_2/12 = -(b_2 - E_2)/12$, since $-(a + c) = E_2/12$ (Lean `mu2_eq`);
- z^4: $\pi_1 - a^2/2$ from the first, $-\pi_1/6$ from the second, so $\mu_4 = \pi_1 - \mathbf{a^2/2} - \pi_1/\mathbf{6} = \mathbf{(c_4 - b_2^2)/288} = \mathbf{-b_4/12}$ by $c_4 = b_2^2 - 24b_4$ — and $\mathbf{E_2}$ **has cancelled** (Lean `mu4_eq`);
- z^6: $\pi_2 - a\pi_1 + a^3/3$ from the first, $-\pi_2/15$ from the second; with $c_4 = b_2^2 - 24b_4$ and $c_6 = -b_2^3 + 36b_2b_4 - 216b_6$ the three terms in b_2^3 cancel exactly (Lean `mu6_no_b2_cube`: $-1/6480 + 1/2880 - 1/5184 = 0$), leaving $\mu_6 = \mathbf{-b_2 b_4/360 - b_6/30}$ (Lean `mu6_eq`).

Only b_2, b_4, b_6 and, through c, E_2 enter. For the denominators, π_k is a polynomial in $g_2 = c_4/12$ and $g_3 = c_6/216$ with denominator dividing $(2k + 2)!$, and the second integration divides the coefficient of z^{2k+2} by $(2k + 1)(2k + 2)$; the logarithm of $1 + az^2 + \pi_1 z^4 + \cdots$ contributes no worse denominator, whence $\mu_\{2m\} \in \mathbb{Z}[1/(2m + 2)!][b_2, b_4, b_6, E_2]$. ■

The coefficients $\mu_\{2m\}$ carry even numerical subscripts throughout; they are not the unit series M(t) of §4.12 nor the homomorphism μ of §7.

Proposition 18 (which terms are visible at the k-th digit) [paper; the valuation arithmetic in Lean. *| In the situation of Theorem N*, reading the right-hand side modulo p^k leaves exactly the terms of valuation ≤ k − 1, and the valuations are | series | j-th (m-th) term | valuation (equality when v_p(Θ) = v_p(F) = 0) | |—|—|—| | p^{−2}·log(1 + p²Θ) | (−1)^{j+1}p^{2j−2}Θ^j/j | **2j − 2 − v_p(j)** | | 2·log(1 + pF) | 2(−1)^{j+1}p^jF^j/j | **j − v_p(j)** | | p^{−2}·ℒ(z) | μ_{2m}z^{2m}/p² | **2me − 2 + v_p(μ_{2m})** | In general the first two entries are **lower bounds**, the excess being j·v_p(Θ) and j·v_p(F); the non-generic case occurs in this paper, since Corollaries N′ and N″ treat boxes with Θ = pΘ₁. A term of valuation v is first visible at the **(v + 1)-th** digit, that is modulo p^{v+1}; in this counting Θ occupies the first digit and F the second, which is Proposition R″_s and Theorem N. In the generic case e = 2 the m-th coefficient of ℒ is first visible at the **(4m − 1 + v_p(μ_{2m}))-th** digit: μ₂ = −c at the **third**, μ₄ = −b₄/12 at the **seventh**, μ₆ at the **tenth** for p = 5 and at the **eleventh** for p ≥ 7.*

Proof. The three lines are the term-by-term valuations of the Iwasawa logarithm, with v_p(Θ) ≥ 0, v_p(F) ≥ 0 and v_p(z) = e; the first two are equalities exactly when Θ and F are units, and lower bounds otherwise. The arithmetic of the subscript is Lean `padicVal_eq_zero_of_lt`, `log_term_depth_small`, `log_term_depth_five`, `log_term_depth_ne_five`, `log_term_factor`, `theta_depth_j5_p5`, `theta_depth_j5_p7`, `log_term_j10_five`, `log_term_j10_seven`; the three digits of the last sentence are `lambda_depth_mono`, `lambda_depth_m2_e2`, `val_thirty`, `val_three_sixty`, `val_thirty_seven`, `mu6_visible_at_ten`. The factor v_p(j) is the whole content of the first two lines: the j-th term of a logarithm is one digit shallower than p^j whenever p | j, which is what puts the term with j = p inside the p^{j−1} digit. The third line is why μ_6, whose denominator is divisible by p for p = 5 alone, is visible one digit earlier at p = 5 than at p ≥ 7. ■

Theorem N* also fixes the coordinate in which the deep digits are to be read. The quantity W of §3.11 uses 1/x(P″) where ℒ uses z^2, and the two are not equal.

Lemma 21 (the two coordinates) [paper. *| As power series in the formal logarithm,*

$$z^2 - 1/x(z) = -(b_2/12)\cdot z^4 + O(z^6), \text{ no odd power occurring.}$$

Hence v_p(z(P′)² − 1/x(P′)) ≥ 4e, and p^{−2}(z(P′)² − 1/x(P′)) has valuation ≥ 4e − 2 ≥ 6 for e ≥ 2: ***1/x(P″) serves in place of z² through the sixth digit and not at the seventh.***

Proof. The proof of Proposition 17 gives $x(z)z^2 = 1 + az^2 + \pi_1 z^4 + \cdots$ with $a = -b_2/12$, a series in z^2. Inverting, $1/x(z) = z^2(1 + az^2 + \pi_1 z^4 + \cdots)^{-1} = z^2 - az^4 + O(z^6)$, again with only even powers, so $z^2 - 1/x(z) = az^4 + O(z^6)$. The valuation follows from v_p(z) = e, $v_p(b_2) \geq 0$ and p ≥ 5. ■

The leading coefficient is b_2 and not a_1: the terms of R linear in a_1 cancel against $a_1 t$ to leading order, so no explanation through a_1 alone can reach the fourth power of z. §5.10 measures the valuation in all 497 boxes with $b_2 \neq 0$, where it is 4e + $v_p(b_2)$ exactly.

3.12 The digits of the height

Every congruence for $n^2 h_p(P)$ in this paper is Theorem N* read modulo a power of p, the terms being selected by Proposition 18. Theorem N is the case k = 2, and the cases k = 3, 4 are the two theorems of the fourth digit; from k = 5 on the truncation is mechanical, and what is stated is what was measured.

Corollary N$^{(3)}$ (the third digit) [paper; the arithmetic in Lean. *] Under the hypotheses of Theorem N,*

$$n^2 \cdot h_p(P) \equiv (1 + p + p^2)\Theta + 2p(1 + p)F - p^2F^2 - (p^2/2)\Theta^2 - c/(p^2 \cdot x(P'')) \pmod{p^3}.$$

Corollary N$^{(4)}$ (the fourth digit) [paper; the arithmetic in Lean. *] In the same situation,*

$$n^2 \cdot h_p(P) \equiv (1 + p + p^2 + p^3)\Theta + 2p(1 + p + p^2)F - p^2(1 + p)F^2 + (2/3)p^3F^3 - (p^2/2)(1 + p)\Theta^2 - c/(p^2 \cdot x(P'')) \pmod{p^4}.$$

***No new local quantity enters**: by Proposition 18 the next coefficient of $\mathcal{L}$ is μ_4, and its digit is the seventh.*

Every term on the right of the two congruences is invariant under (x, y) $\mapsto$ (x, y + sx), since Θ, F, c and x(P″) are: the odd division polynomials ψ_n and φ_n are polynomials in b_2, b_4, b_6, b_8, the b_i are invariant under that substitution, x does not move, and $E_2(E, \omega)$ does not either. So the congruences are statements about invariants of E; the model with p | a_1 is where the proof runs (§4.12), not a hypothesis of the statement.

Corollary N$^{(5)}$ (the fifth digit) [computation: 648 of 648 boxes. *] In the same situation,*

$$n^2 \cdot h_p(P) \equiv (1 + p + p^2 + p^3 + p^4)\Theta - (p^2/2)(1 + p + p^2)\Theta^2 + (p^4/3)\Theta^3 + 2p(1 + p + p^2 + p^3)F - p^2(1 + p + p^2)F^2 + (2/3)p^3(1 + p)F^3 - (1/2)p^4F^4 + (2/5)p^5F^5 - c/(p^2 \cdot x(P')) \pmod{p^5}.$$

The term $(2/5)p^5F^5$ is $2p^4F^5 \equiv 2p^4F$ at $p = 5$, by Fermat, and $\equiv 0 \pmod{p^5}$ for $p \geq 7$: it is the first appearance of the factor v_p(j) of Proposition 18. ***No new local quantity enters.***

Corollary N$^{(6)}$ (the sixth digit) [computation: 648 of 648 boxes. *] In the same situation,*

$$n^2 \cdot h_p(P) \equiv (1 + p + p^2 + p^3 + p^4 + p^5)\Theta - (p^2/2)(1 + p + p^2 + p^3)\Theta^2 + (p^4/3)(1 + p)\Theta^3 + 2p(1 + p + p^2 + p^3 + p^4)F - p^2(1 + p + p^2 + p^3)F^2 + (2/3)p^3(1 + p + p^2)F^3 - (1/2)p^4(1 + p)F^4 + (2/5)p^5(1 + p)F^5 - c/(p^2 \cdot x(P'')) \pmod{p^6}.$$

***No new local quantity enters**, and $1/x(P'')$ may still be used in place of z^2: by Lemma 21 the error $z^2 - 1/x(P'')$ has valuation $\geq 4e \geq 2e + 4$, so it is invisible modulo p^6. The geometric factor of the F^5 term is $(1 + p)$ and not 1, because at $p = 5$ the term $(2/5)p^5F^5$ has valuation 4 and not 5: the factor v_p(j) of Proposition 18 shifts the geometric series as well. For $p \geq 7$ the second summand has valuation 6 and is invisible modulo p^6, so the one display serves both primes.*

Corollary N$^{(7)}$ (the seventh digit) [computation: 648 of 648 boxes. *] In the same situation,*

$$n^2 \cdot h_p(P) \equiv (1 + p + p^2 + p^3 + p^4 + p^5 + p^6)\Theta - (p^2/2)(1 + p + p^2 + p^3 + p^4)\Theta^2 + (p^4/3)(1 + p + p^2)\Theta^3 - (p^6/4)\Theta^4$$

$$+ 2p(1 + p + p^2 + p^3 + p^4 + p^5)F - p^2(1 + p + p^2 + p^3 + p^4)F^2 + (2/3)p^3(1 + p + p^2 + p^3)F^3$$

$$- (1/2)p^4(1 + p + p^2)F^4 + (2/5)p^5(1 + p + p^2)F^5 - (1/3)p^6F^6 + (2/7)p^7F^7$$

$$+ p\hat{}\{-2\}\cdot(\ -c\cdot z(P'')^2 - (b_4/12)\cdot z(P'')^4\ )\ (\mathrm{mod}\ p^7).$$

At this depth Θ^4 and F^6 enter for every p; the term $(2/7)p^7F^7$ is effective at $p = 7$ alone, where its valuation is 6, and the third summand of the geometric factor of F^5 is effective at $p = 5$ alone — both are invisible modulo p^7 at the other prime, so the one display serves both. ***b_4 is the first new local invariant after c, and the seventh digit is where it enters:*** *dropping the term $\mu_4 z^4$ leaves 232 of the 648 boxes, and keeping $1/x(P'')$ in place of z^2 leaves 310 of 648 (§5.10). The congruence is sharp — the two sides differ at valuation exactly 7 — in 499 of the 648 boxes.*

The pattern of the coefficients is Proposition 18 and nothing else, so no further corollary is worth displaying. What is worth recording is how far the closed form was tested and what each ingredient of it is needed for. Over the 648 boxes of §5.10 — conductor ≤ 12000, p $\in \{5, 7\}$, a_p = 1, #E(F_p) = p, p $\nmid \prod_{\ell}$ c_ℓ, ι, split, e ≥ 2:

At k = 3, 4 and 5 the truncation of Theorem N* is the corollary displayed above and holds in 648 of the 648 boxes; since $\mathcal{L}$ contributes the single term $\mu_2 z^2$ there, the truncations of $\mathcal{L}$ do not differ from it. From k = 6 on:

k	Theorem N* truncated by Proposition 18	$\mathcal{L}$ truncated at z^4	$\mathcal{L}$ truncated at z^2	$\mathcal{L}$ without its odd powers
6	**648/648**	648/648	648/648	648/648
7	**648/648**	648/648	232/648	648/648
8	**648/648**	648/648	146/648	648/648
9	**648/648**	648/648	125/648	648/648
10	**648/648**	392/648 (p = 7: **316/316**; p = 5: 76/332)	114/648	648/648

The one entry that two independent runs do not agree on is the third column at k = 10: the run of §5.10 gives 114 and an independent recomputation at p-adic precision 40 and series order 26 gives 113. The difference is one box; it has not been resolved, and nothing in the paper turns on it.

Three readings of the table are the content of Proposition 18. First, the last column: $\mathcal{L}$ **needs no odd power at any depth reached**, which is the first claim of Proposition 17 under test. Second, the third column: truncating $\mathcal{L}$ at z^2 is exact through k = 6 and fails at k = 7, which is μ_4 arriving at its predicted digit. Third, the second column at k = 10: truncating at z^4 still holds for **all** 316 boxes with p = 7 and fails for 256 of the 332 with p = 5, which is v_p(μ_6) = -1 for p = 5 and 0 for p ≥ 7 — a local invariant whose digit depends on p, the same mechanism as the factor v_p(j) on the side of the logarithms.

Remark (where the normalisation of σ becomes visible, and where it stops). *The third digit is the first at which the choice of c — that is, of E_2 — is visible in the height: by Proposition 18 the term $\mu_2 z^2$ has valuation $2e - 2 = 2$ for $e = 2$, hence the third digit. The remark of §2, that another constant in place of E_2 shifts h_p by a multiple of the square of the p-adic elliptic logarithm, is this statement with the digit made explicit; and the square of the elliptic logarithm is exactly z^2, so the remark of §2 is the term $\mu_2 z^2$ itself. By Proposition 17 the third digit is also the* ***last*** *at which the normalisation is visible:* ***E_2 occurs in μ_2***

***alone**, and every deeper coefficient μ_4, μ_6, ... is a polynomial in b_2, b_4, b_6 with rational coefficients — the ring is Z[1/(2m + 2)!] and not Z[1/6], which is what makes the digit of μ_{2m} depend on p (Proposition 17). Unlike Θ, the term $c/(p^2x(P'))$ is invariant under $x \mapsto u^2x$ as well ($c \mapsto u^2c$ and $x(P') \mapsto u^2x(P')$), so it does not share the model dependence recorded in the Remark of §4.12.*

3.13 The minimal global data

Descending to the deeper layers gives two more criteria of the same shape as Corollary N′.

Corollary N″ (the third layer) [paper; the bookkeeping in Lean. *] Let $r = 1$ and $v_p(\mathrm{Reg}_p) \geq 2$, and write $\Theta = p\Theta_1$, $S := (\Theta_1 + 2F)/p$. Then*

$$\boldsymbol{v_p(\mathrm{Reg}_p) \geq 3 \iff S \equiv F^2 + W \pmod p}.$$

Corollary N‴ (the fourth layer) [paper; the bookkeeping in Lean. *] Let $r = 1$ and $v_p(\mathrm{Reg}_p) \geq 3$, and write $S - F^2 - W = pU$. Then*

$$\boldsymbol{v_p(\mathrm{Reg}_p) \geq 4 \iff 6U + 6(S - F^2) + 4F^3 \equiv 0 \pmod p},$$

and together with the three earlier criteria $\min(v_p(\mathrm{Reg}_p), 4)$ is determined (Lean `quintichotomy`).

W is a datum of the formal group. The quantity W is the only new ingredient of the third digit, and it is not a new *global* one.

Proposition 12 (W in the first layer) [paper; the bookkeeping in Lean. *] Let $T := \chi(P)\cdot T_0$ and $t_1 := t(nP - T) \in pZ_p$, with digits $t_1/p = \gamma + p\gamma_2 + p^2\gamma_3 + \cdots$. Then*

$$\boldsymbol{t(P'') = [p](t_1) \equiv p^2\gamma \pmod{p^3}}, \text{ hence } \boldsymbol{W \equiv c\cdot\gamma^2 \pmod p} \text{ and } \boldsymbol{e = 2 \iff \gamma \not\equiv 0 \pmod p}.$$

So Corollary N″ reads $v_p(\mathrm{Reg}_p) \geq 3 \iff S \equiv F^2 + c\cdot\gamma^2 \pmod p$, with c local by the Remark of §2 and γ the first layer of the filtration: W adds no global datum to (χ, γ, F).

Proposition 11 (the x-coordinate to three digits) [paper. *] With $x \circ \tau_T(t) = x(T) + c_1 t + \kappa t^2 + \nu t^3 + \cdots \in Z_p[[t]]$ (Lemma 7) and $\Gamma := \gamma + p\gamma_2 + p^2\gamma_3$, the quantity $\delta := (x(nP) - x(T))/p$ satisfies*

$$\delta \equiv c_1\cdot\Gamma + p\cdot\kappa\cdot\Gamma^2 + p^2\cdot\nu\cdot\Gamma^3 \pmod{p^3};$$

in particular δ mod p^3 depends only on the image of nP in $E(Q_p)/E_4(Q_p)$.

The last piece is the rate at which the layers of the filtration reach the digits of Θ. It needs one non-degeneracy, which is the non-vanishing of the slope of Proposition R⁗.

Theorem G′ (how deep a layer reaches) [paper; the valuation arithmetic in Lean. *] Let $T \in D\backslash\{O\}$, $X(t) := x(\tau_T(R(t))) \in Z_p[[t]]$ (Lemma 7) and $\Theta(t) := Q_2(\varphi_p(X(t)))$. Assume*

(ND) $v_p(\varphi_p'(x(T))) = 1$, equivalently $G(x(T)) \not\equiv 0 \pmod p$, equivalently $s(\bar{T}) \neq 0$.

Then for every $t_1 \in pZ_p$ and every $j \geq 1$

$$\boldsymbol{v_p(\Theta(t_1 + p\hat{}j) - \Theta(t_1)) = j - 1}.$$

In particular $\Theta(t_1)$ mod $p\hat{}\{j-1\}$ is a function of t_1 mod $p\hat{}j$, and no coarser datum will do: the j-th layer of the filtration moves exactly the digits of Θ from $p\hat{}\{j-1\}$ on.

(ND) is a condition on $(\bar{E}, x(\bar{T}))$ alone, since $\varphi_p' = p \cdot G$ with $G \in Z[a_1, \ldots, a_6][x]$ (Lemma 5). It holds whenever $\mathrm{Res}_x(G, \psi_p \bmod p) \neq 0$ in F_p, because for anomalous p the x-coordinate of every point of $E(F_p)\backslash\{O\}$ is a root of $\psi_p \bmod p$ (ker V = $E(F_p)$, §2). **That resultant is non-zero for every $p \geq 5$** (Theorem ND, §3.14, proved in §4.13), so (ND) is a hypothesis no longer; §5.8 records the exhaustive verification for $p \leq 41$ that preceded the proof. The corresponding statement for ψ_p needs no hypothesis either: $v_p(\psi_p'(x(T))) = 1$ exactly, always (§4.12).

Theorem G″ (the minimal global data at the third digit) [paper; the sharpness in Lean. *] Let $p \geq 5$ be good ordinary, anomalous and split, $p \nmid \prod_ \ell\ c_\ell$, ι, $r = 1$. Fix E, p, T_0 and the local invariants x(T), c_1, κ, ν, δ_*, c. Then $\min(v_p(\mathrm{Reg}_p), 3)$ is a function of*

$$(\chi(P),\ \gamma(P),\ \gamma_2(P),\ \gamma_3(P),\ F \bmod p^2)$$

— three layers of the formal filtration of $E(Q_p)$ and two digits of the Fermat quotient of the denominator — and of nothing coarser. The refinement is: $\min(v_p, 1)$ uses (χ, γ); $\min(v_p, 2)$ adds $(\gamma_2, F \bmod p)$; $\min(v_p, 3)$ adds $(\gamma_3, F \bmod p^2)$.

The two halves of Theorem G″ come from the two halves of Theorem G′. *Sufficiency*: by the inequality half, $\Theta \bmod p^3$ is determined by $t_1 \bmod p^4$, that is by $(\gamma, \gamma_2, \gamma_3)$, and by Corollary $N^{(3)}$ and Corollary N″ the three criteria use Θ only modulo p^3 and F only modulo p^2. *Necessity of γ_3*: by the sharp half with j = 3, two points with the same (χ, γ, γ_2) and different γ_3 have Θ differing at valuation exactly 2, hence S mod p differs, hence at most one value of γ_3 satisfies the criterion of Corollary N″. *Necessity of F*: the pair of curves 3888p1 and 3267d1 at p = 7 are isomorphic over Q_p, have the same $\Theta \bmod p^2$ and differ in $v_p(\mathrm{Reg}_p)$ by the term $q_p(d')$ alone (§5.6, §5.8); on one curve, the two points of 4407a1 of §5.13 do the same (Proposition 21 (ii); read pointwise, $v_p(n^2 h_p(P))$ in place of $v_p(\mathrm{Reg}_p)$, since 4407a1 has rank two).

Proposition 19 (the argument of $\mathcal{L}$ is a layer of the filtration) [paper; the valuation step in Lean. *] Let $T := \chi(P) \cdot T_0$ and $t_1 := t(nP - T) \in pZ_p$ as in Proposition 12. Then $P'' = p \cdot (nP - T)$, and since $\log_E$ is a homomorphism of formal groups,*

$$z(P'') = p \cdot \log_E(t_1), \text{ hence } e = 1 + v_p(t_1).$$

So $\mathcal{L}(z(P''))$ is a function of t_1 — that is, of the layers $(\gamma, \gamma_2, \gamma_3, \ldots)$ — and of the invariants $\mu_2, \mu_4, \mu_6, \ldots$ of Proposition 17: ***the third term of Theorem N* brings in no global datum beyond the filtration*** *(Lean `layer_depth`; verified in 40 of 40 boxes, §5.10).*

Proposition 19 makes Proposition 12 the case of the third digit: $W \equiv c\gamma^2 \pmod p$ is the term $\mu_2 z^2$ read modulo the first digit it occupies. It also says which coordinate to use deeper. $W = c/(p^4 x(P''))$ is a quantity **of the digits up to the sixth**: by Lemma 21 the difference $z^2 - 1/x(P'')$ has valuation $\geq 4e$, so $p\hat{}\{-2\}(z^2 - 1/x(P''))$ has valuation $\geq 4e - 2 \geq 6$, and from the seventh digit on the formal logarithm $z(P'') = p \cdot \log_E(t_1)$, not $x(P'')$, is the right coordinate.

The two ingredients of Theorem N* that a layer of the filtration moves — Θ and $\mathcal{L}$ — respond to it at different rates, and the whole of what follows rests on the gap between them.

Proposition 20 (the two rates) [paper; the valuation arithmetic in Lean. *] In the situation of Theorem G′ and Proposition 19, let $e = 1 + v_p(t_1) \geq 2$ and $t_1' := t_1 + p\hat{}j$ with $j \geq 1$, and write $z' := z(t_1') = p \cdot \log_E(t_1')$. Then (i) $v_p(\Theta(t_1') - \Theta(t_1)) = j - 1$ (Theorem G′); (ii) $v_p(z' - z) = 1 + j$; (iii) $v_p(\mathcal{L}(z') - \mathcal{L}(z)) \geq \min_\{m \geq 1\}$ [*

***v_p(μ_{2m}) + (1 + j) + (2m − 1)·min(e, 1 + j)]**, with equality when 1 + j ≠ e and the minimum is attained by one m alone. In particular, if*

***v_p(c) < 2·min(e, 1 + j) + v_p(μ_4)** — which holds whenever p ∤ c, since min(e, 1 + j) ≥ 2 —*

*then the term m = 1 leads and the bound reads **v_p(c) + min(e, 1 + j) + (1 + j)**, with equality for 1 + j ≠ e; whereas when c = 0 — which forces j(E) = 0, and then μ_4 = 0 as well in the models met here — the leading term is m = 3 and the value is **(1 + j) + 5·min(e, 1 + j) + v_p(μ_6)** for 1 + j ≠ e; (iv) hence **v_p(p^{−2}(ℒ(z′) − ℒ(z))) ≥ j + 1 = (j − 1) + 2**, using only e ≥ 2, j ≥ 1, p ≥ 5 and v_p(c) ≥ 0 — c = 0 deletes the m = 1 term and makes the bound deeper, so **(iv) and (v) need no hypothesis on c**; (v) hence the difference of the two sides of Theorem N*, with F held fixed, has valuation **exactly j − 1**. So the ledger lags the digits of Θ by at least two digits, **at every depth**: (iv) puts no upper bound on j.*

Proof. (ii) In log_E(t) = Σ_{n≥1}b_nt^n/n with b_n ∈ Z and $b_1 = 1$, the k-th term of the Taylor expansion at t_1 with increment h = p^j is h^k·L^{(k)}(t_1)/k!, and the coefficients of L^{(k)}/k! are b_n·C(n−1, k−1)/k, since C(n,k)/n = C(n−1,k−1)/k; so its denominator divides k and the k-th term has valuation ≥ kj − v_p(k). For k ≥ 2 and p ≥ 5, p^{v_p(k)} ≤ k gives v_p(k) < k − 1 ≤ (k−1)j, so every term past the first is deeper than the first, which is h·L′(t_1) with L′ ∈ 1 + tZ_p[[t]] a unit, of valuation exactly j. Multiplying by p gives 1 + j.

(iii) Δ(z^2) = (z′ + z)(z′ − z) with z′ + z = 2z + Δz, and v_p(2z) = e since 2 is a unit, v_p(Δz) = 1 + j by (ii); the two differ unless 1 + j = e, and then v_p(z′ + z) = min(e, 1 + j). For every m ≥ 1, Δ(z^{2m}) = Δz·Σ_i z′^i z^{2m−1−i} has valuation **(1 + j) + (2m − 1)·min(e, 1 + j) whenever p ∤ 2m**, and ≥ that in general: the term z′^i z^{2m−1−i} has valuation i(1 + j) + (2m − 1 − i)e, which is ≥ (2m − 1)·min(e, 1 + j) with equality for exactly one i when 1 + j ≠ e — the exponent is min(e, 1 + j) and **not e**, since for 1 + j < e it is z′ and not z that is the shallower factor — while at 1 + j = e all 2m terms are of that valuation and their sum is 2m·z^{2m−1} to leading order. Summing over m and using min(e, 1 + j) ≥ 2 gives the bound of (iii), the terms with m ≥ 2 lying at ≥ (1 + j) + 2(2m − 1); with v_p(μ_{2m}) ≥ −v_p((2m+2)!) ≥ −(2m+2)/(p−1) ≥ −(m+1)/2 (Proposition 17) the inequality 4m − 4 ≥ (m+1)/2 for m ≥ 2 leaves those terms at ≥ j + 3 after the factor p^{−2}. The m = 1 term, when it is present, gives v_p(c) + min(e, 1+j) + (1+j) − 2 ≥ j + 1, which is (iv); when c = 0 that term is absent and the bound is the deeper one from m ≥ 2 (Lean `ledger_lags_any_m`, uniform in m and using neither c ≠ 0 nor v_p(μ_{2m}) ≥ 0). The displayed condition for m = 1 to lead is Lean `m1_leads_iff`, and `m1_leads_of_unit_c` is the case p ∤ c. The equality in (iii) needs a unique minimising m, since two summands of equal valuation may cancel: that is what happens at the boundary row 1 + j = e, where Δ(z^{2m}) = z^{2m}((1 + w)^{2m} − 1) with w := Δz/z a unit, so the m-th leading term drops **exactly when (1 + w)^{2m} = 1 in F_p** — for m = 1 the criterion 2u + 1 ≡ 0 of §5.11, and for (p − 1) | 2m every u with u ≢ −1 (Lean `boundary_cancel_m1`, `boundary_cancel_pow`, `boundary_cancel_of_card`).

(v) (1−p)^{−1}p^{−2}[log(1 + p^2Θ′) − log(1 + p^2Θ)] = (1−p)^{−1}·ΔΘ·[1 − p^2(Θ′ + Θ)/2 + ⋯], where the bracket and (1−p)^{−1} are units, so the Θ side has valuation v_p(ΔΘ) = j − 1 by (i), while the ℒ side is ≥ j + 1 by (iv); j − 1 < j + 1. The subscript arithmetic is Lean `theta_rate_digit`, `ledger_rate_digit`, `gap_is_e`, `ledger_lags`, `ledger_lags_general`, `uniform_in_k`. ■

Translates by p-th powers. Proposition 20 compares two values of the parameter t_1

with F held fixed. On the rational points of one curve F is not a free coordinate: the next proposition computes how F moves along P + p^aΛ, and the one after it says that on rational points F is a function of the height and of t_1.

Proposition 21 (translates by p-th powers) [paper; the arithmetic and the linear algebra in Lean. *] In the situation of Theorem N, let $\Lambda \subseteq E(Q)$ be free, $q := n^2 h_p|_\Lambda$ with polarisation q(P, Z) := (q(P + Z) − q(P) − q(Z))/2, $\chi : \Lambda \to Z/p$ the reduction character, and*

$$\zeta(Z) := t(p \cdot nZ)/p^2 \bmod p,$$

a homomorphism $\Lambda \to Z/p$ (since $p \cdot nZ \in E_2(Q_p)$) with $\zeta(P) \neq 0$ exactly when e(P) = 2. For $\chi(P) \neq 0$ put

$$C(P) := -(\varphi_p'(x(P'))/p) \cdot (2y(P') + a_1 x(P') + a_3)/\varphi_p(x(P')) \bmod p,$$

a unit: $v_p(\varphi_p'(x(P'))) = 1$ by (ND) (Theorem ND), since $x(P') \equiv x(T) \pmod p$; $\varphi_p(x(P'))$ is a unit since $x(P'') = \varphi_p(x(P'))/\psi_p(x(P'))^2$ with $v_p(x(P'')) = -2e = -2v_p(\psi_p(x(P')))$; and $2y + a_1 x + a_3$ does not vanish at P′ modulo p, since $E(F_p)$ has odd order. Then for $Z \in \Lambda$ and $a \geq 1$

$$\boldsymbol{F(P + p\hat{}aZ) \equiv F(P) + p\hat{}\{a-1\} \cdot (\chi(Z)/\chi(P)) \cdot (q(P) - C(P)\zeta(P)/2) \pmod{p\hat{}a}.}$$

In particular: (i) if $\chi(Z) = 0$ then $F(P + p\hat{}aZ) \equiv F(P) \pmod{p\hat{}a}$; (ii) if $\chi(Z) \neq 0$ and $p \mid q(P)$, the (a−1)-st digit of F moves if and only if e(P) = 2 — exactly at valuation a − 1 when e(P) = 2, and at valuation ≥ a when e(P) ≥ 3; (iii) if $k \geq 2$ and $S \in \Lambda \cap \ker \chi$ has $\zeta(S) \neq 0$, then $R := p\hat{}\{k-1\}S$ has depth exactly k, the points P and P + R agree in (χ, γ, …, $\gamma_\{k-1\}$) and in F mod $p\hat{}\{k-1\}$ and differ in γ_k, and $v_p(q(P + R) - q(P)) = k - 1$ exactly (Proposition 20 (v)); so whenever $v_p(q(P)) \geq k$, $\min(v_p(q), k)$ separates the two points. (For $S \in \ker \chi$ the conditions $\zeta(S) \neq 0$ and $\gamma(S) \neq 0$ are equivalent, each saying that S has depth exactly one; Theorem $G^{(k)}$ and §7(9) use the second.)

Proof. Four steps. (1) *The linear term of Θ.* If $nR \in E_j(Q_p)$ with $j \geq 1$, then $\Theta(P + R) - \Theta(P) \equiv C(P) \cdot t(nR)/p \pmod{p\hat{}j}$. Indeed $\Delta := t_1(P + R) - t_1(P) \equiv t(nR) \pmod{p\hat{}\{j+1\}}$ by the formal group law; $X(t) := x(T \oplus \iota(t))$ has $X'(t_1) \equiv (2y + a_1 x + a_3)(P') \pmod p$ from the invariant differential, so $h := X(t_1 + \Delta) - X(t_1) \equiv X'(t_1)\Delta \pmod{p\hat{}\{2j\}}$; and since $\varphi_p \bmod p$ is a polynomial in $x\hat{}p$, the Taylor coefficients $\varphi_p\hat{}\{(k)\}/k!$ with $2 \leq k < p$ are divisible by p, so $\varphi_p(x_0 + h) - \varphi_p(x_0) \equiv \varphi_p'(x_0)h \pmod{p\hat{}\{2j+1\}}$. Taking Q_2 gives $[(\Phi + \delta)\hat{}\{p-1\} - \Phi\hat{}\{p-1\}]/p^2 \equiv (p-1)\Phi\hat{}\{p-2\}\delta/p^2 \pmod{p\hat{}\{2j\}}$ with $(p-1)\Phi\hat{}\{p-2\} \equiv -\Phi\hat{}\{-1\}$, and $v_p(\varphi_p') = 1$ gives the claim. (2) *The kernel of χ.* For $S \in \ker \chi$, $q(S) \equiv 0$ and $2q(P, S) \equiv C(P)\zeta(S) \pmod p$. Modulo p Theorem N* reads $q(Q) \equiv \Theta(Q)$, the F term lying in pZ_p and the $\mathcal{L}$ term at valuation $\geq 2e - 2 \geq 2$; by step (1) with j = 1 and $t(nS)/p \equiv t(p \cdot nS)/p^2 \pmod p$, $q(P \pm S) - q(P) \equiv \pm C(P)\zeta(S)$; the second difference gives $2q(S) \equiv 0$ (Lean `second_difference`), and $2q(P, S) = q(P + S) - q(P) - q(S)$. (3) *A linear form.* $L_P(Z) := 2q(P, Z) - C(P)\zeta(Z) \bmod p$ is additive and vanishes on $\ker \chi$ by step (2), so $L_P(Z) \cdot \chi(P) = L_P(P) \cdot \chi(Z)$ with $L_P(P) = 2q(P) - C(P)\zeta(P)$ (Lean `ratio_of_ker_le`, which needs only the inclusion of kernels). (4) *The translate.* For $R = p\hat{}aZ$ the quadratic identity $q(P + R) - q(P) = p\hat{}a \cdot 2q(P, Z) + p\hat{}\{2a\}q(Z)$ (Lean `translate_quadratic`) gives $D := q(P + R) - q(P) \equiv p\hat{}a \cdot 2q(P, Z) \pmod{p\hat{}\{a+1\}}$ (Lean `shift_quadratic`). Since $nR \in E_\{a+1\}(Q_p)$, step (1) and $t(p\hat{}\{a-1\}W) \equiv p\hat{}\{a-1\}t(W)$ put the Θ side of D at $p\hat{}aC(P)\zeta(Z) \pmod{p\hat{}\{a+1\}}$, and Proposition 20 (iv) puts the $\mathcal{L}$ side at valuation $\geq a + 2$; so the F side of Theorem N* is $p\hat{}a \cdot L_P(Z) \pmod{p\hat{}\{a+1\}}$ (Lean `ledger_digit`), and $2(1 - p)\hat{}\{-1\}[\log(1 + pF') - \log(1 + pF)] \equiv 2p(F' - F)$. Step (3) turns $L_P(Z)$ into $(\chi(Z)/\chi(P)) \cdot L_P(P)$. Part (ii) is Lean `kappa_ne_zero` (with $q \equiv 0$, $C \neq$

0, $\zeta \neq 0$ and $p \neq 2$ the coefficient is non-zero), and the separation in (iii) is Lean `separates`. ■

Step (2) is a second proof, from the linear term of Θ alone, of the statement of Theorem K that M_Λ mod p vanishes on ker χ. The point of Proposition 21 is where q(P) enters: **only modulo p**. Along P + p^aΛ the digit of F at p^{a−1} is a function of the local quantities χ, q(P) mod p, C(P) and ζ, because the (a+1)-st digit of the translate of a quadratic form falls back to the form modulo p.

Proposition 22 (Theorem N* inverted) [paper; the polynomial identity in Lean. *] Since log is an isometry 1 + pZ_p → pZ_p, the term (2/(1 − p))·log(1 + pF) determines F modulo each power of p, and Theorem N* can be solved for F:*

F mod p^k is a function of n^2h_p(P) mod p^{k+1}, of t_1 mod p^{k+2}, and of the local invariants,

Θ mod p^{k+1} being a function of t_1 mod p^{k+2} (Theorem G′) and p^{−2}$\mathcal{L}$ a function of t_1 and μ_2, μ_4, … (Proposition 19). For k = 2, with

$$A := [n^2 h_p(P) - (1 + p + p^2)\Theta + (p^2/2)\Theta^2 + c/(p^2 \cdot x(P''))]/p,$$

$$\boldsymbol{F \equiv A/2 - pA/2 + pA^2/8 \pmod{p^2}}.$$

Proof. Corollary $N^{(3)}$ reads $2p(1+p)F - p^2F^2 \equiv pA \pmod{p^3}$, that is $2(1+p)F - pF^2 \equiv A \pmod{p^2}$. If $8F = 4A - 4pA + pA^2 + 8p^2m$ then $16(1+p)(8F) - p(8F)^2 - 64A \in p^2\mathbb{Z}$ (Lean `inverse_n3`), and $p \nmid 64$. The solution is unique modulo p^2: $F \mapsto 2(1+p)F - pF^2$ is injective modulo p^2, the difference of two values being $(F_1 - F_2)(2(1+p) - p(F_1 + F_2))$ with the second factor a unit. ■

So on the rational points of one curve **the two kinds of datum of Theorem $G^{(k)}$ are not independent coordinates**: F mod p^{k−1} is read off the height and the layers. Theorem $G^{(k)}$ is a statement about the parameter space of Theorem N*, and Proposition 21 is what that parameter space looks like along p-power translates. Both are checked in the form displayed, in §5.13.

With Theorem G′, Proposition 19 and Proposition 20 the description at the third digit extends to every digit, in a shape that says how the global input grows.

Theorem $G^{(k)}$ (the minimal global data at the k-th digit) [paper; the valuation arithmetic in Lean. *] Let $p \geq 5$ be good ordinary, anomalous and split, $p \nmid \prod_\ell c_\ell$, ι, $r = 1$, (ND). Fix E, p, T_0 and the local invariants x(T), c_1, κ, ν, δ_*, and μ_2, μ_4, …, μ_{2m_0}, where m_0 is the largest m with $2me - 2 + v_p(\mu_{2m}) \leq k - 1$. Then $\min(v_p(\mathrm{Reg}_p), k)$ is a function of*

$$\boldsymbol{(\chi, \gamma, \gamma_2, \ldots, \gamma_k) \text{ and } F \bmod p^{k-1}}.$$

*— and of nothing coarser. The global input is therefore of **two kinds at every depth** — the position of Λ in the filtration of E(Q_p), and the digits of the Fermat quotient of the denominator — and each grows by one per digit, so the list is of 2k numbers (Lean `data_count`, `data_count_step`, `data_count_linear`). **Theorem G″ is the case k = 3.** The sense of the minimality. It is minimality as a function on the parameter space that Theorem N* defines — t_1 mod p^{k+1} and F mod p^{k−1}, with the local invariants fixed. **Both kinds of necessity are nevertheless realised by rational points of one curve.** For the layer: if $S \in \Lambda \cap \ker \chi$ has $\gamma(S) \neq 0$, then $R := p^{k-1}S$ has depth exactly k and F(P + R) ≡ F(P) (mod p^{k−1}) by Proposition 21 (i), so whenever $v_p(n^2 h_p(P)) \geq k$ the*

two points agree in (χ, γ, …, γ_{k−1}) and in F mod p^{k−1}, differ in γ_k, and are separated by min(v_p, k) (Proposition 21 (iii)); at 4407a1 and 5956a1 with p = 5 and k = 3 this happens in 32 pairs of 32, and at k = 4 in 2 of 2 (§5.13). These curves have rank two, so the statement is read pointwise: v_p(n²h_p(P)) in place of v_p(Reg_p), Theorem N being an identity for each P with χ(P) ≠ 0. For the digit of F: besides the pair of curves 3888p1, 3267d1 at k = 3, the two points P = −5g₁ + 6g₂ and P + 25(−g₁ + g₂) of 4407a1 at p = 5 share t₁ mod p⁴, differ in the second digit of F alone, and have v₅(n²h_p) = 7 and 2 (Proposition 21 (ii)).* ***What rational points do not give is independence of the two kinds of datum along p-power translates****: on P + p^aΛ the digit of F at p^{a−1} is determined by χ, q(P) mod p, C(P) and ζ (Proposition 21), and on rational points F mod p^{k−1} is a function of the height and of t₁ (Proposition 22).*

The list is empty at a CM cell. At a cell with c = 0 — necessarily j(E) = 0, and then $\mu_4 = 0$ as well — the smallest m with 2me − 2 + v_p(μ_{2m}) ≤ k − 1 is m = 3, so for e = 2 the list $\mu_2, \dots, \mu_{2m_0}$ is **empty for every k ≤ 10**: v_p(p^{−2}$\mathcal{L}$(z)) = 6e − 2 = 10, the ledger contributes nothing before the eleventh digit, and min(v_p(Reg_p), k) is a function of (χ, γ, …, γ_k) and F mod p^{k−1} alone. Corollary N″ reads there v_p(Reg_p) ≥ 3 ⟺ S ≡ F² (mod p), since W ≡ cγ² = 0; the verification over the twenty such cells is in §5.11, and the witness for the necessity of F in Theorem G″ — the pair 3888p1, 3267d1 — is one of them (Lean `cm_ledger_invisible_before_eleventh`, `cm_no_local_invariant_before_eleventh`). The lag there is ten digits and not two.

Proof. *Sufficiency*, uniform in k: by Theorem N*, v_p(Reg_p) ≥ k is a condition on Θ mod p^k, on F mod p^{k−1} and on $\mathcal{L}$(z(P″)) mod p^{k+2}; Θ mod p^k is a function of t₁ mod p^{k+1} by the inequality half of Theorem G′, and $\mathcal{L}$(z(P″)) is a function of t₁ and of $\mu_2, \dots, \mu_{2m_0}$ by Propositions 17, 18 and 19, so the list suffices.

Necessity of γ_k. Take two data agreeing in (χ, γ, …, γ_{k−1}) and in F mod p^{k−1} and differing in γ_k. By Proposition 20 (v) with j = k the two values of n²h_p differ at valuation exactly k − 1, so they are not both ≡ 0 (mod p^k) and min(v_p(Reg_p), k) separates them (Lean `exact_pred_not_at`, `no_cancel`, `at_most_one_vanishes`, `layer_is_needed`). *Necessity of the (k−1)-st digit of F.* Replacing F by F + p^{k−2} changes 2log(1 + pF) by a quantity of valuation exactly k − 1, since 2 and 1 + pF are units (Lean `fermat_digit`, `fermat_digits_needed`), and by Proposition 20 (iv) the ledger cannot cancel it. *Nothing deeper contributes.* By Theorem G′ the layer γ_{k+1} moves Θ only at valuation k, and the k-th digit of F moves the second logarithm only at valuation k; neither is visible modulo p^k.

The two rates are what makes this uniform in k: Proposition 20 (iv) assumes only e ≥ 2, j ≥ 1, p ≥ 5, with no upper bound on j, so there is no depth at which the local term catches up with the layers. The larger e is, the further the ledger lags — its response to γ_j sits at the (e + j)-th digit — and the minimality holds with more room to spare. ■

This closes, at the third digit, the question left open by the Remark of §3.10: there is no third homomorphism into E_2/E_3 that carries the answer, but there *is* a finite list, and the whole of the non-filtration content is the pair of digits F mod p². What happens deeper is Theorem N*: the list grows by one layer and one digit of F per digit of the height, and the local side stays finite — by Proposition 17 the new local invariants are $\mu_4 = -b_4/12$ at the seventh digit, μ_6 at the tenth or eleventh, and so on, all of them classical invariants of E over Q.

3.14 The non-degeneracy (ND) is a theorem

Theorem G′ and the sharp half of Theorem G″ used (ND), the non-vanishing of the slope, and v5 of this paper verified it by exhaustion for $p \le 41$. It is a theorem for every $p \ge 5$. The route is a statement about division polynomials alone, so this subsection works over $\bar{F}_p$ with the short model

$$\bar{E} : y^2 = u(x) = x^3 + a_4 x + a_6,\ \Delta_0 = 4{a_4}^3 + 27{a_6}^2,\ \text{weights } w(x) = 2,\ w(a_4) = 4,\ w(a_6) = 6,$$

and $p \ge 5$, so that $E_4 = -48a_4$ and $E_6 = 864a_6$ identify $F_p[a_4, a_6]$ with the ring of modular forms mod p. Since $[p] = V \circ F$ in characteristic p, $\psi_p \bmod p$ is a polynomial in $x\hat{}p$ (§2): write

$$\psi_p \bmod p = \Psi(x\hat{}p),\ \deg_X \Psi = n_p := (p-1)/2 \text{ (the letter n is the Tamagawa product } \prod_\ell c_\ell \text{ elsewhere)},\ \mathrm{lc}_X \Psi = A \text{ (the Hasse invariant)},\ h := \Psi/A,\ \varphi_p \bmod p = \Phi(x\hat{}p),\ \Phi \text{ monic of degree p in X},\ x \circ V = \Phi/(A^2h^2),\ w\hat{}\{(p)\}(X) := X^3 + a_4\hat{}pX + a_6\hat{}p,$$

so that h is the kernel polynomial of the Verschiebung $V : \bar{E}\hat{}\{(p)\} \to \bar{E}$ and $\Phi/(A^2h^2)$ is its x-coordinate. Let $K := \psi_p'/p \bmod p$ and $S_1 := (\varphi_p - 4uK^2)/\psi_p$, so that (ND) $\iff S_1(x(\bar{T})) \ne 0$ (§4.11). Let

$$D := -6a_6 \cdot \partial/\partial a_4 + (4/3){a_4}^2 \cdot \partial/\partial a_6,$$

the derivation that raises the weight by 2 and kills Δ_0; in the normalisation above D is *minus* the Serre derivative, and on the q-expansion of a form of weight $\equiv 0 \pmod p$ it acts as $-q\, d/dq$.

Lemma 13 (two identities for an odd separable isogeny) [paper; the arithmetic in Lean. *] Let $\lambda : E' \to E$ be separable of odd degree m with kernel polynomial h monic of degree (m−1)/2, and write $x \circ \lambda = P/h^2$, $y \circ \lambda = y'\Omega/h^3$, $\lambda\hat{}*\omega_E = c \cdot \omega_\{E'\}$. Then $P'h - 2Ph' = c\Omega$ and $u_\{E'\}\Omega^2 = P^3 + a_4Ph^4 + a_6h^6$, and at every simple root ξ of h with $P(\xi) \ne 0$*

$$\boldsymbol{c^2 P(\xi) = 4u_\{E'\}(\xi)h'(\xi)^2 \text{ and } c^2 P'(\xi) = 2[u_\{E'\} \cdot (h')^2]'(\xi).}$$

For $\lambda = [p]$ this gives $p^2\varphi_p(x_T) = 4u(x_T)\psi_p'(x_T)^2$ and the identity (I-2) of §4.12. For $\lambda = V$, where $c^2 = A^2$ because the numerator of $x \circ \lambda$ has leading coefficient $c\hat{}\{-2\}$, it gives $\Phi(\xi) = 4w\hat{}\{(p)\}(\xi)h'(\xi)^2$ and $\Phi'(\xi) = 2[w\hat{}\{(p)\}(h')^2]'(\xi)$ at every root ξ of h.

Proposition 14 (the second p-adic digit of ψ_p is read off the first) [paper. *] For $p \ge 5$,*

$$\boldsymbol{A \cdot (\psi_p'/p) = u\hat{}\{(p-1)/2\} \cdot (d\Psi/dX)(x\hat{}p) \text{ in } F_p[a_4, a_6][x],}$$

so $u\hat{}\{(p-1)/2\}$ divides K and $4uK^2 = 4w\hat{}\{(p)\}(x\hat{}p) \cdot h'(x\hat{}p)^2$ is a polynomial in $x\hat{}p$.

Proposition 15 (S_1 is a derivative of ψ_p) [paper. *] For $p \ge 5$,*

$$\boldsymbol{\varphi_p \equiv 4u \cdot (\psi_p'/p)^2 + D(\psi_p^2) \pmod p, \text{ equivalently } S_1 = 2 \cdot D\ \psi_p,}$$

equivalently $A^2\Phi = 4w\hat{}\{(p)\}(\Psi')^2 + A^2 \cdot D(\Psi^2)$. The weight is rigid: $w(D\Psi) = p^2 + 1 = w(S_1 \text{ in } X)$, and there is no monomial of weight 2, so no correction term of the shape $c \cdot \Psi$ can be added.

The two propositions are what (ND) rests on. With them the resultant Res_x(G, ψ_p mod p) factors — not the resultant Res_X($\tilde{u}$, g) of §4.6(3), which is a different one — and can be counted at the cusp by the method of §4.6(3), which gives the theorem:

Theorem ND (the non-degeneracy holds always) [paper. *] For every prime $p \geq 5$ there is $c_S \in F_p\hat{}\times$ with*

$$\textbf{\textit{Res_x(}}S_1\textbf{\textit{, }}\psi\textbf{\textit{_p mod p) = c_S}}\cdot\Delta_0\hat{}\{m_S\}\textbf{\textit{ in F_p[}}a_4\textbf{\textit{, }}a_6\textbf{\textit{]}},\ m_S = p^2(p^2-1)/24,$$

and hence (ND) holds for ***every*** *elliptic curve over $\bar{F}_p$ and every $\bar{T} \in \bar{E}(F_p)\backslash\{O\}$: at an ordinary point the resultant is non-zero, so S_1 and ψ_p mod p have no common root, and at a supersingular point ψ_p mod p is a non-zero constant and (ND) is vacuous. Consequently* ***Theorem G′ and Theorem G″ are unconditional****, and so are Proposition R‴ and Proposition 10.*

Proposition 16 (the three constants in closed form) [paper. *] With $n_p = (p-1)/2$, $M = p(p^2-1)/24$, $m_K = (p^2-1)(p^2-3)/24$ and $m_p = m_K + m_S$, in F_p:*

$$\textbf{\textit{Res_X(D}}\Psi\textbf{\textit{, }}\Psi\textbf{\textit{)}} = \gamma\textbf{\textit{_p}}\cdot\Delta_0\hat{}\{M\},\ \gamma\textbf{\textit{_p}} = (-1)\hat{}\{n_p(n_p+1)/2\}\cdot n_p!\cdot(-16)\hat{}\{M\},\ c_S = 2\hat{}n_p\cdot\gamma_p,$$

$$\textbf{\textit{Res_x(K, }}\psi\textbf{\textit{_p mod p) = c_K}}\cdot\Delta_0\hat{}\{m_K\},\ c_K = \varepsilon\textbf{\textit{_p}}\cdot 2\hat{}\{m_K\}\textit{ with }\varepsilon\textbf{\textit{_p}} = -1 \iff p \equiv 3 \pmod 8,$$

and $c_p = (-1)\hat{}\{p(p-1)/2\}c_Kc_S$ for Res_x(G, ψ_p mod p) = $c_p\cdot\Delta_0\hat{}\{m_p\}$. The sign ε_p is the second supplementary law: it is (2/p) times, when $p \equiv 1 \pmod 4$, a second copy of (2/p).

The closed form for c_K, and with it the Legendre symbol, is the shape of the observation recorded in v5 for $p \leq 53$; it is now a consequence of Proposition 14. Note what Theorem ND does *not* say: it is a statement about the reduced curve, and none of the hypotheses of §3.2 or §3.10 are weakened by it.

3.15 Depth and height: an archimedean input

Every statement so far is local at p: the data are the layers of the filtration of E(Q_p) and the digits of Fermat quotients, and §7(3) records that the local data do not decide whether the Schur complement of Corollary R′_s is a unit. One global datum of a different kind does decide it in part, and it is archimedean: how deep a rational point can lie in the filtration is bounded by its naive height. For $Q \in E(\mathbb{Q})$, $Q \neq O$, write $h(x(Q)) := \log \max(|\alpha(Q)|, d(Q)^2)$ for the naive height of $x(Q) = \alpha/d^2$.

Lemma 23 (depth and height) [paper; the inequality in Lean. *] Let p be a prime of good reduction, the equation minimal at p, and $Q \in E_k(Q_p)$ with $k \geq 1$. Then $p\hat{}k \mid d(Q)$, and hence*

$$\textbf{\textit{h(x(Q))}} \geq \textbf{\textit{2k}}\cdot\textbf{\textit{log p}},$$

with no error term. Equivalently, $k \leq h(x(Q))/(2 \log p)$: the depth of a rational point in the formal filtration is bounded by its naive height.

Proof. For $Q \in E_1(Q_p)$ with $v_p(x(Q)) = -2m$ one has $v_p(y(Q)) = -3m$ and $v_p(t(Q)) = m$, so $Q \in E_k(Q_p)$ exactly when $p\hat{}k \mid d(Q)$; and $d^2 \leq \max(|\alpha|, d^2)$. ■ (Lean `log_denom_ge`, `height_ge_of_depth`, `depth_lt_of_height_lt`.)

The inequality has an exact form, which says where the slack goes.

Lemma 23′ (the slack) [paper; Lean. *] Write d(Q) = p^D·c′ with p ∤ c′. Then*

$$h(x(Q)) = 2 \log c' + 2D \cdot \log p + \max(0, \log|\alpha(Q)| - 2 \log d(Q)).$$

Hence, with s := h(x(Q)) − 2D·log p, one has c′ ≤ e^{s/2}; equality h(x(Q)) = 2D·log p holds if and only if c′ = 1 and |α(Q)| ≤ d(Q)²; and s < 2 log 2 forces c′ = 1.

Proof. h = max(log|α|, 2 log d) = 2 log d + max(0, log|α| − 2 log d), and 2 log d = 2D log p + 2 log c′. ■ (Lean `cofactor_log_le`, `cofactor_eq_one_of_slack_lt`, `max_eq_iff`.)

The slack s splits into the part of the denominator prime to p, 2 log c′, and an archimedean excess max(0, log|x(Q)|); §5.12 measures that the second saturates first. Both lemmas are elementary, and we regard them as known (§8).

Combined with Corollary K′(i), Lemma 23 gives an upper bound for v_p(Reg_p) that no local datum gives.

Proposition 24 (an upper bound from the shortest vector) [paper; the linear algebra in Lean. *] Let p ≥ 5 be good ordinary, #E(F_p) = p, p ∤ ∏_ℓ c_ℓ, E(Q) torsion-free, ι, and E split at p; write Λ = E(Q) and*

$$h_min := min\{\ h(x(Q)) : 0 \neq Q \in \Lambda \cap E_1(Q_p)\ \}.$$

***If h_min < 4·log p, then (χ, γ) : Λ → F_p² is onto**, hence ρ₁ = 2 (Corollary K′(i)).*

*In rank two this gives **v_p(Reg_p) = 0** (Corollary R′_s); in rank r ≥ 3 it gives v_p(Reg_p) = (r − 2) + v_p(det T_s) with T_s ≡ p·Ĉ (mod p²) (Corollary K′(iv)).*

Proof. χ is onto by ι. If (χ, γ) is not onto, its image is a line projecting onto the first factor, that is the graph of γ = cχ, so γ vanishes on ker χ = Λ ∩ E_1(Q_p), that is Λ ∩ E_1(Q_p) ⊆ E_2(Q_p). Then every non-zero Q ∈ Λ ∩ E_1(Q_p) has h(x(Q)) ≥ 4 log p by Lemma 23 with k = 2, against h_min < 4 log p. ■ (Lean `surjective_pair_iff`, `surjective_of_shallow_kernel_element`.) Whether h_min is taken over all of Λ ∩ E_1(Q_p) or over (Λ ∩ E_1(Q_p)) \ pΛ does not change the truth of h_min < 4 log p: in the split case pΛ ⊆ E_2(Q_p), so every non-zero element of pΛ has h(x(Q)) ≥ 4 log p by Lemma 23.

Three remarks place it. (a) *Torsion.* If #E(F_p) = p then the torsion of E(Q) prime to p injects into E(F_p), so p ∤ #E(Q)_tors already makes E(Q) torsion-free. (b) *The split hypothesis cannot be dropped.* In the non-split case E(Q_p) ≅ Z_p and pP has depth exactly one for every P with χ(P) ≠ 0, so Λ ∩ E_1 ⊆ E_2 never happens and the proposition has nothing to say; in the split case pΛ ⊆ E_2 automatically, and for r = 2 the quotient (Λ ∩ E_1)/pΛ has order p, so **(χ, γ) fails to be onto if and only if p² | d(Q) for one — equivalently every — Q ∈ (Λ ∩ E_1) \ pΛ**, a test with no p-adic computation in it. (c) *What it is and is not.* h_min is computed from the archimedean height pairing and an integer search; the condition is sufficient and not necessary, and it bounds v_p(Reg_p) from above by an archimedean datum where §7(3) says that no local datum does. Cut by the naive height of the proposition, the sweep of §5.12 shows that it accounts for **about 68 %** of an observed deficit in the frequency of det(χ, γ) ≡ 0 (207.8 of 305.35; 55.8 % on the second set of boxes). The canonical form below, Proposition 24ᶜ, accounts for **all of it in expectation** (100.6 %, and 99.7 % on the second set): where the deficit ends is explained by a theorem. That above the threshold the frequency is 1/p is an empirical regularity, not a consequence of either proposition.

The canonical height. Let the equation be a global minimal model and ĥ the canonical height in the normalisation of PARI (ĥ ≈ h(x)), so that ĥ = Σ_v λ′_v with λ′_v := 2(λ_v

+ (1/12) log|Δ|_v) (the terms in Δ cancel by the product formula). At the good prime p, $\lambda'_p(Q) = 2D(Q)\cdot\log p$ with $D(Q) = v_p(d(Q))$. Write $R(Q) := \hat{h}(Q) - 2D(Q)\cdot\log p = \Sigma_{v\neq p}\, \lambda'_v(Q) = 2\log c' + \lambda'_\infty(Q) + \Sigma_{\ell \text{ bad}}\, \mathrm{corr}_\ell(Q)\cdot\log \ell$, where $\mathrm{corr}_\ell(Q) \leq 0$ is the component correction (zero when Q has non-singular reduction at ℓ). This decomposition and the table of corrections are the known theory of local heights [known; [SilH]]; we use them and do not prove them. Put

$$\beta(E) := m_\infty + \Sigma_{\ell \text{ bad}}\, m_\ell\cdot\log \ell,\ m_\infty := \min_{E(R)\setminus O} \lambda'_\infty,\ m_\ell := \min \text{ over the components of } E(Q_\ell)/E_0(Q_\ell) \text{ of } \mathrm{corr}_\ell,$$

with $m_\ell = 0$ when $c_\ell = 1$, $-\lfloor N/2\rfloor\lceil N/2\rceil/N$ for split I_N, $-N/4$ for non-split I_N with N even, $-1/2, -2/3, -1, -4/3, -3/2$ for III, IV, I_0^*, IV*, III*, and for I_n^* the smaller of -1 and $-(n+4)/4$.

Proposition 24^c (the canonical threshold) [paper; the arithmetic and the logic in Lean. *] Under the hypotheses of Proposition 24 and with the equation globally minimal, let* $\hat{h}_{min} := min\{\, \hat{h}(Q) : 0 \neq Q \in \Lambda \cap E_1(Q_p) \,\}$.

If $\hat{h}_{min} < 4\cdot\log p + \beta(E)$***, then*** (χ, γ) ***is onto****, hence* $\rho_1 = 2$*, and in rank two* $v_p(Reg_p) = 0$.

Proof. $R(Q) \geq \beta(E)$ termwise: $2\log c' \geq 0$, $\lambda'_\infty \geq m_\infty$, $\mathrm{corr}_\ell \geq m_\ell$ (Lean `beta_le_remainder_of_local_bounds`). If (χ, γ) is not onto then, as in Proposition 24, every non-zero $Q \in \Lambda \cap E_1$ has $D(Q) \geq 2$, so $\hat{h}(Q) = 2D\log p + R(Q) \geq 4\log p + \beta(E)$ (Lean `canonical_ge_of_depth`), against the hypothesis. ■ (Lean `surjective_of_canonical_lt_four_log_add_beta`.) As for Proposition 24, taking $\hat{h}_{min}$ over all of $\Lambda \cap E_1$ or outside $p\Lambda$ does not change the truth of the hypothesis.

Remarks. (d) The two propositions are two sufficient conditions, neither implying the other: $\beta(E)$ can be of either sign (its median over the boxes of §5.12 is $-0.07\cdot\log p$), and 23 + 43 boxes of §5.12 lie below the naive threshold and above the canonical one. (e) The lower bound $R \geq \beta(E)$ holds for every reduction type with a bounded constant; it fails only if a hypothesis is dropped (p a bad prime, a non-minimal model, the component corrections omitted). (f) The local height formulas, the component corrections and the value of m_∞ are not in Lean; they enter the Lean statements as hypotheses. m_∞ is computed numerically, so the box-by-box application in §5.12 is [computation]. (g) The proof uses only that every non-zero $Q \in \Lambda \cap E_1$ has $R(Q) \geq \beta(E)$. Since corr_ℓ depends only on the component of Q at ℓ [known], $\beta(E)$ may be replaced by $\beta_\Lambda(E) := m_\infty + \min\{\, \Sigma_\ell\, \mathrm{corr}_\ell(\kappa)\cdot\log \ell : \kappa$ in the image of $\Lambda \cap E_1$ in $\prod_\ell \Phi_\ell \,\} \geq \beta(E)$, a finite computation; $\beta_\Lambda > \beta$ exactly when that image misses the minimal component at some ℓ. The Lean statements take a bound at each ℓ separately ($m_\ell \leq \mathrm{corr}_\ell$); β_Λ needs instead the joint bound $\Sigma_\ell\, \mathrm{corr}_\ell\cdot\log \ell \geq \beta_\Lambda - m_\infty$, a change of hypothesis that we have not made in Lean [paper].

What the threshold leaves open is the frequency of $\det(\chi, \gamma) \equiv 0$ above it. The next lemma says that the frequency 1/p is a local count, so that any departure from it is a correlation between the local class of the box and the global height.

Lemma 25 (the image modulo E_2) [paper. *] Let $p \geq 5$ be good, $\#E(F_p) = p$, ι, E split at p, and $\varphi : \Lambda \to E(Q_p)/E_2(Q_p)$ the reduction. Then (i)* $E(Q_p)/E_2(Q_p) \cong (Z/p)^2$*; (ii)* ***det*****$(\chi, \gamma) \equiv 0$ if and only if** $\varphi(\Lambda)$ **has order p***, and then* $\varphi(\Lambda)$ *is one of the p lines of* $(Z/p)^2$ *other than* E_1/E_2*; otherwise* φ *is onto; (iii) if* $\Lambda' \subseteq \Lambda$ *is a subgroup of index prime to p, then* $\varphi(\Lambda') = \varphi(\Lambda)$*. In particular the verdict* $\det(\chi, \gamma) \equiv 0$ *depends only on*

the images of the generators of any such Λ' in E(Q_p)/E_2(Q_p) $\cong$ E(Z/p^2Z) [known], so, for generators with p-integral coordinates, only on the equation and those coordinates modulo p^2.

Proof. (i) E(Z_p) is an extension of $\tilde{E}$(F_p) $\cong$ Z/p by E_1(Q_p), and E_1 $\cong$ Z_p with E_2 = pE_1 through the formal logarithm (p $\geq$ 3) [known]. So E(Q_p) is Z_p or Z_p $\oplus$ Z/p; in the first case E_1 = pE(Q_p) and the quotient by E_2 is cyclic of order p^2, in the second the quotient is $(Z/p)^2$, and the second is the split case. (ii) $\varphi(\Lambda \cap E_1) = \varphi(\Lambda) \cap E_1/E_2$, and since χ is onto, $\#\varphi(\Lambda) = p{\cdot}\#\varphi(\Lambda \cap E_1)$. As in the proof of Proposition 24, $\det(\chi, \gamma) \equiv 0$ exactly when $\Lambda \cap E_1 \subseteq E_2$, that is when $\varphi(\Lambda \cap E_1) = 0$. (iii) $\varphi(\Lambda)$ is a p-group and Λ/Λ' has order prime to p. ■

For the family of set C of §5.12 (two points with x-coordinates x_1 and $x_1 + 1$), counting the classes (x_1, y_1, y_2) modulo p^2 uniformly, the split classes with a_p = 1 have $\det(\chi, \gamma) \equiv 0$ in exactly 40 of 200 at p = 5 and 112 of 784 at p = 7, that is 1/p; and in the 6,818 boxes of set C where p does not divide the index of the two points in Λ, the verdict computed from the classes modulo p^2 agrees with the measured one in every box [computation]. Lemma 25 is elementary and should be regarded as known.

When the closure is small. If $\Lambda \cap E_1(Q_p) \subseteq E_3(Q_p)$, then the map $\Lambda \to E(Q_p)/E_3(Q_p)$ factors through $\Lambda/(\Lambda \cap E_1) \cong Z/p$ — no hypothesis on γ is needed for this — so its image has index p^2. If moreover $\gamma(P_1) = 0$ for one P_1 with $\chi(P_1) \neq 0$, then **every layer up to the third is a multiple of** χ: $\gamma \equiv 0$ and $\gamma_2 = c_2\chi$ on Λ (Lean `factor_through_chi`). That γ_2 is a homomorphism, which the Lean statement takes as a hypothesis, holds because $\gamma \equiv 0$ puts t_1 in p^2Z_p, where the formal group law is addition modulo p^4. By Theorem K, $\bar{M} = 2\lambda_\{T,1\}\chi\otimes\chi$; if $\lambda_\{T,1\} \neq 0$ then v_p(h_p(P)) = 0 at every P with $\chi(P) \neq 0$ (Lean `two_ne_zero_of_ne_two`, `diag_ne_zero`), and in rank two $\rho_1 = 1$ and v_p(Reg_p) $\geq$ 1. [paper.] The two boxes of this kind that §5.12 finds, 328563a1 and 477950bg1 at p = 7, have $\rho_1 = 1$ and v_p(Reg_p) = 1 [computation]. Such a box cannot realise the necessity of γ_3 in Theorem G″ on its rational points, since no point with $\chi(P) \neq 0$ has v_p(h_p(P)) $\geq$ 1 there; the realisation of §3.13 uses Proposition 21 instead.

4. Proofs

Remark (the ledger is not an induction). *The congruences of Theorem N and of Corollaries $N^{(3)}$–$N^{(7)}$ are not statements proved one after another: they are the p-adic expansion, digit by digit, of the single exact identity*

$$p^2{\cdot}n^2 h_p(P) = h_p(P'') = 2\,\log_p(\sigma_p(t)/t) + 2\,\log_p(\varphi/\omega) - 2p^2{\cdot}\log_p(d'), \quad t = t(P''),$$

in which $\varphi = \varphi_p(x(P'))$, $\omega = \omega_p(P')$. Theorem N is this identity summed rather than expanded, and Proposition 18 is the table below in closed form. What is visible at digit k of n^2h_p — that is, modulo p^k, equivalently modulo p^{k+2} in h_p(P″) — are exactly the terms of valuation $\leq k + 1$, and the table of valuations settles which those are (e = v_p(t) $\geq$ 2, model with p | a_1): | term | source | valuation | first digit k | |—|—|—|—| | $p^2\Theta$ | log_p(φ/ω) | 2 | 0 | | p^3F | denominator | 3 | 1 | | $p^4\Theta^2$, $p^6\Theta^3$ | log_p(φ/ω), Q_2 | 4, 6 | 2, 4 | | (−1)^{j+1}p^{2j}Θ^j/j, j $\geq$ 2 | log_p(φ/ω) | **2j − v_p(j)** | 2j − 2 − v_p(j) | | p^4F^2, p^5F^3 | denominator | 4, 5 | 2, 3 | | 2(−1)^{j+1}p^{2+j}F^j/j, j $\geq$ 2 | denominator | **2 + j − v_p(j)** | j − v_p(j) | | $a_1t + a_1\omega\psi/\varphi^2$ | σ, Weierstrass | **cancel** (remainder $\geq$ 2v_p(a_1) + 2e $\geq$ 6) | — | | the t^2 terms, combined: $\mu_2 z^2 = -c{\cdot}z^2$ | σ (t^2), η (a_2), the a_1*

remainders | 2e | 2 (for e = 2) | | c_4t^3, $a_3\omega\psi^3/\varphi^3$ | σ (t^3), Weierstrass | 3e, and ***no new invariant*** *(Proposition 17) | — | | μ_{2m}z^{2m}, m ≥ 2 | $\mathcal{L}$ (Proposition 17) |* ***2me + v_p(μ_{2m})*** *| 2me − 2 + v_p(μ_{2m}) | | $a_4\psi^4/\varphi^2$, $a_6\psi^6/\varphi^3$ | Weierstrass (inside log(1 + R), hence inside $\mathcal{L}$) | 4e, 6e | 6, 10 | Two features of the table are corrected and generalised by Proposition 18, and the corrections matter from the fifth digit on. First,* ***the j-th term of a logarithm is one digit shallower than p^j whenever p | j****: the factor v_p(j) is what brings (2/5)p^5F^5 into the fifth digit at p = 5 (Corollary $N^{(5)}$) and what makes the depth of a term depend on p. Second,* ***the individual coefficients of σ_p are not the right unit of account****: by Proposition 17 the combination that survives is the even series $\mathcal{L}$, whose m-th coefficient μ_{2m} is an invariant of E. So the coefficients of σ_p of odd index bring in nothing — the term c_4t^3 of the row above cancels against $a_3\omega\psi^3/\varphi^3$ and a reparametrisation of μ_2z^2 — and the first local invariant after c is $\mu_4 = -b_4/12$, at the* ***seventh*** *digit, not the fifth. The denominator descends at rate p^{k−v_p(k)} and $\mathcal{L}$ at rate p^{2me−2+v_p(μ_{2m})}: the two non-local ingredients of Lemma 3 arrive at different speeds, which is one answer — a structural rather than an intrinsic one — to the question of §7 about q_p(d′).*

4.11 Lemma 7, Proposition 8, Theorems K and N, Propositions 9 and 10

Lemma 7. Translation τ_P is an automorphism of the smooth model over Z_p and τ_P(O) = P has non-zero reduction, so x is regular at P and x ∘ τ_P ∈ Ô_{E,O} = Z_p[[t]]. The linear coefficient is read from the invariant differential: $\omega = dx/(2y + a_1x + a_3)$ is translation-invariant and $\omega = (1 + a_1t + \cdots)dt$, so $d(x \circ \tau_P) = \tau_P^*(dx) = (2y + a_1x + a_3)(\tau_P(\cdot))\cdot\omega$ and the coefficient of t at t = 0 is $2y(P) + a_1x(P) + a_3$. Finally $2y + a_1x + a_3$ is divisible by p only at $\bar{P} \in \bar{E}[2]$, and $\#\bar{E}(F_p) = p$ is odd, so only at $\bar{P} = O$. For κ_j apply this with P = n e_1 and R = n e_j ∈ E_1(Q_p), t = t(n e_j) of valuation ≥ 1; the first digit of the difference is $c_1\cdot(t/p) \bmod p = c_1\cdot\gamma(n\ e_j)$. ■

Proposition 8. Corollary H″, applied to the multiplication [m] with p ∤ m, gives F(mQ) = m^2F(Q) for F(Q) := λ^{(2)}(x(Q)) — Theorem H one digit deeper. By Proposition R‴ and Lemma 7, for Q ∈ E(Z_p) with $\bar{Q} \neq O$,

$$F(Q) = \lambda_T(\chi(Q)) + s(\chi(Q))\cdot c_1(\chi(Q))\cdot\gamma(Q),$$

since $x(Q) = x(\chi(Q)T_0) + p\cdot c_1\gamma(Q) + O(p^2)$ and λ^{(2)} depends on the lift only through its first digit. On the coset χ = 1 the value γ(Q) runs over all of F_p, because t : E_1(Q_p) → pZ_p is a bijection. Under Q ↦ mQ one has χ ↦ mχ and γ ↦ mγ, so comparing the two sides of F(mQ) = m^2F(Q) as affine functions of γ(Q) gives λ_T(m) = m^2λ_T(1) for the constant terms and $m\cdot s(m)c_1(m) = m^2\cdot s(1)c_1(1)$ for the linear ones. ■

Theorem K. Proposition R″_s gives $h_p(P) \equiv 2\lambda^{(2)}(x(nP))/n^2 = 2F(nP)/n^2 \pmod p$. Insert the display of the previous proof with χ(nP) = nχ(P), γ(nP) = nγ(P) and then (L1), (L2):

$$F(nP) = \lambda_{T,1}n^2\chi(P)^2 + B_0\cdot n\chi(P)\cdot n\gamma(P) = n^2(\lambda_{T,1}\chi(P)^2 + B_0\chi(P)\gamma(P)),$$

and n^2 cancels. Polarising gives the bilinear form; the Lean statements `master`, `master_symm`, `master_add_left`, `master_smul_left` record its shape and bilinearity. ■

Corollary K′. (i)–(iii) are linear algebra over F_p on the form χ⊗χ and χ⊗γ + γ⊗χ (Lean `master_mem_radical`, `master_eq_of_gam_eq_smul`, `master_offdiag_ne_iff`,

`master_diag_eq_zero_iff`, `master_bad_class_unique`). For (iv) take a basis e_1 with $\chi(e_1) = 1$, $e_2 \in \ker \chi$ with $\gamma(e_2) \neq 0$, and a basis of Λ_1; then M_Λ has the shape [[a, b, 0 …], [b, pC$_{22}$, p(∗)], [0, p(∗), pĈ]] with b ∈ Z_p^× by Theorem K, the Schur complement of the leading 2×2 block is p·Ĉ modulo p^2, and Corollary R′_s applies (Lean `det_three_layer`, `det_sq_dvd_iff`). ■

Proposition 9. By Lemma 5, φ_p′ = p·G with G ∈ Z[a_1, …, a_6][x], and φ_p″ = p·H. Taylor at x_0 with h = pk gives

$$\varphi_p(x_0 + pk) \equiv \varphi_p(x_0) + p^2 \cdot k \cdot G(x_0) + p^3 \cdot k^2 \cdot H(x_0)/2 \pmod{p^4},$$

the cubic term being $p^4k^3 \cdot (H'/6)(x_0)$ with p ∤ 6 (Lean `eval_shift_second_digit`, in the integral form $2(P(r+pk) - P(r)) - 2p^2kG(r) - p^3k^2H(r) \equiv 0 \bmod p^4$). Write ε for the correction, so $p^2 \mid \varepsilon$, and u := φ_p(x$_0$), u^{p−1} = 1 + $p^2\Theta(x_0)$. Then (u + ε)^{p−1} ≡ u^{p−1}(1 + (p−1)ε/u) (mod p^4), because ε^2 is divisible by p^4 (Lean `fermatQuot_second_move`), whence

$$(u + \varepsilon)^{p-1} - 1 \equiv p^2\Theta(x_0) + (p-1)\varepsilon/u \pmod{p^4}.$$

Dividing by p^2 and using (p−1)p ≡ −p (mod p^2) gives the displayed formula. ■

Proposition 10. By Proposition 9 the function $\delta \mapsto \Theta(x(T) + p\delta)$ is, modulo p, the affine map Θ_T + 2sδ with s ≠ 0 by §4.6 — the quadratic term carries a factor p. An analytic function Z_p → Z_p whose reduction modulo p is a bijection of F_p and whose derivative is a unit at every point is a bijective isometry of Z_p, by Hensel's lemma applied at each residue; in particular 0 is attained exactly once. The equality of valuations is the factorisation f(δ) − f(δ_) = (δ − δ_)·(b + p·q) with b = 2s a unit, which is Lean `root_factor` and `dvd_iff_dvd_sub_root`; the uniqueness of the bad class at each layer without assuming a root is `unique_bad_class`, and its existence is `exists_bad_class`, one step of Hensel written by hand. ■

Lemma 3′. u(P) = σ_p(t)/t = 1 + c_2t + $c_3$$t^2$ + ⋯ with $c_2 = a_1/2$ and v_p(t) = e, so v_p(log_p u) ≥ min(v_p(c_2) + e, 2e). For e = 1 and p | a_1 this is 2 = 2e; for e ≥ 2 the first term is the smaller one. ■

Theorem N. Work in a global minimal model with p | a_1 (§2), and put P′ := nP, P″ := pP′, e := v_p(d(P″)) ≥ 2 (Lemma 4). Four terms have to be carried modulo p^4, where before only their leading behaviour was needed.

(1) *The formal part.* By Lemma 3′, 2Λ_σ ≡ a_1·t(P″) (mod p^4), of valuation v_p(a_1) + e ≥ 3; the next term is $2c_3t^2$ of valuation ≥ 2e ≥ 4.

(2) *The Weierstrass equation.* Multiplying at P″ by ψ_p(P′)^6 and writing φ = φ_p(P′), ω = ω_p(P′), ψ = ψ_p(P′) with v_p(ψ) = e and φ, ω units,

$$\omega^2 + a_1\varphi\omega\psi + a_3\omega\psi^3 = \varphi^3 + a_2\varphi^2\psi^2 + a_4\varphi\psi^4 + a_6\psi^6,$$

the terms $a_3\omega\psi^3$, $a_4\varphi\psi^4$, $a_6\psi^6$ having valuation ≥ 3e ≥ 6 and $a_2\varphi^2\psi^2$ valuation ≥ 2e ≥ 4. Hence $\omega^2 \equiv \varphi^3 - a_1\varphi\omega\psi \pmod{p^4}$. Raising to the power p − 1 and writing φ^{p−1} = 1 + p^2A, ω^{p−1} = 1 + p^2B with A = Θ(x(P′)), one gets $2B \equiv 3A + a_1\omega\psi/(p^2\varphi^2) \pmod{p^2}$, so

$$Q_2(\varphi/\omega) = A - B \equiv -\Theta/2 - a_1\omega\psi/(2p^2\varphi^2) \pmod{p^2}.$$

(3) *The denominator.* By Step 1 of §4.4, $\alpha/\beta = (\varphi_p(P')/\omega_p(P'))\cdot d'^{\{-p^2\}}$, and $(1 + p\,q)^{\{p^2\}} \equiv 1 + p^3 q \pmod{p^4}$ for $p \geq 5$, so $Q_2(d'^{\{p^2\}}) \equiv p\cdot q_p(d') \pmod{p^2}$; equivalently $-p^2 \log_p(d') \equiv p^3 q_p(d') \pmod{p^4}$. **This is the term that Proposition R″_s could drop and Theorem N cannot.**

(4) *The logarithm.* For a unit u with $q_p(u) = 0$, $(p-1)\log_p\, u = \log(1 + p^2 Q_2(u)) \equiv p^2 Q_2(u) \pmod{p^4}$, and $1/(p-1) \equiv -(1 + p) \pmod{p^2}$, so $\log_p\, u \equiv -p^2(1 + p)Q_2(u) \pmod{p^4}$.

Putting (2), (3), (4) into $A = \log_p(\alpha/\beta)$ and adding (1),

$$h_p(P'') = 2\Lambda_\sigma + 2A \equiv a_1 t(P'') + (1 + p)(p^2\Theta + a_1\omega\psi/\varphi^2) + 2p^3 q_p(d') \pmod{p^4}.$$

The two a_1 terms cancel: $t(P'') = -x(P'')/y(P'') = -\varphi\psi/\omega$, so $a_1 t(P'') + a_1\omega\psi/\varphi^2 = a_1\psi\cdot(\omega/\varphi^2 - \varphi/\omega) = a_1\psi\cdot(\omega^2 - \varphi^3)/(\varphi^2\omega)$, which by (2) is $-{a_1}^2\psi^2/\varphi + O(p^4)$ of valuation $\geq 2v_p(a_1) + 2e \geq 6$; and the surviving $p\cdot a_1\omega\psi/\varphi^2$ of $(1 + p)$ has valuation $\geq 1 + 1 + e \geq 4$. Hence

$$h_p(P'') \equiv (1 + p)p^2\Theta(x(P')) + 2p^3 q_p(d') \pmod{p^4},$$

and $h_p(P'') = p^2 n^2 h_p(P)$ gives the statement. The cancellation is the Lean statement `a1_correction_cancels`; the two-digit criteria that Theorem N feeds are `second_digit_zero_iff` and `second_digit_criterion`. ■

Corollary N′. Write $\Theta = p\Theta_1$. By Theorem N, $n^2 h_p(P) \equiv p(\Theta_1 + 2q_p(d')) \pmod{p^2}$, and $p \nmid n$. ■

4.12 Corollaries N(3) and N(4), Theorem G′, Propositions 11, 12 and 19, Corollaries N″ and N‴

Throughout, the model has $p \mid a_1$ (§2), $P' = nP$, $P'' = pP'$, $e = v_p(t(P'')) \geq 2$, $\varphi = \varphi_p(x(P'))$, $\psi = \psi_p(x(P'))$, $\omega = \omega_p(P')$, and the identity of the Remark opening §4 is expanded one digit at a time. Only the terms newly reached are discussed; the rest is §4.11.

The coefficient of t^2 in σ_p. Write $\sigma_p(t)/t = 1 + c_2 t + c_3 t^2 + \cdots$ and $g_1 := 2c_3 - {c_2}^2$. From the defining differential equation $x(t) + c = -(d/\omega)((1/\sigma)(d\sigma/\omega))$ with $D := w^{\{-1\}}d/dt$, $w = 1 + a_1 t + ({a_1}^2 + a_2)t^2 + \cdots$ and $g := \sigma'/\sigma = t^{\{-1\}} + g_0 + g_1 t + \cdots$ (so $g_0 = c_2 = a_1/2$), one has $(g/w)' = -w\cdot(x + c)$. Now $w\cdot(x + c) = t^{\{-2\}} + 0\cdot t^{\{-1\}} + c + (a_3 + a_1 c)t + \cdots$ — the vanishing of the $t^{\{-1\}}$ coefficient is what makes σ_p odd — so $g/w = t^{\{-1\}} + C - ct - (a_3 + a_1 c)t^2/2 + \cdots$, while $1/w = 1 - a_1 t - a_2 t^2 - \cdots$ gives for the coefficient of t^1 in g/w the value $g_1 - a_1 g_0 - a_2$. Comparing,

$$\mathbf{g_1 = a_1 g_0 + a_2 - c = {a_1}^2/2 + a_2 - c},\ c = (b_2 - E_2(E, \omega))/12,$$

and the t^0 coefficient $g_0 - a_1 = C = -a_1/2$ is consistent with $c_2 = a_1/2$.

Corollary N(3). Four stages, each one digit deeper than §4.11.

(1) *The formal part.* $2\Lambda_\sigma = a_1 t + g_1 t^2 + O(t^3)$ with $v_p(a_1 t) \geq 1 + e \geq 3$, $v_p(g_1 t^2) \geq 2e \geq 4$ and $v_p(t^3) \geq 3e \geq 6$, so modulo p^5 the t^2 term survives and the t^3 term does not.

(2) *The Weierstrass equation.* Modulo p^5 the surviving corrections are $a_1\varphi\omega\psi$ and $a_2\varphi^2\psi^2$, so $\eta := (\omega^2 - \varphi^3)/\varphi^3 = -a_1\omega\psi/\varphi^2 + a_2\psi^2/\varphi + O(p^6)$ with $v_p(\eta) \geq 3$. Writing $\varphi^{\{p-1\}} = 1 + p^2\Theta$ and $\omega^{\{p-1\}} = 1 + p^2 B$ and expanding $(1 + p^2 B)^2 = (1 + p^2\Theta)^3(1 + \eta)^{\{p-1\}}$

modulo p^6 (η^2 has valuation ≥ 6) gives $2B + p^2B^2 \equiv 3\Theta + 3p^2\Theta^2 + (p-1)\eta/p^2 + 3(p-1)\Theta\eta \pmod{p^4}$, hence $B \equiv 3\Theta/2 + (3/8)p^2\Theta^2 + (p-1)\eta/(2p^2) \pmod{p^3}$ (Lean `sq_cube_step`, in the integral form $8B \equiv 12\Theta + 3p^2\Theta^2 + 4E$).

(2′) *Q_2 is not additive on quotients.* From $(1 + p^2B)(1 + p^2Q) = 1 + p^2\Theta$ with $Q := Q_2(\varphi/\omega)$,

$$\mathbf{Q = \Theta - B + p^2B(B - \Theta) + O(p^4)}$$

(Lean `q2_quotient`, as the exact identity $Q - (\Theta - B + p^2B(B - \Theta)) = p^4B^2Q$), so with $B \equiv 3\Theta/2 \pmod p$ one gets $Q \equiv -\Theta/2 + (3/8)p^2\Theta^2 - (p-1)\eta/(2p^2) \pmod{p^3}$. **This stage is what an induction on digits omits**: the second digit only needed $Q \equiv \Theta - B$.

(3) *The denominator.* $-2p^2\log_p(d') \equiv 2p^3(1 + p)F - p^4F^2 \pmod{p^5}$, from $1/(p-1) \equiv -(1 + p + p^2) \pmod{p^3}$ and the quadratic term of $\log(1 + pF)$ (Lean `denominator_third_digit`, `neg_inv_pred_cube`).

(4) *The logarithm.* $2\log_p(\varphi/\omega) \equiv -2p^2(1 + p + p^2)Q + p^4Q^2 \pmod{p^5}$. Inserting Q from (2′): the Θ term is $p^2(1 + p + p^2)\Theta$; the Θ^2 terms are $-(3/4)p^4\Theta^2$ from Q and $+(1/4)p^4\Theta^2$ from p^4Q^2, total $-(1/2)p^4\Theta^2$; the η term is $(1 + p + p^2)(p-1)\eta = (p^3 - 1)\eta \equiv -\eta \pmod{p^5}$.

Finally $-\eta = a_1\omega\psi/\varphi^2 - a_2\psi^2/\varphi$, and $a_1t(P'') + a_1\omega\psi/\varphi^2 = a_1\psi(\omega^2 - \varphi^3)/(\varphi^2\omega) = -{a_1}^2\psi^2/\varphi + O(p^7)$ has valuation $\geq 2v_p(a_1) + 2e \geq 6$, so the two a_1 terms cancel modulo p^5 as they did modulo p^4. What is left is $g_1t^2 - a_2\psi^2/\varphi$, and $t^2 = \varphi^2\psi^2/\omega^2 \equiv \psi^2/\varphi \pmod{p^7}$, so it equals $(g_1 - a_2)\psi^2/\varphi = ({a_1}^2/2 - c)\psi^2/\varphi \equiv -c\cdot\psi^2/\varphi \pmod{p^6}$ when $p \mid a_1$. Since $\psi^2/\varphi = 1/x(P'')$, dividing $h_p(P'')$ by p^2 gives Corollary $N^{(3)}$. The intermediate constant ${a_1}^2/2 - c$ belongs to this computation in this model and **not to the statement**: the model-free identity behind the same step is $2c_3 - a_2 - (3/4){a_1}^2 = -c$ in the coefficients c_3, c_4 of $\sigma_p(t)/t$ (Lean `sigma_third_coeff`, which writes c_3 as g_1; its companion `sigma_fourth_coeff` is $a_3 - 2c_4 = -a_1({a_1}^2/2 + a_2 + E_2/8)$), the three further contributions being $-a_2/X$, $({a_1}^2/2)(Y/X^2)^2$ and the a_1 remainder $-{a_1}^2/X$. Both identities are Proposition 17 at $m = 1$ read in the parameter t instead of z. ■

Corollary $N^{(4)}$. The same four stages one digit deeper. What is new: (a) $1/(p-1) \equiv -(1 + p + p^2 + p^3) \pmod{p^4}$ (Lean `inv_pred_quartic`); (b) the cubic term of the logarithm of the denominator, $-2p^2\log_p(d') \equiv 2p^3(1 + p + p^2)F - p^4(1 + p)F^2 + (2/3)p^5F^3 \pmod{p^6}$ (Lean `denominator_fourth_digit`, multiplied by 6 to stay in Z); (c) the two contributions of $\Theta\eta$ — one from the term $(1/4)(p-1)\Theta\eta$ of Q and one from $p^4(1 + p)Q^2$ — cancel, the remainder having valuation ≥ 7 (Lean `theta_eta_cancels`); (d) everything of valuation $3e \geq 6$ dies: the t^3 coefficient of σ_p, the term $a_3\omega\psi^3$ of the Weierstrass equation, and $p^6\Theta^3$ (Lean `cubic_term_vanishes`). No mixed term ΘF appears. ■

Corollaries N'' and N'''. With $\Theta = p\Theta_1$ and $\Theta_1 + 2F = pS$, Corollary $N^{(3)}$ collapses to $n^2h_p \equiv p^2(S - F^2 - W) \pmod{p^3}$, which is Corollary N'' (Lean `cube_dvd_sub_crit`, `cube_dvd_iff_crit`, `quadrichotomy`). With $S - F^2 - W = pU$, Corollary $N^{(4)}$ collapses to $6\cdot n^2h_p \equiv p^3(6U + 6(S - F^2) + 4F^3) \pmod{p^4}$ and $p \nmid 6$, which is Corollary N''' (Lean `fourth_digit_reduction`, `quartic_dvd_iff_crit`, `quintichotomy`). ■

Proposition 12. Good ordinary reduction means the formal group has height one, so $[p](t) \equiv (\text{unit})\cdot t\hat{}p \pmod p$ and the coefficients c_i of $[p](t) = pt + \Sigma_\{i\geq 2\}c_it\hat{}i$ are divisible by p for $2 \leq i < p$. With $t_1 = p\gamma + p^2\gamma_2 + \cdots$ this gives $v_p(c_it_1\hat{}i) \geq 1 + 2 = 3$ for $2 \leq i < p$ and $v_p(c_pt_1\hat{}p) \geq p \geq 5$, hence $t(P'') = [p](t_1) \equiv p^2\gamma \pmod{p^3}$ (Lean `low_terms_vanish`, `p_mult_second_digit`); in particular $e = v_p(t(P''))$ equals 2 exactly when $\gamma \not\equiv 0$. Then $1/x(P'') = t(P'')^2(1 + a_1t(P'') + \cdots)$ with $v_p(a_1t^3) \geq 6$ (Lean `inv_x_correction`), so $W = c\cdot t(P'')^2/p^4 \equiv c\cdot\gamma^2 \pmod p$ (Lean `w_local`, `w_ne_zero_iff`). ■

Proposition 11. Lemma 7 read to order three: $x \circ \tau_T \in Z_p[[t]]$, and $\delta = (x(nP) - x(T))/p = (x \circ \tau_T(t_1) - x(T))/p$ with $t_1 = p\Gamma + O(p^4)$. Substituting and dividing by p gives the display, the coefficients being in Z_p. ■

Theorem G′. Four steps.

(1) *The x-coordinate moves by exactly p^j.* By Lemma 7, $X := x \circ \tau_T \in Z_p[[t]]$ and $X'(0) = c_1(T) = 2y(T) + a_1x(T) + a_3 \in Z_p^\times$; since $X' \in Z_p[[t]]$ and $t_1 \in pZ_p$, $X'(t_1) \equiv X'(0) \pmod p$ is again a unit. Writing $X(t_1 + h) - X(t_1) = \Sigma_\{k \geq 1\} b_k h^k$ with $b_k \in Z_p$ the coefficients of the shifted series and $b_1 = X'(t_1)$, the term $k = 1$ has valuation j and the terms $k \geq 2$ valuation $\geq 2j > j$. Hence $v_p(X(t_1 + p^j) - X(t_1)) = j$.

(2) *All divided derivatives of φ_p below the p-th are divisible by p.* By §2, $\varphi_p(x) \equiv f(x^p) \pmod p$ with $f \in F_p[a_1, \ldots, a_6][X]$; hence, as polynomials in two variables over F_p,

$$\varphi_p(x + y) \equiv f((x + y)^p) = f(x^p + y^p) \equiv \varphi_p(x) + (\text{terms in } y^p \text{ only}),$$

so the coefficient of y^k in $\varphi_p(x + y)$ — which is $\varphi_p^{(k)}/k!$, a polynomial with coefficients in $Z[a_1, \ldots, a_6]$ — is divisible by p for every $1 \leq k \leq p - 1$. For $k = 1$ this is Lemma 5, $\varphi_p' = p \cdot G$.

(3) *The value of φ_p moves by exactly p^{j+1}.* Put $x_0 := X(t_1)$ and $\Delta := X(t_1 + p^j) - X(t_1)$, so $v_p(\Delta) = j \geq 1$ by (1). By (2),

$$\varphi_p(x_0 + \Delta) - \varphi_p(x_0) = p \cdot G(x_0) \cdot \Delta + p \cdot \Delta^2 \cdot B + \Delta^p \cdot C \text{ with } B, C \in Z_p,$$

B collecting the divided derivatives of orders $2 \leq k \leq p - 1$ divided by p and C those of order $\geq p$. The three valuations are $j + 1$, $\geq 2j + 1 \geq j + 2$ and $\geq pj \geq 5j \geq j + 4$, and $G(x_0) \equiv G(x(T)) \pmod p$ is a unit by (ND). Hence the difference has valuation exactly $j + 1$ (Lean `shift_depth_exact`; the inequality half, which needs no non-degeneracy, is `shift_depth_dvd`).

(4) *Raising to the power $p - 1$ and dividing by p^2.* $\Phi := \varphi_p(x_0)$ is a unit (Corollary D), and $\Phi' = \Phi + D$ with $v_p(D) = j + 1 \geq 2$ by (3). From $(\Phi + D)^{p-1} - \Phi^{p-1} = (p - 1)\Phi^{p-2}D + D^2C'$ (Lean `pow_sub_pow_expand`) and $v_p(D^2) \geq 2(j + 1) \geq j + 3$, and since $p \nmid (p - 1)\Phi^{p-2}$, the left-hand side has valuation exactly $j + 1$ (Lean `unit_pow_diff_exact`). Dividing by p^2, which is what Q_2 does, gives $v_p(\Theta(t_1 + p^j) - \Theta(t_1)) = j - 1$ (Lean `quotient_by_p_sq`, `theta_sensitivity_exact`; the determination half is `theta_digit_determined`, the case $j = 3$ is `three_layers_determine` and its sharp companion `layer_three_moves_digit_two`). ■

On (ND), and why the same statement for ψ_p is free. Consider $\Psi := \psi_p \circ X \in Z_p[[t]]$. Since $[p](T + R(t)) = R([p](t))$, the identity $x(mQ) = \varphi_m/\psi_m^2$ at $m = p$ reads

$$\varphi_p(X(t)) = x(R([p](t))) \cdot \Psi(t)^2 = [p](t)^{-2} \cdot u([p](t)) \cdot \Psi(t)^2,\ u \in Z_p[[s]],\ u(0) = 1,$$

and with $[p](t) = pt \cdot M(t)$, $M(0) = 1$, and $\Psi(t) = t \cdot \Psi_1(t)$ this becomes $\varphi_p(X(t)) = p^{-2} M(t)^{-2} u([p](t)) \Psi_1(t)^2$. At $t = 0$, $\varphi_p(x(T)) = p^{-2}\Psi_1(0)^2$, and $\varphi_p(x(T))$ is a unit (Corollary D), so $v_p(\Psi_1(0)) = 1$. Since $\Psi_1(0) = \Psi'(0) = \psi_p'(x(T)) \cdot c_1$ with c_1 a unit,

$v_p(\psi_p'(x(T))) = 1$ exactly — no hypothesis needed.

For φ_p the same route gives nothing: the invariant differential $[p]\hat{}*\omega = p\omega$ says, after clearing denominators, $G\cdot\psi_p - 2\varphi_p\cdot K = B$ with $K := \psi_p'/p$ and B a rational function regular at x(T), and at $x = x(T)$, where ψ_p vanishes, this determines K(x(T)) and not G(x(T)); differentiating once more determines K′(x(T)) and again not G(x(T)), because G enters only in the product $p\cdot G\cdot K$. So (ND) is a statement about φ_p modulo p^2, one order beyond what the invariant differential sees. That extra order is exactly what Proposition 14 supplies — the p^2 digit of ψ_p is a function of the p digit — and §4.13 turns it into a proof of (ND) for every p (Theorem ND). It is the same non-degeneracy that Propositions R′′′′ and 10 use, so those are unconditional as well.

Remark (Theorem N is a statement about a p-minimal model). *Scaling the model by $x' = u^2x$, $y' = u^3y$ with $p \nmid u$ multiplies $\varphi_p(x)$ by $u\hat{}\{2p^2\}$ by the weights of the division polynomials, so*

$$\Theta \mapsto \Theta + 2p\cdot q_p(u) \pmod{p^2},$$

while d' is unchanged if $\gcd(u, d') = 1$. As $n^2h_p(P)$ is intrinsic, the right-hand sides of Theorem N and Corollaries $N^{(3)}$, $N^{(4)}$ depend on the model, and those theorems are statements about a model minimal at p. This is a partial answer to the question of §7 about $q_p(d')$: looking for an intrinsic meaning of $q_p(d')$ by itself is the wrong request, since it is paired with the normalisation of the model — with the normalisation of σ. The third-digit term $c/(p^2x(P'))$ is, by contrast, invariant under scaling as well.

Theorem N and Propositions 17 and 18.* Proved where they are stated (§3.11). Three remarks on how they stand to the four stages above. (i) Stages (2), (2′) and (4) — the Weierstrass equation, the failure of additivity of Q_2 on quotients, and the expansion of the logarithm — are, in the closed form, the single step $\log(\varphi/\omega) = -(1/2)(\log\varphi + \log(1 + R))$; the carries that an induction on digits has to track are the carries of the Iwasawa logarithm, and they are not tracked twice. (ii) Stage (1) and the cancellation of a_1 are, in the closed form, the statement that $\mathcal{L}$ is even: the odd powers of z are exactly the terms that the model with $p \mid a_1$ was chosen to suppress, and Proposition 17 suppresses them in every model. (iii) What the closed form does not contain is any claim about which terms a *later* digit can produce; that is Proposition 18, whose proof is an inspection of the three series and not of the expansion, and which is why the caveat of §8 on the valuation table is weaker in v8 than the table itself would suggest.

Proposition 19. $T := \chi(P)T_0$ reduces to the same point of $E(F_p)$ as $P' = nP$, by the definition of χ, so $P' - T \in E_1(Q_p)$ and $t_1 := t(P' - T) \in pZ_p$. Since $pT = O$, $P'' = pP' = p(P' - T)$. The formal logarithm is a homomorphism $E_1(Q_p) \to Q_p$ with $\log_E([p](t)) = p\cdot\log_E(t)$, whence $z(P'') = p\cdot\log_E(t_1)$. For $t_1 \in pZ_p$ one has $v_p(\log_E(t_1)) = v_p(t_1)$, so $e = v_p(t(P'')) = v_p(z(P'')) = 1 + v_p(t_1)$ (Lean `layer_depth`). ■

4.13 Lemma 13, Propositions 14, 15, 16 and Theorem ND

Lemma 13. Write (∗) $P'h - 2Ph' = c\Omega$ and (∗∗) $u_\{E'\}\Omega^2 = P^3 + a_4Ph^4 + a_6h^6$; both come from $x\circ\lambda = P/h^2$, $y\circ\lambda = y'\Omega/h^3$ and $\lambda\hat{}*\omega_E = c\cdot\omega_\{E'\}$ by clearing denominators in $\omega_E = dx/2y$. At a root ξ of h, (∗) gives $\Omega(\xi) = -2P(\xi)h'(\xi)/c$, and (∗∗) then gives $c^2P(\xi) = 4u_\{E'\}(\xi)h'(\xi)^2$. Differentiating (∗) and (∗∗) once and evaluating at ξ gives $-P'(\xi)h'(\xi) - 2P(\xi)h''(\xi) = c\Omega'(\xi)$ and $u'\Omega^2 + 2u\Omega\Omega' = 3P^2P'$ at ξ; eliminating Ω' and dividing by $P(\xi)^2$ gives the second identity. No formal group and no Tate curve are needed. ■

Proposition 14. Both sides are reductions of elements of $Z[a_4, a_6][x]$, so it suffices to prove the identity at every ordinary point, the ordinary locus being dense and open. Lift $\bar{E}$ to a

curve E over a complete discrete valuation ring R with residue field F, v(p) = 1, ordinary. Let C := $\hat{\mathrm{E}}$[p] ⊂ E[p] be the connected component — the canonical subgroup, of order p. Its non-zero points have v(x) = −2/(p−1) while the remaining p^2 − p points of E[p] have v(x) ≥ 0, so ψ_p = λ·M·N with M := p·(kernel polynomial of C) of degree n_p, N of degree pn and λ ∈ R^×; the valuations of the roots put M ∈ R[x], and the leading coefficient of N is a unit, so the whole drop in degree from $(\mathrm{p}^2-1)/2$ to p(p−1)/2 in ψ_p mod p comes from M. Hence

M ≡ $\mathbf{A}_0$ (mod p) for a constant $\mathbf{A}_0$ ∈ R^×, so M′ ≡ 0 (mod p).

Let π : E → E^ := *E/C, with reduction F, and* $\hat{\pi}$: *E^* → E with reduction V, so $\hat{\pi}$∘π = [p] and c_πc_{$\hat{\pi}$} = p; choose the Weierstrass model of E^ *whose reduction is* $\bar{E}$^*{(p)}. Writing x∘π =* $\tilde{P}/M^2$ *and y∘π =* $y\hat{\Omega}/M^3$ *with* $\tilde{P}$, $\hat{\Omega}$ ∈ *R[x] we have (∗) for π, Z := x∘π ≡ x^p,* $\hat{\Omega}/M^3 \equiv u^{n_p}$*, and c_π ≡ 0 (mod p), say c_π = pϖ. The roots of N are the x-coordinates of the points π(Q) with Q ∈ ker* $\hat{\pi}$*, so up to a unit N = M^{2n_p}·h^(Z) with* h^ *the kernel polynomial of* $\hat{\pi}$*, h^* ≡ h (mod p). Differentiating ψ_p = λMN and using 2n_p + 1 = p,

$$\psi_\mathrm{p}' = \lambda(2\mathrm{n_p}+1)\mathrm{M}^{2\mathrm{n_p}}\mathrm{M}'\mathrm{h}^{(Z)} + \lambda M^{2n_p+1}(h^)'(\mathrm{Z})\cdot \mathrm{c_}\pi\hat{\Omega}/\mathrm{M}^3,$$

so that $\mathbf{K} = \psi_\mathbf{p}'/\mathbf{p} = \lambda\mathbf{M}^{2\mathbf{n_p}}\mathbf{M}'\mathbf{h}^{(Z)} + \lambda\varpi M^{2n_p-2}(h^)'(\mathbf{Z})\hat{\Omega}$: the factor p of c_π and the factor p of 2n_p + 1 = p cancel the denominator. Reducing, the first term dies by M′ ≡ 0 and the second gives K ≡ κ·u^{n_p}h′(x^p) with κ ∈ F^× independent of x. Comparing leading coefficients, lc_x K = $(\mathrm{p}^2-1)/2 \equiv -1/2$ and lc_x(u^nh′(x^p)) = n_p ≡ −1/2, both in degree $(\mathrm{p}^2-3)/2$, so κ = 1. ■ (In particular $\lambda\varpi \mathrm{A}_0{}^{\mathrm{p}} = 1$, whence c_{$\hat{\pi}$} ≡ A: the Hasse invariant is the action of V on the invariant differential, as it must be.)

Proposition 15, the reduction to one congruence. By Proposition 14, $4\mathrm{uK}^2 = 4\mathrm{w}^{(\mathrm{p})}(\mathrm{x}^{\mathrm{p}})\mathrm{h}'(\mathrm{x}^{\mathrm{p}})^2$, so Proposition 15 is the identity $\mathrm{A}^2\Phi = 4\mathrm{w}^{(\mathrm{p})}(\mathrm{h}')^2\mathrm{A}^2 /\mathrm{A}^2 + \mathrm{A}^2\mathrm{D}(\Psi^2)$ in X, that is

$$\Phi - 4\mathrm{w}^{(\mathrm{p})}(\mathrm{h}')^2 = \mathrm{D}(\mathrm{A}^2\mathrm{h}^2) = 2\mathrm{A}(\mathrm{DA})\mathrm{h}^2 + 2\mathrm{A}^2\mathrm{h}\cdot\mathrm{Dh}.$$

By Lemma 13 the left side vanishes at every root of h, and its degree is ≤ p − 1 because the leading coefficients cancel ($4\mathrm{n_p}^2 = (\mathrm{p}-1)^2 \equiv 1$), so it equals h·$\mathrm{C}_1$ with deg_X $\mathrm{C}_1 \le$ n_p. Since h is monic of degree n_p, the displayed identity is equivalent to a congruence modulo h together with the coefficient of X^n_p:

(R′) $[\mathrm{w}^{(\mathrm{p})}(\mathrm{h}')^2]' + \mathbf{A}^2\mathbf{h}'\cdot\mathbf{Dh} \equiv 0$ (mod h) and **(L) $[\mathbf{X}^{\mathbf{p}-1}](\Phi - 4\mathbf{w}^{(\mathbf{p})}(\mathbf{h}')^2) = 2\mathbf{A}\cdot\mathbf{DA}$**,

the first being the residue at each root — using $\mathrm{C}_1(\xi) = -2[\mathrm{w}^{(\mathrm{p})}(\mathrm{h}')^2]'(\xi)/\mathrm{h}'(\xi)$ from Lemma 13 — and the second the top coefficient. In terms of the roots, (R′) says $\mathrm{A}^2\cdot\mathrm{D}\xi_\mathrm{i} = (\mathrm{w}^{(\mathrm{p})})'(\xi_\mathrm{i}) + 4\mathrm{w}^{(\mathrm{p})}(\xi_\mathrm{i})\Sigma_{\mathrm{j}\neq\mathrm{i}}(\xi_\mathrm{i} - \xi_\mathrm{j})^{-1}$: it is the Kodaira–Spencer derivative of the étale p-division points.

Proposition 15, the Tate family. Each coefficient of the identity is weight-homogeneous in $(\mathrm{a}_4, \mathrm{a}_6)$, and the kernel of the q-expansion map $\mathrm{F_p}[\mathrm{E}_4, \mathrm{E}_6] \to \mathrm{F_p}[[\mathrm{q}]]$ is generated by A − 1, which is not homogeneous; so q-expansions are injective in each weight ([Ser73], [SwD73]) and it is enough to prove Proposition 15 for the Tate family. There A ≡ 1, $\bar{E}^{(\mathrm{p})}$ =

E_{q^p}, and the isogeny E_{q^p} → E_q induced by q^{pZ} ⊂ q^Z is V, its kernel being {q^j}. In the short model, with Q := q^p, X_Q(W) = Σ_{m∈Z}Q^mW(1−Q^mW)^{−2} − 2s$_1$(Q) + 1/12 and Y_Q = y_Q + x_Q/2, the invariant differential is dX/2Y = dW/W, so W∂_W X = 2Y and W∂_W Y = u_Q′(X); the roots are ξ_i = X_Q(q^i) and η_i := Y_Q(q^i), i = 1, …, n_p. Three cancellations prove (R′).

(a) Since Q = q^p, the chain rule gives q dξ_i/dq = pQ∂_QX + i·W∂_WX, and pQ = 0 in characteristic p:

q dξ_i/dq = 2i·η_i, hence Dξ_i = −q dξ_i/dq = −2iη_i.

(b) Let θ_Q(V) := (1−V)∏_{m≥1}(1−Q^mV)(1−Q^m/V), so that x_Q(W$_1$) − x_Q(W$_2$) = −c$_0^2$W$_2$·θ_Q(W$_1$W$_2$)θ_Q(W$_1$/W$_2$) · θ_Q(W$_1$)^{−2}θ_Q(W$_2$)^{−2} with c$_0$ = ∏(1−Q^m), the two sides having the same divisor and the same principal part at W$_1$ = 1. Substituting into H(W) := h(X_Q(W)) = ∏_i(X_Q(W) − ξ_i) and using 2n_p + 1 = p,

H(W) = C·G(W)·θ_Q(W)^{−p}, G(W) := ∏_{j=−n_p}^{n_p}θ_Q(Wq^j),

and G is a theta function for q, by a rearrangement of the exponents alone: the factors (1 − q^aW) occurring in G are exactly those with a ≥ −n_p, each once, because every integer ≥ −n_p is uniquely pm + j with m ≥ 0 and

j	≤ n_p; the factors (1 − q^b/W) are exactly those with b ≥ n_p + 1. Comparing with θ_q(W),

G(W) = (−1)^nq^{−n_p(n_p+1)/2}·W^n_p·θ_q(W), hence **H(W) = C′·W^n_p·θ_q(W)·θ_Q(W)^{−p}**.

(c) θ_Q(W)^p is a p-th power, so its logarithmic derivative vanishes in characteristic p. With D := W d/dW,

L := DH/H = n_p + D log θ_q(W), D log θ_q(W) = −W/(1−W) + Σ_{m≥1}[−q^mW/(1−q^mW) + q^m/(W − q^m)].

At W = q^i(1 + s) the term m = i is exactly 1/s, and the value at s = 0 of the rest telescopes: writing q^m/(q^i−q^m) = −(1−q^{i−m})^{−1} for m < i and = q^{m−i}(1−q^{m−i})^{−1} for m > i, the sum equals Σ_{k=1}^{i−1}(q^k−1)/(1−q^k) = −(i−1). So L = 1/s + κ + O(s) with **κ = n_p + 1 − i**, and since D = (1+s)d/ds and H = h$_1$s + h$_2$s^2 + ⋯ one has κ = 1 + h$_2$/h$_1$ and D^{2}H/DH|_{s=0} = 1 + 2h$_2$/h$_1$ = 2κ − 1 = p − 2i ≡ −2i. Finally DH = 2Y·h′(X) and D^{2}H = 4Y^{2}h″(X) + 2h′(X)u_Q′(X), so at W = q^i

η_i·D^{2}H/DH = (w^{(p)})′(ξ_i) + 4w^{(p)}(ξ_i)Σ_{j≠i}(ξ_i−ξ_j)^{−1} = −2iη_i = Dξ_i = A^{2}Dξ_i,

which is (R′). (Replacing i by p − i changes η_i to −η_i and i to −i and leaves both sides unchanged.)

For (L), expand the identity divided by A^2h^2: $\Phi/h^2 = 4w\hat{}\{(p)\}(h'/h)^2 + 2A{\cdot}DA + 2A^2Dh/h$. As $X \to \infty$ the coefficient of X matches because $4n_p^2 \equiv 1$, and the constant term is (L) in the form $\varphi_1 + 6\sigma = 2A{\cdot}DA$ with $\varphi_1 := [X\hat{}\{p-1\}]\Phi$ and $\sigma := \Sigma_i\xi_i$. In the Tate family Φ/h^2 is the x-coordinate of V, so it equals X_q(W) at X = X_Q(W), and lim_{W→1}(X_q − X_Q) = 0 because the m = 0 terms cancel and $\Sigma_\{m\neq 0\}c\hat{}m(1-c\hat{}m)\hat{}\{-2\} = 2s_1(c)$; hence $\varphi_1 = -2\sigma$. Summing over the whole kernel, $\Sigma_\{j=1\}\hat{}\{p-1\}x_Q(q\hat{}j) = 2s_1(q) - 2p{\cdot}s_1(Q) \equiv 2s_1(q)$, so $\sigma = n_p/12 + s_1(q) = -1/24 + s_1(q) = -E_2/24$; and $2A{\cdot}DA \equiv -E_2/6$ because $DA = -(\theta A - ((p-1)/12)E_2A) \equiv -E_2/12$ on q-expansions. So $\varphi_1 + 6\sigma = 4\sigma = -E_2/6 = 2A{\cdot}DA$. This proves (L), hence Proposition 15. ■

Theorem ND. By Proposition 15, $S_1 \bmod p = 2(D\Psi)(x\hat{}p)$, and Res_x(f(x^p), g(x^p)) = Res_X(f, g)^p, so $\mathrm{Res}_x(S_1, \psi_p) = (2\hat{}n_p{\cdot}\mathrm{Res}_X(D\Psi, \Psi))\hat{}p$. Writing $\Psi = A\prod(X-\xi_i)$ and applying D coefficientwise, $D\Psi(\xi_i) = -A{\cdot}(D\xi_i){\cdot}\prod_\{j\neq i\}(\xi_i-\xi_j)$, so with the sign of the resultant convention

$$\mathrm{Res}_X(D\Psi, \Psi) = (-1)\hat{}\{n_p(n_p-1)/2\}{\cdot}A\hat{}\{2n_p\}{\cdot}\prod_i(D\xi_i){\cdot}\prod_\{i<j\}(\xi_i-\xi_j)^2.$$

In the Tate family ord_q A = 0, ord_q(Dξ_i) = i and ord_q(ξ_i − ξ_j) = min(i, j), so ord_q of the resultant is $\Sigma_i\, i + 2\Sigma_\{i<j\}\min(i, j) = n_p(n_p+1)/2 + n_p(n_p^2-1)/3 = M = p(p^2-1)/24$, which is exactly 1/12 of its weight $n_p(p^2+1) + n_p(p^2-1) - 2pn^2 = 12M$. By §4.6(3) the only irreducible weight-homogeneous factor vanishing at the cusp is Δ_0, so $\mathrm{Res}_X(D\Psi, \Psi) = \gamma_p\Delta_0\hat{}\{M\}$ with $\gamma_p \in F_p\hat{}\times$ and $c_S = (2\hat{}n_p\gamma_p)\hat{}p = 2\hat{}n_p\gamma_p \neq 0$. ■

Proposition 16. The leading coefficients in the last computation are $\prod_i(D\xi_i) \to (-1)\hat{}n_p\, n_p!$ (by (a) above, $D\xi_i = -2i\eta_i = -iq\hat{}i(1 + \cdots)$), $\prod_\{i<j\}(\xi_i-\xi_j)^2 \to 1$ and $\Delta_0 \to -q/16$, which gives γ_p and c_S. For c_K, Proposition 14 factors the resultant:

$$\mathrm{Res}_x(K, \psi_p) = \mathrm{Res}_x(u, \psi_p)\hat{}n_p{\cdot}(A\hat{}\{n_p-1\}\mathrm{Res}_X(h', h))\hat{}p,$$
$$\mathrm{Res}_x(u, \psi_p) = (-1)\hat{}nA^3\prod_iw\hat{}\{(p)\}(\xi_i) = (-1)\hat{}nA^3\prod_i\eta_i^2, \mathrm{Res}_X(h', h) = \prod_ih'(\xi_i),$$

and in the Tate family $\eta_i = q\hat{}i/2 + \cdots$ and $\prod_ih'(\xi_i) = (-1)\hat{}\{n_p(n_p-1)/2\}q\hat{}\{n_p(n_p^2-1)/3\}(1 + \cdots)$. Hence the q-order is $n_p{\cdot}n_p(n_p+1) + p{\cdot}n_p(n_p^2-1)/3 = m_K$ — again 1/12 of the weight $(p^2-1)(p^2-3)/2$ — and

$$c_K = (-1)\hat{}\{n_p + n_p(n_p-1)/2 + m_K\}{\cdot}2\hat{}\{4m_K - 2n_p^2\}.$$

Now m_K is even: with $j := (p^2-1)/24 \in Z$ for $p \geq 5$ one has $m_K = (p^2-1)(p^2-3)/24 = j(24j-2) = 2j(12j-1)$. (No divisibility of j is needed, and none holds — j = 1 at p = 5.) So the sign is $(-1)\hat{}\{n_p(n_p+1)/2\} = (-1)\hat{}\{(p^2-1)/8\} = (2|p)$. For the exponent, $2n_p^2 = (p-1){\cdot}n_p \equiv 0 \pmod{p-1}$, while $3m_K = (p-1){\cdot}(p+1)(p^2-3)/8$ and $(p+1)(p^2-3)/8 = ab/2$ with $a = (p+1)/2$ and $b = (p^2-3)/2$ odd: for $p \equiv 3 \pmod 4$ the factor a is even and $3m_K \equiv 0 \pmod{p-1}$, so $2\hat{}\{3m_K\} = 1$; for $p \equiv 1 \pmod 4$ both are odd and $3m_K \equiv (p-1)/2 \pmod{p-1}$, so $2\hat{}\{3m_K\} = (2|p)$. Since $2\hat{}\{4m_K-2n_p^2\} = 2\hat{}\{m_K\}{\cdot}2\hat{}\{3m_K-2n_p^2\}$, this gives $c_K = \varepsilon_p2\hat{}\{m_K\}$ with $\varepsilon_p = (2|p)$ for $p \equiv 3 \pmod 4$ and $\varepsilon_p = (2|p)^2 = 1$ for $p \equiv 1 \pmod 4$, that is $\varepsilon_p = -1$ exactly for $p \equiv 3 \pmod 8$. The formula for c_p is §4.11. ■

5. Computations

The computations of the draft that concern the statements of [A] (its §§5.1–5.6) are not repeated here.

5.7 The closed form modulo p, and the second digit

All sweeps here are over curves of good ordinary reduction with $\#E(\mathbb{F}_p) = p$, $5 \le p \le 19$, $p \nmid \prod_\ell c_\ell$, ι, split, and with M_Λ integral; the ranges differ and are stated each time.

Lemma 7. Over two disjoint sweeps — rank exactly 2 to conductor 60000 (617 boxes) and rank ≥ 3 to conductor 400000 (48 boxes) — the identity $\kappa_j = c_1 \cdot \gamma(n\, e_j)$ held in **713 of 713** pairs, and $\kappa_j \ne 0 \iff v_p(t(n\, e_j)) = 1$ in all 713. The depth distribution was 689 pairs at $v_p(t) = 1$, 23 at 2 and 1 at 3, so the negative side is not vacuous. Controls: the wrong constant x_0 in place of c_1 holds in 125 of 713 by coincidence, and forgetting the factor n breaks 579 of the 600 pairs with $n \not\equiv 1 \pmod p$.

Proposition 8. Purely local, with no global point and no height: over the **372** split anomalous boxes of conductor ≤ 3000, with all $p - 1$ classes in each box (**2,876** classes), (L1) held in 2,876 of 2,876 and (L2) in 2,876 of 2,876. Controls: "s/χ constant" holds in 689 of 2,876 and "$s \cdot \bar{x}/\chi$ constant" in 575, of which 372 each are the base class itself, so 317 and 203 are coincidences; positive control, the non-split constancy of λ/χ^2 (Theorem H), held in **2,613 of 2,613** classes.

Theorem K. Over rank ≥ 1 and conductor ≤ 12000 — 1,051 boxes, 1,143 entries — and over rank ≥ 3 and conductor ≤ 400000 — 48 boxes, **all six entries** of each Gram matrix, 288 entries — the closed form held with **no exception**. Controls: replacing γ by a constant, and dropping the B_0 term, each hold in 0 of the 48 rank-3 boxes (and in 111 of the 1,051 boxes of the first sweep, where the rank is usually 1 and the two forms agree). Corollary K′(iv) was checked on the 48 rank-3 boxes: the equivalence $v_p(\mathrm{Reg}_p) = r - 2 \iff \det \hat{C} \not\equiv 0$ held in **48 of 48**, while the exact formula $v_p(\mathrm{Reg}_p) = (r - 2) + v_p(\det \hat{C})$ fails in one of them (310329c1 at $p = 5$: $v_p(\det M_\Lambda) = 2$ against 3).

Theorem N. Over conductor ≤ 15720 — **1,386 boxes and 1,457 entries** with $\chi \ne 0$, of which 1,313 boxes of rank one, contributing one entry each, and 73 of rank two, contributing 144 — the congruence $n^2 h_p(P) \equiv (1 + p)\Theta(x(nP)) + 2p \cdot q_p(d') \pmod{p^2}$ held with **no exception**. The distribution of $e = v_p(d(pnP))$ was 1,294 / 141 / 20 / 2 for $e = 2, 3, 4, 5$. The controls are informative:

control	holds in
drop the term $2p \cdot q_p(d')$	377 of 1,457 — exactly the entries with $q_p(d') = 0$
drop the factor $(1 + p)$, i.e. the term $p \cdot \Theta$	199 of 1,457
$q_p(n)$ in place of $q_p(d')$	310 of 1,457
$q_p(d' + p)$ in place of $q_p(d')$	**0 of 1,457**

The extension that fails. The congruence **without** the denominator term, $n^2 h_p(P) \equiv \Theta(x(nP)) \pmod{p^2}$, holds in **15 of 113** boxes of conductor ≤ 2000 — the frequency of a coincidence, about $1/p$ — and the equality of valuations $v_p(n^2 h_p(P)) = v_p(\Theta(x(nP)))$ fails in 3 of the 10 boxes there with $v_p \ge 1$ (603e1 and 869a1 at $p = 5$, 1520h1 at $p = 7$). So the second digit of the height is not the second digit of the Fermat quotient, and the local model of Propositions 9 and 10 governs Θ and not h_p.

Corollary N′. Among the rank-one boxes of conductor ≤ 15720 there are **184** with v_p(Reg_p) ≥ 1, of which **30** have v_p(Reg_p) ≥ 2. The criterion v_p(Reg_p) $\ge 2 \iff \Theta/p + 2q_p(d') \equiv 0$ held in **184 of 184**; the control that ignores the q_p(d′) term held in 149, so that term decides **35 of the 184** boxes. The frequencies of v_p ≥ 1 by prime were 87, 61, 9, 13, 6, 8 at p = 5, 7, 11, 13, 17, 19.

Propositions 9 and 10. Over the same 372 local boxes, with every class and 2p + 1 lifts per class — **67,444** pairs — the second-digit expansion of Proposition 9 held in 67,444 of 67,444; the control that drops the k^2 term differs from the truth in 55,040 of them, so the term is not decoration. The bad lift $\delta_$ *was computed digit by digit and verified to* p^3 *in* ***372 of 372*** *boxes, and* $v_p(\Theta(x_0)) = v_p(\delta - \delta_)$ held in **113 of 113** rank-one boxes of conductor ≤ 2000.

$\lambda_$ *T one digit deeper.* Modulo p^2 the identity $\Theta_T(m) = m^2\Theta_T(1)$ with the standard lift of m holds in 282 of 2,504 class pairs, and with the Teichmüller lift m^p in 580 of 2,504. Both frequencies are those of coincidences; no law is claimed (§7(7)).

5.8 The third and fourth digits, the locality of W, and the sensitivity

All sweeps here are over curves of good ordinary reduction with #E(F_p) = p, $5 \le p \le 19$, $p \nmid \prod_\ell c_\ell$, ι, and $\chi(P) \neq 0$; ranges are stated each time. Everything was calibrated before it was run, and the calibrations are reported first because two of them are the arithmetic backbone of §4.12.

Calibration. (i) $\varphi_p(x)/\psi_p(x)^2 = x(pP)$ with deg $\varphi_p = p^2$ and integral coefficients: **36 of 36** (twelve curves of conductor $\le 600 \times p = 5, 7, 11$). (ii) $p \mid \varphi_p'$ and $p \mid \psi_p'$ coefficientwise, the two halves of the shape of §2: **20 of 20** each (five curves $\times$ p = 5, 7, 11, 13); negative control, $p \mid f_\{p-1\}'$ for the even-index division polynomial, **0 of 20**. (iii) the two statements of §4.12 about the first coefficient — $X'(0) = 2y(T) + a_1x(T) + a_3$ and its being a unit: **129 of 129** boxes of conductor ≤ 3000, and (iv) $v_p(\psi_p'(x(T))) = 1$ exactly: **129 of 129** in the same boxes. (v) the σ-normalisation: `ellpadics2(ellinit([17,42]),13,4)` reproduces the one worked example of PARI's documentation. (vi) $E_6(E, \omega) = -c_6$ against the Hasse invariant, **250 of 250** (conductor ≤ 12000, $p \le 19$); with $E_6 = +c_6$ only 150 of 250, and the three smallest cases were checked by hand (p = 5: $c_4 \equiv 2A$ and the Hasse invariant 2A; p = 7: $-c_6 \equiv 3B$ and 3B; p = 11: $c_4(-c_6) \equiv 9AB$ and $20AB \equiv 9AB$).

E_2 modulo p is local (§2). Over conductor ≤ 12000 and $p \le 19$, the congruence $E_\{p+1\}(c_4, -c_6)/E_\{p-1\}(c_4, -c_6) \equiv b_2 - 12\cdot$`ellpadics2`$(E, p) \pmod p$ held in **315,953 of 315,953** boxes. The classical forms were computed from their q-expansions and re-expressed in E_4, E_6 by solving for the isobaric coefficients, with the denominators checked prime to p in every case. Controls: $E_6 \mapsto +c_6$ gives 54,109; dropping b_2 gives 65,189; $E_\{p+1\}$ alone, without dividing by $E_\{p-1\}$, gives 67,566; junk gives 34,020 — the rate of a coincidence at these p. Negative control: over the **58,785** supersingular boxes in the same range, $E_\{p-1\}(c_4, -c_6) \equiv 0$ in all of them and `ellpadics2` returned a p-adic number in **none**, so the two sides agree also about where the quantity is undefined. Finally, over the 315,121 pairs of boxes with the same (p, c_4 mod p, c_6 mod p), the two values of E_2 mod p agreed **without exception**, which is the statement of the Remark of §2 read directly.

Corollaries $N^{(3)}$ and $N^{(4)}$. Over conductor ≤ 20000 and rank 1 — **1,678 boxes**, with e = v_p(t(P″)) distributed 1,482 / 164 / 30 / 2 over e = 2, 3, 4, 5 — both congruences held with **no exception**, Corollary $N^{(3)}$ modulo p^3 and Corollary $N^{(4)}$ modulo p^4. The controls are what make the coefficients load-bearing:

control (same 1,678 boxes)	$N^{(3)}$ holds in	$N^{(4)}$ holds in
drop the term $c/(p^2x(P''))$	378	284
change its sign to +	378	—
coefficient of Θ^2 equal to $+1/4$ (the naive guess)	229	—
c replaced by $b_2/12$, i.e. $E_2 = 0$	377	—
c replaced by a number built from the conductor	409	—
drop the Θ^2 term	246	—
drop the F^3 term	—	410
F^2 without the factor (1 + p)	—	410
Θ^2 without the factor (1 + p)	—	246
Corollary $N^{(3)}$ read modulo p^4	—	280
add $p^3\Theta^3$	—	246
add $p^3\Theta F$	—	582
add $p^3 \times$ junk	—	292

The numbers in the high 200s and 300s are the counts of boxes where the touched term vanishes for a reason ($e \geq 3$ makes $c/(p^2x(P'')) \equiv 0$ modulo p, which is 196 boxes, and $q_p(d') = 0$ accounts for most of the rest); the numbers near 410 are two controls that both remove one term of the p^3 digit and therefore fail on the same boxes. **Every control fails.**

Corollaries N″ and N‴. Among the same boxes there are **41** with $v_p(\mathrm{Reg}_p) \geq 2$, of which **9** have $v_p \geq 3$ and **1** has $v_p \geq 4$ (19779a1 at p = 5). Corollary N″ decided all **41 of 41** correctly in both directions; the control that drops W decided 30, so W changes the verdict in **11** of the 41. Corollary N‴ was correct in **9 of 9**, the controls dropping U or F^3 each in 8 of 9 — with a single positive box that is the absence of a violation and not a test of the criterion (§7(11)).

Proposition 12 and the locality of W. Over conductor ≤ 20000 and rank 1, with γ computed independently through the group law in E(Q_p) — **1,676 boxes** — the congruence $W \equiv c\cdot\gamma^2 \pmod p$ held in **1,676 of 1,676**, and $e = 2 \iff \gamma \not\equiv 0$ in 1,676 of 1,676. Controls: $c\cdot\gamma$ in place of $c\cdot\gamma^2$ holds in 579, $W \equiv 0$ in 378, γ^2 without c in 401. Proposition 11 held in **1,676 of 1,676**; dropping the ν (t^3) term gives 351 and dropping γ_3 gives 237.

Theorem G′, the three steps separately. Over **400** boxes of conductor ≤ 5400 and rank 1, each of the three stages of §4.12 was measured on its own, for $j = 1, \ldots, 5$:

measured quantity	j = 1	2	3	4	5
$v_p(X(t_1 + p\hat{}j) - X(t_1))$ — step (1), predicted j	**1**	**2**	**3**	**4**	**5**
$v_p(\varphi_p(x_0 + p\hat{}j) - \varphi_p(x_0))$ — step (3), predicted j + 1	**2**	**3**	**4**	**5**	**6**
$v_p(\Theta(t_1 + p\hat{}j) - \Theta(t_1))$ — the theorem, predicted j − 1	**0**	**1**	**2**	**3**	**4**

each entry holding in **400 of 400** boxes, with no box off the predicted value. In the same boxes the two hypotheses of step (1) were verified directly: $c_1 = 2y(T) + a_1x(T) + a_3$ is a unit in 400 of 400, and $v_p(\psi_p'(x(T))\cdot c_1) = 1$ in 400 of 400.

Negative controls for the sensitivity. The exponent j + 1 of step (3) comes from Lemma 5 and from nothing else. Replacing φ_p by a polynomial that is **not** a polynomial in $x\hat{}p$ modulo p, the slope drops to j: for $x^2 + x$ the valuation equals j in 346 of the 400 boxes and is deeper in the remaining 54 (there the derivative $2x_0 + 1$ happens to be divisible by p), and for the even-index division polynomial f_{p−1} it equals j in 272 and is deeper in 128 — never j + 1 systematically. So the extra digit is the divisibility of φ_p' by p and not an artefact of the perturbation $p\hat{}j$.

The negative control for the split hypothesis. Over the full sweep to conductor 20000 — **14,168 boxes** — the equivalence of Lemma 4, split $\iff e \geq 2$, held in **14,168 of 14,168**:

the 1,678 split boxes have e ∈ {2, 3, 4, 5} (1,482 / 164 / 30 / 2) and the **12,490** non-split boxes have e = 1 in **every** case. On a sample of 300 boxes of each kind, ψ_p had an integral p-adic root in **300 of 300** split boxes and in **0 of 300** non-split ones. This is the sharp form of the negative control for Theorems G′ and G″: in the non-split case there is no p-torsion point T in E(Q_p), hence no formal parameter $t_1 = t(nP - T)$ and no layers γ, γ_2, γ_3 at all. The sensitivity statement itself is local and survives — φ_p′ is divisible by p whatever the reduction — but the chain that turns a digit of Θ into a layer of E(Q_p), and with it Theorem G″, exists only on the split side.

The non-degeneracy (ND). Two computations. (a) Over the 1,678 split boxes, $G(\bar{x}) = \varphi_p'(\bar{x})/p \not\equiv 0 \pmod p$ in **1,678 of 1,678**. (b) Exhaustively: for each p with $5 \le p \le 19$, for **every** pair $(a_4, a_6) \in F_p^2$ with $4{a_4}^3 + 27{a_6}^2 \neq 0$ — that is for every short Weierstrass model over F_p, hence for every isomorphism class, each class occurring several times, 20 + 42 + 110 + 156 + 272 + 342 = **942 models** — the resultant condition holds: gcd(G mod p, ψ_p mod p) = 1 in **942 of 942**, with no common root, ordinary and supersingular alike (**52** of the 942 are anomalous and **110** supersingular). For the 52 anomalous curves the conclusion was also read off directly: $G(x(\bar{Q})) \neq 0$ at **all 732** non-zero points $\bar{Q}$ of E(F_p). Since Lemma 5 is an identity over Z, this settles (ND) for every curve over Q with $5 \le p \le 19$; the exhaustion was later carried to $p \le 41$ (§5.9), and then replaced by the proof of Theorem ND, which covers every $p \ge 5$. That "942 of 942" is not vacuous was checked in two ways. First, the shape of the divisor: ψ_p mod p has exactly (p−1)/2 roots in F_p, each of multiplicity exactly p, in every anomalous box — which is the statement $\psi_p \equiv (\text{const}) \cdot g(x^p)$ of §2 read off, with g separable and split over F_p because ker V = E(F_p). Second, a random polynomial of the same degree as G, deg_x G = $p^2 - (p+3)/2$, shares a root with ψ_p mod p in **813 of 2,080** trials (39.1%), against the predicted $1 - (1 - 1/p)^{(p-1)/2}$ of 36.0% to 38.5%; and the derivatives of the neighbouring division polynomials f_{p+m}, m = 1, . . . , 4, do share a root with ψ_p mod p in many boxes (for instance 6, 6, 0 and 12 of the 12 anomalous curves over F_13). So the coprimality of G with ψ_p mod p is a property of G and not of degrees.

The 576 pairs, and what separates them. Over conductor ≤ 20000, rank 1, $p \le 19$ there are **576** pairs of curves isomorphic over Q_p with different v_p(Reg_p); the pairs (v_lo, v_hi) are (0,1) 475 times, (0,2) 9, (0,3) 74, (1,2) 2, (1,3) 16, and no pair has v_lo ≥ 2. Hence **all 576 are separated by the ledger of Theorem N**: the difference sits in Θ mod p in 558 pairs, in Θ/p only in 4, in **q_p(d′) only in 2**, in both in 12, and in neither in 0. The two pairs separated by the denominator alone — 3888p1 against 3267d1, and 19044a1 against 3267d3, both at p = 7 — are the shortest exhibition of the content of Theorem N, and the necessity of F in Theorem G″. Both pairs have j = 0, hence c = 0: the first local term of the ledger is absent there and, by Proposition 18, the whole of $p^{-2}\mathcal{L}$ is invisible before the eleventh digit, so the separation by q_p(d′) at the second digit is as clean as it can be. That is not a CM phenomenon, though. **432 of the 576 pairs have j = 0** — the boxes with j = 0 fall into two classes of 38, at p = 7 and at p = 19, and the number of pairs inside a class grows quadratically — and the rate of pairs separated at the second digit alone is 16 of 432 against 2 of 144, which is no difference at all (the expectation apportioned by the number of pairs is 13.5 against 16 observed). Theorem N has no c in it: c enters first at the third digit, through $W \equiv c\gamma^2$, so the way the pairs separate could not have depended on the vanishing of c. Structurally, among curves with the same j-invariant anomalousness forces the twisting factor to be a p-adic square, so every same-j pair surviving the filters is Q_p-isomorphic: **6,295 of 6,295** in this sweep (4,889 with j ≠ 0 and 1,406 with j = 0).

The scaling law. $\Theta \mapsto \Theta + 2p \cdot q_p(u)$ under $x \mapsto u^2x$, $y \mapsto u^3y$ with $p \nmid u$ (the Remark of §4.12): **226 of 226**.

5.9 The identities behind Theorem ND, and the constants

Everything in this subsection is a check of §3.14 and §4.13, in PARI/GP, with the controls stated. None of it is a proof; the proofs are in §4.13, and the numbers are what preceded them and what tested them afterwards.

The exhaustion that preceded the proof. (ND) was verified for **every** elliptic curve over F_p with $5 \le p \le 41$ — 942 curves for $p \le 19$ (§5.8) and **5,220** more for p = 23, 29, 31, 37, 41, so **6,162** in all — by gcd(G mod p, ψ_p mod p) = 1, and, on the anomalous curves, directly at all non-zero points of E(F_p): **732** points for $p \le 19$ (§5.8) and **4,480** for p = 23, ..., 41, so **5,212** in all. For $5 \le p \le 19$ the stronger statement Res_x(G, ψ_p mod p) = c_p·Δ_0^{m_p} was proved as a polynomial identity (symbolically for p = 5, 7; for p = 11, ..., 19 by interpolation in the weight-homogeneous shape Res = a_4^{3m}P(a_6^2/a_4^3), deg P ≤ m, over a Galois ring), with c_p = 2, 2, 1, 6, 13, 11.

The two identities. Proposition 14 holds as an identity in F_p[a_4, a_6][x] for p = 5, 7, 11, 13, 17, 19 (7.0 s at p = 19) and its corollaries u^{(p−1)/2} | K and K/u^n_p ∈ F_p[x^p] for $p \le 19$. Proposition 15 holds as an identity in F_p[a_4, a_6][X] — in the form $A^2\Phi = 4w^{(p)}(\Psi')^2 + A^2D(\Psi^2)$, whose degree in X is p — for **p = 5, ..., 23** (574 s at p = 23, 286 MB of PARI stack) and at 56 sampled points for $p \le 43$, the derivation D being taken by dual numbers over F_p[y]/(y^2). Controls: replacing D by $\partial/\partial a_4$, or w^{(p)} by u, breaks the identity at p = 5; replacing ψ_p by ψ_{p−2} breaks Lemma 13 itself (0 of 96).

The Tate family. With Ψ built from ψ_p over F_p[[q]] and h = Ψ/A: the identification ψ_p mod p = A$\prod$_i(x^p − ξ_i) with ξ_i = X_{q^p}(q^i), and A ≡ 1, hold to q^{30} for p = 5, 7; the congruence (R′) in the form $[w^{(p)}(h')^2]' - A^2h'\cdot(q\,dh/dq) \equiv 0 \pmod h$ holds **identically to q^{30} for p = 5, 7, 11, 13**, while the same expression with the opposite sign is non-zero from q^2 (p = 5) and q^3 (p = 7) on — this is the control that fixes the sign of D on the Tate family. The three steps of §4.13(c) were measured separately: q dξ_i/dq = 2iη_i and (R′) in root form to q^{22} for p = 5, 7, 11 and every i; the product rearrangement G(W) = (−1)^n q^{−n_p(n_p+1)/2}W^n_pθ_q(W) to a relative error 3.4·10^{−71} at q = 1/10, W = 0.37; the constant term κ = n_p + 1 − i, the telescoped sum −(i−1) and the ratio D^2H/DH = p − 2i to 20 digits. For (L): $\varphi_1 + 2\sigma = 0$ and $\varphi_1 + 6\sigma$ = 2A·DA to q^{22} for p = 5, 7.

The constants. At ordinary points of F_{p^2}, for one box per prime, the closed forms of Proposition 16 were compared with the measured resultants for **every p with 5 ≤ p ≤ 151** (34 primes): c_K agreed in **34 of 34** and c_S in **33 of 34**, the single exception being a sample point at p = 19 at which lc(S_1) = 2·DA vanishes, so that deg_x S_1 drops from 171 to 152 and the resultant does not specialise; at the other four sample points of p = 19 the value is the predicted 7. The values c_K = 4, 4, 10, 10, 1, 12, 1, 28, 1, 11, 1, 42 for p = 5, ..., 43 were measured before the closed form was proved, and c_K(47) = 1, c_K(53) = 52 were predicted from the closed form before being measured. The factorisation of the resultant used in §4.13 — Res_x(K, ψ_p) = Res_x(u, ψ_p)^n_p·(A^{n_p−1}Res_X(h′, h))^p, with the two factors as displayed there — holds at sample points for p = 5, 7, 11, 13, and γ_p = 2, 3, 10, 2 there agrees with the closed form.

The constant c_p in one line. Assembling the two counts of Proposition 16 gives, in F_p,

$$\mathbf{c_p = \varepsilon_p\cdot(-1)^{n_p + n_p(n_p+1)/2 + M}\cdot n_p!\cdot 2^{m_K + n_p + 4M}},$$

with n_p = (p − 1)/2, M = p(p^2 − 1)/24, m_K = (p^2 − 1)(p^2 − 3)/24, ε_p = −1 exactly when p ≡ 3 (mod 8), and every exponent read modulo p − 1; the regrouping of the

exponents is Lean `cp_one_line`. It agrees with the measured c_p for all twelve primes $5 \le p \le 43$: the six values 2, 2, 1, 6, 13, 11 for $p \le 19$ recorded above, and **22, 12, 30, 14, 9, 1** for $p = 23, 29, 31, 37, 41, 43$, which were printed from the closed form before being measured. One convention has to be fixed for this to hold: $\Delta_0 = \mathbf{4a_4}^3 + \mathbf{27a_6}^2 = -\Delta/\mathbf{16}$, and not Δ itself — dividing by Δ instead makes eight of the twelve values disagree.

5.10 Theorem N* to the tenth digit

Everything here is a check of §§3.11–3.12, in PARI/GP, over one fixed set of boxes. The boxes are: conductor ≤ 12000, $p \in \{5, 7\}$, a_p = 1 (so #E(F_p) = p), $p \nmid \prod_\ell c_\ell$, ι, split, **rank $\ge$ 1** with P the first generator in Cremona's tables, and $e \ge 2$ — **648 boxes**, 332 with p = 5 and 316 with p = 7, the distribution of e being [551, 82, 13, 2] over e = 2, 3, 4, ≥ 5. The point P″ is computed p-adically: x(P″) = φ_p(x$_0$)/ψ_p(x$_0$)^2 at x$_0$ = x(P′) + O(p^{30}), y(P″) from the Weierstrass equation (the sign of the square root is immaterial, since z(−Q) = −z(Q) and only z^2 and z^4 occur), t = −x/y, and z = log_E(t) by the formal logarithm. For the deep digits $\mathcal{L}$ is built as a series: `ellformallog` and `serreverse` give x(z), the differential equation of Proposition 17 is integrated twice with `intformal`, c is PARI's `ellpadics2`, and $\mathcal{L}$(z) = log(x(z)z^2) + 2log(σ_p(z)/z) is evaluated at z(P″).

Calibration, before anything new. The same generating rule — take Theorem N*, keep the terms of valuation $\le$ k − 1 by Proposition 18, discard the rest — reproduces **verbatim** the right-hand sides of Corollary N$^{(3)}$ and Corollary N$^{(4)}$ as they stand in §3.12, and at k = 5 it produces the term (2/5)p^{5}F^5 by itself, with no term put in by hand. Each of the three holds in **648 of 648** boxes. Beyond k = 5 the generated truncation and the **displayed** congruence are tested separately, the displayed one exactly as it is printed in §3.12: the corrected Corollary N$^{(6)}$ holds in **648 of 648** and the full Corollary N$^{(7)}$ in **648 of 648**, the latter sharply (valuation exactly 7) in 499 of them. Printing the geometric factor of the F^5 term as 1 rather than (1 + p) fails in **195** boxes, all with p = 5 and $5 \nmid$ F, which is how the factor was found: a truncation that the machine generates correctly can still be mistranscribed, so every displayed congruence is now checked in the form in which it is displayed. The counts 648, 332, 316 and the distribution of e agree with the independent sweep of §5.3's filter. Restricting the same filter to rank exactly 1 gives 628 boxes (321 with p = 5, 307 with p = 7, the distribution of e being [534, 80, 13, 1]); the congruences of §3.12 assume no rank, since Theorem N* is a statement about one point with $\chi(P) \ne 0$, so the wider filter is the one used.

The coefficients of $\mathcal{L}$, by an independent route. $\mu_2 = -c$, $\mu_4 = -b_4/12$, $\mu_6 = -b_2b_4/360 - b_6/30$ and the vanishing of every odd coefficient were read off the formal series built as above — which uses the differential equation but not the hand computation of §3.11 — for the four curves [0,0,0,−1,1], [1,−1,0,−79,289], [1,2,3,4,5], [0,1,1,−2,0] with c = 7/3, 11, −1/5, 0: **4 of 4**, to the order z^{12}. The curve [1,2,3,4,5] has $a_1 \ne 0$ and $a_3 \ne 0$, which is what makes the vanishing of the odd coefficients a test and not a consequence of the model.

The table of §3.12. The four columns were measured at k = 6, 7, 8, 9, 10, and the counts are those displayed there. The split by p at k = 10 is the sharp form of the third line of Proposition 18: truncating $\mathcal{L}$ at z^4 holds in **316 of the 316** boxes with p = 7 and fails in 256 of the 332 with p = 5, because v_5(μ_6) = −1 and v_7(μ_6) = 0.

Controls at k = 6. Eight modifications of the right-hand side, on the same 648 boxes: dropping the p^5 tail of the Θ series, 111; dropping the p^3 tail of F^2, 204; dropping the p^2 tail of F^3, 204; dropping the (1 + p) of F^4, 204; dropping the (1 + p) of Θ^3, 111; dropping $\mathcal{L}$ altogether, 41 — and in the 15 boxes with $e \ge 4$, where Proposition 18 says $\mathcal{L}$ is invisible modulo p^6, **15 of 15**; adding the term $(a_3 - 2c_4)/(p^2 y(P''))$, which the expansion by hand

had suggested as the next local ingredient, 398; adding $\mu_4 z^4$, **648**, since its valuation is 6 and it cannot change the congruence modulo p^6 — the last two being a positive and a negative control on the same quantity.

Controls on the logarithms, split by p. At k = 8, dropping the term with j = 5 of the Θ series — valuation 8 − v_p(5), hence 7 at p = 5 and 8 at p = 7 — leaves **316 of 316** at p = 7 and 64 of 332 at p = 5. At k = 10, dropping the term with j = 10 of the F series — valuation 10 − v_p(10), hence 9 at p = 5 and 10 at p = 7 — leaves **316 of 316** at p = 7 and 117 of 332 at p = 5. In both, one of the two primes is untouched and the other collapses, which is the factor v_p(j) measured rather than asserted.

z^2 against 1/x(P″). The valuation of $z^2 - 1/x(P'')$, minus 2e, over the 648 boxes: 4 in 393 boxes, 5 in 38, 6 in 52, 7 in 75, 8 in 39, 9 in 2, 10 in 2, and ≥ **11 in 47** (11 in 26, 12 in 9, 14 in 1, 15 in 1, 16 in 2, 17 in 6, 18 in 1, 20 in 1) — 648 in all. The lower bound 4 is Lemma 21: $z^2 - 1/x(z) = -(b_2/12)z^4 + O(z^6)$, so the valuation is 4e + v_p(b_2) as soon as the tail is deeper, and never less than 4e. In the **497** boxes with $b_2 \neq 0$ the measured valuation is 4e + v_p(b_2) **exactly**, in all 497, and $z^2 - 1/x(P'') + (b_2/12)z^4$ is deeper in every one of them; since the excess over 2e is then 2e + v_p(b_2), the 393 boxes of excess 4 are exactly those with e = 2 and $p \nmid b_2$ (b_2 is a p-adic unit in 453 of the 497, and b_2 = 0 in the remaining 151 boxes, where the leading term is absent and the valuation is deeper still). Hence p^{−2}($z^2 - 1/x(P'')$) has valuation ≥ 4e − 2 ≥ 6: **$1/x(P'')$ may be used through the sixth digit and not at the seventh**, where it leaves 310 of 648. This is the computation behind the statement in §3.13 that W is a quantity of the digits up to the sixth. The explanation through a_1 that an earlier version gave is not the right one — it reaches only 2e + 2 — and the terms of R linear in a_1 cancel against $a_1 t$ to leading order, so a_1 cannot be the source of the fourth power.

Proposition 19. In the first 40 boxes of conductor ≤ 2500, T ∈ E(Q_p)[p] was built by Hensel from the p-adic roots of ψ_p, $t_1 = t(P' - T)$ taken for each of the four choices of sign, and z(P″) compared with p·log_E(t_1): **40 of 40**, and e = 1 + v_p(t_1) in **40 of 40**. The sign of T changes z(P″) by a sign and z^2, z^4 not at all.

5.11 The two rates, over the cells of §5.10

Proposition 20 was measured on the filter of §5.10 — the same boxes, called cells here — split by conductor and by e into three sweeps. **Sweep a is the part of sweep b with conductor ≤ 3000, run separately and at a lower working precision; the distinct cells are those of sweeps b and c, which are disjoint (e = 2 against e ≥ 3).** In each cell T ∈ E(Q_p)[p] was built by Hensel from the p-adic roots of ψ_p, $t_1 = t(nP - T)$ by the group law in E(Q_p), X(t) = x(τ_T(R(t))) as a series, $\Theta(t) = Q_2(\varphi\text{_}p(X(t)))$, z(t) = p·log_E(t), and $\mathcal{L}$ as the series obtained by integrating the differential equation of Proposition 17 twice — which uses the equation but not the hand computation of §3.11.

sweep	conductor	e	cells	cells with c = 0	cells with c ≠ 0
a	≤ 3000	= 2	**103** (p = 5: 47; p = 7: 56)	8	**95**
b	≤ 6000	= 2	**259** (126; 133)	14	**245**
c	≤ 12000	≥ 3	**97** (e = 3: 82; e = 4: 13; e = 5: 2)	6	**91**

The cell counts are the counts of the filter of §5.10 restricted to the stated range and to nothing else: they are read off the list of 648 boxes mechanically. Sweeps b and c together

are **356 cells**, of which **336** have $c \neq 0$. The filter fixes P as the first generator at any rank, and the six cells of rank two in this range — three in sweep b, three in sweep c — behave exactly like the rest, so nothing below depends on a restriction to rank one.

The cells where c vanishes. Among the 356 cells the constant c of Proposition 17 — PARI's `ellpadics2` — vanishes in exactly **20**, and those 20 are exactly the cells with **j = 0**: 20 of 20 and 336 of 336, the positive and the negative control on the same criterion. This is the vanishing of the p-adic weight-two Eisenstein series at a CM point, here for complex multiplication by $\mathbb{Q}(\sqrt{-3})$. **All 20 lie at p = 7, and the reason is the trace and not the splitting of p.** For $j = 0$ the curve has complex multiplication by $\mathbb{Q}(\sqrt{-3})$, so at a prime of good ordinary reduction $a_p^2 - 4p = -3f^2$ for some $f \in \mathbb{Z}$; anomalousness $a_p = 1$ therefore forces $\mathbf{4p = 1 + 3f^2}$, hence $p \equiv 1 \pmod 3$, and among $5 \leq p \leq 19$ only $p = 7$ ($f = 3$) and $p = 19$ ($f = 5$) qualify — $4 \cdot 13 - 1 = 51 = 3 \cdot 17$ is not of that shape, so $p = 13$ is excluded although it splits in the field, while $p = 5, 11, 17$ are $\equiv 2 \pmod 3$ and supersingular (Lean `cm_prime_mod_three`, `cm_no_anomalous_of_mod_two`, `cm_seven_nineteen`, `cm_not_thirteen`). Over conductor ≤ 20000 the sweep of §5.8 has **76 boxes with j = 0, 38 at p = 7 and 38 at p = 19**, and **all 76 are split and all 76 have c = 0** (76 of 76 and 1,602 of 1,602); the 20 cells here are the part of those 76 that the filter of §5.10 keeps. Since the corrected Proposition 20 (iii) predicts a value at $c = 0$ as well, these cells are no longer set aside in the $\mathcal{L}$ rows: they are the rows where the leading term is $m = 3$.

measured	Proposition 20 says	result
$v_p(\Theta(t_1 + p^j) - \Theta(t_1))$, $j = 1..7$	exactly $j - 1$	**356 of 356**, every row
$v_p(z(t_1 + p^j) - z(t_1))$	exactly $1 + j$	**356 of 356**
$e = 1 + v_p(t_1)$ (Proposition 19)	—	**356 of 356**
$v_p(\mathcal{L}(z') - \mathcal{L}(z))$, $1 + j \neq e$	$\min_m [\, v_p(\mu_{2m}) + (1 + j) + (2m - 1) \cdot \min(e, 1+j) \,]$	**2,136 of 2,136 rows**, all 356 cells, $j = 1..7$
— the same rows against the old wording $v_p(c) + \min(e, 1+j) + (1 + j)$		**2,016 of 2,016** where $c \neq 0$; **false in all 120 rows** where $c = 0$
$v_p(\mathcal{L}(z') - \mathcal{L}(z))$, $1 + j = e$	at least the same, with the cancellation of the boundary row	**356 of 356 rows** at the boundary value — see below
v_p of the difference of the two sides of Theorem N*, F fixed	exactly $j - 1$	**356 of 356**
$v_p(2\log(1 + p(F + p^i)) - 2\log(1 + pF))$, $i = 0..3$	exactly $i + 1$	**356 of 356**

Controls. The **positive** control is the row $j = 3$, which reproduces the sharpness used by Theorem G″ — Θ moving at valuation exactly 2 — with the same instruments. The **negative** controls are the rows $j = 5, 6, 7$, where nothing moves modulo p^4: the fifth and deeper layers do not enter the verdict at $k = 4$, and at $j = 4$ the count is 0 of 356, as it must be. A second **independent route** measures $\mathcal{L}$ by the single term $-cz^2$ instead of the whole series and gets the same valuation in all **336** cells where $c \neq 0$.

(iii) row by row, and the unconditional form. Over the 356 cells and $j = 1, \ldots, 7$ — **2,492 rows** — the measured $v_p(\mathcal{L}(z') - \mathcal{L}(z))$ was compared with the displayed right-hand side of Proposition 20 (iii) itself, the minimum over m of $v_p(\mu_{2m}) + (1 + j) + (2m - 1) \cdot \min(e, 1 + j)$, the coefficients μ_{2m} read off the series for $m \leq 8$ and the prediction formed from $(e, j, v_p(\mu_{2m}))$ by a separate program. Equality in **2,136 of the 2,136 rows** with $1 + j \neq e$, and the minimum is attained by one m alone in all 2,492 of them: $m = 1$ in the 2,352 rows with $c \neq 0$ and $m = 3$ in the 140 rows with $c = 0$. The **unconditional**

form — v_p(c) + min(e, 1+j) + (1 + j) with no hypothesis on c — is false in exactly the **140 rows with c = 0**, where its right-hand side is infinite while the measured valuation is finite and equals (1 + j) + 5·min(e, 1 + j): 13, 13, 14, ..., 18 at the eleven cells with e = 2 that cancel at the boundary row, 12, 13, 14, ..., 18 at the three that do not, and 12, 19 or 21, 19, 20, ..., 23 at the six with e = 3 — v_p(μ_6) = 0 in all twenty. Its hypothesis v_p(c) < 2·min(e, 1+j) + v_p(μ_4) holds in all 2,352 rows with c ≠ 0, the narrowest margin being **one digit** (2115g1 and 2502e1, where v_p(c) = 3 against the threshold 4).

A box with c ≠ 0 where the hypothesis on c is needed. The positive control for the hypothesis needs a box with c deep, so the search was widened to conductor ≤ 40000 over 5 ≤ p ≤ 19: **3,214 split boxes** (1,009 at p = 5, 976 at p = 7, 301, 484, 151, 293), of which **61** have v_p(c) ≥ 3 — v_p(c) = 3 in 8 boxes, 4 in 49 and 6 in 4. In 60 of the 61 the hypothesis still holds, because v_p(μ_4) rises with v_p(c) in this sample (all 49 boxes with v_p(c) = 4 have v_p(μ_4) = 2). The exception is **27729a1 at p = 7**, with e = 3, $v_7(c) = 6$ and $p \nmid b_4$: at j = 1 the threshold is 2·min(3, 2) + 0 = 4, the m = 2 term leads, and the measured valuation is **8 = v_p(μ_4) + (1 + j) + 3·min(e, 1 + j)**, while the unconditional form gives ≥ 6 + 2 + 2 = 10. **The unconditional lower bound is false there by two digits, and Proposition 20 (iii) gives 8 exactly.** Over all 111 boxes measured outside the cells of §5.10 — the 61 deep ones and the 50 boxes with j = 0 at p = 19 up to conductor 40000 — Proposition 20 (iii) holds in **666 of 666** rows with 1 + j ≠ e and the boundary formula in **111 of 111**, while the unconditional form fails in 351 rows: 350 at c = 0 and the one at 27729a1. The six rows where two values of m tie (27729a1 at j = 2, ..., 7) showed no cancellation, so equality held there as well — the statement claims it only for a unique minimiser.

The boundary row. When 1 + j = e the two factors of $\Delta(z^2) = (2z + \Delta z)\Delta z$ have equal valuation and the leading terms can cancel. Writing $t_1 = p^{e-1}u$ with u a unit, z = p^e·u + O(p^{e+1}) and Δz = p^{1+j} + O(p^{2+j}), so at 1 + j = e the sum is 2z + Δz ≡ p^e(2u + 1), and the cancellation occurs **exactly when u ≡ −1/2 (mod p)**. Over the 336 cells with c ≠ 0 that criterion is exact: the $\mathcal{L}$ row is deeper than v_p(c) + min(e, 1+j) + (1 + j) in **67 cells, and those 67 are precisely the cells with u ≡ −1/2 (mod p)** — 67 of 67 and 269 of 269, and the same at two working precisions. Split by e the count is 44 of 245 at e = 2, 22 of 76 at e = 3, 0 of 13 at e = 4 and 1 of 2 at e = 5. Since u is a unit the heuristic frequency is **1/(p − 1) and not 1/p**: it predicts 51.3, 16.2, 2.8 and 0.4 respectively, and 70.8 over the 336 cells against 67 observed. Every deviation from the general bound is on the slow side — the ledger lagging more, not less — so the conclusion of Proposition 20 is strengthened.

The boundary row for a general m, and how deep the cancellation goes. Write w := Δz/z, a unit at the boundary row, with w ≡ u^{−1} (mod p). Then Δ(z^{2m}) = z^{2m}((1 + w)^{2m} − 1), so the leading term of the m-th summand of $\mathcal{L}$ cancels **exactly when (1 + w)^{2m} = 1 in F_p**, that is when (u + 1)^{2m} = u^{2m}. For m = 1 this is 2u + 1 ≡ 0, the criterion above; when (p − 1) | 2m — for instance p = 7 and 2m = 6, which is the leading term at a cell with c = 0 — **every** u with u ≢ −1 cancels. Over the 20 cells with c = 0, all at p = 7 with leading term m = 3, the cancellation happens in exactly the 17 cells with u ≢ −6 and fails in the three with u = 6 (1521a1, 1521a2, 3600bd1): **20 of 20**, the positive and the negative control on one criterion. The depth of the cancellation is v_p((1 + w)^{2m} − 1) − v_p(w) as well, which for m = 1 is v_p(2z + Δz) − e: over the 336 cells with c ≠ 0 the excess of the measured $\mathcal{L}$ row over the general bound equals that quantity in **336 of 336**, with the excess 0 in 269 cells, 1 in 54, 2 in 12 and 3 in 1. So the boundary row is not an exception with an unknown size: **which** cells lag more is one residue condition, and **by how much** is the valuation of one explicit unit. With w in hand the value at the boundary row is min_m [v_p(μ_{2m}) + 2m·e + v_p((1 + w)^{2m} − 1)], and

that predicted the measured valuation in **356 of 356** boundary rows, the 20 cells with c = 0 included (Lean `boundary_cancel_m1`, `boundary_cancel_pow`, `boundary_cancel_of_card`, `boundary_p7_m3`, `boundary_p7_exception`).

The criterion of the boundary row at p = 19, as a negative control. At p = 7 the criterion (1 + w)^{2m} = 1 is satisfied by every u ≠ −1 when 2m = 6, because 6 = p − 1; at p = 19 it is not, and the same formula must then make the cancellation the exception. The sweep of §5.8 extended to conductor 40000 has **50 boxes with j = 0 at p = 19** (38 of them within conductor 20000, which reproduces the count there), all split, all with c = 0 and all with $b_4 = 0$, so the leading term is m = 3 in every one. The congruence (u + 1)^6 ≡ u^6 (mod 19) has exactly five solutions, u ≡ 2, 7, 9, 11, 16, and the boundary row is deeper than the general value in exactly the **12 of the 50 boxes whose u is one of those five — 50 of 50**, against 17 of 20 at p = 7. The heuristic frequency 5/18 = 27.8% sits against 24% observed. One criterion, two primes, opposite verdicts, and right both times.

The residue u is a datum of the generator, not of (E, p). Replacing P by −P replaces T by −T and t_1 by $-t_1 + a_1 t_1^2 + \cdots$, so u = t_1/p^{e-1} is replaced by −u modulo p: measured in **356 of 356** cells, with no exception. Hence (E, p) determines only the class {±u}, and the sign is the convention by which the generator is chosen in `elldata`. The criterion of the boundary row is unaffected: it is stated for an ordered pair t_1, $t_1 + p^j$, and if both increments ±p^j are allowed it reads u ≡ ±1/2, which is sign-free, so the heuristic frequency 1/(p − 1) stands. It does mean that a count of raw residues is a count of a convention: at p = 7 the 178 cells give $\chi^2 = 12.85$ on 5 degrees of freedom for the raw u — an apparent excess at u = 5 — while the classes {±u} give 58, 67, 53 against 59.3, $\chi^2 = 1.70$ on 2 degrees of freedom, and at p = 5 the classes give 87, 91 against 89.0. The whole of the raw excess is the split 21 against 46 **inside** the class {2, 5}, and it survives counting isogenous curves once (39 against 24.7 at p = 7, with the classes then 53, 51, 44): it is a property of how generators are tabulated and there is nothing p-adic left in it.

5.12 Depth and height over the split boxes of rank two

The boxes. `elldata`, conductor ≤ 200000, p ∈ {5, 7, 11, 13, 17, 19}, rank exactly 2, a_p = 1, #E(F_p) = p, p ∤ $\prod_\ell$ c_ℓ, p ∤ #E(Q)_tors, ι — **25,374 curves and 27,355 boxes**, of which **3,056 are split** (988, 830, 321, 448, 138, 331 by p) and 24,299 non-split. E(Q) is torsion-free in all 27,355, as remark (a) of §3.15 requires. Nothing p-adic is computed: by Lemma 23 a rational point Q lies in E_k(Q_p) \ E_{k+1}(Q_p) exactly when v_p(d(Q)) = k, splitting is read as $D_1 := v_p(d(pP_1)) \geq 2$ for a generator P_1 with $\chi(P_1) \neq 0$, and det(χ, γ) ≡ 0 as D_Q := v_p(d(Q)) ≥ 2 for Q ∈ (Λ ∩ E_1) \ pΛ (remark (b) of §3.15). The sweep takes 85 seconds.

Calibration. On the six boxes whose ρ_1 had been measured from the p-adic height matrix, the test agrees in 6 of 6 (det(χ, γ) ≡ 0 at 8577a1 and 16011e1 at p = 5 and 12208i1 at p = 13; ≢ 0 at 2343c1, 3333b1, 4407a1). Three elements of (Λ ∩ E_1) \ pΛ — the first, third and fifth candidates in order of norm — give the same verdict D_Q ≥ 2 in **3,056 of 3,056** split boxes; the value of D_Q itself can differ (at 16011e1, p = 5, another choice of three elements has depths 2, 2 and 3), and it is the verdict, not the value, that remark (b) makes a property of the box. The negative control, since the argument of remark (b) needs splitting: the three elements disagree in 4,291 of the 24,299 non-split boxes, a number that depends on the choice of elements (7,279 for another choice).

The frequency, and its cause. If γ were a random multiple of χ on the two generators, det(χ, γ) ≡ 0 would occur with probability 1/p, and Σ1/p = 405.35 over the split boxes. It occurs in **100**.

measured	observed	naive expectation	ratio
$\det(\chi, \gamma) \equiv 0$ (rank 2, split, 3,056 boxes)	**100**	405.35	**0.247**
$\gamma(\mathrm{P}) \equiv 0$ (rank 1, split, 13,926 boxes)	1,805	1,847.2	0.977
$\gamma(\mathrm{P}_1) \equiv 0$ (rank 2, split)	420	405.4	1.036
$\chi(\mathrm{P}) = 0$ (rank 1, all 133,079 boxes)	2,975	14,559.7	0.204

The statistics that fall short are exactly those that ask for a power of p in the denominator of a rational point; $\gamma(\mathrm{P}) \equiv 0$ asks it of pP and not of P, and shows no deficit. Stratified by u := h_min/log p, with h_min the smallest canonical height $\hat{\mathrm{h}}$ on $(\Lambda \cap \mathrm{E}_1) \setminus \mathrm{p}\Lambda$ found in the box $[-2\mathrm{p}, 2\mathrm{p}]^2$ of coefficients, in the normalisation of PARI — which is not the naive height of Proposition 24, nor the threshold of Proposition 24^c, which adds $\beta(\mathrm{E})$: over the 3,056 boxes the naive minimum minus the canonical minimum ranges from -1.72 to $+7.61$ times log p, and is negative in about one box in six. The last column cuts the same boxes by the naive minimum h(x(Q)) instead (boxes, $\det(\chi, \gamma) \equiv 0$, $\Sigma 1/\mathrm{p}$, ratio):

u	split boxes	$\det(\chi, \gamma) \equiv 0$	$\Sigma 1/\mathrm{p}$	ratio	cut by the naive height
$0 \le \mathrm{u} < 2$	506	**0**	74.3	**0.000**	0 boxes (Lemma 23, k = 1)
$2 \le \mathrm{u} < 4$	1,732	15	236.5	0.063	1,528 · **0** · 207.8 · **0.000**
$4 \le \mathrm{u} < 6$	510	59	62.4	**0.946**	1,031 · 52 · 140.0 · **0.371**
$6 \le \mathrm{u} < 8$	178	18	20.1	0.895	307 · 36 · 37.1 · 0.971
$8 \le \mathrm{u} < 12$	100	5	9.4	0.531	144 · 7 · 16.1 · 0.433
$12 \le \mathrm{u} < 20$	27	3	2.2	1.363	43 · 5 · 3.9 · 1.281
$\mathrm{u} \ge 20$	3	0	0.4	0.000	3 · 0 · 0.4 · 0.000

Cut by the canonical height, the ratio returns to 1 at $\mathrm{u} \approx 4$. Counting only the boxes with $\mathrm{u} \ge 4$ gives the expectation 94.5 against 100 observed (ratio 1.058); moving the threshold to 2, 3, 4.5 or 6 gives 0.30, 0.57, 1.36 and 3.11, so on that scale the agreement is not a fit. **Proposition 24 is a statement about the naive height, and cut by the naive height the picture is different.** Below 4 log p there is no box with $\det(\chi, \gamma) \equiv 0$ among 1,528, against 207.8 expected; the 15 boxes with $\det(\chi, \gamma) \equiv 0$ in the canonical stratum $2 \le \mathrm{u} < 4$ all have naive height between 4.92 and 7.65 times log p. Between 4 and 6 log p the ratio is 0.37, and above the naive threshold 100 are observed against 197.5 (ratio 0.51). So Proposition 24 accounts for 207.8 of the deficit of $405.35 - 100 = 305.35$, about 68 %. **Proposition 24^c accounts for the rest.** The 15 boxes all have $\beta(\mathrm{E}) < 0$ and $(\hat{\mathrm{h}}_\mathrm{min} - \beta)/\log \mathrm{p}$ between 4.000 and 5.154: they lie above the canonical threshold $4 \log \mathrm{p} + \beta(\mathrm{E})$, as Proposition 24^c requires, and the return of the ratio to 1 near $\mathrm{u} \approx 4$ is the median of $\beta(\mathrm{E})/\log \mathrm{p}$, -0.07, seen on the scale $\beta = 0$. The naive column was computed twice, by separate programs: once from the shortest candidates in $\hat{\mathrm{h}}$ with a second pass, in the boxes first found at $\mathrm{u} \ge 4$, over every element of $\hat{\mathrm{h}} \le 4 \log \mathrm{p} + 12$, and once by enumerating every element of Λ with $\hat{\mathrm{h}} \le 6 \log \mathrm{p} + 14$, recognising $\ker \chi$ by $\mathrm{p} \mid \mathrm{d(Q)}$ with no discrete logarithm, and doubling the bound in the 33 boxes where none was found. Each naive minimum is therefore an upper bound, exact whenever the minimising element lies within the bound; the two computations agree in every entry, the boundaries at 4 and 6 are fixed within the observed range of $\hat{\mathrm{h}} - \mathrm{h}$, and the strata above 6 are approximate. On the side of χ: over the 66,013 boxes of rank one with $\mathrm{h(x(P))} < 2 \log \mathrm{p}$ there is no box with $\chi(\mathrm{P}) = 0$ (Lemma 23 with k = 1), and above u = 6 the ratio is 0.98 to 1.12.

The next table cuts the same boxes, and the 5,572 split boxes of the extension below, by the threshold of each proposition (boxes below the threshold, $\det(\chi, \gamma) \equiv 0$ among them, $\Sigma 1/\mathrm{p}$ over them, share of the deficit) [computation, with m_∞ a numerical minimum]. We call it table 5.12^c.

cut	set A: conductor $\leq$ 200000, 3,056 boxes, 100 with det $\equiv$ 0, deficit 305.35	set B: conductor (200000, 500000], 5,572 boxes, 283, deficit 458.15	A + B: 8,628 boxes, 383, deficit 763.50
naive, h_min $<$ 4 log p (Prop. 24)	1,528 · **0** · 207.8 · 68.05 %	1,874 · **0** · 255.6 · 55.78 %	3,402 · **0** · 463.4 · 60.7 %
canonical, $\hat{\text{h}}$_min $<$ 4 log p + β(E) (Prop. 24^c)	2,192 · **0** · 307.1 · **100.6 %**	3,251 · **0** · 456.8 · **99.70 %**	5,443 · **0** · 763.9 · **100.05 %**
canonical with β_Λ (remark (g))	2,221 · **0** · 311.7 · 102.07 %	3,320 · **0** · 467.8 · 102.11 %	5,541 · **0** · 779.5 · 102.09 %
either	2,215 · 0 · 310.1 · 101.6 %	3,294 · 0 · 462.8 · 101.0 %	5,509 · 0 · 772.9 · 101.2 %
naive only	23 · 0 · 3.0 · 1.0 %	43 · 0 · 6.0 · 1.3 %	66 · 0 · 9.0 · 1.2 %
control: $\hat{\text{h}}$_min $<$ 4 log p ($\beta := 0$)	2,239 · **15** · 311.0	3,366 · **28** · 471.6	5,605 · **43** · 782.6
above the canonical threshold: observed / $\Sigma 1/\text{p}$	100 / 98.2 = 1.018	283 / 284.4 = 0.995	383 / 382.6 = 1.001

The same canonical cut on the two sets C and D described below, which lie outside Cremona's tables (the naive cut was not computed there):

cut	set C: 5,782 boxes, 812 with det $\equiv$ 0, $\Sigma 1/\text{p}$ 830.3	set D: 7,861 boxes, 788, $\Sigma 1/\text{p}$ 1,087.4
canonical, $\hat{\text{h}}$_min $<$ 4 log p + β(E) (Prop. 24^c)	695 · **0** · 97.8	2,039 · **0** · 317.3 · 106.0 % of the deficit 299.4
canonical with β_Λ (remark (g))	715 · **0** · 101.0	2,063 · **0** · 321.3
above the canonical threshold: observed / $\Sigma 1/\text{p}$	812 / 732.5 = 1.109	788 / 770.1 = 1.023

(On set C the deficit, 18.3, is too small for a share to mean anything: there the frequency above the threshold is above 1/p, at p = 5 alone, 527 against 445.2.)

Controls: with β(E) replaced by 0 the canonical cut lets 15 and 28 boxes with $\det(\chi, \gamma) \equiv 0$ below the threshold; permuting β(E) among the boxes (1,000 times) lets 8 to 24 and 26 to 50, never 0; a constant β (the median, 0, or $-0.5 \cdot \log$ p) gives 15, 15 and 4 on set A and 26, 28 and 17 on set B. Only the curve's own β(E) gives 0. On every enumerated non-zero element of Λ in all 8,628 boxes $\text{R(Q)} \geq \beta$(E) holds; dropping the component corrections breaks it in 2,833 and 5,176 boxes. The minimum m_∞ is taken on a grid with golden-section refinement and checked by a second route through the x-coordinate (all of set A; 412 boxes of set B, all those within 0.05 log p of the threshold): the second route never goes below the first. Since a sampled minimum is at least the true one, the computed β(E) is at least the true one, so "no box with $\det(\chi, \gamma) \equiv 0$ below the threshold" cannot be produced by the grid missing a narrow valley; what the error could inflate is the share. The smallest margins are $7.5 \cdot 10^{-5}$ log p on the side of $\det(\chi, \gamma) \equiv 0$ (434832b1, p = 5) and $1.6 \cdot 10^{-4}$ log p on the explained side (106947h1). At I_n* with c_ℓ = 2 (205 and 524 boxes) the rational component was not determined and the smaller correction was used, which can only lower β(E). By p, the ratio above the canonical threshold over A + B is 0.91, 1.02, 0.96, 1.10, 1.46, 1.16 for p = 5, 7, 11, 13, 17, 19 (139/153.0, 117/115.3, 32/33.4, 49/44.6, 18/12.3, 28/24.1).

Just above the threshold: an effect whose cause is not known. Stratified by ($\hat{\text{h}}$_min $-$ β)/log p over A + B: [4, 4.5) 118/97.2 = 1.21, [4.5, 5) 51/63.0 = 0.81, [5, 6) 81/84.3 = 0.96,

[6, 8) 76/80.3 = 0.95, [8, 12) 37/43.8 = 0.85, ≥ 12 20/14.1 = 1.41. The excess just above the threshold is 1.46 on set A and 1.10 on set B (Poisson z = +2.6 and +0.8). To decide whether it is noise we fixed, before computing, two further sets and a rule (Poisson z > 2 on the set alone: an effect; z < 1: noise; for the last set also the premise that above the threshold the frequency is within z = ±2 of 1/p at each p). Set C: rank-two curves $y^2 = x^3 + Ax + B$ through two small points whose x-coordinates differ by 1. Set D: rank-two curves $y^2 = x^3 + Ax + B$ with random $|A| \le 1000$, $|B| \le 10^4$ on which two independent points of naive height at most 10^4 were found. On C and D the rank is fixed by ellrank with equal bounds, and the generators are saturated at all primes ≤ 200, which we assume to be enough; neither set meets A, B or the other. In the stratum [4, 4.5) (observed $\det(\chi, \gamma) \equiv 0$ against Σ1/p, Poisson z):

set	conductor	split boxes	[4, 4.5), threshold with β(E)	with β_Λ(E)	above the threshold
A	≤ 200000 (Cremona)	3,056	45 / 30.8 = 1.46 (+2.6)	46 / 30.1 = 1.53 (+2.9)	1.02
B	(200000, 500000] (Cremona)	5,572	73 / 66.4 = 1.10 (+0.8)	80 / 65.4 = 1.22 (+1.8)	0.99
A + B	≤ 500000	8,628	118 / 97.2 = 1.21 (+2.1)	126 / 95.5 = 1.32 (+3.1)	1.00
C	512,144 to $5.8 \cdot 10^{12}$, median $1.1 \cdot 10^{10}$	5,782	99 / 50.7 = 1.95 (+6.8)	100 / 50.7 = 1.97 (+6.9)	1.11
D	508,528 to $1.0 \cdot 10^{11}$, median $3.5 \cdot 10^{9}$	7,861	94 / 51.8 = 1.82 (+5.9)	96 / 50.2 = 1.91 (+6.5)	1.02

On D the premise holds (|z| ≤ 1.3 at each p). On C the frequency above the threshold is 1.18 times 1/p at p = 5, which is not seen on D (0.98); on C the excess in [4, 4.5) remains when the positions above the threshold are permuted within each p (z = +6.6). By the rule the excess is an effect on C and on D. It is not traced to the archimedean place, nor to the component corrections once the curves are matched. By Lemma 25 the frequency 1/p is exact for the local classes, so the effect is a correlation between the class of a box modulo E_2 and its heights.

Whether the size of the effect — 1.21 on A + B against 1.95 and 1.82 on C and D — follows the conductor, β(E)/log p or the way the set was built was tested by a second rule fixed before counting: strata by the tertiles of the conductor within D, the halves of the conductor within A + B, and the tertiles of β(E)/log p over all four sets; a score test for a log-linear trend in a Poisson model within D and within the β-strata (summed over the sets), a conditional binomial test for the two halves of A + B, and a Mantel–Haenszel-type stratified conditional binomial test for C + D against A + B within the β-strata. In [4, 4.5) the ratios are 1.60, 1.93, 1.95 over the conductor tertiles of D (trend z = +0.8) and 1.22, 1.21 over the halves of A + B (z = −0.0); the β-trend gives z = +1.5 and C + D against A + B within the β-strata z = +0.8 (with β_Λ: +0.7, +0.2, +0.9, +1.2). None reaches |z| = 2, so by that rule the size of the effect is attributed neither to the conductor nor to the construction of the sets. The rule has little power, and we do not read the outcome as an absence: on synthetic data in which the ratio moves log-linearly in log N from 1.21 at the median conductor of A + B to 1.88 at that of C + D, it detects the conductor in 14 % of runs, since within D the conductor moves by a factor of about 6.6 between the tertile boundaries and between the medians of A + B and C + D by about $1.8 \cdot 10^4$; and the β-strata hardly overlap (82 % of the boxes of A + B lie in the lowest, 98 % of those of C + D in the two upper ones). Matched on β(E)/log p ∈ [0.4, 1.3] instead — all 1,378 such boxes from Cremona's tables, which end at conductor 500000 (Σ1/p = 20.3 in [4, 4.5)), against curves

built as for D until $\Sigma 1/p$ reached 50 there (10,727 boxes) — the ratios are $30/20.3 = 1.48$ and $76/50.1 = 1.52$, and a further rule fixed before counting (a Mantel–Haenszel-type test within three sub-strata of $\beta(E)/\log p$) gives $z = -0.1$, rate ratio 0.97 with an approximate 95 % interval from 0.58 to 1.62 (with β_Λ, $z = +0.1$), which on synthetic data detects a ratio of 1.2 between the two constructions in at most 9 % of runs and so decides nothing; no box of either set has $\det(\chi, \gamma) \equiv 0$ below the threshold. Within the curves built as for D alone, which can be multiplied at will, a third rule fixed before counting divided the boxes with $\beta(E)/\log p \in [0.4, 2.0]$ into four layers (boundaries 1.0, 1.3, 1.6) and tested for a trend with a Cochran–Armitage-type score test stratified by p and by the range of (A, B), after adding 10,355 boxes from curves with $10^4 < |B| \le 2{\cdot}10^4$ that had not been looked at: the ratios in [4, 4.5) are 1.31, 1.58, 1.61 and 1.84 ($\Sigma 1/p = 26.7, 43.8, 27.9, 46.3$), the trend gives $z = +1.2$ (with β_Λ, +1.0) and the ratio of the top layer to the bottom one is estimated at 1.26, with a profile-likelihood 95 % interval from 0.86 to 1.88. On synthetic data a ratio moving linearly from 1.4 to 1.9 across $\beta(E)/\log p \in [0.4, 2.0]$ is detected in 10 % of runs at this size, and 80 % would need $\Sigma 1/p \approx 1{,}300$ in the band, so this rule decides nothing either: the excess is not shown to follow $\beta(E)/\log p$, nor to be independent of it. On the new curves alone the excess recurs (93 against $\Sigma 1/p = 59.0$, a ratio of 1.58) and the trend is $z = +0.1$; none of the 6,156 boxes built as for D (2,328 of them new) has $\det(\chi, \gamma) \equiv 0$ below the threshold. A fourth rule fixed before counting compared two predictions without free parameters on 11,008 further boxes built as for D from curves with $10^4 < |B| \le 2{\cdot}10^4$ whose (A, B) had not been drawn before. The first is a constant multiple of $\Sigma 1/p$, with the constant 1.62 taken from the earlier boxes built as for D. The second splits the band into boxes with $\det(\chi, \gamma) \equiv 0$ whose shortest non-zero point Q of $\Lambda \cap E_1(Q_p)$ has p-adic depth 2 and an excess $s < 0.5$ outside p, and boxes with $\det(\chi, \gamma) \not\equiv 0$ whose Q has depth 1 and $s \in [2, 2.5)$ (on all the boxes measured, $\det(\chi, \gamma) \not\equiv 0$ has always come with depth 1), and predicts the first kind from the boxes outside the band, assuming that $s < 0.5$ is equally frequent in both classes and that depth 2 has frequency $1 - 1/p$. The band holds 111 boxes with $\det(\chi, \gamma) \equiv 0$ against 91.3 and 116.9 predicted. The Poisson log-likelihood ratio is -14.6, in favour of the constant multiple; the split prediction fails its own resampling ($p < 10^{-4}$) and its fit by layers of $\beta(E)/\log p$ ($p = 0.03$), while the constant multiple fits by layers ($p = 0.19$) but falls short in total ($z = +2.1$; the ratio on these curves is 1.97). So the split does not account for the excess. The ratio in [4, 4.5) differs between sets of curves (1.37 to 1.97), but over the six sets — A + B, C, and four sets built as for D, two with $|B| \le 10^4$ and two with $10^4 < |B| \le 2{\cdot}10^4$ — it is consistent with a single constant (pooled ratio 1.73, Poisson $\chi^2 = 7.4$ on 5 degrees of freedom, $p = 0.19$) and shows no trend in $|B|$ ($p = 0.41$) [computation, outside the rules]; none of the 2,449 new boxes below the threshold has $\det(\chi, \gamma) \equiv 0$. A fifth rule fixed before counting tested the frequency $1 - 1/p$ of depth 2 among the boxes with $\det(\chi, \gamma) \equiv 0$ at $p = 5$, which had come out higher on every earlier set: on 8,181 further boxes at $p \in \{5, 7\}$ from curves whose (A, B) had not been drawn before, 391 of the 454 boxes with $p = 5$ and $\det(\chi, \gamma) \equiv 0$ have depth 2 (0.861 against 0.800; one-sided exact binomial test, $p = 4.3{\cdot}10^{-4}$), and at $p = 7$ 461 of 511 (0.902 against 0.857). The frequency $1 - 1/p$ rests on two assumptions about the map $\theta : \Lambda \cap E_2(Q_p) \to E_2(Q_p)/E_3(Q_p) \cong F_p$, through which depth 2 is decided: that θ is uniformly distributed, and that it is independent of the line of Q modulo p. On these boxes the first is not contradicted ($\theta = 0$ in 17 of 454 against 1/25, and no dependence on (A, B) mod 25 is seen, χ^2 $p = 0.49$), while the second fails: the kernel of θ is the line of Q in 46 of the 437 boxes with $\theta \neq 0$, against 1/6. A point of depth 3 carries at least 2 log p more in $\hat{h}$ than one of depth 2, so choosing Q by $\hat{h}$ pushes the kernel of θ off the shortest line; subtracting that amount raises the count from 46 to 58, against 72.8 expected [computation, outside the rules]. None of the 2,330 new boxes below the threshold has $\det(\chi, \gamma) \equiv 0$. A sixth rule fixed before counting chose instead the element

Q' of $(\Lambda \cap E_1(Q_p)) \setminus \{0\}$ that minimises $\hat{h} - 2(d-2)\log p$, with d the p-adic depth — the canonical height with the contribution $2d \log p$ of p to the denominator removed, up to a constant — and asked whether depth at least 3 then has frequency $1/p$ among the boxes with $\det(\chi, \gamma) \equiv 0$ at $p = 5$. The depth was read from the p-adic valuation of the formal logarithm on a basis of $\Lambda \cap E_1$ and checked against the denominators, and the minimum was taken over all elements of depth below 40. On 8,048 further boxes at $p \in \{5, 7\}$ from curves whose (A, B) had not been drawn before, 80 of the 433 boxes with $p = 5$ and $\det(\chi, \gamma) \equiv 0$ have Q' of depth at least 3 (0.185 against 0.200; one-sided exact binomial tests: $P(K \leq 80) = 0.23$ under $1/p$, and $P(K \geq 80) = 0.005$ under the frequency 0.139 of the earlier boxes with Q chosen by $\hat{h}$), and the verdict of the rule is that the frequency is consistent with $1/p$; at $p = 7$ it is 71 of 535 (0.133 against 0.143). On these boxes Q chosen by $\hat{h}$ itself has depth at least 3 in 69 of 433 (0.159), so the rule separates the two choices by 11 boxes. Over all six sets built as for D the frequency with Q' is 347 of 1,973 (0.176, $P(K \leq 347) = 0.004$ under $1/p$) against 287 with Q: removing the depth from the height recovers about half of the deviation, and the rest is not explained [computation, outside the rules; the earlier sets include the one on which the explanation was found]. The threshold and the band do not depend on this choice. When $\det(\chi, \gamma) \equiv 0$ every non-zero element of $\Lambda \cap E_1$ lies in E_2, so the argument of Proposition 24^c applies to every such element and none of them, Q' included, lies below $4 \log p + \beta(E)$ [paper, under the hypotheses of Proposition 24^c]; and over the six sets Q' differs from Q in 97 of the 5,351 boxes with $\det(\chi, \gamma) \equiv 0$ and $\beta(E)/\log p \in [0.4, 2.0]$, and in one of the 48,168 boxes with $\det(\chi, \gamma) \not\equiv 0$, three boxes leave $[4, 4.5)$ and none enters, and the ratio there is 1.81 with either choice (593/327.3 and 590/326.7) [computation]. So the excess just above the threshold is not an artefact of choosing the shortest element by $\hat{h}$. A seventh rule fixed before counting looked at the shape of $\Lambda \cap E_1$ instead. With b_1, b_2 a Gauss-reduced basis of $\Lambda \cap E_1(Q_p)$ for the quadratic form $\hat{h}$ (b_1 is the element Q above) and $r = \hat{h}(b_2)/\hat{h}(b_1)$, the boxes with $\det(\chi, \gamma) \equiv 0$ at $p = 5$ were divided into five classes of r with boundaries 1.1, 1.3, 1.7 and 2.5, and in each class the frequency of $\theta(b_1) = 0$, that is of depth at least 3 for b_1, was compared with $1/p$ by an exact two-sided binomial test, with Holm's correction over the five classes. The hypothesis written down beforehand, from the shape of the remaining deviation on the earlier sets, was that only the classes with r near 1 would deviate. On 23,927 further boxes at $p \in \{5, 7\}$ from curves whose (A, B) had not been drawn before, the 1,334 boxes with $p = 5$ and $\det(\chi, \gamma) \equiv 0$ give 0.161, 0.193, 0.203, 0.168 and 0.050 from the first class to the last, and only the last ($r \geq 2.5$; 26 of 525, $p = 5 \cdot 10^{-23}$) is rejected: by the rule the deviation is not confined to r near 1, and it sits at large r. The frequency of $\theta(b_2) = 0$ is consistent with $1/p$ in every class (247 of 1,334 in all), and among the boxes with $\theta \neq 0$ the kernel of θ is the line of b_2 as often as $1/(p+1)$ predicts in every class (86 against 86.3 in the last), so the deviation is not an exchange of the two shortest lines. At $p = 7$ the pattern is the same (only the last class deviates, 68 of 707). A secondary rule, added before these boxes were measured, reads this through Proposition 24^c: under its hypotheses every non-zero element v of $\Lambda \cap E_1$ has $\hat{h}(v) \geq \beta(E) + 2d(v) \log p$, so depth at least 3 for b_1 requires $X(b_1) := (\hat{h}(b_1) - \beta(E))/\log p \geq 6$ [paper, under the hypotheses of Proposition 24^c], and a null model in which θ is independent of the lattice is not compatible with that inequality. On these boxes no box with $X(b_1) < 6$ has $\theta(b_1) = 0$ (0 of 495 at $p = 5$, 0 of 332 at $p = 7$), and on the boxes with $X(b_1) \geq 6$ the frequency is 174 of 839 (0.207, two-sided $p = 0.60$; at $p = 7$, 178 of 1,225), with no class rejected; the deficit against $1/p$, 92.8 boxes, is of the size of the mass $495/5 = 99.0$ that the inequality excludes. Since the determinant of $\hat{h}$ on $\Lambda \cap E_1$ is $p^2 \cdot \mathrm{Reg}(\Lambda)$, a large r goes with a small $\hat{h}(b_1)$, which is why the deficit sits in the last class (361 of its 525 boxes have $X(b_1) < 6$) [computation; a rule fixed before counting and a secondary rule fixed before measuring; this reading accounts for the counts and is not a theorem]. As an observation outside the rules,

the element Q' of the sixth rule has depth at least 3 in 207 of these 1,334 boxes (0.155), so the agreement with 1/p found by the sixth rule did not recur; on the boxes with $X(b_1) \geq 6$ it is 205 of 839 (0.244) [computation]. An eighth rule fixed before counting asked whether the boxes with $\det(\chi, \gamma) \equiv 0$ are pressed against their lower bound more than the others. Let $s := X(b_1) - 2d(b_1)$, with d the p-adic depth; under the hypotheses of Proposition 24^c, $s \geq 0$ [paper, under the hypotheses of Proposition 24^c], and the bound sits at $X(b_1) = 4$ (depth 2) when $\det(\chi, \gamma) \equiv 0$ and at $X(b_1) = 2$ (depth 1) when $\det(\chi, \gamma) \not\equiv 0$. The rule compared $\rho := \#\{s \in [0, 0.5)\}/\#\{s \in [0.5, 1)\}$ between the two classes by Fisher's exact test (two-sided, level 0.05), with $p = 5$ and $p = 7$ together. On 13,920 further boxes at $p \in \{5, 7\}$ from curves $y^2 = x^3 + Ax + B$ with $|A| \leq 1000$ and $|B| \leq 10^4$ whose (A, B) had not been drawn before ($\beta(E)/\log p \in [0.4, 2.0]$: 1,709 boxes with $\det(\chi, \gamma) \equiv 0$ and 12,133 with $\det(\chi, \gamma) \not\equiv 0$), ρ is 2.57 (296 against 115) and 2.44 (2,017 against 826), $p = 0.68$, and the verdict of the rule is that there is no difference: both classes crowd just above their own bound to the same degree. The same holds at $p = 5$ and at $p = 7$ separately (2.70 against 2.77, 2.32 against 2.06) and after dividing s by the width of the window up to Hermite's bound $\sqrt{(4/3 \cdot \det)}$ for $\hat{h}(b_1)$ (2.02 against 2.40, $p = 0.07$), and a split of the boxes with $\det(\chi, \gamma) \not\equiv 0$ by the parity of the draw is not rejected either; over the whole range $s < 4$ the two distributions do differ (χ^2 $p = 0.012$, after the fact). Outside the rules: every box with $s < 0.5$ has a point b_1 whose x-coordinate has a power of p as denominator, which the hypotheses of Proposition 24^c force, since a further prime in the denominator adds at least $2 \log 2/\log p > 0.5$ to s [paper, under the hypotheses of Proposition 24^c]; near 0 the fraction of the boxes with $s < 0.5$ that have $s < a$ is 0.005, 0.013, 0.044 and 0.13 for $a = 10^{-5}$, 10^{-4}, 10^{-3} and 10^{-2} ($\det(\chi, \gamma) \not\equiv 0$), close to $\sqrt{(2a)}$, as when λ'_∞ has a non-degenerate minimum on E(R) near which the point is smoothly distributed; the smallest s is $5 \cdot 10^{-8}$, so the threshold is approached to that precision; in [4, 4.5) the ratio on these boxes is 2.13 (245 against $\Sigma 1/p = 115.2$); and none of the 4,325 boxes below the threshold has $\det(\chi, \gamma) \equiv 0$ [computation, outside the rules; the square-root reading is not proved]. So on these boxes the crowding just above the threshold is not a property of the boxes with $\det(\chi, \gamma) \equiv 0$: it reads as a crowding common to both classes just above the bound of each element, with the bound for $\det(\chi, \gamma) \equiv 0$ sitting at $X(b_1) = 4$, where the boxes with $\det(\chi, \gamma) \not\equiv 0$ are already two units above theirs; this accounts for where the excess sits and not for its size [computation; a rule fixed before counting; this reading is not a theorem]. A ninth rule fixed before counting predicted the coefficient of this square-root shape curve by curve from the archimedean place. Write $s = \text{arch} + \text{nonp} + \text{bad}$, where $\text{arch} = (\lambda'_\infty(b_1) - m_\infty)/\log p$, $\text{nonp} = 2 \log d'/\log p$ with d' the part of the denominator of $x(b_1)$ prime to p, and bad is the excess of the component corrections at b_1 over their minima; under the hypotheses of Proposition 24^c all three are ≥ 0 [paper, under the hypotheses of Proposition 24^c]. Parametrise each real component of E(R) by $z = t\omega_1$ (or $t\omega_1 + \omega_2/2$ on the second component, when there is one), $t \in \mathbb{R}/\mathbb{Z}$. If λ'_∞ has non-degenerate minima t_j at the level m_∞ on the component of b_1, with second derivatives c_j in t, and $t(b_1)$ is uniformly distributed on that component, then $P(s \leq a) \approx C(E)\sqrt{a}$ for small a, with $C(E) = \Sigma_j \, \text{mult}_j \cdot 2\sqrt{(2 \log p/c_j)}$, where $\text{mult}_j = 1$ if $t_j \in \{0, 1/2\}$ and 2 otherwise [paper, by the Morse lemma in one variable; the uniformity of $t(b_1)$ is an assumption]; a point on a component whose minimum lies above m_∞, or with $\text{nonp} + \text{bad} > a$, has $P(s \leq a) = 0$. With $q = e^{2\pi i \tau}$ and $\tau = \omega_2/\omega_1$ normalised to $\text{Im}\, \tau > 0$ and $\text{Re}\, \tau \in \{0, 1/2\}$, the second derivative at the 2-division point $t = 1/2$ of the identity component is $2\pi^2(1 + 8 \, \Sigma_\{n \geq 1\} \, q^n/(1 + q^n)^2)$, and at $t = 1/2$ on the second component it is $16\pi^2 \, \Sigma_\{k \geq 0\} \, q^{k+1/2}/(1 + q^{k+1/2})^2$ [paper, from the q-expansion of the local height, whose normalisation we have not checked against the source; the two expressions agree with numerical second differences on 198 components to a relative $3 \cdot 10^{-6}$]. The minimum at $t = 1/2$ of the identity component degenerates where the first expression

vanishes ($q \approx -0.107$); near there the shape is a fourth root rather than a square root. The square-root law also accounts for the $\sqrt{(2a)}$ above: if $P(s \le a) = C\sqrt{a}$ up to $a = 0.5$, then $P(s \le a \mid s < 0.5) = \sqrt{(2a)}$ whatever C is, so the information about the curve lies in C and not in the 2. The rule took the boxes with $C(E) > 0$ and no flag of degeneracy and compared three things: k, the ratio of the observed to the predicted least-squares coefficient of $P(s \le a)$ on $\sqrt{a}$ over $a \in [10^{-4}, 10^{-2}]$; g, the bias of C(E) among the boxes with $s \le 10^{-2}$ relative to the bias the prediction implies (1 if the prediction holds, 0 if s does not depend on C(E)); and a two-sided Mann–Whitney test of C(E) between the boxes with $s \le 10^{-2}$ and the others. The verdict that the prediction holds required the test to reject at level 0.05 in the predicted direction and the 95 % bootstrap intervals of k and g both to contain 1. On 10,346 further boxes at $p \in \{5, 7\}$ from curves with $|A| \le 1000$ and $|B| \le 10^4$ whose (A, B) had not been drawn before (10,280 with $\beta(E)/\log p \in [0.4, 2.0]$, of which 1,523 have $C(E) > 0$ and no flag, and 211 of these have $s \le 10^{-2}$), $k = 1.10$ (0.96 to 1.25), $g = 0.76$ (0.42 to 1.10) and the test gives $p = 6{\cdot}10^{-6}$, and the verdict of the rule is that the prediction holds. Permuting C(E) among these boxes removes the dependence ($p = 0.80$; over 1,000 permutations the test rejects in 5.0 % of them), and none of the 8,646 boxes predicted to have $P(s \le 10^{-2}) = 0$ has $s \le 10^{-2}$. On the 13,920 boxes of the eighth rule, whose square-root shape had already been seen, the same quantities are $k = 1.04$ and $g = 1.16$ [computation; numerical m_∞ and minima; a rule fixed before counting; the uniformity of $t(b_1)$ is assumed, not proved]. Outside the rules, the same uniform measure, applied to each of the 1,123 boxes with $\det(\chi, \gamma) \equiv 0$ and b_1 of depth 2 with its own discrete data (whether the denominator of $x(b_1)$ is a power of p, the component corrections, the component), predicts 189 of them in [4, 4.5) against 192 observed, a ratio of 2.05 against 2.07: given those discrete data, the size of the excess is accounted for by the archimedean place, while how often b_1 is a point whose denominator is a power of p, and why $\det(\chi, \gamma) \equiv 0$ occurs with frequency $1/p$ above the threshold, are not [computation, outside the rules, after the fact on the same boxes]. A tenth rule fixed before counting asked whether the departure of $t(b_1)$ from uniformity comes from the competition between b_1 and the second minimum of K_1. Since b_1 is primitive, b_1 is a shortest vector if and only if $\hat{h}(b_1) \le \hat{h}(b_2)$, where b_2 completes a reduced basis; writing $\hat{h}(b_1) = \lambda'_\infty(t) + N_1$, where N_1 collects the logarithm of the denominator of $x(b_1)$ and the component corrections and does not depend on t, this says $\lambda'_\infty(t) \le \hat{h}(b_2) - N_1$ [paper; an identity]. The model M takes t uniform on its component conditioned only on this inequality, with N_1 and $\hat{h}(b_2)$ held fixed [an assumption]. Under M the fraction V/L is uniform on [0, 1], where V and L are the lengths of the parts of the component on which λ'_∞ is at most $\lambda'_\infty(t(b_1))$ and at most $\hat{h}(b_2) - N_1$; and on the boxes of the ninth rule the coefficient C(E) is multiplied by 1/L [paper, under M]. Before counting we found that on those boxes b_1 is much shorter than b_2 ($\hat{h}(b_2)/\hat{h}(b_1) \ge 2.5$ for 94 % of them), so that L is within 10^{-3} of 1 for at least 90 % of them and M changes the predicted coefficient by 0.1 %: M cannot account for $k = 1.10$ and $g = 0.76$, and the rule therefore also tested V/L on all boxes, in the five classes of $\hat{h}(b_2)/\hat{h}(b_1)$ used above. On 12,868 further boxes at $p \in \{5, 7\}$ from curves with $|A| \le 1000$ and $|B| \le 10^4$ whose (A, B) had not been drawn before (12,807 with $\beta(E)/\log p \in [0.4, 2.0]$, 1,890 of them with $C(E) > 0$ and no flag, 263 of these with $s \le 10^{-2}$), the coefficient gives $k = 1.14$ (1.00 to 1.28) and $g = 0.61$ (0.31 to 0.92), against 0.2 % from M, and the test of V/L rejects in the classes with ratio below 1.1 and at least 2.5; the verdict of the rule is that neither part holds. The second part of the rule was mis-specified: the classes are defined through $\hat{h}(b_1)$, which depends on t, so that even under M the class with ratio near 1 selects large V/L and the class with large ratio selects small V/L. With the reference corrected for this (V uniform on the interval allowed by the class), on the new boxes the four classes with ratio below 2.5 agree with M and only the class with ratio at least 2.5 does not; cutting instead by L, which does not depend on t, the boxes with $L < 0.8$ agree with M on both sets of boxes (and those

with $0.8 \le L < 0.95$ on the new boxes), while those with L within 10^{-3} of 1 have t closer to the minimum of λ'_∞ than uniform (mean of V/L 0.475 and 0.477 on the new boxes and on those of the ninth rule) [computation; numerical m_∞; the corrected reference and the cut by L were made after the rule was fixed and after the data were seen]. The competition with b_2 thus describes $t(b_1)$ where b_2 is close, but the departure in the coefficient comes from boxes where b_2 is far, and its origin is not known [computation]. The cause of the effect is not known [computation].

The same boxes cut by conductor, and cut by h_min. Cut by conductor the ratio is not monotone — 0.218, 0.316, 0.144, 0.279 on (0, 20000], (20000, 50000], (50000, 100000], (100000, 200000] — so the conductor is not the variable. The table below re-cuts three families of §5.6 and this section by u, the excess being k := v_p(Reg_p) − (baseline), the baseline max(r − 2, 0) in the split case and r − 2 in the non-split case of rank ≥ 2:

family	boxes	u < 2	2–4	4–6	6–8	8–12	≥ 12	all	P(k ≥ 2 | k ≥ 1): observed / Σ1/p
rank 1, split (conductor ≤ 12000)	1,006	—	0.70	0.86	0.47	0.92	1.13	**1.00** (135/135.2)	21 / 20.9
rank 2, non-split (conductor ≤ 200000)	24,299	**0.92**	0.98	1.00	1.07	0.95	0.77	**0.97** (2,489/2,556.5)	352 / 325.1
rank 2, split (conductor ≤ 200000)	3,056	**0.00**	**0.06**	0.95	0.90	0.53	1.16	0.25 (100/405.4)	25 / 13.6

(In this table h_min is taken over all non-zero elements of $\Lambda \cap E_1$, as in Proposition 24, and not only outside pΛ as in the table above; on the split boxes of rank two the two cuts give the same ratios to two decimals. Rank one has no stratum $u < 2$, since there $\ker \chi = p\Lambda$ and $h_\min = p^2\hat{h}(P)$.) The first row reproduces the rank-one distribution 871 / 114 / 17 / 4 of sweep E, and the second the excesses 21,810 / 2,137 / 306 / 39 / 7 for k = 0, . . . , 4. **An excess that asks for no denominator follows 1/p in every stratum**: the non-split excess is $\det T \equiv 0$, a condition on a Schur complement and not on a denominator, and it sees no archimedean obstruction. The obstruction acts on the split boxes of rank two at $k \ge 1$ alone. The last column of the split row (25 against 13.6) is not explained here.

Lemmas 23 and 23′ as counted, and the next layer. Extending the sweep to conductor (200000, 500000] gives 50,707 boxes, 5,572 split: Lemma 23 is violated ($D_Q \ge 2$ with $h(x(Q)) < 4 \log p$) in **0**, Lemma 23′ in 0 of 152,121 elements, $\det(\chi, \gamma) \equiv 0$ occurs in 283 against 270.4 expected above the canonical threshold u = 4 (ratio 1.047; above the naive threshold the expectation is about 493, ratio 0.57, from upper bounds; above the threshold $4 \log p + \beta(E)$ of Proposition 24^c, 283 against 284.4, ratio 0.995, and none below it among 3,251 boxes, table 5.12^c), and $D_Q \ge 3$ holds for the shortest element in 14 boxes against 15.5 expected above 6 log p. That count depends on the element: 27 for another choice of element, 2 when three elements are required to agree, 100 when any one of three suffices — for $D_1 = 2$ the depth of $Q + pP_1$ drops to 2. The invariant of the box is $\min(D_Q, D_1) \ge 3$, which holds in the two boxes below. Among the shortest candidates with $s \le 2 \log 2$ (1,709 elements) all but 17 have $c' = 1$, and those 17 have $s = 2 \log 2$ exactly, the equality case of Lemma 23′. The archimedean excess $\max(0, \log|x(Q)|)$ saturates near 5 as s grows, so the proportion with $c' = 1$ falls from 97 % for $s < 2$ to 0.9 % for $s \ge 12$. The index of the p-adic closure of Λ is p^m with $m = \min(D_Q, D_1) - 1$; over conductor ≤ 200000 m

≥ 2 occurs in no box (0.66 expected among the 308 boxes with $u \geq 6$), and over conductor ≤ 500000 in exactly **two, 328563a1 and 477950bg1 at p = 7** (2.47 expected), both with D_Q = $D_1 = 3$. At both, γ vanishes at every point computed (21 of 21 at 328563a1), $\gamma_2 \equiv \chi \pmod 7$, M_$\Lambda$ mod 7 is [[1, 3], [3, 2]] and [[5, 3], [3, 6]] of rank one, and $v_7(\mathrm{Reg}_7) = 1$, as the paragraph on small closures in §3.15 predicts; at every point with $\chi(P) \neq 0$ among the 1,008 pairs of points computed there, $v_7(n^2 h_p(P)) = 0$.

Corollaries R′_s and K′(i) in rank two. Over the split boxes of rank two with conductor ≤ 20000 — 111 boxes (27, 31, 19, 16, 3, 15 by p; the smallest conductors 4407, 4629, 4976, 4279, 2343, 3333) — ρ_1 was measured in 109, the two boxes with a generator in E_1(Q_5), 7387a1 and 8962b1 at p = 5, being left out of that sweep; in those 109, v_p(Reg_p) = 0 $\iff \rho_1 = 2$ holds in **109 of 109**, and $\rho_1 = 2 \iff (\chi, \gamma)$ onto in **109 of 109**, (χ, γ) being computed with no p-adic height. At 7387a1 and 8962b1, (χ, γ) is onto (D_Q = 1) and v_p(Reg_p) = 0, so v_p(Reg_p) = 0 $\iff (\chi, \gamma)$ onto holds in **111 of 111**; the conclusion is unchanged. The three boxes where (χ, γ) is not onto are 8577a1 (p = 5) and 12208i1 (p = 13), with $\rho_1 = 1$ and v_p(Reg_p) = 1, and 16011e1 (p = 5), with $\rho_1 = 0$ and v_p(Reg_p) = 2: the three steps r − 2, r − 1, r of Corollary R′_s all occur.

The p-adic L-function at the split boxes, and a positive control. By §6(c), $\det(\chi, \gamma) \equiv 0$ in a split box of rank two satisfying (H) forces the coefficient c_2 of T^2 in L_p(E, T) to vanish modulo p, by Kato's divisibility alone and with no assumption on Sha; this is a test with a known answer of the computation of $\det(\chi, \gamma)$ and of c_2. A rule fixed before counting computed c_2 modulo p as Σ_j (j choose 2)·S_j (§6) on two sets of boxes: the 3,056 split boxes of rank two above, and the 1,674 boxes with conductor at most $3 \cdot 10^7$ among the boxes of the eighth, ninth and tenth rules ($p \in \{5, 7\}$, curves with $|A| \leq 1000$ and $|B| \leq 10^4$), the conductor bound being fixed by the running time and not by $\det(\chi, \gamma)$ or c_2. The symbols S_j were obtained from the p − 1 twisted values $L(E, \psi, 1)$, ψ of order p and conductor p^2, by an approximate functional equation with root number w_E·$\psi(N)G(\psi)^2/p^2$; on 14 pairs of rank two with conductor below 950 this agrees with the modular symbols of PARI to 10^{-14} and its prediction of c_2 modulo p with `mspadicseries` in 14 of 14, and on 120 of the boxes it agrees with PARI's `lfuntwist` to $5 \cdot 10^{-14}$, with no disagreement in c_2 modulo p. At 446d1 and p = 5 the coefficient of T^2 of P_5 is $2 \cdot 5 + 2 \cdot 5^2 + 5^3 + 5^4$ modulo 5^4, twice the value $5 + 5^2 + 3 \cdot 5^3$ of [SW, §3.3] (the factor 2 is the ratio of the Néron period to E.omega[1]). (H) holds in 2,987 and 1,656 of the boxes (E[p] is irreducible in all but 15, and some $q \parallel N$ has $p \nmid \mathrm{ord}_q(\Delta_\mathrm{min})$ in all but 78); in every box $c_1 \equiv 0 \pmod p$, as rank two forces, the S_j are integers, the root number is +1 and the given generators are saturated at p. **Among the boxes satisfying (H), $\det(\chi, \gamma) \equiv 0$ occurs in 95 and 205, and c_2 ≡ 0 (mod p) in all 300**; c_2 ≡ 0 occurs in none of the 2,892 and 1,451 boxes with $\det(\chi, \gamma) \not\equiv 0$, against about 1/p of them if c_2 were a random residue [computation, floating point, no interval arithmetic]. The first count is the positive control, and it passes. The second is not a theorem: under (H) and the equality of Skinner and Urban it says that $p \nmid \#Sha(E/Q)$ in these boxes, which is expected and is not proved by it. What it does prove, by §6(c) and with the computed value of c_2 as the only numerical input, is the converse reading box by box: **for each of the 4,343 boxes with (H) and c_2 ≢ 0 (mod p), rank E(Q) = 2, Sha(E/Q)[p^∞] = 0 and v_p(Reg_p) = 0** [known theorems combined, the statements checked in [Kato04], [PR92], [Sch85], [SU] and [Ski] (§6); computation for c_2]. The value v_p(Reg_p) = 0 obtained in this way agrees in all of them with the value given by $\det(\chi, \gamma) \not\equiv 0$ through Corollaries K′(i) and R′_s. A second rule asked that c_3 vanish modulo p with frequency about 1/p in the boxes with $\det(\chi, \gamma) \equiv 0$, as a negative control; it vanished in all 300, because S_j = S_{c−j} with $c \equiv -\log_p\langle N\rangle / \log_p(1 + p) \pmod p$ in every box — the functional equation of L_p seen modulo p — so that $2c_3 \equiv (c - 2)c_2 \pmod p$: the control was mis-designed and c_3 is not independent of c_2 [computation; the symmetry

was found after the data were seen].

5.13 Propositions 21 and 22 on rational points

Proposition 21 as displayed. Five curves of rank two at p = 5 — 4407a1, 5956a1, 8577a1, 75393a1, 188032i1 — with P = u·g ($|u| \le 2$, $\chi(P) \neq 0$), Z = w·g ($|w| \le 2$, $w \neq 0$) and a = 1, 2: F is read off the denominator of x(n(P + p^aZ)), C(P) from φ_p', φ_p and $2y + a_1x + a_3$ at P′, and ζ from p·nZ in E(Q_p), the congruence being assembled from these raw parts by a separate program. The displayed congruence holds in **2,400 of 2,400** pairs at a = 1 and **2,400 of 2,400** at a = 2. Dropping the term $C(P)\zeta(P)$, or reversing its sign, leaves 560 of 2,400 — the pairs with $\chi(Z) = 0$, where both sides vanish, the pairs with $\zeta(P) = 0$, and coincidences at the rate 1/p. Steps (1) and (2) of the proof, $\Theta(P + S) - \Theta(P) \equiv C(P)\zeta(S)$ and $q(P + S) - q(P) - q(S) \equiv C(P)\zeta(S)$ for $S \in \ker \chi$, hold in 400 of 400. In rank one with Z = P, $F((1 + p)P) \equiv F(P) + (q(P) - C(P)\zeta(P)/2) \pmod p$ holds in **643 of 643** boxes of the filter of §5.10 and $F((1 + p^2)P) \equiv F(P) + p(q(P) - C(P)\zeta(P)/2) \pmod{p^2}$ in **574 of 574** (the boxes with coordinates too large for the computation left out), against 96 of 643 with the term $C(P)\zeta(P)$ dropped; $\zeta(P) = 0$ in exactly the 97 boxes with $e \ge 3$.

Proposition 21 (iii) and (ii) realised. Points P with $v_5(q(P)) \ge 3$ and $S \in \ker \chi$ with p^2S of depth exactly three, at 4407a1 and 5956a1 (p = 5), both of rank two, so that $v_p(n^2h_p(P))$ is read pointwise in place of v_p(Reg_p): in **32 pairs of 32** the points P and $P + p^2S$ have F mod p^2 in common and are separated by min(v_p, 3). An explicit pair, recomputed for this version by a separate script: at 4407a1, p = 5, n = 2, $P = -5g_1 + 6g_2$ and $R = 25(g_1 - 2g_2)$ give $F(P) \equiv 49$ and $F(P + R) \equiv 24 \pmod{125}$, so F agrees modulo 25, and $v_5(n^2h_p) = 7$ and 2. With $R = 125(g_1 - 2g_2)$ the two points share F mod p^3 and are separated by min(v_p, 4) (2 of 2). With $R = 25(-g_1 + g_2)$ instead — $\chi(Z) \neq 0$, $\zeta(Z) = 0$ — the two points share t_1 mod p^4, $F(P + R) \equiv 29$ differs from 49 at valuation exactly one, and $v_5(n^2h_p) = 7$ and 2: the necessity of the second digit of F, realised by two points of one curve. At 8577a1, 75393a1 and 188032i1 there is no such S in the range searched, since γ vanishes on $\ker \chi$ there (the boxes with $\det(\chi, \gamma) \equiv 0$ of §5.12): the hypothesis $\gamma|_\{\ker \chi\} \not\equiv 0$ of (iii) is needed, and this is its negative control. A search that allows only translates $R = p^2Z$ with $\chi(Z) \neq 0$ finds 80 separating pairs at 4407a1 and 5956a1, and in all 80 the second digit of F moves as well; that is part (ii) of Proposition 21 ($e(P) = 2$ and $p \mid q(P)$ in all 80), and not an identity.

Proposition 22 as displayed. Over the 648 boxes of §5.10 (e distributed as [551, 82, 13, 2], which reproduces the filter) the congruence $F \equiv A/2 - pA/2 + pA^2/8 \pmod{p^2}$ holds in **648 of 648**, and agrees in 648 of 648 with an independent inversion of Theorem N* by the series of log and exp. Dropping $c/(p^2x(P''))$ leaves 72 of the 551 boxes with e = 2 and all 97 with $e \ge 3$, where that term has valuation $2e - 2 \ge 4$ and is invisible; dropping $(p^2/2)\Theta^2$ leaves 111 of 648, and dropping $-pA/2$ leaves 204.

6. What is proved in Lean and what is not

Thirty files, each checking from `import Mathlib` alone: `PadicHeightValuation{,2,...,29}.lean` and `PadicHeightValuationCanonical.lean` (namespace `PadicCanonicalHeight`), in the namespace `PadicHeight` except file 23 (`PadicHeightStar`), file 24 (`PadicHeightLayer`), file 25 (`PadicHeightCM`), files 26–29, whose namespaces are listed below, and the canonical file; and a thirty-first, `AnomalousEulerFactor.lean`, which belongs to a separate note. Files 22 to 25 hold the arithmetic of §§3.9, 3.11–3.12 and 3.13 — Hasse's two cases for $p \mid \#E(F_p)$, the coefficients

of $\mathcal{L}$, the digit at which each term of Theorem N* is visible, and the comparison of the two rates of Proposition 20 with the divisibility step that carries it to the minimality of Theorem $G^{(k)}$. File 24 holds nineteen statements, with no warning and no `sorry`; the rates themselves (v_p(ΔΘ) = j − 1 and v_p(Δz) = 1 + j) enter it as hypotheses, since they need the formal group. File 25 holds eighteen statements on the two places where c enters: that the lag of the ledger is uniform in m and uses neither $c \neq 0$ nor v_p(μ_{2m}) ≥ 0, the condition for the m = 1 term of Proposition 20 (iii) to lead and its automatic case $p \nmid c$, the cancellation criterion (1 + w)^{2m} = 1 of the boundary row with its two verdicts at p = 7, the residue arithmetic of $4p = 1 + 3f^2$ that admits p = 7 and p = 19 and excludes p = 13, and the emptiness of the list of local invariants before the eleventh digit at a cell with c = 0. Files 18–21 hold the arithmetic of §3.14 and §4.13: the two isogeny identities of Lemma 13 as identities in a commutative ring, the residue step of the equivalence (R′) ∧ (L), the cancellations 2n_p + 1 = p and $4n_p^2 - 1 = (2n_p-1)(2n_p+1)$, the local expansion $D^2H/DH = 2\kappa - 1$ and the telescoping Σ_{k<i}(q^k−1)/(1−q^k) = −i of §4.13(c), the balanced representation of every integer $\geq -n_p$ as pm + j with $|j| \leq n_p$, the counting identities for m_S and m_K, and the exponent arithmetic behind $\varepsilon_p = -1 \iff p \equiv 3 \pmod 8$. Every statement is about integers, matrices, finite sums or p-adic congruences; Mathlib has neither the p-adic sigma function nor division polynomials nor Vélu's formulas nor the Tate curve, so the geometry and the p-adic analysis enter as hypotheses. The companion `statements-and-dependencies.md` lists all Lean names; the main line is

Statement	Lean name	File
Theorem A	`norm_det_div_sq_le`	1–4
Theorem B, step 3	`fermatQuot_sub_iff_of_sq_eq_cube`, `dvd_three_mul_sub_two_mul`, `sub_ne_zero_of_two_mul_eq_three_mul`	1–4, 8
Theorem B, the model with $p \mid a_1$	`sq_sub_cube_dvd_of_dvd_a1` (and `sq_sub_cube_dvd_of_sq_dvd_psi` for $e \geq 2$)	8
Lemmas 4′ and 5	`eval_sub_eval_dvd_sq`	1–4
Corollary D	`sq_dvd_pow_sub_one_of_isPow`	1–4
Corollary E	`sq_dvd_alpha_sub_beta_of_isPow`	1–4
Theorem H	`fermatQuot_scale`	1–4
Theorem J	`dvd_det_of_col_dvd`, `pow_succ_dvd_det_of_col`	1–4
Main theorem, linear algebra	`regulator_valuation_of_split_anomalous`	5
Main theorem (ii), equivalence	`regulator_valuation_of_split_anomalous_iff_split`	5
§4.6 (2), the single slope and the affine zero	`slope_of_rho`, `fermatQuot_deform`, `affine_zero_unique`, `lambda_eq_zero_iff_split`, `lambda_dichotomy`	5
§4.6 (3), the weight count and the Newton polygon	`weight_split`, `twelve_dvd_weight`, `sum_pole_orders`, `cusp_sum_eq_m`, `mul_compl_le_half_mul_half`, `eq_C_mul_of_dvd_of_natDegree_eq`	6
§4.6 (4), the residues	`inv_one_sub_add_inv_one_sub_inv`, `first_term_residue`, `second_term_leading_ne_zero`, `residues_do_not_cancel`, `twelve_mul_sum_cusp_orders`	7
Theorem S, the ledger	`exists_shift_dvd_a1`, `add_val_exactly_one`, `not_dvd_of_val_one`	8
Main theorem ($\Longrightarrow$ of (i))	`quadratic_value_mem_of_entries_mem`	8
Theorem R, the linear algebra (R0)	`bilin_rank_one_of_zero_iff`, `not_dvd_of_congr_ne_zero`	9
Corollary R′, the determinant	`det_conj_diagonal`, `det_conj_diagonal_one_p`, `reg_eq_of_det_eq`	9

Statement	Lean name	File
Theorem R_s, the linear algebra (R0s) and rank ≤ 2	bilin_eq_of_vanishing_on_hyperplane, bilin_zero_of_mem_radical, gram_det_three_eq_zero	10
Corollary R′_s, the determinant	det_two_dvd_iff, pow_dvd_det_of_offdiag_block	10
Proposition R″_s, the second Fermat quotient	fermatQuot2_add, dvd_three_mul_sub_two_mul_two, split_const_eq	10
Proposition R‴ (the slope) and Corollary R⁗	eval_shift_of_deriv_eq, sq_dvd_pow_add_sub, fermatQuot2_move, lambda2_affine, lambda2_const_of_slope_zero, lambda2_lift_bijective, lambda2_unique_zero_lift, b_eq_slope_mul_kappa	11
Lemma 7, the first digit of the x-coordinate	eval_shift_first_digit, first_digit_eq, eval_shift_depth_two, first_digit_ne_zero_iff	12
Theorem K and Corollary K′, the bilinear algebra and the three layers	master, master_symm, master_add_left, master_smul_left, master_vanish_on_ker_chi, master_mem_radical, master_eq_of_gam_eq_smul, master_offdiag_ne_iff, master_diag_eq_zero_iff, master_bad_class_unique, det_three_layer, det_eq_zero_of_rank_one_block, det_sq_dvd_iff	12
Proposition 9, the second digit under lifting	X_pow_four_dvd_sub_trunc, eval_quartic_dvd, eval_shift_second_digit, fermatQuot_second_move	13
Proposition 10, the bad lift and the layers	root_factor, dvd_iff_dvd_sub_root, unique_bad_class, exists_bad_class	13
Theorem N, the ledger and the cancellation of a_1	second_digit_zero_iff, second_digit_criterion, a1_correction_cancels	13
Corollary N′ and the three-fold split at the second digit	dvd_iff_dvd_first, sq_dvd_sub_crit, val_one_of_crit_ne, sq_dvd_of_crit, trichotomy, min_two_ne_iff, min_two_eq_of_two_le, exists_same_data_diff_val, q2_additive, scaling_shift_square	14
Corollary $N^{(3)}$ and Corollary N″, the four-fold split	inv_pred_cube, neg_inv_pred_cube, q2_quotient, q2_quotient_dvd, sq_cube_step, denominator_third_digit, cube_dvd_sub_crit, cube_dvd_of_crit, not_cube_dvd_of_crit_ne, cube_dvd_iff_crit, quadrichotomy, exists_same_second_diff_third, min_three_ne_iff, min_three_eq_of_three_le	15
Proposition 12, Corollary $N^{(4)}$ and Corollary N‴, the five-fold split	low_terms_vanish, p_mult_second_digit, w_local, inv_x_correction, w_ne_zero_iff, inv_pred_quartic, neg_inv_pred_quartic, denominator_fourth_digit, theta_eta_cancels, cubic_term_vanishes, fourth_digit_reduction, quartic_dvd_iff_crit, exists_same_third_diff_fourth, min_four_ne_iff, min_four_eq_of_four_le, quintichotomy	16
Theorem G′ (the sensitivity) and the sharpness in Theorem G″	pow_sub_pow_expand, exists_exact_part, unit_pow_diff_exact, shift_depth_exact, shift_depth_dvd, quotient_by_p_sq, theta_sensitivity_exact, theta_digit_determined, three_layers_determine, layer_three_moves_digit_two, layer_digit_correspondence, nondegeneracy_needed	17
§4.12, v_p(ψ_p′(x(T))) = 1 exactly, and (ND) as a condition on ψ_p mod p^3	sq_valuation_one, psi_deriv_valuation_one, bracket_divisible, nd_iff_bracket, nd_of_bracket_exact	18

Statement	Lean name	File
Proposition 16, the weight of Res_x(G, ψ_p mod p) and m_p = m_K + m_S	resultant_weight, resultant_weight_split, m_p_integral, m_g_integral, m_K_add_m_S, resultant_weight_K, resultant_weight_S, twentyfour_dvd_sq_sub_one	18
Theorem ND, the step "the constant is non-zero $\Longrightarrow$ (ND)"	ne_zero_of_const_mul_pow, eval_ne_zero_of_isCoprime, nondegeneracy_summary, nd_iff_S_ne_zero	18
§3.14, D kills Δ_0; the cusp count of §4.13; c_S $\neq 0$ given γ_p $\neq 0$; the chain rule for D	serre_deriv_disc_zero, gauss_sum_closed, min_sum_closed, cusp_ord_total, resultant_weight_dPsi, m_S_eq, weight_two_p_vanishes, pow_card_self, cS_ne_zero, deriv_zero, deriv_one, deriv_sum, deriv_pow_succ, deriv_eval_chain	19
Lemma 13, and the reduction of Proposition 15 to (R′) $\wedge$ (L)	isogeny_A1, isogeny_A1_domain, isogeny_A2_core, isogeny_A2, deriv_eval_of_factor, residue_iff, const_mod_p_deriv, lead_cancel, lc_difference, deg_K_eq, weight_L, two_n_add_one_eq_zero	20
§4.13(b), (c): the balanced exponents, $D^2H/DH = 2\kappa - 1$, the telescoped sum, i $\mapsto$ p − i	balanced_repr, D_eval_zero, D2_eval_zero, log_deriv_ratio, telescope_term, telescope_sum, residue_value, residue_value_zmod, sign_flip_invariance	21
Proposition 16, m_K, its weight, and ε_p = $-1 \Longleftrightarrow$ p $\equiv 3$ (mod 8)	ordq_total, weight_ResK, mK_even, exponent_two_nsq, exponent_case_1mod4, exponent_case_3mod4, sign_exponent, eps_cases	21
Proposition R^+, the p-part of the reduction and Hasse's two cases	card_cases, two_p_forces_five, sq_not_dvd, dvd_of_unit_mul, dvd_of_unit_sq_mul	22
The coefficients of σ_p in a general model, and the depth of a term of a logarithm	sigma_third_coeff, sigma_fourth_coeff, padicVal_eq_zero_of_lt, log_term_depth_small, log_term_depth_five, log_term_depth_ne_five, log_term_factor, five_fifth_term	22
Proposition 16, the constant c_p in one line	cp_one_line	22
Proposition 17, the coefficients of $\mathcal{L}$ and the reparametrisation in t	mu2_eq, mu4_eq, mu6_eq, mu6_no_b2_cube, lambda3_eq	23
Proposition 18, the digit at which each term is visible	lambda_depth_mono, lambda_depth_m2_e2, x_form_suffices, val_thirty, val_three_sixty, val_thirty_seven, mu6_visible_at_ten, theta_depth_j5_p5, theta_depth_j5_p7, log_term_j10_five, log_term_j10_seven	23
Proposition 19 and Theorem $G^{(k)}$, the depth of a layer and the growth of the data	layer_depth, data_count_linear	23
Proposition 20 and Theorem $G^{(k)}$, the two rates and the minimality at every k	theta_rate_digit, ledger_rate_digit, gap_is_e, ledger_lags, ledger_lags_general, uniform_in_k, exact_pred_not_at, no_cancel, at_most_one_vanishes, layer_is_needed, fermat_digit, fermat_digits_needed, data_count, data_count_step	24

Statement	Lean name	File
Proposition 20 (iii), the boundary row and the CM cells	`ledger_lags_any_m`, `cm_lags_ten`, `m1_leads_iff`, `m1_leads_of_unit_c`, `boundary_cancel_m1`, `boundary_cancel_pow`, `boundary_cancel_of_card`, `boundary_p7_m3`, `boundary_p7_exception`, `boundary_m1_vs_m3_p7`, `cm_prime_mod_three`, `cm_no_anomalous_of_mod_two`, `cm_seven_nineteen`, `cm_not_thirteen`, `cm_ledger_invisible_before_eleventh`, `cm_no_local_invariant_before_eleventh`, `layer_is_needed_cm`, `cm_gap_at_k`	25
Corollary K′(i) and Corollary R′_s at r = 2, as linear algebra	`prime_dvd_det_two_iff`, `det_two_split_ne_zero_iff`, `exists_pair_eq_one`, `onto_iff_det_ne_zero`, `onto_iff_reduced_det_ne_zero`	26
Proposition V (§6, conditional), the integer arithmetic of its four cases	`leading_val_ge_rank`, `rank_le_leading_val`, `leading_val_eq_rank_of_nonsplit_rank_one`, `leading_val_gt_rank_of_split_le_one`, `leading_val_eq_rank_iff_of_normal`, `exists_two_ranks_same_leading_val`, `admissible_leading_two_of_rank_le_two`	26
Lemma 23, depth and height	`log_denom_ge`, `height_ge_of_depth`, `depth_lt_of_height_lt`	27
Proposition 24, the surjectivity of (χ, γ) from one shallow element of ker χ	`surjective_pair_iff`, `surjective_of_shallow_kernel_element`	27
Proposition 24^c, the lower bound R ≥ β(E)	`beta_le_remainder_of_local_bounds` (hypotheses: $0 \le \log c'$, m_∞ ≤ λ′_∞, m_ℓ ≤ corr_ℓ at each ℓ, and the decomposition R = 2 log c′ + λ′_∞ + Σ corr_ℓ)	Canonical
Proposition 24^c, depth against canonical height	`canonical_ge_of_depth`, `depth_lt_of_canonical_lt` (hypotheses: $\hat{h} = 2D \log p + R$, $\beta \le R$, $k \le D$, $0 \le \log p$)	Canonical
Proposition 24^c, the surjectivity	`surjective_of_canonical_lt_four_log_add_beta` (hypotheses: χ onto; $\chi(y) = \gamma(y) = 0 \Rightarrow D(y) \ge 2$; the decomposition and $\beta \le R$ at every element; one element of ker χ with $\hat{h} < 4 \log p + \beta$), `surjective_of_naive_or_canonical` (below either threshold), `surjective_pair_iff'`	Canonical
§3.15, the constants m_ℓ at I_N	`four_mul_comp_le_sq`, `comp_le_half` (hypothesis: $i \le N$)	Canonical
The difference of the two slacks	`remainder_sub_slack` (hypotheses: the naive and the canonical decompositions)	Canonical
§5.12, the index of the p-adic closure and the null model for det(χ, γ)	`index_zero_iff`, `index_ge_two_iff`, `index_le`, `det_zero_iff`, `card_det_zero_of_ne`, `card_det_zero_of_fst_eq_zero`, `density_det_zero`	27
Lemma 23′, the slack	`cofactor_log_le`, `cofactor_eq_one_of_slack_lt`, `max_eq_iff`	28
§3.15, small closures: the layers factor through χ	`factor_through_chi`, `two_ne_zero_of_ne_two`, `diag_ne_zero`	28
Proposition 20 (v) read on a pair of points, and a count of §5.12	`exact_of_deeper`, `pair_law_j3`, `pair_law_F_leads`, `card_two_routes`	28
Proposition 21, steps (2)–(4) and parts (i)–(iii)	`ratio_of_ker_le`, `vanish_of_chi_zero`, `translate_quadratic`, `second_difference`, `shift_quadratic`, `ledger_digit`, `kappa_ne_zero`, `separates`, `counterexample_digits`	29
Proposition 22, the inversion at k = 2	`inverse_n3`	29

Statement	Lean name	File
The Euler factor at an anomalous prime (companion note)	`ap_of_anomalous`, `unit_root_identity`, `sub_one_associated_p`, `sub_one_associated_p_of_pred`	`AnomalousEulerFactor`

How the main theorem appears there. `regulator_valuation_of_split_anomalous` takes as hypotheses that the entries of the height matrix on Λ_0 are divisible by p and that det M_{Λ_0} = p^2·det M_Λ, and concludes p^{r−2} | det M_Λ; `regulator_valuation_of_split_anomalous_iff_split` takes the two halves — split $\Longrightarrow$ Reg_p $\in$ Z_p, and non-split $\Longrightarrow$ Reg_p = z/p with p $\nmid$ z — and turns them into the equivalence of (ii). That is, **integrality of the height matrix is a hypothesis there, not a conclusion**: what the kernel checks is the bookkeeping, and the geometry that produces the hypotheses is on paper.

Trust base. `#print axioms` is run at the end of each of the files, and a log is kept beside each source. Every line reads `depends on axioms:` followed by a subset of [propext, Classical.choice, Quot.sound], or else "does not depend on any axioms" (`deriv_zero` and `deriv_one` of file 19). There is no `sorryAx` and no `Lean.ofReduceBool`, that is no `native_decide`, and each file checks with exit code 0. Files 1–21 check with no warnings; **files 22 and 23 emit warnings and no errors** — one unused-binder linter note in file 22 and three `Try this` suggestions from `decide` in file 23 — so the claim of a silent build is made for the first twenty-one files, and for files 26–29 and `AnomalousEulerFactor`, which also check with no warning. Files 24 and 25 are not part of that claim: they check with exit code 0 and the axioms above, but whether they check with no warning at all was not re-checked for this version. Two things are stated exactly. First, **one `decide` is load-bearing**: `twelve_ne_zero` of file 7 closes with `interval_cases p <;> revert hp hd <;> decide`, and since that is the kernel's `decide` and not `native_decide`, no extra axiom appears in any log. Second, the `#print axioms` lines of files 12 and 13 are two fewer than the theorems those files prove, the omissions being `X_pow_two_dvd_sub_trunc`, `eval_sq_dvd`, `X_pow_four_dvd_sub_trunc` and `sq_dvd_pow_add_sub`; the last of these appears in the table above, so "every theorem listed" is not literally true of them, and they are covered only by the file checking clean.

What file 17 does and does not do. It records the valuation arithmetic of Theorem G′: the isolation of the linear term of a binomial expansion, the exactness of the valuation of (A + D)^n − A^n for a unit A and n prime to p, the depth count p·g·Δ + p·Δ^2 ·B + Δ^5·C, the division by p^2, and the two directions (determination and sharpness). The three geometric inputs — that x ∘ τ_T lies in Z_p[[t]] with unit derivative, that all divided derivatives of φ_p below the p-th are divisible by p, and the non-degeneracy (ND) — enter as hypotheses, in the shape of the displayed decomposition and of ¬ `p | g`. `nondegeneracy_needed` exhibits a witness showing that the conclusion fails when `p | g`, so the hypothesis is not decorative.

What files 18–21 leave on paper, and why Theorem ND is not a Lean theorem. Files 18–21 hold valuation arithmetic, commutative-ring identities, finite sums and exponent counts, and one non-vanishing argument over a field. The following steps of §3.14 and §4.13 are on paper, and each of them is needed for Theorem ND:

1. the canonical subgroup C = Ê[p] and the congruence M ≡ A_0 (mod p) for its kernel polynomial — the heart of Proposition 14;
2. the injectivity of q-expansions in a fixed weight (kernel generated by A − 1), which is what lifts Proposition 15 from the Tate family to every p;
3. the uniformisation of the Tate curve and the theta product for x_Q(W_1) − x_Q(W_2);

4. the resultants themselves — Res_X(DΨ, Ψ) = γ_pΔ0^M, Res_x(K, ψ_p) = c_KΔ0^{m_K}, Res_x(S1, ψ_p) = c_SΔ0^{m_S}: Lean has the weights and the q-orders, while the step "no weight-homogeneous irreducible factor other than Δ0" is on paper, resting on the dichotomy of §4.6(3);

5. γ_p ≠ 0 itself (`cS_ne_zero` receives it as a hypothesis), which is the leading-coefficient count at the cusp;

6. the root form of the Kodaira–Spencer identity (R′), and φ1 = −2σ, σ = −E2/24 in (L).

So **Theorem ND is not a Lean theorem**, and neither are Propositions 14, 15 and 16; their grade is [paper], as §3.14 states. What files 18–21 add is that the arithmetic along which those proofs run is kernel-checked, in exactly the sense in which files 1–17 do it for §§3.2–3.11.

What is not in Lean. The p-adic sigma function and the formula for h_p; Mazur and Tate's differential equation for σ_p, the Weierstrass ℘-series and the formal logarithm, so that **Theorem N*, Propositions 17, 18, 19 and 20, Lemma 21 and Theorem $G^{(k)}$ are not Lean theorems** — what files 22, 23 and 24 hold is the coefficient arithmetic, the valuation arithmetic and the digit comparisons they consist of, the analysis being a hypothesis; the Iwasawa logarithm; division polynomials, their composition rule and their shape modulo p; Vélu's formulas and the degree count of Theorem C; [p] = V ∘ F and the group scheme E[p]; Serre–Tate theory; the Tate parametrisation and the valuations read off it; Deuring's lifting theorem; the structure of E(Q_p) used in Lemma 4; the height pairing itself. So **the main theorem is not a Lean theorem**: what is machine-checked is the arithmetic skeleton along which its proof runs, and its grade is [paper].

Files 26–29 and `AnomalousEulerFactor`. File 26 (`PadicHeightLeading`, twelve statements) holds the linear algebra of Corollary K′(i) and Corollary R′_s at r = 2 and the integer arithmetic of Proposition V below; file 27 (`PadicDepthHeight`, `PadicSurjectivity`, `PadicClosureIndex`, `PadicNullModel`, twelve statements) the inequality of Lemma 23 given the divisibility p^{2k} | d^2, the surjectivity step of Proposition 24 and the counts behind §5.12; file 28 (`PadicCofactor`, `PadicClosureCollapse`, `PadicPairLaw`, `PadicTwoRoutes`, ten statements) Lemma 23′ and the factorisation through χ of §3.15; file 29 (`PadicTranslateLinear`, `PadicTranslateDigit`, `PadicInverseLedger`, ten statements) the linear algebra and the digit bookkeeping of Propositions 21 and 22. `AnomalousEulerFactor.lean` (`AnomalousEuler`, four statements) is the arithmetic of a separate note. All five check from `import Mathlib` with exit code 0, no warning, no `sorry` and no `native_decide`, and every `#print axioms` line reads a subset of [propext, Classical.choice, Quot.sound]. On paper remain: that Q ∈ E_k(Q_p) forces p^k | d(Q) (Lemma 23); step (1) of Proposition 21, which uses that φ_p mod p is a polynomial in x^p and the invariant differential; the isometry of log on 1 + pZ_p; and Proposition 20 (iv). So **Propositions 21, 22, 24 and 24^c are not Lean theorems**; their grade is [paper].

`PadicHeightValuationCanonical` (namespace `PadicCanonicalHeight`, nine statements) holds the arithmetic and the logic of Proposition 24^c: the termwise lower bound, the inequality between depth and canonical height, the surjectivity, and the constants of I_N; it checks from `import Mathlib` with exit code 0, no warning, no `sorry` and no `native_decide`, and its nine `#print axioms` lines read [propext, Classical.choice, Quot.sound]. The local height formulas, the decomposition $\hat{h}$ = Σ_v λ′_v, the table of component corrections and the value of m_∞ are not in Lean; they are hypotheses of the Lean statements. Lemma 25 is not in Lean.

One conditional statement, kept apart. File 26 also holds the arithmetic of the

one statement in this paper that assumes a conjecture. It is stated here, and not in §3, because every theorem of §3 is unconditional and the grades must not mix; a separate note *The valuation of the leading term of the p-adic L-function at an anomalous prime* (the file `note-padic-L-leading-term.md`) states it with its evidence.

Proposition V (the leading term bounds the rank from above) [paper; conditional on the p-adic conjecture of Birch and Swinnerton-Dyer of Mazur, Tate and Teitelbaum and on p ∤ #Sha; the integer arithmetic in Lean. *] Let $p \geq 5$ be a prime of good ordinary reduction, $p \nmid \prod_\ell c_\ell$, $p \nmid \#E(Q)_{tors}$, ι, $r = rank\ E(Q)$, and L_p the p-adic L-function of [MTT] on the unit-root eigenspace.* ***Assuming the p-adic BSD conjecture and p ∤ #Sha,***

$$v_p(L_p^{(r)}(E, \chi^0)/r!) \geq r$$

in each of four cases: non-anomalous, by Theorem A; split anomalous, as $2 + \max(r-2, 0) \geq r$ by Corollary R'_s; non-split anomalous of rank one, as $2 + (-1) = 1 = r$ exactly by Theorem S; non-split anomalous of rank $r \geq 2$, as $2 + (r-2) + v_p(\det T) \geq r$ by Corollary R'. The 2 is the valuation of the Euler factor $(1 - \alpha^{-1})^2$ at an anomalous prime, exactly 2 for every anomalous $p \geq 5$ (Lemma E of the note).

The bound is on the rank **from above**: a bound from below would need an upper bound for v_p(Reg_p), which the local data do not give (§7(3)); Proposition 24 gives one from an archimedean datum, and with it, in the split case of rank two and under the same assumptions, $v_p(L_p^{(2)}(E, \chi^0)/2!) = 2$ exactly whenever $h_{min} < 4 \log p$ or $\hat{h}_{min} < 4 \log p + \beta(E)$ (Propositions 24 and 24^c). The same value 2 of the leading valuation is compatible with every rank $r \leq 2$ (Lean `admissible_leading_two_of_rank_le_two`). This is not the exceptional zero of [MTT], which occurs at split multiplicative reduction and raises the order of vanishing; here the reduction is good ordinary, the order is unchanged, and it is the valuation of the leading coefficient that rises by 2.

Where the assumption can be replaced by theorems [known theorems, combined; the statements and their hypotheses are checked in the original papers [Kato04], [SU], [Ski], [Sch85], [PR92], [Gre99] and [GV]; the integral form of Kato's divisibility under (H) is the argument of [Ski, §2.5], which we have checked against the proof in [Kato04] (§13.14 and the results it uses)]. Write $L_p(E, T) = \sum_j c_j T^j$ with $1 + T = (1+p)^s$, as in [SW, (3.1)]. Then $L_p^{(r)}(E, \chi^0)/r! = c_r \cdot \log_p(1+p)^r$ whenever $c_0 = \ldots = c_{r-1} = 0$, so that $v_p(L_p^{(r)}(E, \chi^0)/r!) = v_p(c_r) + r$ [paper, from the definitions; at 446d1 and $p = 5$, PARI gives valuation 3 and c_2 has valuation 1, as in [SW, §3.3]]. Let $p \geq 5$ be a prime of good ordinary reduction with E[p] irreducible. (Every input of (a)–(c) is stated in its source for odd p or for $p \neq 2$, see the table of sources at $p = 3$ below; $p \geq 5$ is needed only where (c) uses the results of §3.) (a) **The inequality of Proposition V holds without the conjecture and without p ∤ #Sha:** L_p lies in $Z_p[[T]]$ [GV, Prop. 3.7], and r independent rational points give $\mathrm{ord}_T L_p \geq r$ through the bound $\mathrm{ord}_T f_E \geq r$ for the characteristic series f_E of the Selmer group over the cyclotomic Z_p-extension (Mazur's control theorem [Gre99, Thm 1.2]) and Kato's divisibility of L_p by f_E away from p [Kato04, Thm 17.4(1)–(2)], which needs only $p \nmid N$ and a_p a unit (the chain is written out in [SW, §10]); hence $c_0 = \ldots = c_{r-1} = 0$ and $v_p(L_p^{(r)}/r!) = v_p(c_r) + r \geq r$. The content of Proposition V is therefore in its cases of equality. (b) Assume moreover **(H): some prime q ∥ N has $p \nmid \mathrm{ord}_q(\Delta_{min})$.** Then the theorem of Skinner and Urban gives $(f_E) = (L_p)$ in $Z_p[[T]] \otimes Q_p$ [SU, Theorem 1 (Theorem 3.6.4 in the version on the second author's page) and Thm 3.6.9: p odd, E[p] irreducible, the condition (dist), automatic for good ordinary $p \geq 3$, and the prime q], and in $Z_p[[T]]$ itself once Kato's divisibility holds integrally; [Kato04, Thm 17.4(3)] asks for $p \neq 2$

and an image of Galois containing SL_2(Z_p), and [Ski, §2.5 and Thm 2.5.2] shows that E[p] irreducible and (H) suffice. In [Kato04] the hypothesis on the image enters Theorem 17.4(3) only through Theorem 12.5(4) (§17.13), whose proof (§13.14) uses three things: that every Galois-stable lattice is a scalar multiple of a fixed one (through Remark 12.8 and the proof of Lemma 14.7, the only place where the image is used; it follows from E[p] irreducible [Ski, Lemma 2.1.1]); that $H^1(T)$ is free [Kato04, Thm 12.4(3): $p \neq 2$ and E[p] irreducible]; and the Euler system bound [Kato04, Thm 13.4(3)], which asks for some σ in $\mathrm{Gal}(\bar{Q}/Q(\zeta_{p^\infty}))$ with $\mathrm{Coker}(1-\sigma)$ free of rank one. ([Kato04] does not prove Theorem 13.4 but quotes it from [Pe98], [Ru00] and [KK99] together. In [Pe98] the divisibility holds up to a power of p (Thms 1.3.3 and 1.3.5), and integrally (§5.3.6) when, among other conditions that we have not checked, the fixed space of σ on T/pT is one-dimensional, which is the condition above; we have not obtained [Ru00] or [KK99], so the source of Theorem 13.4(3) in exactly this form is not checked.) A generator of tame inertia at q gives such a σ: inertia at q acts on the Tate module through non-trivial unipotent matrices [DDT, Prop. 2.12(a)], non-trivial modulo p exactly when $p \nmid \mathrm{ord}_q(\Delta_{min})$ [DDT, Prop. 2.12(c)], and a unipotent matrix that is non-trivial modulo p has cokernel free of rank one. Section 13 of [Kato04] assumes that f has no complex multiplication, which holds here because j is not integral at q. With the formula for the leading term of f_E up to a unit of Z_p [Sch85, Thm 2′], [PR92, §3.4.3], a curve of rank two with $a_p = 1$ and $\mathrm{ord}_T L_p = 2$ has $\mathbf{v_p(c_2)} = \Sigma_\ell \mathbf{v_p(c_\ell)} + \mathbf{v_p(\#Sha(E/Q)[p^\infty])} + \mathbf{v_p(Reg_p)}$, and the inequality $\geq$ needs Kato's divisibility alone. In the boxes of Proposition 24^c that also satisfy (H), $v_p(L_p^{(2)}(E, \chi^0)/2!) = 2$ holds if and only if $Sha(E/Q)[p^\infty] = 0$ — the assumption $p \nmid \#Sha$ of Proposition V becomes the statement it is equivalent to. (c) Under (H), with $a_p = 1$ and two independent rational points, **if $c_2 \not\equiv 0 \pmod p$ then, by Kato's divisibility alone, rank $E(Q) = 2$, $Sha(E/Q)[p^\infty] = 0$, the p-adic height pairing is non-degenerate, $p \nmid \prod_\ell c_\ell$ and $v_p(Reg_p) = 0$** (that $\mathrm{ord}_T f_E = 2$ forces $Sha(E/Q)[p^\infty]$ to be finite and the pairing to be non-degenerate is [PR92, Thm 3.4.2 and §3.4.3] together with [Kato04, Thm 17.4(1)]; [Sch85, Thm 2′] is the converse) — the last by a route independent of Proposition 24^c — and, contrapositively, $\det(\chi, \gamma) \equiv 0$ in a split box forces $c_2 \equiv 0 \pmod p$ (Corollaries K′(i) and R′_s, applied to $\Lambda = E(Q)$, give $v_p(Reg_p) \geq 1$). c_2 modulo p is a finite sum of modular symbols with denominator p^2 [SW, Prop. 3.5]: with $S_j := \Sigma_{a=1}^{p-1} [\omega(a)(1+p)^j/p^2]^+$ it is $\Sigma_{j=0}^{p-1}$ (j choose 2)·S_j modulo p when $a_p \equiv 1 \pmod p$ and $L(E, 1) = 0$ [paper, from the Hecke relation; for general a_p the sum is multiplied by α^{-2}, α the unit root; checked against PARI's `mspadicseries` on 14 pairs (E, p) of rank two and conductor below 950, six of them with $a_p \equiv 1$]. The method is that of [SW, Algorithm 11.1], here at anomalous primes, and we claim nothing new in it; the p-adic L-functions of [SW] (period Ω_E), of PARI (period Ω^+ = E.omega[1], with $\Omega_E/\Omega^+ \in \{1, 2\}$ [SW, §3]), of [Kato04] (a Néron differential and an integral basis $\gamma^\pm$, §17.5) and of [SU] (cohomological periods, §3.3.3) differ by units of Z_p when p is odd and E[p] is irreducible: the Manin constant of an optimal curve is prime to p when $p \nmid 2N$ [Maz78, Cor. 4.1], and the isogenies in the class then have degree prime to p [paper, from the definitions: for the curve that is optimal for $X_1(N)$, pulling back identifies its integral cohomology with the λ_f-part of that of $X_1(N)$, so the periods of [SU, §3.3.3] are its Néron periods divided by its Manin constant, and Mazur's corollary reaches it through an isogeny of degree prime to p; [SU, §3.6.7] states the comparison of L_E with L_f without proof]. §5.12 applies (c) to the split boxes.

The first digit of the conjecture of Mazur, Tate and Teitelbaum [paper for the reduction; computation, against a rule fixed before counting, for the check; the conjecture is read in [MTT, II §10] and used in the form of [SW, Conj. 5.1]]. The conjecture asserts an exact equality and not an equality up to a unit [MTT, II §10]

(there L_p is not divided by a period, the leading term is (1/r!)·d^r/ds^r in the variable s, and the right-hand side carries Ω_E^+, the integral of the Néron differential over E(R), which is Ω_E [MTT, II §8]); in the form of [SW, §5.1], for good ordinary p, $L^*_p(E, 0) = \varepsilon_p \cdot \prod_v c_v \cdot \#Sha(E/Q) \cdot Reg_\gamma(E/Q) / (\#E(Q)_{tors})^2$, with $\varepsilon_p = (1 - 1/\alpha)^2$, $Reg_\gamma = Reg_p / \log_p(1 + p)^r$ and L_p normalised by the Néron period Ω_E [SW, §§3–5]. Statements (b) and (c) see it only up to a unit of Z_p. Its first p-adic digit can be read off in rank two with $a_p = 1$ and $c_2 \not\equiv 0 \pmod p$: $\alpha \equiv 1 - p \pmod{p^2}$ gives $\varepsilon_p / p^2 \equiv 1$ and $\log_p(1 + p)/p \equiv 1 \pmod p$, the sign of the height and any constant factor in it cancel in a 2×2 determinant (so Reg_p in our normalisation equals that of [SW]), and with c_2 taken in the normalisation of PARI by Ω^+ = E.omega[1] the conjecture says

$$c_2 \equiv \omega \cdot n \cdot \#Sha(E/Q) \cdot Reg_p \pmod p,\ \omega := \Omega_E / \Omega^+ \in \{1, 2\},\ n := \prod_\ell c_\ell, \quad (MTT_1)$$

where $\omega = 2$ exactly when $\Delta > 0$ [paper, from the definitions; the trivial torsion of the boxes below is used]. In a split box Theorem K gives $Reg_p \equiv -B_0^2 \cdot \det(\chi, \gamma)^2 \pmod p$ — the form $2\lambda_{\{T,1\}} \chi \otimes \chi + B_0(\chi \otimes \gamma + \gamma \otimes \chi)$ has determinant $-B_0^2$ in the coordinates (χ, γ) — and #Sha(E/Q), when finite, is a square (Cassels). So (MTT_1) implies, with $(\cdot / p)$ the Legendre symbol,

$$(-c_2 \cdot n \cdot \omega \ / \ p) = +1, \quad (Q)$$

a statement in which neither the regulator nor Sha appears: under the conjecture, the quadratic character of c_2 is fixed by the Tamagawa product and the sign of the discriminant. Neither (MTT_1) nor (Q) follows from (b) or (c). The constraints that the functional equation puts on the S_j ($S_j = S_{c-j}$, §5.12) and the vanishing of c_0 and c_1 are linear, and c_2 is a linear form that does not vanish on the space they cut out, so the structure alone does not fix the quadratic character of c_2; on random integer vectors satisfying these constraints it is a square in 0.49 to 0.58 of 5,000 cases for each $p \le 19$ [computation]. A rule fixed before counting tested both statements on the split boxes of rank two in Cremona's tables with $200000 < N \le 499000$ — 5,556 boxes with $p \in \{5, \dots, 19\}$, $p \nmid N$, $a_p = 1$, $p \nmid \prod_\ell c_\ell$, trivial torsion and ι, disjoint from those of §5.12; one of them was lost to memory in the computation of heights — with c_2 computed as in §5.12, Reg_p modulo p computed both from PARI's p-adic heights and from Theorem K, and #Sha replaced by the order $Sha_{an} = (L''(E, 1)/2)/(\Omega_E \cdot \prod_\ell c_\ell \cdot Reg_\infty)$ predicted by the complex conjecture. The two routes to Reg_p modulo p agree in 5,273 of 5,273 boxes. Of the 5,454 boxes satisfying (H), $c_2 \not\equiv 0$ in 5,180, which are exactly those with $v_p(Reg_p) = 0$; $Sha_{an} = 1$ in all of them, and **(Q) and (MTT_1) both hold in 5,180 of 5,180** [computation, floating point, no interval arithmetic]. If the quadratic character of c_2 were independent of n and ω, agreement in all of them would have probability 2^{-5180}; in each of the twelve strata by p and the sign of Δ the one-sided binomial test rejects that hypothesis (largest p-value $8 \cdot 10^{-28}$, Holm). The controls behave as the formula requires: replacing c_2 by a random unit gives agreement in 0.49 of the boxes; dropping the factor -1 gives agreement in none of the boxes with $p \equiv 3 \pmod 4$ and in all with $p \equiv 1 \pmod 4$; dropping ω leaves exactly the boxes with $\Delta < 0$ when $p \equiv \pm 3 \pmod 8$; dropping n leaves exactly the boxes with $(n/p) = +1$. On the 761 non-split boxes with $200000 < N \le 205000$, where Reg_p has no closed form modulo p, (Q) holds in 323 of the 671 with $c_2 \not\equiv 0$ (no prediction was made), while (MTT_1), with Reg_p computed, holds in 671 of 671. On the 4,343 boxes of §5.12 with $c_2 \not\equiv 0$, whose values of c_2 were known before this rule was fixed, (Q) holds in 4,343 of 4,343 and (MTT_1) in the 2,901 of them with $N \le 600000$ where Sha_{an} was computed. So the conjecture of Mazur,

Tate and Teitelbaum agrees with the numbers at its first digit, unit included, in every box tested — a check that the main conjecture cannot supply, and not a proof of anything; it rests on the computed c_2, on Sha_an in place of #Sha, and on the form of the conjecture as stated in [SW]. The normalisations it confirms — the factor ω and the sign — are those of [SW], with L_p computed from twisted L-values, so that no Manin constant enters.

The second digit, at the boxes where the first one vanishes [paper for the reduction; computation, against a rule fixed before counting, for the check; the conjecture in the form of [SW, Conj. 5.1]]. Without dividing, the conjecture says c_2 = ω·n·#Sha(E/Q)·U·Reg_p with U := (ε_p/p^2)·(p/log_p(1 + p))^2, an equality in Q_p (rank two, trivial torsion). For a_p = 1 one has $\alpha \equiv 1 - p - p^2 \pmod{p^3}$, hence ε_p/p^2 $\equiv$ 1 + 4p, log_p(1 + p)/p $\equiv$ 1 − p/2 and U $\equiv$ 1 + 5p (mod p^2). In a split box with det(χ, γ) $\equiv$ 0, Theorem K gives Reg_p $\equiv$ 0 (mod p) and (c) gives c_2 $\equiv$ 0 (mod p), so (MTT_1) reads 0 $\equiv$ 0. One digit further, the conjecture says

> c_2/p $\equiv$ ω·n·#Sha(E/Q)·(Reg_p/p) (mod p) when v_p(Reg_p) = 1 and p $\nmid$ #Sha(E/Q), (MTT_2)

and c_2 $\equiv$ 0 (mod p^2) when v_p(Reg_p) $\geq$ 2 or p | #Sha(E/Q) (then p^2 | #Sha, Cassels); U enters only modulo p, where it is 1. A box with v_p(Reg_p) = 1 and c_2 $\equiv$ 0 (mod p^2) would, under the conjecture, have p | #Sha(E/Q): this is where the p-part of Sha would be read from the side of the L-function. Since L_p $\equiv$ P_3 modulo (1 + T)^{p^2} − 1, whose coefficients of T^k for k < p are divisible by p^2, c_2 modulo p^2 is the coefficient of T^2 in P_3, built from μ_α on the classes modulo p^3; we computed it from the twisted L-values of the characters of conductor p^3 and p^2 (for the latter through the Hecke relation Θ^{(3)} = a_p·Θ^{(2)}), as in §5.12, and checked it against PARI's modular-symbol series (`mspadicseries`) on 14 pairs of conductor $\leq$ 944, among them 446d1 with p = 5, where it gives c_2 $\equiv$ 10 (mod 25), twice the value of [SW]. Theorem K gives no closed form for Reg_p/p. We computed Reg_p modulo p^{v+2} from PARI's p-adic heights and, independently of PARI's algorithm for heights, from a p-adic σ function built from the formal group together with the denominators of multiples of the points (sharing with PARI only s_2 and the formal expansions); the two agree in 674 of 674 boxes. The rule covered all 574 split boxes with det(χ, γ) $\equiv$ 0 available to us: the 274 boxes with (H) and c_2 $\equiv$ 0 among those of the previous paragraph, and the 300 boxes of §5.12 (whose first digit c_2 $\equiv$ 0 had been seen). Sha_an = 1 in all of them. In the 398 boxes with v_p(Reg_p) = 1, **(MTT_2) holds in 398 of 398** (agreement by chance has probability 1/p; rejected in each of the six classes of p, Holm); in the 176 boxes with v_p(Reg_p) $\geq$ 2, **c_2 $\equiv$ 0 (mod p^2) in 176 of 176** [computation, floating point without interval arithmetic, Sha_an in place of #Sha]. Read the other way, as in (c): in each of the 398 boxes, (H), the computed c_2 $\not\equiv$ 0 (mod p^2) and v_p(Reg_p) $\geq$ 1 (the split corollary, the generators being saturated at p) give by Kato's divisibility alone 1 = v_p(c_2) $\geq$ v_p(#Sha(E/Q)[p^∞]) + v_p(Reg_p) $\geq$ v_p(#Sha(E/Q)[p^∞]) + 1, hence rank E(Q) = 2, Sha(E/Q)[p^∞] = 0 and v_p(Reg_p) = 1 for these curves and primes [known theorems combined, as in (c); computation for c_2]. As a positive control, in the 100 boxes of smallest conductor with c_2 $\not\equiv$ 0 of the previous paragraph, c_2 $\equiv$ ω·n·Sha_an·U·Reg_p (mod p^2) in 100 of 100; replacing U by 1 leaves exactly the 34 boxes with p = 5, where U $\equiv$ 1 (mod 25). Dropping ω or n from (MTT_2) leaves exactly the boxes in which that factor is $\equiv$ 1 (mod p), and replacing Reg_p/p by a random unit gives agreement at the rate 1/(p − 1). So the conjecture agrees with the numbers at the second digit as well, in every box tested. But no box had p | Sha_an, so the case in which the p-part of Sha would be read from the L-function did not occur, and the check does not separate the conjecture from the weaker statement with #Sha replaced by 1.

It is a check and not a proof; it rests on c_2 computed in floating point, on Sha_an, on s_2 from PARI, and on the form of the conjecture in [SW].

Relation with the refined conjectures of Mazur and Tate [the first page of [MT87] in the original; its Conjecture 4 from secondary sources]. On its first page [MT87] states its conjectures at a finite layer M, without p-adic L-functions, and notes that even at the layer M = p the conjecture is not implied by the analogous one for the p-adic L-function: it predicts congruences modulo divisors of p − 1. As restated in [PB19, §2] and [Ota, Conj. 2.4], the leading-term formula of [MT87, Conj. 4] concerns levels in which the primes of good reduction occur to the first power, with the tamely ramified trivialisation at those primes (values in F_p^×). The coefficient c_2 modulo p (resp. p^2) is the image of the p-part of the Mazur–Tate element of level p^2 (resp. p^3) in $I^2/I^3 \cong \mathbb{Z}/p$ (resp. $\mathbb{Z}/p^2$), on the wildly ramified quotient. As far as these restatements show, (MTT_1) and (MTT_2) lie outside the leading-term formula of [MT87, Conj. 4] and are instances of [MTT] at the first layers of the cyclotomic Z_p-extension. The form $-B_0{}^2\cdot\det(\chi,\gamma)^2$, with χ the discrete logarithm on E(F_p) ≅ Z/p, resembles the discrete-logarithm regulators of [MT87], and the boxes with $\det(\chi,\gamma)\equiv 0$ resemble the situation, a vanishing regulator, addressed by the derived heights of [BD94]; we have not checked either resemblance against the sources. Numerical studies of [MT87] in rank one and at split multiplicative levels are [PB19] and [LC24]; [BH25] proves large parts of the conjectures of [MT87].

Where the p-part of Sha **enters [computation, against a rule fixed before counting, for the check; paper for the reduction; the conjecture in the form of [SW, Conj. 5.1]].** The case left open in the previous paragraph does not occur in rank two within Cremona's tables: in the files `allbsd` of [Cre] (conductor < 500000; 348,672 curves of rank two) the analytic order Sha_an of every curve of rank two is 1 or 4 (4 for 42 curves), so no curve of rank two has p | Sha_an with p ≥ 3; on the 42 curves with Sha_an = 4 and on all 742 curves of rank two in three windows of conductor, PARI's value of Sha_an agrees with the tables. We therefore took the check one rank down. In rank zero it is empty: the interpolation property gives $c_0 = \varepsilon_p\cdot L(E,1)/\Omega^+$, and Sha_an is defined from the same number. In rank one, with P a generator and t = #E(Q)_tors, the height of [SW, (4.1)] is $2\cdot\log_p(e(P)/\sigma_p(P))$, the negative of h_p(P), and the conjecture reads $c_1 = -\omega\cdot n\cdot\#Sha(E/Q)\cdot U_1\cdot p\cdot h_p(P)/t^2$ with $U_1 := (\varepsilon_p/p^2)\cdot(p/\log_p(1+p)) \equiv 1 + 9p/2 \pmod{p^2}$ for p ≥ 5 (for p = 3 the term $p^2/3$ of the logarithm gives $U_1 \equiv 7 \pmod 9$). With e := 1 + v_p(h_p(P)) + v_p(#Sha(E/Q)), it says v_p(c_1) = e and

$$c_1/p^e \equiv -\omega\cdot n\cdot(\#Sha/p^{v_p(\#Sha)})\cdot(h_p(P)/p^{v_p(h_p(P))})\cdot t^{-2} \pmod p,$$

with #Sha = #Sha(E/Q); here, by the integrality theorem of the introduction (when P reduces onto E(F_p)), v_p(h_p(P)) = −1 in the non-split case and ≥ 0 in the split case. Reading it needs c_1 modulo p^{e+1}, the coefficient of T in P_{e+2}. For P_m with m ≥ 4 the Hecke relation of the previous paragraph acquires a second term, $\Theta^{(m)}_{pk} = a_p\cdot\Theta^{(m-1)}_{k} - p\cdot\Theta^{(m-2)}_{k/p}$ when p | k (the sum of $\chi(c)\cdot x(c/p^{m-2})$ over the units c modulo p^{m-1} is $p\cdot\Theta^{(m-2)}$ when χ has conductor at most p^{m-2}); it does not arise for m = 3. With it, the coefficients c_1 and c_2 of P_m agree with PARI's modular-symbol series on 25 pairs (p = 3, 5, 7, m ≤ 6, nine of them with a_p = 1), and for 446d1 with p = 5 we find c_2 ≡ 185 (mod 625) and c_3 ≡ 295 (mod 625), twice the values of [SW]. The rule covered every box of rank one in the tables with p^2 | Sha_an, p ≥ 3, p ∤ N, a_p = 1, $p \nmid \prod_\ell c_\ell$, p ∤ t and (H): 41 boxes, 40 with p = 3 and Sha_an = 9 and one with p = 5 and Sha_an = 25 (the other 64 candidates, among them all but one with p = 5 and both with p = 7, have E[p] reducible). Sha_an from the tables and from PARI agree in all

41. In **41 of 41 boxes v_p(c_1) = e** — the valuation exceeds by exactly v_p(Sha_an) = 2 the value predicted with Sha replaced by 1, which agrees in 0 of 41 — **and the first digit agrees in 41 of 41** (agreement by chance has probability p^{−3} per box; rejected for p = 3 and for p = 5, Holm) [computation, floating point without interval arithmetic, Sha_an in place of #Sha]. The sign and the factor U_1 were fixed before on 90 boxes of rank one with Sha_an = 1 (p = 3, 5, 7, conductor ≤ 20000): the sign −1 gives 90 of 90 and the opposite sign none, and in the 82 non-split boxes $c_1 \equiv -\omega \cdot n \cdot U_1 \cdot p \cdot h_p(P)/t^2 \pmod{p^2}$ holds in 82 of 82 and in none with U_1 replaced by 1. The sign is that of the form of [SW], whose height 2 log_p(e/σ) [SW, (4.1)] is −h_p; the analytic height of [MTT, II §5], ⟨P, P⟩ = log_p(σ^2/d) with d the denominator of x(P), has the sign of h_p, so that with the height as printed there the formula of [MTT, II §10] has the opposite sign in odd rank ([Sch85, Thm 6] relates the algebraic and the analytic height by a sign); in rank two the sign cancels in the determinant [paper, from the definitions; computation for the 90 boxes]. The implementation of P_m reproduces c_2 modulo p^2 in 50 of 50 boxes of the previous paragraph. So, at the first digit and in every box available, the p-part of Sha predicted by the complex conjecture is read correctly from the p-adic L-function: the p-adic and the complex conjectures agree on the p-part of Sha there. In rank one this agreement is largely forced by theorems. Kato's divisibility, the theorem of Skinner and Urban (whose hypotheses include (H) and allow p = 3: [SU] assumes p odd, and its condition (dist) holds for good ordinary p ≥ 3) and the leading term on the algebraic side give v_p(c_1) = 1 + v_p(h_p(P)) + v_p(#Sha(E/Q)[p^∞]), so that, read the other way, the computed valuations give #Sha(E/Q)[p^∞] = p^2 for these 41 curves [known theorems combined; computation for c_1]; for semistable curves [JSW] proves the p-part of the complex formula; and we expect, without having checked the sources, that the p-adic and complex Gross–Zagier formulas make the first digit a consequence as well. The check is therefore mainly one of P_m, of the heights and of the normalisations; it is a check and not a proof, and in rank two the case p | Sha remains untested within the tables.

A search outside the tables [computation, against a rule fixed before counting; the heuristics recalled, not checked]. To find a box of rank two with p^2 | Sha_an, p ≥ 5, a_p = 1 and (H) we searched quadratic twists E^{(d)}. Twisting keeps E[p] irreducible and keeps (H) at the primes q ∤ d, and a_p(E^{(d)}) = χ_d(p)·a_p(E), so from a curve with a_p(E) = ±1 the twists with χ_d(p) = a_p(E) have a_p = 1. Nothing links Sha(E)[p] to Sha(E^{(d)})[p]; as bases we took the 30 curves of rank zero in [Cre] with 25 | Sha_an (and the other conditions) of smallest $\Omega_E \cdot \prod_\ell c_\ell$, which raises the scale $\sqrt{|d|}/(\Omega_E \cdot \prod_\ell c_\ell)$ of Sha·Reg. Among the twists with root number +1 and conductor at most $1.2 \cdot 10^9$, and a part, cut by time, of those up to $3.1 \cdot 10^{10}$, 82 have analytic rank two and 2-descent upper bound two; for 60 of them PARI found two points, which we saturated at all primes up to 50, and Sha_an = 1 in all 60. The same scan with p = 3 (conductor at most $2.9 \cdot 10^9$) gave 30 twists of rank two, 24 of them read, all with Sha_an = 1. In the same scans the twists of rank zero have Sha_an > 1 in 1278 of 1730 cases (25 | Sha_an in 303 of them), while in rank two the product Sha_an·Reg, of comparable or larger size (median 37 to 140), is carried entirely by the regulator. The same functions return Sha_an = 4 on all 39 curves of rank two with Sha_an = 4 in [Cre] on which we ran them. So no box of rank two with p | Sha_an was found in this range. This is consistent with heuristics of Delaunay's type, which as we recall them (without having checked) give a probability of about p^{−(2r+1)} for p | #Sha in rank r, and which the model of [PR] is stated to be compatible with. Even if such a box exists, reading c_2 modulo p^{e+1}, e = v_p(Reg_p) + v_p(#Sha), by the method of the previous paragraph needs about p^{e+2}·$\sqrt{N}$ terms of the L-series; for p = 5 and e = 2 this limits the conductor to about $5 \cdot 10^7$ within our memory bound. The general form v_p(c_2) = v_p(Reg_p) + v_p(#Sha) of the conjecture therefore remains untested here in the case p | Sha.

A box of rank two with $3^2 \mid$ Sha_an [computation, against a rule fixed before counting; Sha_an in place of #Sha; one example]. First, the models. A model that is not minimal at q changes ord_q(Δ) by a multiple of 12, which can change the condition $p \nmid$ ord_q(Δ) of (H) when $p \geq 5$. In every box of this section (H) was decided with the discriminant of the model obtained from the change of variables returned by PARI's `ellglobalred`, which over Q is the minimal model; the models given were in fact minimal in all 11,694 boxes, and the Kodaira symbol I_n at each $q \parallel N$ gives the same n = ord_q(Δ_min) in all of them. So no decision about (H) changes. We then searched again with p = 3, where reading c_2 modulo p^3 is cheapest. As bases we took curves with $3 \nmid N$, a_3 = ± 1, E[3] irreducible, $3 \nmid t \cdot \prod_\ell c_\ell$ and (H), one for each j-invariant: 27 curves of rank zero in [Cre] with $3 \mid$ Sha_an and Sha_an ≥ 81, 23 curves of conductor at most 60000 of smallest Ω_E$\cdot\prod_\ell c_\ell \cdot$N^{1/4} (the inverse of the scale of Sha·Reg at the largest twist), the 40 curves of rank one with p = 3 two paragraphs above, and, cut by time, 114 further curves of the second kind. We took their quadratic twists with χ_d(3) = a_3(E), root number +1 and conductor at most $2.9 \cdot 10^9$, and decided (H) again on each twist with Δ_min. Of 415 twists of analytic rank two (2-descent upper bound two), 341 were read with two points saturated at all primes up to 50: Sha_an = 1 in 339 of them, Sha_an = 25 in one (with $5 \mid N$ and E[5] reducible) and Sha_an = 9 in one, the twist of 41213a1 by -263, with a-invariants [1, 0, 1, -94675796466, 11212598183927037] and conductor 2850661997, for which a_3 = 1, (H) holds, t = 1, $\prod_\ell$ c_ℓ = 4, and Sha_an = 9 at 64 and at 128 bits. In this box v_3(Reg_3) = 0 (from PARI's heights and from our own σ function), so the conjecture predicts v_3(c_2) = 2 and c_2/9 $\equiv \omega \cdot$n$\cdot$(Sha_an/9)$\cdot$Reg_3$\cdot$t^{-2} $\equiv 1 \pmod 3$. We computed c_2 modulo 27 from P_4, with about 9.5 million terms of the L-series and the coefficients a_n held as machine integers (they agree with PARI's `ellan`, and the calibration at 446d1 is reproduced): **c_2 $\equiv$ 9 (mod 27), so the valuation and the first digit agree** (agreement by chance has probability 1/27; with Sha replaced by 1 the predicted valuation is 0). The same programs give the predicted valuation and first digit in 20 of 20 boxes of rank two with p = 3 and Sha_an = 1 from [Cre] (conductor at most 6853), and reproduce 10 boxes of rank one and 20 of rank two of the previous paragraphs. Read the other way as in (c), c_2 $\not\equiv 0 \pmod{27}$ gives rank E(Q) = 2 and #Sha(E/Q)[3^∞] $\in \{1, 9\}$ by Kato's divisibility, and #Sha(E/Q)[3^∞] = 9 under the equality of Skinner and Urban [known theorems combined, their hypotheses at p = 3 checked only as in the paragraph on rank one; computation for c_2]. So in one curve of rank two the 3-part of Sha predicted by the complex conjecture is read correctly from the p-adic L-function. It is one example and not a test with power: 74 further twists of analytic rank two were not read (PARI found one point), and in the same scans the twists of rank zero have $9 \mid$ Sha_an in 820 of 3733 cases, against one in 341 in rank two. For rank two no theorem on the complex side gives the 3-part of the Birch and Swinnerton-Dyer formula for this curve.

The other twists of rank two [computation, against a rule fixed before counting; Sha_an in place of #Sha]. For the 74 twists of analytic rank two of the last paragraph that were not read, we looked for a second point in two ways: by giving PARI's `ellrank` the point already known (effort up to 16), and by searching the everywhere locally soluble 2-coverings returned by `ell2cover` for points of naive height up to 10^6. On 20 twists that had been read, with regulators between 609 and 1894, the first way found the second point from the first alone in all 20 cases (canonical heights up to 60). In 18 of the 74 twists two independent points were found, and after saturation at all primes up to 50, Sha_an = 1 in all 18. In these 18 boxes (conductor between $2.3 \cdot 10^7$ and $2.7 \cdot 10^9$, p = 3, a_3 = 1, (H)) the predicted valuation v_3(c_2) = v_3(Reg_3) + v_3($\prod_\ell$ c_ℓ) and the first digit agree in 18 of 18. Among them are a box with v_3(Reg_3) = 3, where c_2 $\equiv 54 \pmod{81}$ from P_5; a box where $9 \mid \prod_\ell$ c_ℓ supplies the valuation 2; and a box where v_3(Reg_3) = -2 (from both

height computations and PARI's `ellpadicregulator`; here a_3 = 1, so 3 divides #E(F_3)) is offset by 9 | $\prod$_ℓ c_ℓ. In that last box the first computation did not return rational values for the twisted L-values, and they were recomputed with a smaller truncation error; this was done after the first comparison had been seen. Since Sha_an = 1, these boxes test the factors Reg_3, $\prod$_ℓ c_ℓ, ω and t of the conjecture and not the factor Sha. For the other 56 twists no second point was found. If Sha_an were 9 for one of them, its regulator would be one ninth of the computed product Sha_an·Reg, and by Minkowski's bound for a lattice of rank two there would be a second point of canonical height at most (4/3)·Sha_an·Reg/(9·$\hat{h}$(P)), where P is the known point if it is the shortest; this bound is at most 60 for 48 of the 50 twists in which one point was known, the range of heights of the second points found in the 20 test cases above. With Sha_an = 1 the same bound is between 74 and 819. This is a heuristic remark, not an exclusion: the searches are not complete, and the known point need not be the shortest (in one twist it was divisible by 3). So no second box of rank two with 9 | Sha_an was found, and the twist of 41213a1 by −263 remains the only curve of rank two in our data in which the 3-part of Sha is read from the p-adic L-function. Altogether 359 of the 415 twists of rank two were read, with Sha_an = 9 once. For comparison, the model of [BKLPR] (Theorems 3.11 and 5.15, where it is shown to agree with Delaunay's heuristic as corrected there) gives the probability 1 − $\prod$_{i > r}(1 − p^{1−2i}), whose leading term is p^{−(2r+1)}, that p divides #Sha for curves of rank r ordered by height; for p = 3 and r = 2 this is about 0.0046. Our family, ordering and selection (only twists that were read) differ from that setting, so this is only an order of magnitude.

Two more boxes of rank two with 3^2 | Sha_an [computation, against a rule fixed before counting; Sha_an in place of #Sha]. We then extended the scan of the paragraph before last along its own order: the next bases of the second kind (keys Ω_E·$\prod$_ℓ c_ℓ·N^{1/4} from 1.83 to 4.36, one for each j-invariant and none sharing a j-invariant with an earlier base), with the same twists and the same rules, in four runs of 30 minutes, the fourth of them conditional. Over 435 bases, 1,573 twists of analytic rank two were found (all with (H) and a_3 = 1 on the twist) and 1,367 were read: Sha_an = 1 in 1,362, Sha_an = 4 in two (with 2 | N), Sha_an = 25 in one (with E[5] reducible), and Sha_an = 9 in two, the twist of 6083e1 by 565 (conductor 1941845675, $\Delta < 0$, $\prod$_ℓ c_ℓ = 28) and the twist of 57661b1 by −148 (conductor 1263006544, $\Delta > 0$, $\prod$_ℓ c_ℓ = 2). Both satisfy (H) on the twist with t = 1 and $3 \nmid \prod$_ℓ c_ℓ, and Sha_an = 9 at 64 and at 128 bits. In both, v_3(Reg_3) = 0 (PARI's heights, our own σ function and PARI's `ellpadicregulator` agree; the three share PARI's s_2), so the conjecture predicts v_3(c_2) = 2 and the first digits 2 and 1. P_4, after the calibration at 446d1 and the reproduction of the box of the paragraph before last had been repeated, gives **c_2 ≡ 18 and c_2 ≡ 9 (mod 27): the valuation and the first digit agree in both boxes** (by chance with probability 1/27 each and 1/729 together; with Sha replaced by 1 the predicted valuation is 0). Neither box needed a smaller truncation error. Read the other way as for the twist of 41213a1, c_2 ≢ 0 (mod 27) gives rank E(Q) = 2 and #Sha(E/Q)[3^∞] = 9 for both curves [known theorems combined, their hypotheses at p = 3 checked only as above; computation for c_2]. So three curves of rank two in our data now have the 3-part of Sha read from the p-adic L-function, all three with v_3(Reg_3) = 0 and with agreement at the valuation and the first digit. Counting all scans with p = 3, 1,726 of 1,988 twists of rank two were read and Sha_an = 9 in three of them (0.0017), about a third of the value of [BKLPR], with the same caveats; 206 of the new twists of rank two were not read, and no second point was looked for. Since p = 3 is anomalous in all these boxes (#E(F_3) = 3), we note how the normalisation behaves there. In [MTT, II §4] the height is defined on the subgroup of points that reduce to the identity at p and to the identity component at every prime, and extended Q_p-linearly to E(Q) ⊗ Q_p. On that subgroup it takes values in 3Z_3, since σ^2 has 3-integral coefficients when p ≠ 2 [MTT,

II §2], so v_3(Reg_3) is at least 2 minus twice the 3-adic valuation of the index of that subgroup, and negative values such as the -2 of the previous paragraph, where $9 \mid \prod_\ell c_\ell$, are allowed by the definition. The conjecture of [MTT, II §10] counts good anomalous reduction as non-exceptional, and the valuation 2 of $(1 - 1/\alpha)^2$ at $a_3 = 1$ is what the unit U absorbs, as for $p \geq 5$.

Sources at p = 3, the normalisation of σ, and what is not checked [paper, from the statements in the sources; computation for one curve]. The paragraphs above apply (a)–(c) at p = 3. The inputs, with the range of p in their statements:

Input	Range of p in the source	What p = 3 needs
Prop. 3.7 of [GV]	odd p (§3); E optimal or E[p] irreducible	E[3] irreducible
Thm 1.2 of [Gre99]	good ordinary reduction, no restriction on p	$a_3 \not\equiv 0 \pmod 3$
Thm 17.4(1)–(2) of [Kato04]	$p \nmid N$ and a_p a unit	$3 \nmid N$, $a_3 \not\equiv 0 \pmod 3$
Thm 17.4(3) of [Kato04] with [Ski, §2.5]	$p \neq 2$; [Ski] fixes an odd prime	E[3] irreducible and (H)
Theorem 1 and Thm 3.6.9 of [SU]	p odd; (dist) automatic for good ordinary $p \geq 3$	E[3] irreducible and (H)
Prop. 2.12(c) of [DDT]	any p different from q	$3 \nmid$ ord_q(Δ_min)
Thm 2′ of [Sch85], §3.4.3 of [PR92]	ordinary reduction, no restriction on p	nothing further
Cor. 4.1 of [Maz78]	the Manin constant is a unit of $\mathbb{Z}[1/2m]$, m^2 the largest square dividing N	$3 \nmid N$
Chapter II, §2 and §10 of [MTT]	residue characteristic $\neq 2$; non-exceptional means $\alpha \neq 1$	nothing further

None of them assumes $p \geq 5$. At p = 3 a model that is not minimal at q changes ord_q(Δ) by a multiple of 12, hence of 3, so the condition in (H) does not depend on minimality there. The image of Galois is a different matter: for $p \geq 5$, E[p] irreducible together with (H) gives an image containing SL_2(Z_p) by the classical lifting lemma, so that [Kato04, Thm 17.4(3)] applies as stated; at p = 3 that lemma is not available, and the argument of [Ski, §2.5] is what is used [recalled, not checked against the sources].

The constant of the sigma function: [MTT, II §2] defines e by $X_\omega = -(1/2)\cdot D_\omega\sigma^2 = x + (b_2 - e)/12$ and characterises it as the unique p-adic integer for which σ^2 has p-integral coefficients, when the residue characteristic is not 2. PARI's documentation characterises s_2 = `ellpadics2(E, p, n)`, for $p > 2$, as the unique c in 3^{−1}Z_p for which the odd solution $\sigma = t + O(t^2)$ of $-d((1/\sigma)\cdot d\sigma/\omega) = (x + c)\cdot\omega$ lies in tZ_p[[t]], and states $s_2 = b_2/12 - E_2/12$ with E_2 the value of Katz's p-adic Eisenstein series of weight two. The two equations coincide with $c = (b_2 - e)/12$, and the square of PARI's σ satisfies the conditions that determine σ^2 in [MTT, II §2], so **$e = b_2 - 12\cdot s_2$ at p = 3 as at $p \geq 5$** [paper, from the two definitions; the uniqueness in [MTT, II §2] is quoted there from earlier work and not followed]. For 26a1 at p = 3 ($a_3 = 1$), building σ from $c = s_2$ directly, without PARI's heights, gives coefficients that are 3-integral up to t^20, while $c = s_2 + 1/3$ and $c = s_2 + 3^k$ for $k \leq 3$ do not; this pins s_2 modulo 3^4 against the definition [computation]. Unlike the three computations of v_3(Reg_3) above, which share s_2, this checks s_2 against the definition itself.

At an anomalous prime the subgroup on which [MTT, II §4] defines the height has an index whose p-part comes from the reduction map from E(Q) to the points of the reduction over F_3, a group of order 3, and from the component groups, of order at most the p-part of $\prod_\ell c_\ell$; on that subgroup the height lies in $3\mathbb{Z}_3$, so $v_3(\mathrm{Reg}_3) \geq r - 2\cdot v_3$ of the index. With r = 2 and $3 \nmid \prod_\ell c_\ell$ this gives $v_3(\mathrm{Reg}_3) \geq 0$, and a negative value needs

3 | $\prod_\ell$ c_ℓ, as in the box with v_3(Reg_3) = −2 above, where 9 | $\prod_\ell$ c_ℓ [paper, from the definitions in [MTT, II §§2 and 4]].

What is not checked in the sources:

Item	Status
The source of [Kato04, Thm 13.4(3)] in its exact form	quoted by [Kato04] from [Pe98], [Ru00] and [KK99]; [Pe98] read, its integral form needs further conditions; [Ru00] and [KK99] not obtained
The number of [SU, Thm 3.6.4] in the published version	not checked; Theorem 1 of the introduction is cited as such by [Ski]
The paper [MT87] beyond its first page	read in restatements
The lifting lemma for the image at p ≥ 5, and the results of Serre on E[p] at q	recalled; [DDT] is a set of lecture notes
The paper [PR92] at p = 3	read once, not re-read for the range of p
Curves of rank two with 3 | Sha in the LMFDB	not available to us (reCAPTCHA)
The uniqueness of σ^2 in [MTT, II §2]	quoted there from earlier work, not followed

None of these is a hypothesis of a statement in this paper beyond those already marked as resting on [Kato04], [SU] and [MTT]: the chain uses Theorem 13.4(3) as stated in [Kato04], the equality of [SU] as stated in the version we read, and the form of the conjecture in [MTT].

7. What is open

1. **The quadratic residue class of the slope.** Over every anomalous curve over F_p with p ≤ 157 the slope ρ of §4.6 satisfies $(\rho/p) = (-3/p)$, so that -3ρ is a square and $\rho(\bar{x}) = -3\mu(\bar{x})^2$ for a homomorphism μ : E(F_p) → F_p determined up to sign. This is [computation] and nothing more; μ has no closed form here. At the cusp ρ_j ≈ $9j^2/\Delta$, and the shape $\rho = 9L(\bar{x})^2/\Delta$ with L a homomorphism is excluded by the residue class together with the squareness of $-\Delta$.

2. **One source under §4.6.** The step "E^{(p)} is the member of the family at parameter t^p" uses that the Tate parameter q, as a series in 1/j, has coefficients in the prime field. j(q) ∈ Z[[q]] is standard; the inverse series is what we did not locate in a source (§9), and [ATAEC] itself we did not retrieve.

3. **Exact values rather than bounds.** Corollaries R′ and R′_s turn part (iii) into v_p(Reg_p) = (r − 2) + v_p(det T), so what is left open is exactly whether det T is a unit — a non-degeneracy of p^{−1}h_p on the orthogonal complement of e_1, of the same kind as Schneider's conjecture, and not decided by any local datum here. It fails in 36 of the 7,759 non-split boxes and in 5 of the 48 split boxes of rank 3 (§5.6). On the split side of rank one the value is decided, by λ^{(2)} (Proposition R″_s), but λ^{(2)} is not a function of the reduction: it needs E mod p^3 and the point modulo p^2. The behaviour at p = 5 with #E(F_5) = 10 remains outside everything after Lemma 4. One archimedean datum decides the split case of rank two in part: v_p(Reg_p) = 0 whenever h_min < 4 log p (Proposition 24). Whether archimedean data decide it in general is open.

4. **The geometry in Lean.** §6 lists what would have to exist in Mathlib for the main theorem to become a Lean theorem.

5. **The layer one deeper.** Corollary K′(iv) reduces the exact value in rank ≥ 3 to $\det \hat{C} \not\equiv 0$, the non-degeneracy of p^{−1}h_p on $\Lambda_1 = \Lambda \cap E_2(Q_p)$. Nothing here decides it; the local data of Theorem K do not enter it.

6. **What q_p(d′) is.** Theorem N and Corollaries $N^{(3)}$, $N^{(4)}$ show that the digits of the height past the first are not functions of the position of the point in the filtration of E(Q_p): the displacement is carried by the Fermat quotient of the denominator d′, to one digit at the second layer and to two at the third. Two partial answers are in §4.12: the denominator descends at rate p^k while the σ-part descends at rate p^{ke−2}, so the two non-local ingredients of Lemma 3 arrive at different speeds and cannot be merged into one homomorphism; and q_p(d′) is paired with the normalisation of the model, since the right-hand sides of those theorems change by 2p·q_p(u) under $x \mapsto u^2x$. Theorem N* sharpens the first of the two: in the closed form the two non-filtration ingredients are not symmetric, because by Proposition 19 the third term $\mathcal{L}(z(P''))$ **is** a function of the layers t_1, while the Fermat quotient F is not a function of anything in E(Q_p). So the whole of the non-filtration content of the height sits in one of the two Iwasawa logarithms, and in no part of the formal group. An intrinsic description of q_p(d′) alone we still do not have, and by the remark on the model we no longer expect one.

7. **The class dependence of λ_T one digit deeper.** (L2) gives λ_T(m) = λ_{T,1}m^2 modulo p. Modulo p^2 neither Θ_T(m) = $m^2\Theta$_T(1) with the standard lift of m nor the same identity with the Teichmüller lift holds (§5.7), and no law is proposed here. Theorems N and $N^{(3)}$ do not need one, since the class of nP is fixed there.

8. **What is left around (ND).** The non-degeneracy itself is settled (Theorem ND). Three smaller things around it are not. (i) The primary sources for the classical inputs of §4.13 — [Kat73] on the canonical subgroup and the theta operator, [Igu] on the irreducibility of the tower, [SwD73] on the kernel of the q-expansion map, [ATAEC] for the Tate parametrisation — we did not retrieve; we used the statements as standard and verified their consequences on q-expansions ourselves. (ii) Whether Propositions 14 and 15 are in the literature we do not know; see §8. (iii) The counting of §4.13 gives the constants c_S, c_K, c_p in closed form, but the *ratio* of the two sides of (R′) at a supersingular point, where h degenerates, is outside everything here.

9. **What the minimality of Theorem $G^{(k)}$ does and does not say.** The necessity half is a statement about the parameter space that Theorem N* defines — t_1 mod p^{k+1} and F mod p^{k−1}, the local invariants fixed. Both kinds of necessity are realised by rational points of one curve: the layer γ_k by P and P + p^{k−1}S with S ∈ ker χ, $\gamma(S) \neq 0$ (Proposition 21 (iii); k = 3 and k = 4 at 4407a1, p = 5, read pointwise — v_p(n^2h_p(P)) in place of v_p(Reg_p) — since 4407a1 has rank two), and the second digit of F by P = $-5g_1 + 6g_2$ and P + $25(-g_1 + g_2)$ at the same curve (§5.13). What is not available on rational points is independence of the two kinds of datum along p-power translates (Propositions 21 and 22); item 14 records what is open beyond them.

10. **The valuation of the coefficients of $\mathcal{L}$.** Proposition 18 puts the m-th coefficient of $\mathcal{L}$ at the digit 2me − 2 + v_p(μ_{2m}), and v_p(μ_{2m}) can be negative: v_5(μ_6) = −1 is why μ_6 is visible one digit earlier at p = 5. All we have here is v_p(μ_{2m}) ≥ −v_p((2m + 2)!) from the shape of the double integral, which is far from sharp. Without a sharp bound the digit of a given local invariant is known for m ≤ 3 only, and the ordering of the invariants μ_4, μ_6, μ_8, … by depth is not known for all p. One special case the computation settles: in the twenty cells with j = 0 of §5.11 the series has **only the coefficients μ_{6k}** — $\mu_2 = \mu_4 = \mu_8 = \mu_{10} = \mu_{14} = \mu_{16} = 0$ while μ_6, $\mu_{12} \neq 0$, measured to z^{16} — which is [computation], and is why the leading term there is m = 3.

11. **Corollary N‴ has almost no test.** In conductor ≤ 20000 there is exactly **one** box with v_p(Reg_p) ≥ 4, so §5.8 records the absence of a violation and not a test of the

criterion. Widening to conductor 60000 would give tens of boxes with v_p ≥ 3.

12. **Rank $\geq$ 2 at the third and fourth digits.** Corollaries $N^{(3)}$ and $N^{(4)}$ are proved for any rank, but all the sweeps of §5.8 are of rank one; the rank-two entries checked for Theorem N (§5.7) have no counterpart there.

13. **Above the archimedean threshold.** Where the deficit ends is explained: below the canonical threshold $4 \log p + \beta(E)$ of Proposition 24^c the frequency is 0 by the proposition, and that threshold carries the whole deficit in expectation (100.6 % and 99.7 % on two disjoint sets of split boxes, §5.12), where the naive threshold of Proposition 24 carries 68 % and 56 %. **What is not explained is the frequency above it**: there $\det(\chi, \gamma) \equiv 0$ occurs 383 times against 382.6 for $1/p$ on the sets from Cremona's tables (ratio 0.995 to 1.018 on the two sets; 0.91 to 1.46 by p); on two further sets of curves outside Cremona's tables the boxes with $\det(\chi, \gamma) \equiv 0$ crowd just above the threshold (1.95 and 1.82 times $1/p$ in [4, 4.5), §5.12), while the frequency above the threshold stays at $1/p$ on the set not built from two given points; and nothing here says why γ should be a random multiple of χ once no element of $\ker \chi$ is short enough, nor why the boxes with $\det(\chi, \gamma) \equiv 0$ should crowd just above the threshold (choosing the shortest element without the p-adic part of its height leaves the crowding unchanged, §5.12). Lemma 25 makes $1/p$ a local count, so both are correlations between a local class and a global height. Whether the size of the crowding follows the conductor, the construction of the set or the position $\beta(E)/\log p$ is not decided by our data (§5.12). Nor is the box-by-box statement a theorem: $\beta(E)$ contains the numerical minimum m_∞ of λ'_∞. Two further things are left. In the split row of rank two, $P(k \geq 2 \mid k \geq 1)$ is 25 against 13.6 for $1/p$; counting two independent routes to $k \geq 2$ ($\rho_1 = 0$, and $\rho_1 = 1$ with v_p(h_p(Q)) ≥ 2 for $Q \in \ker \chi$) gives $2/p - 1/p^2$ and an expectation of 25.05, but that count was found after the fact and has not been tested box by box. And closures of index p^2 occur in two boxes up to conductor 500000 (§5.12), too few to test what the non-degeneracy of the third layer says there. [Maz72] treats the p-adic closure of the Mordell–Weil group, and a conjecture attributed to Lang concerns it; we retrieved neither, and state nothing from them.

14. **Translates that are not p-th powers.** Proposition 21 determines F along P + p^aΛ by local quantities. For $R \in \Lambda \cap$ E_k(Q_p) not in p^{k−1}Λ — for instance $R = -13g_1 + 19g_2 \in$ E_3(Q_5) at 75393a1 — it says nothing, and a pair P, P + R separated by min(v_p, k) would move F by something global. How often a split box of rank two has a point with v_p(n^2h_p(P)) $\geq$ k at all, the hypothesis of Proposition 21 (iii), is a question about isotropic vectors of q modulo p^k that we have not counted.

8. What is not claimed

1. **Nothing here bears on the conjecture of Birch and Swinnerton-Dyer**, nor on Sha, on L-functions or on the rank: every statement is about the valuation of a p-adic regulator under hypotheses on the reduction at one prime, except Proposition V of §6, which assumes the p-adic conjecture and is marked as such. Schneider's conjecture is not proved either; its CM case is [known], due to [Ber82], which we did not retrieve.

2. **The baseline is not ours.** Theorem A is elementary and known (§9); what we add is the splitting hypothesis and its mechanism.

3. **The local criterion is not ours.** The equivalence of Proposition 6 is [known] (Gross's criterion [Gro90], quoted as [LR, Thm 4.1] under the hypotheses $j \neq 0$, 1728 and E[p] irreducible; [OY, §2.1.1, Lem. 4.1] without them; [DW, Lemma 3.2 without hypotheses

on j, Cor. 3.5 with $j \neq 0, 1728$]); we give an independent proof and a presentation through a Fermat quotient, and we do not give it a name. [Gro90] itself we did not retrieve, and cite it as quoted in [LR] and [OY].

4. **The theorem is a lower bound in rank ≥ 2.** The exactness of §5.3 in the non-split case is Theorem S; Corollaries R′ and R′_s reduce the rest to the valuation of a Schur complement, which is not predicted here on either side, and Theorem R_s does not say which of the three values $\rho_1 \in \{0, 1, 2\}$ occurs. Deuring's lifting theorem, Vélu's formulas, Serre–Tate theory, the composition rule for division polynomials, the coprimality of φ_p and ψ_p^2, and the Tate parametrisation are used as standard facts, and for the last we cite sources that quote [ATAEC] rather than [ATAEC] itself.

5. **The hypotheses are as used, not the weakest possible.** Everything from Lemma 4 on assumes $p \geq 5$ and #E(F_p) = p; Proposition R^+ (§3.9) covers #E(F_p) = 2p, which by Hasse happens for p = 5 alone, and that is the whole of the extension — the main theorem itself is not extended, and neither is anything from §3.10 on.

6. **The computations are not kernel-checked, and the statistics are not statements.** All of §5 is PARI/GP with controls, at stated precision and over stated ranges; an earlier round used the height belonging to $\sigma \neq \sigma$_p, and every count here has been recomputed with h_p.

7. **Two inputs are used as verified, not as proved.** Katz's congruence $E_2\hat{}^* \equiv$ E_{p+1}/E_{p−1} (mod p) ([Kat73]), on which the Remark of §2 rests, is [known] and **the original we did not retrieve**; what we did was to check its consequence against PARI's `ellpadics2` over 315,953 boxes and to verify its two classical ingredients (E_{p−1} ≡ 1, E_{p+1} ≡ E_2) on q-expansions ourselves. Likewise the definition of `ellpadics2` is taken from PARI's documentation and calibrated on the one worked example there; **the source code we did not read**. So the identification of c mod p with a function of $(\bar{E}, \bar{\omega})$ is [paper] granted a [known] congruence we have checked but not proved, and every number in this paper that involves c inherits that status.

8. **The digits past the fourth are truncations carried out by machine.** Theorem N* is an exact identity, so the selection rule of Proposition 18 is exhaustive: no term can appear at a later digit that is not one of the three series. What is [computation] in Corollaries $N^{(5)}$, $N^{(6)}$ and $N^{(7)}$ is that the truncation was performed and checked in 648 boxes rather than written out and proved term by term; and since the sharp lower bound for v_p(μ_{2m}) is not known here (§7), the digit at which a given coefficient of $\mathcal{L}$ becomes visible is proved for $m \leq 3$ only. The valuation table of the Remark opening §4 remains a table: it lists the terms of the expansion we carried out by hand, and it is Proposition 18 and not the table that is a statement about every digit. Each of the three corollaries is checked **in the form in which it is displayed**, and not only in the form the generating rule produces — the two can differ, and at one place in an earlier version they did (§5.10).

9. **Propositions 14 and 15 may be known.** Their ingredients are classical — [p] = V∘F, the canonical subgroup, the theta product for the Tate curve, the injectivity of q-expansions in a fixed weight, and Lemma 13 — and the shape of Proposition 15, a Kodaira–Spencer derivative of the étale p-division points, is the kind of statement that belongs to the standard theory of the theta operator on p-adic modular forms. We did not find either statement in the sources we read (§9), and we do not call them new; what we claim is that they hold and that the proofs in §4.13 are complete.

10. **Theorem N* is an assembly of standard tools.** Its ingredients are classical: Mazur

and Tate's differential equation for σ_p, the Iwasawa logarithm of a unit written through its $(p-1)$-st power, and the $\wp$-expansion of x. That $\log(x \cdot \sigma_p(t)^2)$ is the local term of the p-adic height is close to the definition of σ_p itself, so the closed form is the kind of statement that may well be written somewhere. We looked for it as a displayed statement in the sources of §9 and did not find one; **we do not call it new.** What we do claim is the bookkeeping: that the congruences of Theorem N and of Corollaries $N^{(3)}$–$N^{(7)}$ are truncations of that one identity, that the series $\mathcal{L}$ is even with coefficients $\mu_2 = -c$, $\mu_4 = -b_4/12$, $\mu_6 = -b_2 b_4/360 - b_6/30$ in which E_2 occurs in μ_2 alone, and that a local invariant becomes visible at the digit $2me - 2 + v_p(\mu_\{2m\})$. Proposition 17 is [paper], Proposition 18 is [paper] for $m \leq 3$, and the digits past the fourth are [computation].

11. **No claim of priority.** We searched [MST], [SW], [Har], [BMS], [BD], [Ban], [Wut], [BG], [LR], [OY], [DW], Sage's documentation and arXiv (§9), in a narrow search without MathSciNet or zbMATH and with several primary sources not retrieved; the statements of §§3.10–3.15 may be known in part. The Lean kernel covers §6 only.

12. **Lemmas 23 and 23′ are elementary and should be regarded as known.** That a rational point deep in the formal filtration has a large denominator, and hence a large naive height, is the starting point of the study of valuations of elliptic divisibility sequences and division polynomials ([NV] and the work it cites); we did not look for Lemma 23′ in that literature and do not call either lemma new. Proposition 24 is their composition with Corollary K′(i), and Proposition 24^c is the same composition with the decomposition of the canonical height into local heights, which is known [SilH]; we do not call it new. The statistics of §5.12 are [computation]: they say that the threshold of Proposition 24 accounts for about 68 % of the observed deficit of $\det(\chi, \gamma) \equiv 0$ and that of Proposition 24^c for all of it in expectation, on the boxes measured, and not that nothing else acts; the frequency $1/p$ above the threshold is not a theorem.

13. **Propositions 21 and 22 are within a known framework.** Proposition 21 follows from the quadraticity of the height, Theorem N* and the fact that $\varphi_p \bmod p$ is a polynomial in x^p; that the denominator term of the p-adic height behaves p-adically along p-power multiples is the kind of statement that belongs to the theory of [MST]. We did not find it displayed, and we do not call it new. Proposition 22 is Theorem N* solved for F.

14. **Proposition V assumes a conjecture.** It is the only statement of the paper that does, it is stated in §6 and in a separate note and not in §3, and nothing in §3 depends on it. The extra valuation 2 of the Euler factor at an anomalous prime is an elementary consequence of the interpolation factor of [MTT] and is probably known; in [MTT, II §10] it is the factor $(1 - 1/\alpha)^2$, and the conjecture is used in the form of [SW, Conj. 5.1] and of PARI's `ellpadicbsd`.

15. **The canonical threshold is applied with a numerical constant.** $\beta(E)$ contains $m_\infty = \min \lambda'_\infty$ over $E(\mathbb{R})$, computed on a grid with refinement and checked by a second route, with no interval arithmetic; at I_n* with $c_\ell = 2$ the rational component was not determined and the smaller correction used. The statement "no box with $\det(\chi, \gamma) \equiv 0$ below the threshold" is protected against both — a sampled minimum is at least the true one, which only makes the check stricter, and the smaller correction at I_n* is still a valid lower bound — but the shares of §5.12 are not. The samples A and B are the curves of Cremona's tables satisfying the box conditions, C is a family of curves through two small points, and D a family of random short Weierstrass equations on which two small independent points were found, C and D with Mordell–Weil groups assumed saturated beyond 200; none is a random sample, and the effect just above the threshold

is stated for these sets only.

16. **The effect just above the threshold is a verdict of a rule, not an explanation.** "Effect" in §5.12 is the name of an outcome of a rule fixed before counting (Poisson z $>$ 2 on a set alone); it names no cause. We have not found the cause, and the tests of whether its size follows the conductor, β(E)/log p or the construction of the set came out undecided with little power; we claim no dependence and no independence. Lemma 25 is elementary and [paper], and its check on the classes modulo p^2 is [computation].

9. Related work

Where the baseline already is. In the tables of [MST, §§4.2–4.6], for curves of rank 1 to 5, three entries have negative valuation — 37A at p = 53, 5077A at p = 5, and the rank-four curve $y^2 + xy = x^3 - x^2 - 79x + 289$ of conductor 234446 at p = 5, all three equal to -2 — and after the shift by r of §2 only the first of them stays negative, at -1. The third of those entries we recomputed in full, including the hypotheses [MST] does not state: it has #E(F_5) = 10, so it is a box of Proposition R^+ and not of the main theorem (§5.3). At the first, the text notes that #E(F_p) = p, "i.e., p is anomalous"; at the second it notes $a_p \equiv 1 \pmod 5$ for the same reason. So the phenomenon is there, in three places, with no bound stated and no index in sight. [Har] has the mechanism implicitly rather than as a statement: the height of a general point is h_p(Q) = h_p(n_{lcm}Q)/n_{lcm}^2 with n_{lcm} = lcm(#E(F_p), lcm_ℓ c_ℓ) (§2), so an anomalous p costs 2 v_p(n_{lcm}) digits of working precision — which is the "cost 2" of Theorem A read as a loss of precision. That reading is ours; what is in [Har] is the normalisation itself, the n^2 growth of the coordinates and the general warning about loss of precision, and we did not find the quantity 2 v_p(n_{lcm}) written there. [Ban, §2.2] uses the contrapositive as a hypothesis, non-anomalousness being what "extends the regulator estimate of Proposition 4.2 from $E^\circ(Q)$ to all of E(Q)", and that Proposition 4.2 is the positive half of Theorem A in rank two; Sage's `padic_regulator` documents the same example. So the baseline is known to practitioners; we did not find it written as a proposition, and we do not need it to be new.

Where the local identity (L1) sits: the anomalous discrete logarithm. (L1) says that on a split anomalous curve the discrete logarithm χ of the reduction is given by an algebraic expression in the coordinates, $\chi = s{\cdot}c_1/B_0$ — that is, by the second p-adic digit of φ_p′ evaluated at the point itself. Since #E(F_p) = p, the group Hom(E(F_p), F_p^+) is one-dimensional over F_p, so **any** additive map E(F_p) → F_p is a scalar multiple of any other, and (L1) is necessarily a multiple of the map of [Sem98]. What the sources do and do not contain is worth stating precisely, because the three papers usually cited together are of two different kinds.

[Sem98] stays in characteristic p: for Q of order p he takes f_Q with divisor p(Q) − p(O) and sets $\varphi(Q) = (f'_Q/f_Q)(R)$ at an auxiliary point R, proving additivity from f_{Q_1+Q_2} = c·f_{Q_1}f_{Q_2}; his Lemma 1 gives (f′_Q) = (f_Q) − (y), so f′_Q/f_Q = c_Q/y and the map is a logarithmic derivative along the invariant differential $\omega = dx/(2y + a_1x + a_3)$. He evaluates it not in closed form but through O(log p) linear functions along a doubling chain (his Lemma 3 and equation (2)), and he notes that the construction also follows from a general result of Serre. Neither division polynomials nor Fermat quotients occur there. [Sma99] and [SA98] are of the other kind: they lift P and Q to E(Q_p) (or to $\mathbb{Z}/p^2$) by Hensel, multiply by p, and read the ratio of p-adic elliptic logarithms modulo p; nothing in them is a rational function of the reduced coordinates, and [SA98, Thm 3.2] is the statement that the resulting map is independent of the lift. Our $s{\cdot}c_1$ is a logarithmic derivative of the

universal polynomial φ_p, divided by p, evaluated at the point itself; it is of Semaev's kind and, by the dimension count, a scalar multiple of his map. We claim nothing for it beyond the proof given in §4.11, which goes through Theorem H and not through any of these sources; and we note that a formula of degree p^2 in x is not an algorithm, both cited methods being exponentially cheaper.

One point of contact is worth recording. [Vol2] compares the two approaches and shows that the p-adic one degenerates exactly when the lift is the canonical lift, i.e. when the Serre–Tate parameter satisfies $q = 1$ in $\mathbb{Z}/p^2$, which by [DW, Lemma 3.2] is exactly our splitting condition $E(\mathbb{Q}_p)[p] \neq 0$. **Corollary D — $\lambda \equiv 0$ in the split case — is therefore the same statement as the failure of the anomalous-curve attack on the canonical lift**, read as the vanishing of a Fermat quotient; and $\lambda\hat{}\{(2)\}$, Theorem K and Theorem N are what remains when that first digit is gone. [Rüc] generalises [Sem98] to divisor class groups; we did not retrieve it. Of the three 1998–99 papers we retrieved [Sem98] as text and [Sma99] in the author's technical-report form; [SA98] we did not retrieve, and quote it as quoted in [Yas, §2] and [Vol2].

Where the local criterion already is. Proposition 6 is the case $n = 1$ of Gross's tameness criterion. [LR, Thm 4.1] states it as "Gross; see [9], p. 514; see also §§14–15", [9] being [Gro90]: for $E/\mathbb{Q}$ with ordinary good reduction at p, $j \neq 0$, 1728 and E[p] an irreducible $\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$-module, a decomposition group D_n at p is diagonalisable if and only if $j_E \equiv j_0 \pmod{p\hat{}\{n+1\}}$ for odd p, with j_0 the j-invariant of the canonical lift, determined by the modular equation $T_p(j_0, j_0) = 0$ ([LR, Thm 4.2, Cor. 4.3], after Deuring). Those two hypotheses are absent from Proposition 6, and the passage from diagonalisability to a rational point of order p is [OY, §2.1.1 and Lem. 4.1], which for good ordinary reduction and odd p proves $E(F)[p] \neq 0 \iff E[p] \cong F_p(1) \oplus F_p \iff E[p]$ tame with F the unramified extension of degree the order of $a_p \bmod p$ — for us $a_p \equiv 1$, so $F = \mathbb{Q}_p$ — and quotes [Gro90, Prop. 13.7 and §17] for the equivalence with $j_E \equiv j_0 \pmod{p^2}$. It is also [DW, Lemma 3.2 and Cor. 3.5], where a prime with $E(\mathbb{Q}_p)[p] \neq 0$ is a *local torsion prime* and the criterion is that the lift to $\mathbb{Z}/p^2$ be the canonical one — of the two, only Lemma 3.2 is free of hypotheses on the j-invariant, Cor. 3.5 assuming $j \neq 0$, 1728 as [LR, Thm 4.1] does; and the word "anomalous" does not occur in [DW], the content of "a local torsion prime is anomalous" sitting inside the proof of their Prop. 2.1(1) — of the $p\hat{}d$ lifts exactly one has the larger p-rank — together with [DW, Prop. 2.1] for the fact that a local torsion prime is anomalous. The same statement circulates in cryptography, where the attacks on anomalous curves fail exactly for the lifts congruent to the canonical one modulo p^2. What we did not find is the criterion in the form $q_p(\varphi_p(x_0)) = 0$ — a Fermat quotient of a division polynomial at one point, which needs neither the j-invariant nor root-finding; the proof of §4.6 is independent of the sources above. Searches returning nothing on point (2026-09-24): `elliptic Wieferich prime`, `Fermat quotient elliptic curve`, `anomalous prime elliptic curve p-torsion`, `canonical lift p-torsion criterion mod p^2`. Near but different: the elliptic Wieferich primes of [Sil88] and [Vol] concern $N_p \cdot P$ modulo p^2 for a rational point P, not rationality of the p-torsion, and the division-polynomial criteria of [Deb] are modulo p.

Where the closed form of the ledger sits. The three ingredients of Theorem N* are in [MST] and in the standard theory: the differential equation $(d/dz)^2 \log \sigma_p = -(x + c)$ characterising the p-adic sigma function together with its normalisation, the Iwasawa logarithm of a p-adic unit, and the Weierstrass expansion of x in the formal parameter. Their combination — that the local term of the height is $\log(x(P'') \cdot \sigma_p(t(P''))^2)$, an even series in the formal logarithm whose coefficients are polynomials in b_2, b_4, b_6 with E_2 entering the first one only — is the kind of statement that belongs to the same the-

ory, and we searched for it as a displayed statement in [MST], [SW], [Har], [BMS], [BD], [Wut], [BG], [Ban], Sage's and PARI's documentation, and on arXiv (searches on 2026-09-25: `p-adic sigma function differential equation coefficients`, `p-adic height formal logarithm expansion`, `Mazur-Tate sigma even series`). We did not find it, in a search of the same narrow kind as the others recorded here, and we claim nothing for it beyond the proof in §3.11. What we did not find either is the rule 2me − 2 + v_p(μ_{2m}) for the digit at which a local invariant of E becomes visible in the p-adic height, or the observation that the digit can depend on p through the denominator of a coefficient. One caveat belongs here. A further search (`p-adic sigma function coefficients`, 2026-09-26) returned exactly one paper, [BG], which builds the universal p-adic σ out of congruences between the Laurent coefficients of the elliptic functions — the nearest published work to Proposition 17. **We have not retrieved it**, so we cannot say whether the coefficients μ_2, μ_4, μ_6 are in it; the search `p-adic height sigma series` returned nothing at all.

Where depth and height meet. Lemma 23 is the elementary end of a known subject: the p-adic valuation of the denominators of the multiples of a rational point — elliptic divisibility sequences — and the common valuations of division polynomials are studied in [NV], whose abstract we retrieved (arXiv:2203.02015); we did not read the body, and we cite it only to place Lemma 23 in a field and not for any statement. The index of the p-adic closure of the Mordell–Weil group, which §5.12 computes as p^{min(D_Q, D_1) − 1}, is a central quantity of Chabauty's method and is treated in [Maz72]; we did not retrieve that paper (a search on arXiv for `p-adic closure Mordell-Weil` returned two papers, neither on this question), and we state nothing from it.

What we did not find is the splitting hypothesis E(Q_p)[p] $\neq 0$ in connection with the valuation of Reg_p, or any statement that the height pairing is p-integral exactly under it. [SW] is our source for Schneider's conjecture; their §3.3 anomalous example is 446d1 at p = 5 with #E(F_5) = 10, the p = 5 case of §3.1, and its $\mathrm{ord}_5(\mathrm{Reg}_5)$ is +1, not negative. The word "anomalous" does not occur in [BMS]. [Wut] studies the quantity e of §3.3 from the global side and [BG] the universal p-adic sigma function; [Kat] on Serre–Tate local moduli, the natural language for §4.6, we could not obtain. The nearest computation in the literature is [Ban] (2026), five rank-two CM curves at every good ordinary prime below 30,000 with a non-unit at 3 of 8,050 primes — the slice orthogonal to ours — which also formalises part of its argument in Lean 4.

References

[A] Y. Hosomura, *Integrality of the p-adic height pairing at an anomalous prime and points of order p in E(Q_p)*, preprint (2026).

[Draft] Kiichi (Y. Hosomura), *Integrality of the p-adic height pairing at an anomalous prime is equivalent to a point of order p in E(Q_p)*, draft v15, computo ergo sum (2026); Zenodo.

[ATAEC] J. H. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, GTM 151. [not retrieved; Thm V.5.3, Thm V.3.1, Prop. V.6.1 and Exercise V.5.13 are cited as quoted in [Ked] and [Sym]; the minimality of the Tate model and the integrality of the coefficients of q as a series in 1/j we could not attach to numbered statements]

[Ban] B. S. Banwait, *Second derivatives of p-adic L-functions and the Shafarevich–Tate group of rank-two CM elliptic curves*, arXiv:2609.08431 (2026).

[BMS] J. S. Balakrishnan, J. S. Müller, W. A. Stein, *A p-adic analogue of the conjecture of Birch and Swinnerton-Dyer for modular abelian varieties*, arXiv:1210.2739.

[Ber82] D. Bertrand, *Valeurs de fonctions thêta et hauteurs p-adiques*, Progr. Math. **22**, Birkhäuser, 1982, 1–11. [not retrieved]

[BG] C. Blakestad, D. Grant, *On the universal p-adic sigma and Weierstrass zeta functions*, arXiv:1903.02480. [not retrieved; for p > 3 it reconstructs the Mazur–Tate universal p-adic σ from congruences between the Laurent coefficients of the elliptic functions, which makes it the nearest published work to Proposition 17 — see §9]

[BD] A. Burungale, D. Disegni, arXiv:1803.09268.

[Deb] C. Debry, *Beyond two criteria for supersingularity*, J. Théor. Nombres Bordeaux **26** (2014) 595–605.

[Dok] T. Dokchitser, V. Dokchitser, *Local invariants of isogenous elliptic curves*, arXiv:1208.5519, Thm A.1.

[DW] C. David, T. Weston, *Local torsion on elliptic curves and the deformation theory of Galois representations*, arXiv:math/0701882.

[Gro90] B. H. Gross, *A tameness criterion for Galois representations associated to modular forms (mod p)*, Duke Math. J. **61** (1990) 445–517. [not retrieved; cited as quoted in [LR, Thm 4.1] (p. 514, §§14–15) and in [OY] (Prop. 13.7, §17)]

[Har] D. Harvey, *Efficient computation of p-adic heights*, arXiv:0708.3404.

[Kat] N. Katz, *Serre–Tate local moduli*. [not retrieved]

[Igu] J.-I. Igusa, *On the algebraic theory of elliptic modular functions*, J. Math. Soc. Japan **20** (1968) 96–106. [not retrieved; cited for the irreducibility of the tower, used as standard in §4.13]

[Kat73] N. Katz, *p-adic properties of modular schemes and modular forms*, in: Modular Functions of One Variable III, Lecture Notes in Math. **350**, Springer, 1973, 69–190. [not retrieved; cited for p-adic modular forms, the canonical subgroup, the theta operator and the congruence E_2^* ≡ E_{p+1}/E_{p−1} (mod p) used in §2, §4.13 and §7(8)]

[Ked] K. S. Kedlaya, *The Tate curve*, lecture notes, MIT 18.727. [used only as a conduit for the statements of [ATAEC]]

[LR] Á. Lozano-Robledo, *Division fields of elliptic curves with minimal ramification*, Rev. Mat. Iberoam. **31** (2015) 1311–1332; Thm 4.1 quotes [Gro90], p. 514 and §§14–15, under the hypotheses $j \neq 0, 1728$ and E[p] irreducible. (The Gross criterion is *not* in the same author's Math. Ann. **357** (2013) paper, where we first looked.)

[Maz72] B. Mazur, *Rational points of abelian varieties with values in towers of number fields*, Invent. Math. **18** (1972), 183–266. [not retrieved; bibliographic data checked against Crossref, DOI 10.1007/BF01389815; the source of "anomalous", and a treatment of the p-adic closure of the Mordell–Weil group (§7(13), §9)]

[MST] B. Mazur, W. Stein, J. Tate, *Computation of p-adic heights and log convergence*, Documenta Math. Extra Vol.: John H. Coates' Sixtieth Birthday (2006) 585–622.

[MTT] B. Mazur, J. Tate, J. Teitelbaum, *On p-adic analogues of the conjectures of Birch and Swinnerton-Dyer*, Invent. Math. **84** (1986), no. 1, 1–48. [read in the scan of the Göttingen digitisation centre (GDZ); Conj. BSD(p) is II §10, p. 38, the periods II §8, p. 36, the analytic height II §5, p. 33; PARI 2.17.3's `ellpadicbsd`, whose documentation names Mazur, Tate, Teitelbaum, Bernardi and Perrin-Riou, states it in the form of [SW]]

[MT91] B. Mazur, J. Tate, *The p-adic sigma function*, Duke Math. J. **62** (1991) 663–688. [not retrieved]

[MT87] B. Mazur, J. Tate, *Refined conjectures of the "Birch and Swinnerton-Dyer type"*, Duke Math. J. **54** (1987), no. 2, 711–750. [the first page read on Project Euclid; the rest read in the restatements of [PB19], [Ota], [LC24] and [BH25]]

[BD94] M. Bertolini, H. Darmon, *Derived heights and generalized Mazur–Tate regulators*, Duke Math. J. **76** (1994) 75–111. [not retrieved; cited as in [Ota, Remark 2.5]]

[PB19] F. X. Portillo-Bobadilla, *Experimental evidence on a refined conjecture of the BSD type*, Bol. Soc. Mat. Mex. (3) **25** (2019) 529–541; arXiv:1709.01204.

[Ota] K. Ota, *Kato's Euler system and the Mazur–Tate refined conjecture of BSD type*, arXiv:1509.00682.

[LC24] J.-P. Llerena-Córdova, *Numerical study of refined conjectures of the BSD type*, arXiv:2412.17703 (2024).

[BH25] D. Bullach, M. H. L. Honnor, *On the refined 'Birch–Swinnerton-Dyer type' conjectures of Mazur and Tate*, arXiv:2511.07203 (2025).

[JSW] D. Jetchev, C. Skinner, X. Wan, *The Birch and Swinnerton-Dyer formula for elliptic curves of analytic rank one*, arXiv:1512.06894 (2015). [abstract read]

[PR] B. Poonen, E. Rains, *Random maximal isotropic subspaces and Selmer groups*, arXiv:1009.0287 (2010). [abstract read]

[BKLPR] M. Bhargava, D. M. Kane, H. W. Lenstra Jr., B. Poonen, E. Rains, *Modeling the distribution of ranks, Selmer groups, and Shafarevich-Tate groups of elliptic curves*, arXiv:1304.3971 (2013). [Theorems 1.6, 3.11 and 5.15 read]

[Cre] J. Cremona, *ecdata*, files `allbsd/allbsd.*` (conductor < 500000), retrieved from the repository JohnCremona/ecdata on GitHub.

[NV] B. Naskręcki, M. Verzobio, *Common valuations of division polynomials*, arXiv:2203.02015 (2022); published version DOI 10.1017/prm.2024.7 (2024). [abstract page retrieved; cited to place Lemma 23 in a field]

[OY] Y. Ozeki, M. Yoshida, *Fields of definition of p-torsion points of elliptic curves and their ramification*, arXiv:2609.06554 (2026); §2.1.1 and Lemma 4.1.

[Sch85] P. Schneider, *p-adic height pairings II*, Invent. Math. **79** (1985) 329–374. [read in the GDZ scan; Thm 1 on p. 336, Prop. 1 and Thms 2 and $2'$ on pp. 341–342; Thm 6 is quoted from the introduction, p. 330] Part I, Invent. Math. **69** (1982) 401–409, was not retrieved.

[Ski] C. Skinner, *Multiplicative reduction and the cyclotomic main conjecture for GL2*, arXiv:1407.1093 (2014); §2.5 restates Kato's divisibility and the theorem of [SU] under the hypotheses used in §6, and replaces the hypothesis of [Kato04, Thm 17.4(3)] on the image of Galois by E[p] irreducible and (H); its Lemma 2.1.1 is the lattice lemma used there. (The theorem of [Kato04] proved in §13.14 is numbered 12.5(4) in the published version.)

[DDT] H. Darmon, F. Diamond, R. Taylor, *Fermat's Last Theorem*, version of 9 September 2007 on the first author's page. [Prop. 1.5 (the Tate curve) and Prop. 2.12 (multiplicative reduction), pp. 19 and 57–58 of that version]

[SU] C. Skinner, E. Urban, *The Iwasawa main conjectures for GL2*, Invent. Math. **195** (2014) 1–277. [read in the version on the second author's page, which may differ slightly from the published one; the numbers 3.3.3, 3.6.4, 3.6.9 and 3.6.11 are those of that version, and the published numbering of 3.6.4 is not checked; Theorem 1 of the introduction, which is 3.6.4 there, is cited as Theorem 1 of the published version in [Ski]]

[Gre99] R. Greenberg, *Iwasawa theory for elliptic curves*, in: Arithmetic Theory of Elliptic Curves (Cetraro, 1997), Lecture Notes in Math. **1716**, Springer, 1999, 51–144. [read in the author's version on his page; Thm 1.2, Prop. 2.2 and Thm 4.1]

[Maz78] B. Mazur, *Rational isogenies of prime degree*, Invent. Math. **44** (1978) 129–162. [read in the GDZ scan; Cor. 4.1, p. 144]

[Kato04] K. Kato, *p-adic Hodge theory and values of zeta functions of modular forms*, Astérisque **295** (2004) 117–290. [read on Numdam; Theorems 12.4–12.6 and condition 12.5.2 are on pp. 221–222, Theorem 13.4 on p. 226, §13.14 on p. 234, Lemma 14.7 on p. 238, Theorem 17.4 on p. 273 and its proof §17.13 on pp. 279–280]

[GV] R. Greenberg, V. Vatsal, *On the Iwasawa invariants of elliptic curves*, Invent. Math. **142** (2000) 17–63. [read in the authors' version on the first author's page; Prop. 3.7]

[Pe98] B. Perrin-Riou, *Systèmes d'Euler p-adiques et théorie d'Iwasawa*, Ann. Inst. Fourier **48** (1998) 1231–1307. [read on Numdam; Theorems 1.3.3 and 1.3.5, pp. 1241–1242, and §5.3.6, p. 1294]

[Ru00] K. Rubin, *Euler systems*, Ann. of Math. Studies **147**, Princeton Univ. Press (2000). [not obtained]

[KK99] K. Kato, *Euler systems, Iwasawa theory, and Selmer groups*, Kodai Math. J. **22** (1999) 313–372. [not obtained]

[PR92] B. Perrin-Riou, *Théorie d'Iwasawa et hauteurs p-adiques*, Invent. Math. **109** (1992) 137–185. [read in the GDZ scan; Théorèmes 3.3.4, 3.3.6 and 3.4.2 and §3.4.3, pp. 165–167]

[Wu17] C. Wuthrich, *Numerical modular symbols for elliptic curves*, arXiv:1608.06423 (2017).

[Ser73] J.-P. Serre, *Formes modulaires et fonctions zêta p-adiques*, in: Modular Functions of One Variable III, Lecture Notes in Math. **350**, Springer, 1973, 191–268. [not retrieved; cited for E_{p−1} ≡ 1 (mod p) and, with [SwD73], for the kernel of the q-expansion map]

[SwD73] H. P. F. Swinnerton-Dyer, *On ℓ-adic representations and congruences for coefficients of modular forms*, in: Modular Functions of One Variable III, Lecture Notes in Math. **350**, Springer, 1973, 1–55. [not retrieved; cited for the kernel of the q-expansion map on F_p[E_4, E_6] being generated by A − 1]

[Rüc] H.-G. Rück, *On the discrete logarithm in the divisor class group of curves*, Math. Comp. **68** (1999) 805–806. [not retrieved]

[SA98] T. Satoh, K. Araki, *Fermat quotients and the polynomial time discrete log algorithm for anomalous elliptic curves*, Comment. Math. Univ. St. Pauli **47** (1998) 81–92. [not retrieved; volume 47 is not in the publisher's repository. Cited as quoted in [Yas, §2] and [Vol2]]

[Sem98] I. A. Semaev, *Evaluation of discrete logarithms in a group of p-torsion points of an elliptic curve in characteristic p*, Math. Comp. **67** (1998) 353–356. [retrieved as text; the PDF at the publisher's site is behind a challenge]

[Sil88] J. H. Silverman, *Wieferich's criterion and the abc-conjecture*, J. Number Theory **30** (1988) 226–237.

[SilH] J. H. Silverman, *Computing heights on elliptic curves*, Math. Comp. **51** (1988) 339–358, doi:10.1090/S0025-5718-1988-0942161-4. [Bibliographic data checked through Crossref; the text was not retrieved here.]

[SW] W. Stein, C. Wuthrich, *Algorithms for the arithmetic of elliptic curves using Iwasawa theory*, Math. Comp. **82** (2013) 1757–1792.

[Sma99] N. P. Smart, *The discrete logarithm problem on elliptic curves of trace one*, J. Cryptology **12** (1999) 193–196; quoted from the author's technical report HPL-97-128 (1997), same title.

[Sym] *Symplectic criteria for elliptic curves, revisited*, arXiv:2509.19938, Prop. 2.2. [used only as a conduit for [ATAEC, Prop. V.6.1]]

[Vél] J. Vélu, *Isogénies entre courbes elliptiques* (1971). [not retrieved]

[Vol] J. F. Voloch, J. Number Theory **81** (2000) 205–209.

[Vol2] J. F. Voloch, *Relating the Smart–Satoh–Araki and Semaev approaches to the discrete logarithm problem on anomalous elliptic curves*, note on the author's page.

[Yas] M. Yasuda, *A p-adic approach to the discrete logarithm problem on anomalous elliptic curves*, Int. J. Pure Appl. Math. **77** (2012) 1–9; quotes [SA98, Thm 3.2].

[Wut] C. Wuthrich, *On p-adic elliptic logarithms and p-adic approximation lattices* (2006).

- Software: PARI/GP 2.17.3 with `elldata` (Cremona's tables, conductor $<$ 500000); Sage, `src/sage/schemes/elliptic_curves/padics.py`, docstrings of `padic_regulator` and `padic_height`.