

General Topology and its Relations
to Modern Analysis and Algebra VI
Proc. Sixth Prague Topological Symposium 1986
Z. Frolík (ed.)
Copyright Heldermann Verlag Berlin 1988
323–330

On universal metric spaces

M. Katětov

We present the following theorems (for the terms involved, see 1.1 and 1.2 below).

Theorem I. There is exactly one (up to isometry) complete ω -homogeneous ω -universal metric space of weight ω .

Theorem II. If $\alpha > \omega$ is a cardinal and $\alpha^\alpha = \alpha$, then there exists exactly one (up to isometry) Urysohn universal space of weight α . Every Urysohn universal space of uncountable weight is complete.

Theorem III. If α is a cardinal, $\alpha^\alpha > \alpha$, then there is no α -homogeneous α -universal space of weight α .

Theorem IV. There exists a meager Urysohn universal space of weight ω .

Theorem I has been proved by P. Urysohn, see [2] and [3]. In [3], P. Urysohn posed the question whether there exist non-complete ω -homogeneous strongly ω -universal spaces; this question is answered by Theorem IV. Theorem II can be easily deduced from Jónsson's theorems concerning relational systems (for the pertinent theorem and references to Jónsson's articles, see [1], §4).

Theorems II–IV are immediate consequences of propositions proved below: Theorem II follows from 2.8, 2.9, 2.7 and 2.10, Theorem III from 2.9 and 2.11, Theorem IV from 3.2.

1

1.1. Notation and conventions. A) As a rule, the same symbol denotes a metric space and its underlying set, or a mapping $f : A \longrightarrow B$ and the set $\{(a, b) : a \in A, b = f(a)\}$, etc. – B) If $f : A \longrightarrow B$ is a mapping, we often write fa instead of $f(a)$. – C) The class of all ordinals (all cardinals) will be denoted by Ord (respectively, by Card). The letters x, y and z (respectively, α, β and γ) will stand for ordinals (for cardinals). The cardinality of a set A is denoted by $\text{card } A$. We have $\text{Card} \subset \text{Ord}$, and $x = \{y : y < x\}$ for all $x \in \text{Ord}$. – D) If $x \in \text{Ord}$, $\alpha \in \text{Card}$, we put $\alpha^x = \sup(\alpha^\beta : \beta < x)$. – E) “Space” always means a metric space. The class of all non-void spaces is denoted by $\mathcal{M}$. The weight of a space S is denoted by $\text{w}S$. – F) If S and T are spaces, then $[S, T]$ denotes the set of all isometric mappings $f : S \longrightarrow T$, and $\text{Is}(S, T)$ denotes the set of isometries $g : S \longrightarrow T$, i.e. of bijective mappings $g \in [S, T]$. A mapping $h \in \text{Is}(S, S)$ will be called an isometry of S . If $\text{Is}(S, T) \neq \emptyset$, we write $S \cong T$.

1.2. Basic definitions. Let $\alpha \geq \omega$ be a cardinal. A space S will be called (1) α -homogeneous if, for any $f \in \text{Is}(A, B)$, where $A \subset S$, $B \subset S$, $\text{card } A < \alpha$, there is a $g \in \text{Is}(S, S)$ such that $g \upharpoonright A = f$, (2) α -universal (strongly α -universal) if $[T, S] \neq \emptyset$ whenever $T \in \mathcal{M}$ and $\text{card } T \leq \alpha$ (respectively, $\text{w}T \leq \alpha$). A space of weight α will be called Urysohn universal if it is α -homogeneous strongly α -universal.

1.3. Fact and notation. Let $(S, \varrho) \in \mathcal{M}$. If $p \in S$, let $\hat{p}$ (or merely p) denote the function $s \longmapsto \varrho(p, s)$. Let $E'(S)$ denote the set of all $f : S \longrightarrow R$ satisfying $|fp - fq| \leq \varrho(p, q) \leq fp + fq$ for all $p, q \in S$ and distinct from all $\hat{s}$, $s \in S$. Put $E(S) = S \cup E'(S)$. If $p, q \in E(S)$, put $\varrho^E(p, q) = \sup\{|p(s) - q(s)| : s \in S\}$. Then ϱ^E is a metric on $E(S)$. The space $(E(S), \varrho^E)$ will be denoted by $E(S, \varrho)$ or by $E(S)$.

1.4. Fact and notation. Let $S \in \mathcal{M}$, $\emptyset \neq T \subset S$. If $f \in E'(T)$, let f^* denote the function $s \longmapsto \inf\{\varrho(s, t) + ft : t \in T\}$ defined on S . If $f^* = \hat{p}$ for some $p \in S$, put $f_S = p$; if not, put $f_S = f^*$. If $t \in T$, put $t_S = t$. Then the mapping $r \longmapsto r_S$, defined on $E(T)$, is in $[E(T), E(S)]$.

Proof. It is easy to see that either $f^* \in E'(S)$ or $f^* = \hat{p}$ for some $p \in S$. We are going to show that if $f, g \in E'(T)$ and $f^*, g^* \in$

$E'(S)$, then $\sup\{|f^*s - g^*s| : s \in S\} \leq \varrho^E(f, g)$; the remaining assertions are easy to prove. Suppose there are $s \in S$ and $\varepsilon > 0$ such that $|f^*s - g^*s| > \varrho^E(f, g) + \varepsilon$. We can assume $f^*s > g^*s$. Choose $t \in T$ such that $gt + \varrho(s, t) < g^*s + \varepsilon$. Since $ft + \varrho(s, t) \geq f^*s$, we get $ft - gt \geq f^*s + g^*s - \varepsilon > \varrho^E(f, g)$, which is a contradiction.

1.5. Notation. Let $S \in \mathcal{M}$ and let $\alpha > 1$. The subspace of $E(S)$ consisting of all t_S , where $t \in \bigcup(E(T) : T \subset S, 0 < \text{card } T < \alpha)$, will be denoted by $E(S, \alpha)$.

1.6. Fact and notation. Let $S \in \mathcal{M}$ and let $\varphi \in \text{Is}(S, S)$. Then there is exactly one $\Phi \in \text{Is}(E(S), E(S))$ such that $\Phi \upharpoonright S = \varphi$. The isometry Φ will be denoted by $E(\varphi)$. For any $\alpha > 1$, $E(S, \alpha)$ is invariant under Φ .

Proof. If $s \in S$, put $\Phi(s) = \varphi s$. If $f \in E'(S)$, put $\Phi(f) = f \circ \varphi^{-1}$. It is easy to see that Φ is an isometry of $E(S)$. – Let Ψ be an isometry of $E(S)$ and let $\Psi \upharpoonright S = \varphi$. Let $f \in E'(S)$, $g = \Psi(f)$. It is easy to see that, for any $h \in E'(S)$ and any $p \in S$, $\varrho^E(h, p) = h(p)$. For any $s \in S$, we have $\varrho^E(g, \varphi s) = \varrho^E(f, s)$, hence $g(\varphi s) = f(s)$, $(g \circ \varphi)(s) = f(s)$. This implies $g \circ \varphi = f$, $g = f \circ \varphi^{-1}$.

1.7. Fact. Let $(S, \varrho) \in \mathcal{M}$, $A \subset S$, $0 < \text{card } A < \omega$, $B \subset S$. Let $\varphi : A \rightarrow B$ be a bijection. Let $\varepsilon > 0$ and let $\varrho(a, \varphi a) \leq \varepsilon$ for each $a \in A$. If $f \in E'(A)$, $g \in E'(B)$ and $f = g \circ \varphi$, then $\varrho^E(f_S, g_S) \leq \varepsilon$.

Proof. Let $s \in S$. We have $f_S(s) = \inf\{ft + \varrho(s, t) : t \in A\}$, $g_S(s) = \inf\{g(\varphi t) + \varrho(s, \varphi t) : t \in A\} = \inf\{ft + \varrho(s, \varphi t) : t \in A\}$. Since $|\varrho(s, t) - \varrho(s, \varphi t)| \leq \varrho(t, \varphi t) \leq \varepsilon$ for each $t \in A$, we get $|f_S(s) - g_S(s)| \leq \varepsilon$.

1.8. Lemma. Let $1 < \alpha \leq \omega$. Let $S \in \mathcal{M}$. Then $w(E(S, \alpha)) = \omega \cdot wS$, $\text{card } E(S, \alpha) = 2^\omega \cdot \text{card } S$.

Proof. Since $E(S, \omega) = \bigcup(E(S, n) : 1 < n < \omega)$, we can assume that $\alpha = n < \omega$. If $A \subset S$, $0 < \text{card } A < n$, put $P(A) = \{u_S : u \in E(A)\}$. Since $w(E(A)) = \omega$, $\text{card } E(A) = 2^\omega$, we get, by 1.4, $w(P(A)) = \omega$, $\text{card } P(A) = 2^\omega$. Let $T \subset S$, $\text{card } T = wS$. By 1.7, the union of $P(A)$, where $A \subset T$, $0 < \text{card } A < n$, is dense in $E(S, n)$. This proves $wE(S, n) \leq \omega \cdot wS$; evidently, $wE(S, n) \geq \omega \cdot wS$. Finally, we have $E(S, n) = \bigcup(E(T) : T \subset S, 0 < \text{card } T < n)$, hence $\text{card } E(S, n) \leq 2^\omega \cdot \text{card } S$.

1.9. Fact and notation. Let $S \in \mathcal{M}$. There exists exactly one indexed class $(S_x : x \in \text{Ord})$ such that (1) $S_0 = S$, (2) if $x \in \text{Ord}$, then $S_{x+1} = E(S_x, \omega \cdot (\text{card } x)^+)$, (3) if x is a limit ordinal, then $S_x = \bigcup (S_y : y < x)$. – The spaces S_x will be denoted by $U_x(S)$.

1.10. Lemma. Let $S \in \mathcal{M}$, $\text{card } S \leq 2^\omega$. Then, for any ordinal $x > \omega$, $\text{card } U_x(S) \leq 2^x$.

Proof. By 1.8, $\text{card } U_\omega(S) = 2^\omega$. Put $V = \{x : x > \omega, \text{card } U_x(S) > 2^x\}$. Suppose $V \neq \emptyset$ and put $y = \min V$. Put $\beta = \text{card } y$. If y is a limit ordinal, then $U_y(S) = \bigcup (U_x(S) : x < y)$, hence $\text{card } U_y(S) \leq \beta \cdot 2^y$, which contradicts $y \in V$. If $y = z + 1$, then $U_y = E(U_z(S), \beta^+)$, hence $\text{card } U_y(S) = \alpha^\beta$, where $\alpha = 2^z$. If $z \in \text{Card}$, then, clearly, $\alpha^\beta = \alpha^z = 2^z = 2^y$. If $z \notin \text{Card}$, then $\alpha = 2^\beta$, $\alpha^\beta = 2^\beta = 2^y$. Hence $\text{card } U_y(S) \leq 2^y$, which contradicts the supposition.

2

2.1. Notation. If α and β are cardinals, then $\mathcal{E}(\alpha, \beta)$ will denote the class of all $(T, S) \in \mathcal{M} \times \mathcal{M}$ such that $T \subset S$ and if $M \in \mathcal{M}$, $A \subset M$, $\text{card } A < \alpha$, $\text{card}(M \setminus A) < \beta$, $f \in [A, T]$, then $g \upharpoonright A = f$ for some $g \in [M, S]$.

2.2. Lemma. Let $\alpha \geq \omega$ be a cardinal. Let $S \in \mathcal{M}$. Let S_x , $x < \alpha$, be subspaces of S . Assume that (1) $\bigcup S_x = S$, (2) if $x < \alpha$, then $(S_x, S_y) \in \mathcal{E}(\alpha, 2)$ for some $y < \alpha$, (3) if $K \subset S$, $\text{card } K < \alpha$, then $K \subset S_z$ for some $z < \alpha$. Then $(S, S) \in \mathcal{E}(\alpha, \alpha^+)$.

Proof. Put $\beta = \alpha^+$. Let M, A and f be as in 2.1. Put $\gamma = \text{card}(M \setminus A)$. Choose a bijection $h : \gamma \rightarrow M \setminus A$. For $x \leq \gamma$ let G_x consist of all $g \in [A \cup h\{y : y < x\}, S]$ such that $g \upharpoonright A = f$. Put $G = \bigcup G_x$. It is easy to see that G , ordered by inclusion (cf. 1.1 A), contains a maximal element, say k . Suppose that $k \in G_x$, where $x < \gamma$. Then $\text{dom } k = A \cup h\{y : y < x\}$, hence $\text{card}(\text{dom } k) < \alpha$ and therefore $\text{dom } k \subset S_y$ for some $y < \alpha$. Choose a $z < \alpha$ such that $(S_y, S_z) \in \mathcal{E}(\alpha, 2)$. Then there exists a $k^* \in [(\text{dom } k) \cup \{hx\}, S]$ such that $k^* \upharpoonright \text{dom } g = g$. This is a contradiction, for $k^* \in G_{x+1} \subset G$. We have shown that $k \in G_\gamma$, which proves the lemma.

2.3. Corollary. For any cardinal $\alpha \geq \omega$, $\mathcal{E}(\alpha, 2) = \mathcal{E}(\alpha, \alpha^+)$.

2.4. Notation. If $\alpha \geq \omega$, we put $\mathcal{E}(\alpha) = \{S : (S, S) \in \mathcal{E}(\alpha, 2)\}$.

2.5. Lemma. Let $\alpha \geq \omega$. Let $S_i \in \mathcal{E}(\alpha)$, $i = 1, 2$. Let $A_i \subset S_i$, $\text{card } A_i < \alpha$ and let $f \in \text{Is}(A_1, A_2)$. Let $K_i \subset S_i$, $\text{card } K_i \leq \alpha$. Then there exist subspaces $T_i \subset S_i$, and an isometry $g : T_1 \rightarrow T_2$ such that $A_i \cup K_i \subset T_i$ and $g \upharpoonright A_1 = f$.

Proof. Clearly, we can assume that $K_i \cap A_i = \emptyset$, $\text{card } K_1 = \text{card } K_2 \geq \omega$. Put $\beta = \text{card } K_1$ and choose bijections $u : \{x : x < \beta\} \rightarrow K_1$, $v : \{x : x < \beta\} \rightarrow K_2$. For $x \leq \beta$ put $U_x = u\{z : z < x\}$, $V_x = v\{z : z < x\}$. If $x, y \leq \beta$, let $G(x, y)$ consist of all $g : [M, P_2]$ such that $A_1 \subset M \subset S$, $g \upharpoonright A_1 = f$, $\text{dom } g \setminus A_1 = U_x \cup g^{-1}(V_y)$. Put $G = \bigcup \{G(x, y) : x \leq \beta, y \leq \beta\}$. It is easy to see that G , ordered by inclusion (cf. 1.1 A), contains a maximal element, say h . Put $x_0 = \sup\{x \leq \beta : h \in G(x, y) \text{ for some } y\}$, $y_0 = \sup\{y \leq \beta : h \in G(x, y) \text{ for some } x\}$. It is easy to show that $h \in G(x_0, y_0)$. Suppose that $x_0 < \beta$. Then $u(x_0) \notin \text{dom } h$. Hence, due to $(S_2, S_2) \in \mathcal{E}(\alpha, 2)$ and $\text{card}(\text{dom } h) < \alpha$, there is an isometric mapping $h' : (\text{dom } h \cup \{u(x_0)\}) \rightarrow S_2$ such that $h' \upharpoonright \text{dom } h = h$. We have $h' \in G(x_0 + 1, y_0) \subset G$. This contradicts the fact that h is maximal, and proves that $x_0 = \beta$. The proof of $y_0 = \beta$ is analogous. Hence, $\text{dom } h \supset K_1$, $\text{dom } h^{-1} \supset K_2$, which proves the lemma.

2.6. Proposition. Let $\alpha \geq \omega$. If $S \in \mathcal{E}(\alpha)$ is complete and $\text{ws} S \leq \alpha$, then S is Urysohn universal.

Proof. By 2.3, S is α -universal, hence, being complete, strongly α -universal. Let $A_1 \subset S$, $A_2 \subset S$, let $f : A_1 \rightarrow A_2$ be an isometry and let $\text{card } A_i < \alpha$. By 2.5, there are dense sets $T_1 \subset S$, $T_2 \subset S$ and an isometry $g : T_1 \rightarrow T_2$ such that $A_i \subset T_i$ and $g \upharpoonright A_1 = f$. Since T_i are dense and S is complete, there is an isometry $h : S \rightarrow S$ such that $h \upharpoonright \text{dom } g = g$, hence $h \upharpoonright A_1 = f$.

2.7. Lemma. If $\alpha > \omega$ and $S \in \mathcal{E}(\alpha)$, then S is complete.

Proof. Let $(x_n : n \in N)$ be a Cauchy sequence in S ; assume that $x_m \neq x_n$ if $m \neq n$. Let $M = N \cup \{\omega\}$; put $\sigma(i, j) = \varrho(x_i, x_j)$, $\sigma(\omega, i) = \sigma(i, \omega) = \lim_{m \rightarrow \infty} \varrho(x_i, x_m)$. Then (M, σ) is a metric space. Due to $(S, S) \in \mathcal{E}(\alpha, 2)$, there exists a $g \in [M, S]$ such that $g(i) = x_i$ for $i \in N$. Clearly, $x_n \rightarrow g(\omega)$.

2.8. Proposition. Let $\alpha > \omega$ be a regular cardinal and let $\alpha^\alpha = \alpha$. Let $T \in \mathcal{M}$, $\text{card } T \leq 2^\omega$. Then $U_\alpha(T)$ is a complete Urysohn universal space of weight α .

Proof. Put $S = U_\alpha(T)$. By 2.2, $(S, S) \in \mathcal{E}(\alpha, \alpha^+)$, hence S is α -universal. By 2.7, S is complete. By 1.10, $\text{card } S \leq 2^\alpha$, hence $\text{ws} S \leq \alpha$. By 2.6, this proves the proposition.

2.9. Lemma. Let $\alpha \geq \omega$. If S is α -homogeneous α -universal, then $S \in \mathcal{E}(\alpha)$.

Proof. Let $M \in \mathcal{M}$, $A \subset M$, $M \setminus A = \{q\}$, $\text{card } A < \alpha$. Let $f \in [A, S]$. Since S is α -universal, there exists a $g \in [M, S]$. For $p \in g(A)$ put $h(p) = f(g^{-1}p)$. Then $h : g(A) \rightarrow f(A)$ is an isometry. Since S is α -homogeneous, there exists an isometry $v : S \rightarrow S$ such that $v \upharpoonright g(A) = h$. Clearly, $v \circ g \in [M, S]$, $(v \circ g) \upharpoonright A = f$. We have shown that $(S, S) \in \mathcal{E}(\alpha, 2)$.

2.10. Proposition. Let $\alpha \geq \omega$. Let both S_1 and S_2 be complete Urysohn universal spaces of weight α . Then $S_1 \cong S_2$.

Proof. Since $\text{w}S_i = \alpha$, there exist, by 2.5, dense sets $T_i \subset S_i$ and an isometry $g : T_1 \rightarrow T_2$. Since S_i are complete, there is an isometry $h : S_1 \rightarrow S_2$.

2.11. Lemma. If $\alpha \geq \omega$ and $(S, \varrho) \in \mathcal{E}(\alpha)$, then $\text{w}S \geq \alpha^{\mathfrak{G}}$.

Proof. I. Let $\beta < \alpha$. Let A be a set of cardinality α . Put $T = \bigcup (B^x : x < \beta)$. If $\xi \in A^x$, $\eta \in A^y$, $x < y \leq \beta$, and $\eta \upharpoonright x = \xi$, put $\xi < \eta$. Let G consist of all $g : M \rightarrow S$ such that (1) $M \subset T$, (2) if $\xi < \eta$, $\eta \in M$, then $\xi \in M$, $\varrho(g\xi, g\eta) = 1$, (3) if $x < \beta$, $\xi \in A^{x+1} \cap M$, $\eta \in A^{x+1}$, $\eta \upharpoonright x = \xi \upharpoonright x$, $\eta \neq \xi$, then $\eta \in M$, $\varrho(g\xi, g\eta) = 2$.

II. It is easy to see that G contains a maximal element, say h . We will suppose that $\text{dom } h \neq T$ and derive a contradiction. Let x be the least ordinal such that $A^x \setminus \text{dom } h \neq \emptyset$. Choose a $\xi \in A^x \setminus \text{dom } h$. Clearly, each $\eta < \xi$ is in $\text{dom } h$. – If x is a limit ordinal, then, due to $(S, S) \in \mathcal{E}(\alpha, 2)$, there is a point $q \in S$ such that $\varrho(q, h\eta) = 1$ for all $\eta < \xi$. Put $h'(\zeta) = h(\zeta)$ if $\zeta \in \text{dom } h$, and put $h(\xi) = q$. It is easy to see that $h' \in G$, which contradicts the maximality of h . – Consider the case $x = z + 1$. Since, by 2.3, $(S, S) \in \mathcal{E}(\alpha, \alpha^+)$, there exists a set $K \subset S$ of cardinality α such that $\varrho(p, q) = 2$ whenever $p, q \in K$, $p \neq q$, and $\varrho(p, h\eta) = 1$ whenever $\eta < \xi$. Let $f : A \rightarrow K$ be a bijection, put $h'(\zeta) = h(\zeta)$ if $\zeta \in \text{dom } h$, and $h'(\zeta) = f(\zeta(z))$ whenever $\zeta \in A^x$, $\zeta \upharpoonright z = \xi \upharpoonright z$. It is easy to prove that $h' \in G$, which contradicts the fact that h is maximal. Thus, we have shown that $\text{dom } h = T$.

III. If $\xi \in A^\beta$, choose a point $p(\xi) \in S$ such that $\varrho(p(\xi), h\eta) = 1/2$ for all $\eta < \xi$ (this is possible, due to $(S, S) \in \mathcal{E}(\alpha, 2)$). Put $P = \{p(\xi) : \xi \in A^\beta\}$. If $\xi, \zeta \in A^\beta$, $\xi \neq \zeta$, let x be the least ordinal such that $\xi(x) \neq \zeta(x)$. Put $z = x + 1$,

$\xi' = \xi \upharpoonright z$, $\zeta' = \zeta \upharpoonright z$. By (3), we have $\varrho(h\xi', h\zeta') = 2$. Since $\xi' < \xi$, $\zeta' < \zeta$, we have $\varrho(p(\xi), h\xi') = 1/2$, $\varrho(p(\zeta), h\zeta') = 1/2$, and therefore $\varrho(p(\xi), p(\zeta)) \geq 1$. This proves that P is a discrete subspace of cardinality α^β . Hence, $\beta < \alpha$ being arbitrary, we have $wS \geq \alpha^\alpha$.

2.12. Proposition. Let $\alpha \geq \omega$. If S is an α -homogeneous α -universal space, then $wS \geq \alpha^\alpha$.

This follows at once from 2.9 and 2.11.

3

3.1. Fact. Let $T \in \mathcal{M}$ and let $\varphi \in \text{Is}(T, T)$. Then there is an isometry Φ of $U_\omega(T)$ such that $\Phi \upharpoonright T = \varphi$ and all $U_n(T)$ and $E(U_n(T), m)$, where $n \in N$, $m > 1$, are invariant under Φ .

Proof. Put $T_n = U_n(T)$. Define φ_n as follows: $\varphi_0 = \varphi$; if $n \in N$, then $\varphi_{n+1} = E(\varphi_n) \upharpoonright T_{n+1}$. For $p \in U_\omega(T)$, let $\Phi(p) = \varphi_n(p)$, where n satisfies $p \in T_n$. It is easy to see that we obtain a mapping Φ which has the properties stated above.

3.2. Proposition. Let $T \in \mathcal{M}$, $wT \leq \omega$. Let $T_0 = T$; for $n \in N$, let T_{n+1} be a completion of $U_\omega(T_n)$. Put $P = \bigcup T_n$. Then (1) P is Urysohn universal, $wP = \omega$, (2) if $n \in N$ and $\varphi \in \text{Is}(T_n, T_n)$, then $\Phi \upharpoonright T_n = \varphi$ for some isometry Φ of P .

Proof. I. Put $S_n = U_\omega(T_n)$. From 1.8, it follows, by induction, that $wT_n = \omega$ for all $n \in N$, hence $wP = \omega$. For each $n \in N$, by 2.2, $S_n \in \mathcal{E}(\omega)$, hence S_n is ω -universal. This implies that each T_{n+1} , hence also P , is strongly ω -universal. – II. We are going to prove the assertion (2). Put $\varphi_n = \varphi$. For $k \leq n$, put $\Psi_{k+1} = E(\varphi_k) \upharpoonright S_{k+1}$ and let φ_{k+1} be the isometry of T_{k+1} such that $\varphi_{k+1} \upharpoonright S_k = \Psi_{k+1}$. In this way, we get isometries $\varphi_k \in \text{Is}(T_k, T_k)$, where $k \geq n$, such that $\varphi_{k+1} \upharpoonright T_k = \varphi_k$. For any $p \in P$, put $\Phi(p) = \varphi_m(p)$, where $m \geq n$, $p \in T_m$. Clearly, Φ is an isometry of P , $\Phi \upharpoonright T_n = \varphi$. – III. It remains to show that P is ω -homogeneous. Let $A \subset P$, $B \subset P$, $0 < \text{card } A < \omega$, and let $f \in \text{Is}(A, B)$. Choose $n \in N$ such that $A \cup B \subset S_n$. Since $S_n \in \mathcal{E}(\omega)$, there are, by 2.5, dense subsets K and M of S_n and $g \in \text{Is}(K, M)$ such that $g \upharpoonright A = f$. Hence, there is an isometry φ of T_{n+1} such that $\varphi \upharpoonright A = f$. By II, there is an isometry Φ of P such that $\Phi \upharpoonright T_{n+1} = \varphi$.

3.3. Lemma. If S is a complete space, $wS = \omega$, T is dense in S and $T \in \mathcal{E}(\omega)$, then $S \in \mathcal{E}(\omega)$.

We omit the proof, since its main idea is that of P. Ury-

sohn's proof of Theorem I in [3].

3.4. Proposition. Let S be a complete space, let $wS = \omega$ and let $T \subset S$ be dense in S . If T is Urysohn universal, then so is S .

Proof. By 2.9, $T \in \mathcal{E}(\omega)$. By 3.3, $S \in \mathcal{E}(\omega)$. By 2.6, S is Urysohn universal.

3.5. Theorem I is an immediate consequence of 3.2 and 3.4.

References

- [1] W. W. Comfort, S. Negrepointis: The theory of ultrafilters. Springer-Verlag, 1974.
- [2] P. Urysohn: Sur un espace métrique universel, C. R. Acad. Sci. Paris 180 (1925), 803–806.
- [3] P. Urysohn: Sur un espace métrique universel, Bull. Sci. Math. (2)51 (1927), 43–64.

Math. Institute, Charles University
Sokolovská 83, 18600 Praha 8
Czechoslovakia