





```
STEP GROUPS (discovery masks: AND parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=75):
    Baseline (WT), n=75 cell lines

Step 1:
    MTAP_homdel | MTAP_ko | MTAP_altered
    discovery=MTAP_homdel;MTAP_ko
```





```
STEP GROUPS (discovery masks: AND parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
------------------------
WT (n=64):
    Baseline (WT), n=64 cell lines

Step 1:
    CDKN2B_homdel | CDKN2B_ko |
    CDKN2B_altered | CDKN2A_homdel     disco
    very=CDKN2B_homdel;CDKN2B_ko;CDKN2B_alte
    red
```




```
STEP GROUPS (discovery masks: AND parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=82):
    Baseline (WT), n=82 cell lines

Step 1:
    MTAP_lowexp     discovery=MTAP_lowexp
```




```
STEP GROUPS (discovery masks: AND parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=91):
    Baseline (WT), n=91 cell lines

Step 1:
    Q7Z417_lowsangerprot
    discovery=Q7Z417_lowsangerprot
```