




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=588):
    Baseline (WT), n=588 cell lines


Step 1:
    MTAP_lowexp     discovery=MTAP_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=530):
    Baseline (WT), n=530 cell lines


Step 1:
    MTAP_homdel | MTAP_ko | MTAP_altered
    discovery=MTAP_homdel

Step 2:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=493):
    Baseline (WT), n=493 cell lines

Step 1:
    CDKN2B_altered | CDKN2B_homdel |
    CDKN2B_ko     discovery=CDKN2B_altered
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=623):
    Baseline (WT), n=623 cell lines

Step 1:
    IFNE_ko | IFNE_homdel | IFNE_altered |
    IFNA8_homdel | IFNA8_ko
    discovery=IFNE_ko


Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel

Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=596):
    Baseline (WT), n=596 cell lines

Step 1:
    DMRTA1_altered | DMRTA1_homdel |
    DMRTA1_ko     discovery=DMRTA1_altered
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=626):
    Baseline (WT), n=626 cell lines

Step 1:
    IFNA8_altered | IFNA2_homdel |
    IFNA13_homdel | IFNA6_homdel | IFNA13_ko
    | IFNA6_ko | IFNA13_altered |
    IFNA6_altered | IFNA2_altered |
    KLHL9_altered
    discovery=IFNA8_altered

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel

Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=634):
    Baseline (WT), n=634 cell lines

Step 1:
    KLHL9_homdel | IFNA2_ko | KLHL9_lowexp |
    KLHL9_ko | IFNA14_homdel | IFNA14_ko |
    IFNA16_homdel | IFNA16_ko |
    IFNA21_homdel     discovery=KLHL9_homdel

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel


Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
------------------------
WT (n=634):
    Baseline (WT), n=634 cell lines

Step 1:
    KLHL9_homdel | IFNA2_ko | KLHL9_lowexp |
    KLHL9_ko | IFNA14_homdel | IFNA14_ko |
    IFNA16_homdel | IFNA16_ko |
    IFNA21_homdel     discovery=KLHL9_homdel

Step 2:
    Q13126_lowdepmapprot
    discovery=Q13126_lowdepmapprot
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=645):
    Baseline (WT), n=645 cell lines

Step 1:
    IFNB1_homdel | HACD4_homdel |
    IFNW1_homdel | IFNW1_ko | IFNA21_ko |
    IFNB1_ko | IFNB1_altered | IFNW1_altered
    | HACD4_ko | HACD4_altered |
    FOCAD_homdel     discovery=IFNB1_homdel

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel


Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=645):
    Baseline (WT), n=645 cell lines

Step 1:
    IFNB1_homdel | HACD4_homdel |
    IFNW1_homdel | IFNW1_ko | IFNA21_ko |
    IFNB1_ko | IFNB1_altered | IFNW1_altered
    | HACD4_ko | HACD4_altered |
    FOCAD_homdel     discovery=IFNB1_homdel

Step 2:
    ELAVL2_homdel | ELAVL2_ko |
    IZUMO3_homdel
    discovery=ELAVL2_homdel
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=645):
    Baseline (WT), n=645 cell lines

Step 1:
    IFNA16_altered | IFNA14_altered |
    IFNA21_altered
    discovery=IFNA16_altered

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel


Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
------------------------
WT (n=645):
    Baseline (WT), n=645 cell lines

Step 1:
    IFNA16_altered | IFNA14_altered |
    IFNA21_altered
    discovery=IFNA16_altered


Step 2:
    ELAVL2_homdel
    discovery=ELAVL2_homdel
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
------------------------
WT (n=645):
    Baseline (WT), n=645 cell lines

Step 1:
    IFNA16_altered | IFNA14_altered |
    IFNA21_altered
    discovery=IFNA16_altered

Step 2:
    Q13126_lowdepmapprot
    discovery=Q13126_lowdepmapprot
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=635):
    Baseline (WT), n=635 cell lines

Step 1:
    ELAVL2_homdel | ELAVL2_ko |
    IZUMO3_homdel | ELAVL2_altered |
    IZUMO3_altered | IZUMO3_ko
    discovery=ELAVL2_homdel

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel


Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
------------------------
WT (n=635):
    Baseline (WT), n=635 cell lines

Step 1:
    ELAVL2_homdel | ELAVL2_ko |
    IZUMO3_homdel | ELAVL2_altered |
    IZUMO3_altered | IZUMO3_ko
    discovery=ELAVL2_homdel

Step 2:
    FOCAD_ko     discovery=FOCAD_ko
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
------------------------
WT (n=661):
    Baseline (WT), n=661 cell lines

Step 1:
    FOCAD_ko     discovery=FOCAD_ko

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel


Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=661):
    Baseline (WT), n=661 cell lines

Step 1:
    FOCAD_ko     discovery=FOCAD_ko

Step 2:
    Q13126_lowdepmapprot
    discovery=Q13126_lowdepmapprot
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=668):
    Baseline (WT), n=668 cell lines

Step 1:
    Q13126_lowdepmapprot
    discovery=Q13126_lowdepmapprot

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel


Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=668):
    Baseline (WT), n=668 cell lines

Step 1:
    Q13126_lowdepmapprot
    discovery=Q13126_lowdepmapprot


Step 2:
    MLLT3_homdel     discovery=MLLT3_homdel
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=668):
    Baseline (WT), n=668 cell lines

Step 1:
    Q13126_lowdepmapprot
    discovery=Q13126_lowdepmapprot

Step 2:
    TUSC1_altered
    discovery=TUSC1_altered
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=655):
    Baseline (WT), n=655 cell lines

Step 1:
    TUSC1_altered | TUSC1_homdel | TUSC1_ko
    discovery=TUSC1_altered

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel

Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=649):
    Baseline (WT), n=649 cell lines

Step 1:
    FOCAD_altered
    discovery=FOCAD_altered

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel

Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=673):
    Baseline (WT), n=673 cell lines

Step 1:
    MLLT3_homdel | MLLT3_ko
    discovery=MLLT3_homdel

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel

Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```





```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
------------------------
WT (n=659):
    Baseline (WT), n=659 cell lines

Step 1:
    FOCAD_lowexp     discovery=FOCAD_lowexp

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel

Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```




```
STEP GROUPS (discovery masks: OR parent subset)
Red points: sensitive. Blue points: resistant.
p-values are the discovery Wilcoxon stats used for selection.
-------------------------
WT (n=671):
    Baseline (WT), n=671 cell lines

Step 1:
    CAAP1_altered
    discovery=CAAP1_altered

Step 2:
    CDKN2A_homdel
    discovery=CDKN2A_homdel

Step 3:
    CRIP2_lowexp     discovery=CRIP2_lowexp
```