

Exercises for Week 13

The work handed in should be entirely your own. You can consult Dummit and Foote, Artin and/or the class notes but nothing else. To receive full credit, justify your answer in a clear and logical way. Due Dec. 6 (Note the special date!).

Reading. Read Dummit and Foote Sections 7.1, 7.2, 10.1. Alternatively, read Artin Sections 11.1, 11.2, 14.1.

1. Let R and S be two rings.
 - (a) Show that the product $R \times S$ is a ring under componentwise addition and multiplication.
 - (b) Show that $R \times S$ is commutative iff both R and S are commutative.
 - (c) Prove that, if $R = S$, $\{(r, r) | r \in R\} \subset R \times R$ is a subring.
2. The center of a ring R is $\{z \in R | z \cdot r = r \cdot z, \forall r \in R\}$. Prove that the center of a ring is a subring. Find the center for the $n \times n$ -matrix ring $M(n, \mathbb{R})$.
3. Use the definition of ring isomorphism to show that the group ring $F[\mathbb{Z}]$ is isomorphic to the ring of Laurent polynomials over F :

$$F[x, x^{-1}] := \left\{ \sum_{i \in \mathbb{Z}} a_i x^i \mid \text{all but finitely many } a_i \text{ is zero} \right\}.$$

4. Show that if M is an R -module, then $0 \cdot m = 0$, and $(-1) \cdot m = -m$ for all $m \in M$.
5. Prove the statement we made in class: Modules over \mathbb{Z} coincide with abelian groups.
6. Product modules. Consider a ring R and two given modules M, N over R . We define their *direct sum*

$$M \oplus N := \{(m, n) | m \in M, n \in N\},$$

with the R -action $r \cdot (m, n) := (rm, rn), \forall r \in R, m \in M, n \in N$. Show that $M \oplus N$ is an R -module.

7. Let R be a ring, and M be an R -module. Let $R^* := \{x \in R | x \text{ has a left and right inverse}\}$.
 - (a) Show that $R^* \times M \rightarrow M, (r, m) \mapsto rm$ defines a group action.
 - (b) When $R = FG$ is the group ring of a group G , and M is an F -vector space with a G action. Show that $G \subset (FG)^*$, and the action in (a), when restricted to G , agrees with the G -action on M .
8. Let F be a field, and X the following 2×2 -matrix

$$X = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Consider the $F[x]$ module $M := F^2$ by letting x act on M through the matrix X . Find a non-zero proper $F[x]$ -submodule of M .