Cairo University
Faculty of Engineering
Electronics and Electrical Communications Engineering

Assignment 1

ELC 4028 - Artificial Neural Networks

ID	*Sec*	*Name*
9220399	**2**	**Shehda Gamal Mahmoud Hussein**
9220411	**2**	**Salah Eldin Hassan Hassan Mohamed**
9220984	**4**	**Yousef Khaled Omar Mahmoud**

Supervised by:
Dr. Mohsen Rashwan
Dr. Mohamed Motaz

Contents

PART1 : Regressing Problems **3**
Problem 1: Insurance Company . . . 3
1.1 Dataset . . . 3
1.2 Case A: WITH Outlier . . . 3
1.3 Case B: WITHOUT Outlier . . . 11
1.4 Best-Fit Model Selection . . . 19
1.5 Prediction of Insured People . . . 21
Problem 2: Heating Oil . . . 21
2.1 Dataset . . . 22
2.2 Case A: WITH Outlier . . . 22
2.3 Case B: WITHOUT Outlier . . . 24
2.4 Comparison of Models and Best-Fit Selection . . . 27
2.5 Prediction of Oil . . . 28

List of Figures

1 Scatter plot of original dataset identifying the outlier at x = 3 (y = 1050) 4
2 Linear Model of Original Data with Outlier 6
3 Simulation Results of Linear Model of Original Data with Outlier 6
4 Quadratic Model of Original Data with Outlier 8
5 Simulation Results of Quadratic Model of Original Data with Outlier 8
6 Cubic Model of Original Data with Outlier 10
7 Simulation Results of Cubic Model of Original Data with Outlier 10
8 Comparison between three Models of Original Data with Outlier 11
9 Scatter Plot of Cleaned Data without Outlier 12
10 Linear Model of Cleaned Data without Outlier 14
11 Simulation Results of Linear Model of Cleaned Data without Outlier 14
12 Quadratic Model of Cleaned Data without Outlier 16
13 Simulation Results of Quadratic Model of Cleaned Data without Outlier 16
14 Cubic Model of Cleaned Data without Outlier 18
15 Simulation Results of Cubic Model of Cleaned Data without Outlier 18
16 Comparison between three Models of Cleamed Data without Outlier 19
17 R-square Comparison between three Models 19
18 Best Fit Model without Outlier 20
19 Simulation Results of Best Fit Model with Outlier 20
20 Best Fit Model with Outlier 21
21 SImulation Results of Best Fit Model without Outlier 21
22 Simulation Results of Predection Insured People at year 1997 21
23 3D Scatter Plot to Detect Outlier (outlier in red: Insulation=40) 22
24 3D Linear Regression Surface (With Outlier), R2 = 0.6967 23
25 Simulation Results of Linear Regression Surface (With Outlier) 23
26 3D Quadratic Regression Surface (With Outlier), R2 = 0.9742 24
27 Simulation Results of Quadratic Regression Surface (With Outlier) 24
28 3D Scatter Plot (Cleaned Data, Outlier Removed) 25
29 3D Linear Regression Surface (Cleaned Data), R2 = 0.9606 25
30 Simulation Results of Linear Regression Surface (Cleaned Data) 26
31 3D Quadratic Regression Surface (Cleaned Data), $R^2 = 0.9769$ 26
32 Simulation Results of Quadratic Regression Surface (Cleaned Data) 27
33 Best Fit Model of Cleaned Data (Linear Model) 27
34 Best Fit Model of Original Data (Quadratic Model) 28
35 Simulation Results of Best Fit Model 28
36 Simulation Results of Prediction Oil of Cleaned Data 29
37 Simulation Results of Prediction Oil of Original Data 29

PART1 : Regressing Problems

Problem 1: Insurance Company

The numbers of insured persons y with an insurance company for the years 1987 to 1996 are given. We let x represent the number of years since 1987 (x = 0 for 1987, x = 9 for 1996).

1.1 Dataset

x (years since 1987)	Year	y (insured persons)
0	1987	12,400
1	1988	10,900
2	1989	10,000
3	1990	**1,050 (OUTLIER)**
4	1991	9,500
5	1992	8,900
6	1993	8,000
7	1994	7,800
8	1995	7,600
9	1996	7,200

** Value 1050 at x = 3 (year 1990) is the outlier — likely a data entry error or anomaly.*

1.2 Case A: WITH Outlier

1. Scatter Plot & Outlier Identification A scatter plot of y (number of insured persons) versus x (years since 1987) reveals a generally decreasing trend from ~12,400 down to ~7,200. However, one data point stands out sharply:

- x = 3 (year 1990): y = 1050

This value is dramatically lower than the surrounding data , which are all within the range of approximately **7,200 to 12,400**. The sharp drop at this point breaks the otherwise smooth downward trend.

Therefore, this observation is identified as a clear outlier, most likely due to a data entry error (for example, a missing digit such as *10,500* instead of *1,050*).



Figure 1: Scatter plot of original dataset identifying the outlier at x = 3 (y = 1050)
dataset identifying the outlier at x = 3 (y = 1050)

2. Regression Models (WITH Outlier) We apply both OLS and matrix-form methods to compute linear model parameters, then use the matrix form for quadratic and cubic models.

2.1 Linear Model

•

The equation for the linear model is given as follows:

$$\widehat{y_i} = \beta_0 + \beta_1 x_i$$

The key formulas used are:

$$\widehat{\beta_1} = \frac{\sum_{i=i}^{n} x_i y_i \; - \; \frac{\left(\sum_{i=1}^{n} x_i\right)\left(\sum_{i=1}^{n} y_i\right)}{n}}{\sum_{i=1}^{n} x_i^2 \; - \; \frac{\left(\sum_{i=1}^{n} x_i\right)^2}{n}} \; , \; \widehat{\beta_0} = \widehat{y} \; - \; \widehat{\beta_i} x$$

Data Table for Hand Computation:

year	$\mathbf{x}_i$	$\mathbf{y}_i$	${\mathbf{x}_i}^2$	${\mathbf{y}_i}^2$	$\mathbf{x}_i\mathbf{y}_i$
1987	0	12400	0	153,760,000	0
1988	1	10900	1	118,810,000	10,900
1989	2	10000	4	100,000,000	20,000
1990	3	1050	9	1,102,500	3,150
1991	4	9500	16	90,250,000	38,000
1992	5	8900	25	79,210,000	44,500
1993	6	8000	36	64,000,000	48,000
1994	7	7800	49	60,840,000	54,600

year	$\mathbf{x}_i$	$\mathbf{y}_i$	${\mathbf{x}_i}^2$	${\mathbf{y}_i}^2$	$\mathbf{x}_i\mathbf{y}_i$
1995	8	7600	64	57,760,000	60,800
1996	9	7200	81	51,840,000	64,800
$\Sigma =$	**45**	**83,350**	**285**	**777,572,500**	**344,750**

$n \ = \ 10$, $y = \frac{83350}{10} = 8335$, $x = \frac{45}{10} = 4.5$

$\widehat{\beta_1} = \frac{344{,}750 \ - \ \frac{45 \ * \ 83{,}350}{10}}{285 - \ \frac{45^2}{10}} \ = \ - \ 367.57$

$$\widehat{\beta_0} = 8335 \ + 367.57 \ * \ \ 4.5 = 9989.09$$

•

$$\beta = (X^T X)^{-1} X^T Y$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \end{pmatrix} \ , \ \ X = \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 1 & 4 \\ 1 & 5 \\ 1 & 6 \\ 1 & 7 \\ 1 & 8 \\ 1 & 9 \end{pmatrix} \ , \ Y \ = \begin{pmatrix} 12400 \\ 10900 \\ 10000 \\ 1050 \\ 9500 \\ 8900 \\ 8000 \\ 7800 \\ 7600 \\ 7200 \end{pmatrix}$$

$$\begin{pmatrix} \beta_0 \\ \beta_1 \end{pmatrix} = \begin{pmatrix} 9989.09 \\ -367.576 \end{pmatrix}$$

So ,linear equation is:

$$\widehat{y_i} = \ 9989.09 - 367.576 \ x$$

•

$$\overline{y} = \frac{\sum_{i=1}^{n} y_i}{n} = \frac{83350}{10} = 8335$$

year	$\mathbf{x}_i$	$\mathbf{y}_i$	$\hat{y}_i = \beta_\mathbf{0} + \beta_\mathbf{1}\mathbf{x_i}$	$\mathbf{SSE} = (\mathbf{y}_i - \hat{y}_i)^2$	$\mathbf{SSyy} = (\mathbf{y}_i - \bar{y})^2$
1987	0	12400	9,989.1	5,812,483	16,524,225
1988	1	10900	9,621.5	1,634,524	6,579,225
1989	2	10000	9,253.9	556,606	2,772,225
1990	3	1050	8,886.4	61,408,600	53,071,225
1991	4	9500	8,518.8	962,777	1,357,225
1992	5	8900	8,151.2	560,683	319,225
1993	6	8000	7,783.6	46,813	112,225
1994	7	7800	7,416.1	147,410	286,225
1995	8	7600	7,048.5	304,169	540,225
1996	9	7200	6,680.9	269,455	1,288,225
$\Sigma =$	**45**	**83,350**	**83,350**	**71,703,515**	**82,850,250**

$$R^2_{linear\ model} = \frac{SS - SSE}{SS} = \frac{82,850,250 - 71,703,515}{82,850,250} = 0.13454$$

•



Figure 2: Linear Model of Original Data with Outlier

•

```
Linear Model:
  Equation: y = 9989.09 + (-367.58)*x^1
  R^2: 0.1345
```

Figure 3: Simulation Results of Linear Model of Original Data with Outlier

2.2 Quadratic Model The equation for the quadratic model is given as follows:

$$\widehat{y_i} = \beta_0 + \beta_1 x_i + \beta_2 x_i^2$$

•

$$\beta = (X^T X)^{-1} X^T Y$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \end{pmatrix} \quad , \quad X = \begin{pmatrix} 1 & 0 & 0^2 \\ 1 & 1 & 1^2 \\ 1 & 2 & 2^2 \\ 1 & 3 & 3^2 \\ 1 & 4 & 4^2 \\ 1 & 5 & 5^2 \\ 1 & 6 & 6^2 \\ 1 & 7 & 7^2 \\ 1 & 8 & 8^2 \\ 1 & 9 & 9^2 \end{pmatrix} \quad , \; Y \; = \begin{pmatrix} 12400 \\ 10900 \\ 10000 \\ 1050 \\ 9500 \\ 8900 \\ 8000 \\ 7800 \\ 7600 \\ 7200 \end{pmatrix}$$

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} 11627.73 \\ -1596.55 \\ 136.55 \end{pmatrix}$$

So ,linear equation is:

$$\widehat{y_i} = 11627.73 - 1596.55x + 136.55x^2$$

•

$$\overline{y} = \frac{\sum_{i=1}^{n} y_i}{n} = \frac{83350}{10} = 8335$$

year	$\mathbf{x}_i$	$\mathbf{y}_i$	$\hat{y}_i = \beta_0 + \beta_1 \mathbf{x_i} + \beta_2 \mathbf{x_i^2}$	**SSE** $= (\mathbf{y}_i - \hat{y}_i)^2$	**SSyy** $= (\mathbf{y}_i - \bar{y})^2$
1987	0	12400	**11,627.7**	**596,405**	**16,524,225**
1988	1	10900	10,167.7	536,223	6,579,225
1989	2	10000	8,980.8	1,038,701	2,772,225
1990	3	1050	8,067.0	49,238,925	53,071,225
1991	4	9500	7,426.4	4,299,969	1,357,225
1992	5	8900	7,058.8	3,390,064	319,225
1993	6	8000	6,964.3	1,072,639	112,225
1994	7	7800	7,143.0	431,710	286,225
1995	8	7600	7,594.7	28	540,225
1996	9	7200	8,319.5	1,253,377	1,288,225
$\Sigma =$	**45**	**83,350**	83,350	61,858,042	82,850,250

$$R^2_{quad\ model} = \frac{SS - SSE}{SS} = \frac{82,850,250 - \mathbf{61,858,042}}{82,850,250} = 0.2534$$

•

•



Figure 4: Quadratic Model of Original Data with Outlier

```
Quadratic Model:
  Equation: y = 11627.73 + (-1596.55)*x^1 + (136.55)*x^2
  R^2: 0.2534
```

Figure 5: Simulation Results of Quadratic Model of Original Data with Outlier

2.3 Cubic Model The equation for the cubic model is given as follows:

$$\widehat{y_i} = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3$$

•

$$\beta = (\mathbf{X^T X})^{-1} \mathbf{X^T Y}$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}, \quad X = \begin{pmatrix} 1 & 0^2 & 0^3 \\ 1 & 1^2 & 1^3 \\ 1 & 2^2 & 2^3 \\ 1 & 3^2 & 3^3 \\ 1 & 4^2 & 4^3 \\ 1 & 5^2 & 5^3 \\ 1 & 6^2 & 6^3 \\ 1 & 7^2 & 7^3 \\ 1 & 8^2 & 8^3 \\ 1 & 9^2 & 9^3 \end{pmatrix}, \quad Y = \begin{pmatrix} 12400 \\ 10900 \\ 10000 \\ 1050 \\ 9500 \\ 8900 \\ 8000 \\ 7800 \\ 7600 \\ 7200 \end{pmatrix}$$

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 13113.15 \\ -4313.93 \\ 932.31 \\ -58.95 \end{pmatrix}$$

So ,linear equation is:

$$\widehat{y_i} = 13113.15 - 4313.93\ x_i\ +\ 932.31\ x_i^2\ -\ \ 58.95\ x_i^3$$

•

$$\overline{y} = \frac{\sum_{i=1}^{n} y_i}{n} = \frac{83350}{10} = 8335$$

year	$\mathbf{x}_i$	$\mathbf{y}_i$	$\hat{y}_i = \beta_0 + \beta_1 \mathbf{x_i} + \beta_2 \mathbf{x_i^2} + \beta_3 \mathbf{x_i^3}$	**SSE** $= (\mathbf{y}_i - \hat{y}_i)^2$	**SSyy** $= (\mathbf{y}_i - \bar{y})^2$
1987	0	12400	**13,113.1**	**508,578**	**16,524,225**
1988	1	10900	9,672.6	1,506,542	6,579,225
1989	2	10000	7,743.0	5,094,123	2,772,225
1990	3	1050	6,970.7	35,054,266	53,071,225
1991	4	9500	7,002.0	6,240,214	1,357,225
1992	5	8900	7,483.2	2,007,341	319,225
1993	6	8000	8,060.7	3,684	112,225
1994	7	7800	8,380.8	337,334	286,225
1995	8	7600	8,089.8	239,940	540,225
1996	9	7200	6,834.1	133,864	1,288,225
$\Sigma =$	**45**	**83,350**	83,350	51,125,885	82,850,250

$$R^2_{cubic\ model} = \frac{SS\ -\ SSE}{SS} = \frac{82,850,250\ \mathbf{-51,125,885}}{82,850,250} = 0.3829$$

•

•



Figure 6: Cubic Model of Original Data with Outlier

```
Cubic Model:
  Equation: y = 13113.15 + (-4313.93)*x^1 + (932.31)*x^2 + (-58.95)*x^3
  R^2: 0.3829
```

Figure 7: Simulation Results of Cubic Model of Original Data with Outlier

3. Comparison between three models (WITH Outlier):

Model	Equation (MATLAB)	R^2
Linear	$\widehat{y}_i = 9989.09 - 367.576\ x_i$	0.1345
Quadratic	$\widehat{y}_i = 11627.73 - 1596.55\ x_i + 136.55\ x_i^2$	0.2534
Cubic	$\widehat{y}_i = 13113.15 - 4313.93\ x_i + 932.31\ x_i^2 - 58.95\ x_i^3$	0.3829



Figure 8: Comparison between three Models of Original Data with Outlier

Comment:

All R^2 values are low (< 0.50) due to the heavy influence of the outlier at x = 3 (y = 1050). The cubic model yields the highest R.

1.3 Case B: WITHOUT Outlier

1. Outlier Removal and Scatter Plot of Cleaned Data



Figure 9: Scatter Plot of Cleaned Data without Outlier

2. Regression Models (WITHOUT Outlier) After removing the outlier at x = 3 (y = 1,050), the cleaned dataset contains 9 data points. We now repeat the regression analysis using the same methods.

2.1 Linear Model

•

The equation for the linear model is given as follows:

$$\widehat{y_i} = \beta_0 + \beta_1 x_i$$

Data Table for Hand Computation:

year	$\mathbf{x}_i$	$\mathbf{y}_i$	${\mathbf{x}_i}^2$	${\mathbf{y}_i}^2$	$\mathbf{x}_i\mathbf{y}_i$
1987	0	12,400	0	153,760,000	0
1988	1	10,900	1	118,810,000	10,900
1989	2	10,000	4	100,000,000	20,000
1991	4	9,500	16	90,250,000	38,000
1992	5	8,900	25	79,210,000	44,500
1993	6	8,000	36	64,000,000	48,000
1994	7	7,800	49	60,840,000	54,600
1995	8	7,600	64	57,760,000	60,800
1996	9	7,200	81	51,840,000	64,800
$\Sigma =$	**42**	**82,300**	**276**	**776,470,000**	**341,600**

$n = 9$, $y = \frac{82300}{9} = 9,144.44$, $x = \frac{42}{9} = 4.667$

$\widehat{\beta_1} = \frac{341{,}600 \ - \ \frac{42*82300}{9}}{276 - \ \frac{42^2}{9}} \ = \ - \ 530.83$

$$\widehat{\beta_0} = 9,144.44 \ + 530.83 \ * \ 4.667 = 11,621.67$$

•

$$\beta = (X^T X)^{-1} X^T Y$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \end{pmatrix} \ , \ X = \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 2 \\ 1 & 4 \\ 1 & 5 \\ 1 & 6 \\ 1 & 7 \\ 1 & 8 \\ 1 & 9 \end{pmatrix} \ , \ Y \ = \begin{pmatrix} 12400 \\ 10900 \\ 10000 \\ 9500 \\ 8900 \\ 8000 \\ 7800 \\ 7600 \\ 7200 \end{pmatrix}$$

$$\begin{pmatrix} \beta_0 \\ \beta_1 \end{pmatrix} = \begin{pmatrix} 11,621.67 \\ -530.83 \end{pmatrix}$$

So ,linear equation is:

$$\widehat{y_i} = 11,621.67 \ - 530.83 \ x$$

•

$$\widehat{y_i} = \frac{82300}{\mathbf{9}} = 9,144.44$$

year	$\mathbf{x}_i$	$\mathbf{y}_i$	$\hat{y}_i = \beta_0 + \beta_1 \mathbf{x_i}$	$\mathbf{SSE} = (\mathbf{y}_i - \hat{y}_i)^2$	$\mathbf{SSyy} = (\mathbf{y}_i - \bar{y})^2$
1987	0	12400	11621.7	605,803	10,598,642
1988	1	10900	11090.8	36,417	3,081,975
1989	2	10000	10560.0	313,600	731,975
1991	4	9500	9498.3	3	126,420
1992	5	8900	8967.5	4,556	59,753
1993	6	8000	8436.7	190,678	1,309,753
1994	7	7800	7905.8	11,201	1,807,531
1995	8	7600	7375.0	50,625	2,385,309
1996	9	7200	6844.2	126,617	3,780,864
$\Sigma =$	**42**	**82,300**	**82300.0**	**1,339,500**	**23,882,222**

$$R^2_{linear\ model} \ = \ \frac{SS \ - \ SSE}{SS} \ = \ \frac{23,882,222 \ - \ 1,339,500}{23,882,222} \ = \ 0.9439$$

•



Figure 10: Linear Model of Cleaned Data without Outlier

•

```
Linear Model:
  Equation: y = 11621.67 + (-530.83)*x^1
  R^2: 0.9439
```

Figure 11: Simulation Results of Linear Model of Cleaned Data without Outlier

2.2 Quadratic Model The equation for the quadratic model is given as follows:

$$\widehat{y_i} = \beta_0 + \beta_1 x_i + \beta_2 x_i^2$$

•

$$\beta = (\mathbf{X^T X})^{-1} \mathbf{X^T Y}$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \end{pmatrix} \quad , \quad \mathbf{X} = \begin{pmatrix} 1 & 0 & 0^2 \\ 1 & 1 & 1^2 \\ 1 & 2 & 2^2 \\ 1 & 4 & 4^2 \\ 1 & 5 & 5^2 \\ 1 & 6 & 6^2 \\ 1 & 7 & 7^2 \\ 1 & 8 & 8^2 \\ 1 & 9 & 9^2 \end{pmatrix} \quad , \quad \mathbf{Y} = \begin{pmatrix} 12400 \\ 10900 \\ 10000 \\ 9500 \\ 8900 \\ 8000 \\ 7800 \\ 7600 \\ 7200 \end{pmatrix}$$

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} \mathbf{12,024.17} \\ \mathbf{-863.13} \\ \mathbf{37.44} \end{pmatrix}$$

So ,linear equation is:

$$\widehat{\mathbf{y_i}} = \mathbf{12,024.17} \ - \mathbf{863.13\ x} + \mathbf{37.44\ x^2}$$

•

$$\overline{y} = \frac{\sum_{i=1}^{n} y_i}{n} = \frac{\mathbf{82,300}}{9} = 9,144.44$$

year	$\mathbf{x}_i$	$\mathbf{y}_i$	$\hat{y}_i = \beta_0 + \beta_1 \mathbf{x_i} + \beta_2 \mathbf{x_i^2}$	$\mathbf{SSE} = (\mathbf{y}_i - \hat{y}_i)^2$	$\mathbf{SSyy} = (\mathbf{y}_i - \bar{y})^2$
1987	0	12400	12024.2	141,248	10,598,642
1988	1	10900	11198.5	89,090	3,081,975
1989	2	10000	10447.7	200,411	731,975
1991	4	9500	9170.7	108,429	126,420
1992	5	8900	8644.6	65,249	59,753
1993	6	8000	8193.3	37,362	1,309,753
1994	7	7800	7816.9	286	1,807,531
1995	8	7600	7515.4	7,156	2,385,309
1996	9	7200	7288.8	7,884	3,780,864
$\Sigma =$	**42**	**82,300**	**82300.0**	**657,116**	**23,882,222**

$$\mathbf{R^2_{quad\ model}} = \frac{\mathbf{SS\ -\ SSE}}{\mathbf{SS}} = \frac{\mathbf{23,882,222 - 657,116}}{\mathbf{23,882,222}} = \mathbf{0.9725}$$

•



Figure 12: Quadratic Model of Cleaned Data without Outlier

Cleaned Data without Outlier

•

```
Quadratic Model:
  Equation: y = 12024.17 + (-863.13)*x^1 + (37.44)*x^2
  R^2: 0.9725
```

Figure 13: Simulation Results of Quadratic Model of Cleaned Data without Outlier

2.3 Cubic Model The equation for the cubic model is given as follows:

$$\widehat{y_i} = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3$$

•

$$\beta = (\mathbf{X^T X})^{-1} \mathbf{X^T Y}$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}, \quad X = \begin{pmatrix} 1 & 0^2 & 0^3 \\ 1 & 1^2 & 1^3 \\ 1 & 2^2 & 2^3 \\ 1 & 4^2 & 4^3 \\ 1 & 5^2 & 5^3 \\ 1 & 6^2 & 6^3 \\ 1 & 7^2 & 7^3 \\ 1 & 8^2 & 8^3 \\ 1 & 9^2 & 9^3 \end{pmatrix}, \quad Y = \begin{pmatrix} 12400 \\ 10900 \\ 10000 \\ 9500 \\ 8900 \\ 8000 \\ 7800 \\ 7600 \\ 7200 \end{pmatrix}$$

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 12,204.42 \\ -1,221.74 \\ 140.17 \\ -7.46 \end{pmatrix}$$

So ,linear equation is:

$$\widehat{y_i} = \mathbf{12,204.42} \ \mathbf{-1,221.74} \ x_i + \mathbf{140.17} \ x_i^2 - \ \mathbf{7.46} \ x_i^3$$

•

$$\overline{y} = \frac{\sum_{i=1}^{n} y_i}{n} = \frac{82,300}{9} = 9,144.44$$

year	$\mathbf{x}_i$	$\mathbf{y}_i$	$\hat{y}_i = \beta_0 + \beta_1 \mathbf{x_i} + \beta_2 \mathbf{x_i^2} + \beta_3 \mathbf{x_i^3}$	**SSE** $= (\mathbf{y}_i - \hat{y}_i)^2$	**SSyy** $= (\mathbf{y}_i - \bar{y})^2$
			12204.4	38,251	
1987	0	12400			10,598,642
			11115.4	46,392	
1988	1	10900			3,081,975
			10261.9	68,595	731,975
1989	2	10000			
			8666.9		126,420
1991	4	9500		174,339	
			8307.8	54,323	59,753
1992	5	8900			
			7960.3	94,746	
1993	6	8000			1,309,753
			7579.7	25,702	
1994	7	7800			1,807,531
			7121.1	413	
1995	8	7600			2,385,309
			82300.0	6,227	
1996	9	7200			3,780,864
			12204.4		
$\Sigma =$	**42**	82,300		**508,988**	**23,882,222**

$$R^2_{cubic\ model} = \frac{SS - SSE}{SS} = \frac{\mathbf{23,882,222} - \mathbf{508,988}}{\mathbf{23,882,222}} = 0.9787$$

•



Figure 14: Cubic Model of Cleaned Data without Outlier

•

```
Cubic Model:
  Equation: y = 12204.42 + (-1221.74)*x^1 + (140.17)*x^2 + (-7.46)*x^3
  R^2: 0.9787
```

Figure 15: Simulation Results of Cubic Model of Cleaned Data without Outlier

1.3 Comparison Between Three Models (Cleaned Data):

Model	Equation (MATLAB)	R^2
Linear	$\widehat{y}_i = 11,621.67 \ - 530.83\ x$	0.9439
Quadratic	$\widehat{y}_i = 12,024.17 \ - 863.13\ x + 37.44\ x^2$	0.9725
Cubic	$\widehat{y}_i = 12,204.42 \ - 1,221.74\ x_i + 140.17\ x_i^2 - \ 7.46\ x_i^3$	0.9787



Figure 16: Comparison between three Models of Cleamed Data without Outlier

Comment:

All three R^2 values are very high (> 0.94) after removing the outlier, confirming that it was the primary source of distortion. The linear model alone explains 94.4% of the variance.

1.4 Best-Fit Model Selection



Figure 17: R-square Comparison between three Models

R^2 Comparison

between three Models

Model	Parameters	R^2 (Original) → R^2 (Cleaned)	Gain in R^2 (Original) → Gain in R^2 (Cleaned)
Linear	2	0.1345 → 0.9439	—
Quadratic	3	0.2534 → 0.9725	+0.1189 → +0.0286
Cubic	4	0.3829 → 0.9787	+0.1295 → +0.0062

Conclusion (with outlier):

- Linear → Quadratic: R^2 gain = 0.1189 (significant) → Quadratic is better.
- Quadratic → Cubic: R^2 gain = 0.1295 (significant) → Cubic is better.

Best-fit model (with outlier): Cubic $\hat{y} = 13113.15 - 4313.93\,x + 932.31\,x^2 - 58.95\,x^3$ ($R^2 = 0.3829$)

Conclusion (without outlier):

- Linear → Quadratic: R^2 gain = 0.0286 (significant) → Quadratic is better.
- Quadratic → Cubic: R^2 gain = 0.0062 → gain is NOT significant enough.

Best-fit model (without outlier): Quadratic $\hat{y} = 12{,}024.17 - 863.13\,x + 37.44\,x^2$ ($R^2 = 0.972$)



Figure 18: Best Fit Model without Outlier

```
>>  Model Selection: Cubic recommended
   dR2 (Linear -> Quadratic) = +0.1188
   dR2 (Quadratic -> Cubic)  = +0.1295
```

Figure 19: Simulation Results of Best Fit Model with Outlier



Figure 20: Best Fit Model with Outlier

```
>>  Model Selection: Quadratic recommended
   dR2 (Linear -> Quadratic) = +0.0286
   dR2 (Quadratic -> Cubic)  = +0.0062
```

Figure 21: SImulation Results of Best Fit Model without Outlier

1.5 Prediction of Insured People

At year 1997 (x=10) :

- Using the best-fit model from the original data with outlier (Cubic model as selected):

$$\widehat{y_i} = 13113.15 - 4313.93\ x_i + 932.31\ x_i^2 - \ 58.95\ x_i^3$$

$\hat{y} = 13113.15 - 4313.93(10) + 932.31(10^2) - 58.95(10^3) = 4{,}254.85$

Insured People ≈ 4,254.85 insured person

- Using the best-fit model from the cleaned data without outlier (Quadratic model as selected):

$$\widehat{y_i} = 12,024.17\ - 863.13\ x + 37.44\ x^2$$

$\hat{y} = 12{,}024.17 - 863.13(10) + 37.44(10^2) = 7{,}136.87$

Insured People ≈ 7,136.87 insured person

```
============================================================
  PREDICTIONS FOR 1997  (year = 1997 , x = 10)
============================================================
  Original Data with-outlier (Cubic Model) -> 4,260.00 insured persons

  Cleaned Data without-outlier (Quadratic Model) -> 7,137.06 insured persons
```

Figure 22: Simulation Results of Predection Insured People at year 1997

Predection Insured People at year 1997

Problem 2: Heating Oil

We develop a regression model to estimate heating oil usage for a single-family home in January, based on two predictors: average temperature (Temp, in °F) and amount of insulation (Insulation, in inches).

2.1 Dataset

Oil (gallons)	Temp (°F)	Insulation (in)
270	40	4
362	27	4
162	40	10
45	73	6
91	65	7
233	**65**	**40 ← OUTLIER**
372	10	6
305	9	10
234	24	10
122	65	4
25	66	10
210	41	6
450	22	4
325	40	4
52	60	10

2.2 Case A: WITH Outlier

1. Scatter Plot & Outlier Identification A 3D scatter plot of Oil Usage vs. Temperature and Insulation reveals that one data point is far outside the cluster formed by the other observations: the point (Temp=65°F, Insulation=40 in, Oil=233 gallons). All other insulation values range from 4 to 10 inches; an insulation thickness of 40 inches is physically unreasonable for a typical single-family home and is almost certainly a data entry error (e.g., the value 4 recorded as 40). This point is identified as the outlier and is highlighted in red in the scatter plot below.



Figure 23: 3D Scatter Plot to Detect Outlier (outlier in red: Insulation=40)

Detect Outlier (outlier in red: Insulation=40)

2. Regression Models

2.1 Linear Model



Figure 24: 3D Linear Regression Surface (With Outlier), R2 = 0.6967

The multivariate linear model has the form:

$$Oil = \beta_0 + \beta_1 * \ T \ + \beta_2 * \ I$$

where T = Temp, I = Insulation

$$Oil = 426.97 \ - \ 5.274 * T \ + 1.966 * I$$

```
Linear model:
  Intercept : 426.967656
  Coef      : Temp=-5.273465,  Insulation=1.966053
  Equation  : y = 426.9677  +  (-5.2735)*Temp  +  (1.9661)*Insulation
  R2        : 0.696686
```

Figure 25: Simulation Results of Linear Regression Surface (With Outlier)

So, the linear fit equation for the original dataset is:

$$\widehat{y_i} = 426.97 - \ 5.27x_{1i} + 1.966 \ \ x_{2i}$$

The Coefficient of Determination (R^2):

R^2 (Linear, With Outlier) = 0.6967

2.2 Quadratic Model



Figure 26: 3D Quadratic Regression Surface (With Outlier), R2 = 0.9742

Surface (With Outlier), $R^2 = 0.9742$

The multivariate linear model has the form:

$$Oil = \beta_0 + \beta_1\ T\ + \beta_2\ I\ + \beta_3 T^2 + \beta_4 TI + \beta_5\ I^2$$

```
Quadratic model:
  Intercept : 680.025958
  Temp                : -6.167784
  Insulation          : -41.716060
  Temp^2              : -0.012475
  Temp Insulation     : 0.239147
  Insulation^2        : 0.658593
  Equation  : y = 680.0260  +  (-6.1678)*Temp  +  (-41.7161)*Insulation  +  (-0.0125)*Temp^2  +  (0.2391)*Temp
  Insulation  +  (0.6586)*Insulation^2
  R2        : 0.974152
```

Figure 27: Simulation Results of Quadratic Regression Surface (With Outlier)

where T = Temp, I = Insulation

$$Oil = 680.03\ -\ 6.168\ T\ -\ 41.716\ I\ -\ 0.01248\ T^2 + 0.2392\ TI + 0.6586\ I^2$$

So, the quadratic fit equation for the original dataset is:

$$\widehat{y_i} = 680\ -\ 6.17x_{1i} - 41.72\ x_{2i} -\ 0.012\ x_{1i}^2 + 0.24\ x_{1i}x_{2i} + 0.66\ x_{2i}^2$$

The Coefficient of Determination (R^2):

R^2 (Quadratic, With Outlier) = 0.9742

2.3 Case B: WITHOUT Outlier

1. Scatter Plot & Outlier Identification (WITHOUT Outlier) The data point (Temp=65, Insulation=40, Oil=233) is removed. All remaining 14 data points have insulation values between 4 and 10 inches, which is physically reasonable. The cleaned scatter plot is shown below.



Figure 28: 3D Scatter Plot (Cleaned Data, Outlier Removed)

2. Regression Models (WITHOUT Outlier)

2.1 Linear Model



Figure 29: 3D Linear Regression Surface (Cleaned Data), R2 = 0.9606

Surface (Cleaned Data), $R^2 = 0.9606$

The multivariate linear model has the form:

$$Oil = \beta_0 + \beta_1 * \ T \ + \beta_2 * \ I$$

where T = Temp, I = Insulation

$$Oil = 601.6 \ - \ 5.482 \ T \ - \ 23.232 \ I$$

```
Linear model:
  Intercept : 601.601412
  Coef      : Temp=-5.481747,  Insulation=-23.232031
  Equation  : y = 601.6014  +  (-5.4817)*Temp  +  (-23.2320)*Insulation
  R2        : 0.960609
```

Figure 30: Simulation Results of Linear Regression Surface (Cleaned Data)

So, the linear fit equation for the original dataset is:

$$\widehat{y_i} = 601.6 - \ 5.482 \ x_{1i} + 23.232 \ \ x_{2i}$$

The Coefficient of Determination (R^2):
R^2 (Linear, With Outlier) = 0.96061

2.2 Quadratic Model



Figure 31: 3D Quadratic Regression Surface (Cleaned Data), R^2 = 0.9769

Surface (Cleaned Data), R^2 = 0.9769

The multivariate quadratic model has the form:

$$Oil = \beta_0 + \beta_1 \ T \ + \beta_2 \ I \ + \beta_3 T^2 + \beta_4 TI + \beta_5 \ I^2$$

where T = Temp, I = Insulation

$$Oil = 785.52 - \ 6.963 \ T \ - \ 72.159 \ I \ - \ 0.00348 \ T^2 + 0.2567 \ TI + 2.7438 \ I^2$$

```
Quadratic model:
  Intercept : 785.516872
  Temp                : -6.962462
  Insulation          : -72.158742
  Temp^2              : -0.003478
  Temp Insulation     : 0.256701
  Insulation^2        : 2.743800
  Equation  : y = 785.5169  +  (-6.9625)*Temp  +  (-72.1587)*Insulation  +  (-0.0035)*Temp^2  +  (0.2567)*Temp
  Insulation  +  (2.7438)*Insulation^2
  R2        : 0.976889
```

Figure 32: Simulation Results of Quadratic Regression Surface (Cleaned Data)

So, the linear fit equation for the original dataset is:

$$\widehat{y_i} = 785.52 \; - \; 6.963 \; x_{1i} - 72.159 \; x_{2i} - \; 0.00348 \; x_{1i}^2 + 0.2567 \; x_{1i}x_{2i} + 2.7438 \; x_{2i}^2$$

The Coefficient of Determination (R^2):

R^2 (Quadratic, With Outlier) = 0.9769

2.4 Comparison of Models and Best-Fit Selection

Model	R^2 (With Outlier)	$R^{2\ 2}$ (Without Outlier)	Gain (With Outlier)	Gain ^{2} (Without Outlier)
Linear	0.6967	0.9606	—	—
Quadratic	0.9742	0.9769	+0.2775	+0.0163

Conclusion (Part a — With Outlier):

Linear → Quadratic gain = +0.2775

Best-fit model (with outlier): Quadratic (R^2 = 0.9742).

Conclusion (Part b — Without Data):

Linear → Quadratic gain = +0.0163 (NOT significant enough).

Best-fit model (without outlier): Linear (R^2 = 0.9606).



Figure 33: Best Fit Model of Cleaned Data (Linear Model)



Figure 34: Best Fit Model of Original Data (Quadratic Model)

After removing the outlier, the linear model already achieves $R^2 = 0.9606$. The gain from linear to quadratic is only +0.0163, which is NOT significant enough as specified in Note 4 of the assignment.

- **Simulation Results:**

```
Part a  (With Outlier):
  dR2 (Linear -> Quadratic) = +0.2775  -->  significant gain
  Best-fit model: Quadratic  (R2 = 0.9742)

Part b  (Without Outlier):
  dR2 (Linear -> Quadratic) = +0.0163  -->  not significant
  Best-fit model: Linear     (R2 = 0.9606)
```

Figure 35: Simulation Results of Best Fit Model

2.5 Prediction of Oil

Temp =15°F, Insulation = 5 inches

The linear model could be abstracted as follows:

$\hat{y}_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}$

Linear Model	**Data with Outlier**	**Data without Outlier (Best Fit)**
β_0	427	601.6
β_1	- 5.273	- 5.482
β_2	1.966	- 23.23
$\mathbf{R}^2$	0.6966	0.9606
$\hat{y}$**(temp = 15, insulation = 5)**	357.7	403.22

- Using the best-fit model from the cleaned data without outlier (Linear model as selected):

Oil = 601.60 − 5.482 * 15 − 23.232 * 5 = 403.21

Predicted Oil Usage ≈ 403.2 gallons

- **Simulation Results:**

```
Prediction at (15F,5in) using CLEANED Linear best-fit: 403.22 units
```

Figure 36: Simulation Results of Prediction Oil of Cleaned Data

Prediction Oil of Cleaned Data

The quadratic model could be abstracted as follows:

$\hat{y}_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \beta_3 {x_1}^2{}_i + \beta_4 x_{1i} x_{2i} + \beta_5 {x_2}^2{}_i$

Linear Model	**Data with Outlier (Best Fit)**	**Data without Outlier**
β_0	680	785.5
β_1	- 6.17	- 6.96
β_2	- 41.72	- 72.16
β_3	- 0.012	- 0.0034
β_4	0.24	0.2567
β_5	0.66	2.744
$\mathbf{R}^2$	0.9741	0.9768
$\hat{y}$**(temp = 15, insulation = 5)**	410.52	407.35

- Using the best-fit model from the original data with outlier (Quadratic model as selected):

Oil = 680 + $(-6.17)(15)$ + $(-41.72)(5)$ + $(-0.012)(15^2)$ + $(0.24)(15)(5)$ + $(0.66)(5^2)$ = 410.65

Predicted Oil Usage ≈ 410.65 gallons

- **Simulation Results:**

```
Prediction at (15F,5in) using ORIGINAL Quadratic best-fit: 410.52 units
```

Figure 37: Simulation Results of Prediction Oil of Original Data

Prediction Oil of Original Data