ξ-Bundle Curvature as a Critical-Line Discriminator

Universality Across L-Function Families and an Analytic Detection Theorem

Kim Young-Hu

Independent Researcher

April 20, 2026

(Companion paper to "Phase Quantisation from Optical Lattices to the Zeta Landscape," 2026)

Abstract

Extending the ξ-bundle framework of Kim (2026), we demonstrate that the curvature invariant $\kappa\delta^2$ satisfies four universal properties—functional-equation verification, zero enumeration, $\kappa\delta^2 \approx 1$, and monodromy $\pm\pi$—across an enlarged family of L-functions: elliptic-curve L-functions (rank 0–3, conductor 11–10099), Rankin-Selberg GL(4), Dedekind ζ-functions (7 imaginary quadratic fields), and Epstein ζ-functions including three *non-Euler* forms (class number $h > 1$). The Epstein result establishes that the ξ-bundle depends only on the functional equation structure, not on the presence of an Euler product (**Theorem 6**).

A systematic investigation of the Davenport–Heilbronn (DH) function, which satisfies a functional equation yet possesses off-critical zeros, reveals a sharp discriminator: on-critical zeros satisfy $\kappa\delta^2 = 1 + O(\delta^2)$, whereas off-critical zeros satisfy $\kappa\delta^2 = 1 + c_1\delta + O(\delta^2)$ with $c_1 \neq 0$. We prove analytically (**Theorem 7**) that $c_1 = 0$ if and only if $\sigma_0 = \frac{1}{2}$, and verify numerically that $c_1 = \mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$ to within 0.5% for all four known DH off-critical zeros. A δ-sweep across eight values confirms the log-log scaling exponents 2.00 ± 0.003 (on-critical) vs. 0.96 ± 0.04 (off-critical), with complete separation at every tested δ. These results yield an ξ-bundle-based discriminator for the Riemann Hypothesis: any zero off the critical line produces a detectable $O(\delta)$ correction to $\kappa\delta^2$.

Four additional results (#125–#127, #200) establish *weight and spectral universality*: the $\kappa\delta^2$ log-log slope equals 2.0 ± 0.001 for the unique normalised Hecke eigenforms of modular weights $k = 12$ (Ramanujan's Δ, Result #125, slope 2.0008 ± 0.0006), $k = 16$ ($\Delta \cdot E_4$, Result #126, slope 1.9989 ± 0.0035), and $k = 20$ ($\Delta \cdot E_8$, Result #127, slope 1.9984 ± 0.0055). Result #200 extends this to a *non-holomorphic* Maass cusp form (spectral parameter $R = 9.5337$, slope 2.0003 ± 0.0003—the most precise measurement to date). Combined with the $k = 0$ and $k = 2$ results from Paper 1, the slope 2 is confirmed across all tested weights and spectral parameters, demonstrating that the $\kappa\delta^2$ universality is independent of both modular weight and holomorphicity—a numerical consequence of the functional equation alone (Observation 8).

Fifteen numerical results support the framework's universality and the discriminator theorem.

Keywords: ξ-bundle, $\kappa\delta^2$ curvature, elliptic-curve L-functions, Epstein ζ-function, Davenport–Heilbronn function, functional equation, critical-line discriminator, monodromy, Riemann Hypothesis

Contents

1 Introduction **4**

2 Framework Recap **5**

3 Elliptic-Curve L-Functions **5**
3.1 Sign of the Functional Equation and the ε-Correction 5
3.2 κ_{near} Universality Across Rank and Conductor 6
3.3 Monodromy Precision 7
3.4 Conductor Scaling 7

4 Beyond the Euler Product **7**
4.1 Rankin–Selberg GL(4) 7
4.2 Dedekind ζ-Functions 8
4.3 Epstein ζ-Functions: Functional-Equation Sufficiency 8

5 Critical-Line Discrimination: The DH Function **9**
5.1 The Davenport–Heilbronn Function 9
5.2 Four-Property Test on DH Zeros 9
5.3 Theorem 7: Analytic Discrimination 10
5.4 δ-Sweep Validation 10
5.5 The c_1 Coefficient: Analytic Formula and Asymptotics 11

6 Weight and Spectral Universality **12**
6.1 Ramanujan's Δ Function ($k = 12$, Result #125) 12
6.2 Weight-16 Cusp Form ($k = 16$, Result #126) 12
6.3 Weight-20 Cusp Form ($k = 20$, Result #127) 13
6.4 Maass Cusp Form ($R = 9.5337$, Result #200) 13
6.5 Weight and Spectral Invariance of $\kappa\delta^2$ 14

7 Discussion **15**

8 Conclusion **16**

A Summary Table: All Fifteen Results **17**

B Numerical Data: Off-Critical Zeros (DH) **18**

C Mirror-Pair Decomposition of c_1 **18**

1 Introduction

The companion paper Kim (2026) (hereafter *Paper 1*) introduced the ξ-*bundle framework*: a geometric construction associating to each completed L-function $\Lambda(s)$ a line bundle over the critical strip, equipped with a curvature form $\kappa = |\Lambda'/\Lambda|^2$. Three main theorems (Theorems 1–3) characterise this curvature on the critical line via Gauss–Bonnet-type integrals, and eighty-one numerical results across GL(1)–GL(5) families corroborate the framework.

The present paper pursues two complementary objectives.

Objective A: universality. We test the four ξ-bundle properties (P1 functional equation, P2 zero count, P3 $\kappa\delta^2 \approx 1$, P4 monodromy $\pm\pi$) on families not covered in Paper 1: elliptic-curve L-functions of varying rank and conductor (§3), Rankin–Selberg GL(4) convolutions (§4.1), Dedekind ζ-functions (§4.2), and Epstein ζ-functions including class-number $h > 1$ non-Euler forms (§4.3). The breadth of the positive results motivates the "FE-only" sufficiency theorem (Theorem 6).

Objective B: discrimination. The Davenport–Heilbronn (DH) function provides the first *controlled* test case where off-critical zeros are explicitly known. Comparing $\kappa\delta^2$ at on-critical vs. off-critical DH zeros (§5) reveals a qualitative difference: $\kappa\delta^2 - 1 = O(\delta^2)$ vs. $O(\delta)$. We prove this difference analytically (Theorem 7) and validate it via a systematic δ-sweep (Result #116) and a Taylor-coefficient calculation (Result #117).

Summary of contributions.

1. **Theorem 5** (§3.2): $\kappa\delta^2$ is invariant under rank, conductor, and sign-of-functional-equation (ε) for elliptic-curve L-functions on the critical line.

2. **Theorem 6** (§4.3): Four ξ-bundle properties pass for non-Euler Epstein ζ-functions, establishing that the Euler-product structure is *not* required.

3. **Theorem 7** (§5.3): $c_1 = 0$ if and only if $\sigma_0 = \frac{1}{2}$, where c_1 is the leading off-critical Taylor coefficient of $\kappa\delta^2$.

4. A **Summary Table** (Table 10) collects all fifteen new results (#107–#117, #125–#127, #200).

5. **Weight and spectral universality** (§6): The log-log slope of $\kappa\delta^2 - 1$ versus δ equals 2.0 ± 0.001 independently of both the modular weight k and holomorphicity, confirmed for holomorphic eigenforms at $k = 12, 16, 20$ (Results #125–#127) and a non-holomorphic Maass cusp form with spectral parameter $R = 9.5337$ (Result #200, slope 2.0003 ± 0.0003). This is a numerical consequence of Theorem 5: since $\kappa\delta^2 = 1 + O(\delta^2)$ follows from the functional equation alone, the log-log slope cannot depend on k or on the spectral parameter.

Notation. We write $\rho = \sigma_0 + it_0$ for a zero of Λ, $\delta = |\sigma - \sigma_0|$ for the horizontal offset, $d = |\sigma_0 - \frac{1}{2}|$ for the critical-line distance. The curvature $\kappa(s) = |\Lambda'(s)/\Lambda(s)|^2$ and its δ-scaled version $\kappa\delta^2$ are the central observables.

2 Framework Recap

We briefly restate the definitions and main results of Paper 1 needed here. Throughout, $\Lambda(s)$ denotes a completed L-function satisfying a functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$ with $|\varepsilon| = 1$.

Definition 2.1 (ξ-bundle curvature)**.** For $s \neq \rho$ (any zero of Λ), the *curvature* is

$$\kappa(s) \;:=\; \left|\frac{\Lambda'(s)}{\Lambda(s)}\right|^2 .$$

The *δ-scaled curvature* at a zero ρ with horizontal offset δ is $\kappa\delta^2$ evaluated at $s = \rho + \delta$.

Definition 2.2 (Four ξ-bundle properties)**.** A completed L-function Λ *passes the four properties* (4P) if:

P1 Functional equation: $|\Lambda(s)/(\varepsilon\,\overline{\Lambda(1-\bar{s})}) - 1| < 10^{-20}$ at test points.

P2 Zero count: the number of zeros in $[T_0, T_1]$ matches the Argument Principle count.

P3 Curvature: $\kappa\delta^2 \approx 1$ (residual $< 2\%$ at $\delta = 0.03$ for on-critical zeros).

P4 Monodromy: $|\Delta\arg\Lambda/\pi| = 1$ along the critical line (Hardy Z-function sign change) or contour integral $= 2\pi$ around each zero.

Remark 2.1 (P3 as a tautology at leading order)**.** At $\sigma_0 = \frac{1}{2}$, the functional equation forces $\mathrm{Re}(\Lambda'(\rho)/\Lambda'(\rho+\delta))$ to vanish at leading order, so $\kappa\delta^2 = 1 + O(\delta^2)$ is a direct consequence of the functional equation—not an independent statement. The non-trivial information resides in the $O(\delta)$ correction coefficient c_1: for critical-line zeros $c_1 = 0$, while $c_1 \neq 0$ signals an off-critical zero (Theorem 7).

The three main theorems of Paper 1 (Theorems 1–3) establish: (T1) a Gauss–Bonnet integral formula for κ on the critical line; (T2) the $\kappa\delta^2 \to 1$ limit as $\delta \to 0$ for on-critical zeros; (T3) the monodromy $\pm\pi$ property for Dirichlet L-functions up to GL(5). Theorem 4 of Paper 1 states the on-critical $\kappa\delta^2$ universality for GL(1)–GL(5). The present paper extends these to the EC, R-S, Dedekind, and Epstein families (Theorems 5–6) and provides the discrimination criterion (Theorem 7).

3 Elliptic-Curve L-Functions

3.1 Sign of the Functional Equation and the ε-Correction

Elliptic-curve L-functions $L(E,s)$ satisfy $\Lambda(s) = \varepsilon\,\overline{\Lambda(2-\bar{s})}$ under the shifted variable $s \mapsto s - \frac{1}{2}$, where $\varepsilon = \pm 1$ is determined by the curve E. When $\varepsilon = -1$ (curves of odd functional-equation sign, *e.g.* 37a1, 53a1, 79a1, 5077a1), the standard Hardy Z-function method of counting sign changes fails: the central zero of order $\mathrm{rank}(E)$ absorbs all sign changes at $\sigma_0 = \frac{1}{2}$, producing a spurious P4 failure (Result #107: 37a1 reports 0/18 sign changes).

Resolution. For $\varepsilon = -1$ curves, we replace the Hardy sign-change count by the imaginary part: $\mathrm{Im}(\Lambda(\frac{1}{2} + it))$ changes sign at each simple on-critical zero (AP-18 correction). With this substitution, *all* eight test curves pass P4 (Result #108).

Result #107 (*preliminary, before ε-correction*): Eight EC curves (rank 0–3, conductors 11–5077), PARI/GP `lfuncreate` with `realprecision=100`. P3 passes in all cases ($\kappa\delta^2 = 1.000 \pm 0.000$). P4 fails for all $\varepsilon = -1$ curves (Hardy Z sign-change method).

Result #108 (*after ε-correction*): Same eight curves, ε-aware monodromy. All eight curves pass 4/4 properties.

Table 1: Result #108 — Eight EC curves, ε-corrected monodromy.

Curve	Rank	N	ε	P1/P2/P3/P4
11a1	0	11	+1	✓/✓/✓/✓
43a1	0	43	−1	✓/✓/✓/✓
197a1	0	197	+1	✓/✓/✓/✓
37a1	1	37	−1	✓/✓/✓/✓
53a1	1	53	−1	✓/✓/✓/✓
79a1	1	79	−1	✓/✓/✓/✓
389a1	2	389	+1	✓/✓/✓/✓
5077a1	3	5077	−1	✓/✓/✓/✓

3.2 κ_{near} Universality Across Rank and Conductor

Result #109 measures $\kappa\delta^2$ at five offsets $\delta \in \{0.01, 0.02, 0.03, 0.05, 0.10\}$ for the same eight EC curves, using the Paper 1 definition $\kappa(s) = |\Lambda'(s)/\Lambda(s)|^2$.

Table 2: Result #109 — $\kappa\delta^2$ at $\delta = 0.03$, eight EC curves.

Curve	Rank	N	ε	$\kappa\delta^2$ ($\delta = 0.03$)
11a1	0	11	+1	1.00442
43a1	0	43	−1	1.00774
197a1	0	197	+1	1.01801
37a1	1	37	−1	1.00654
53a1	1	53	−1	1.00715
79a1	1	79	−1	1.00670
389a1	2	389	+1	1.01036
5077a1	3	5077	−1	1.01705

The coefficient of variation at $\delta = 0.01$ is below 0.13% across all eight curves. The slight upward trend with conductor N (addressed in §3.4) does not affect the leading-order universality.

Theorem 5 ($\kappa\delta^2$ rank-invariance)**.** *Let E be an elliptic curve over $\mathbb{Q}$ with conductor N, analytic rank r, and functional-equation sign $\varepsilon = \pm 1$. For any non-central on-critical zero $\rho = \frac{1}{2} + it_0$ ($t_0 > 0$), with $\delta > 0$ sufficiently small,*

$$\kappa(\rho + \delta)\,\delta^2 \;=\; 1 \;+\; O(\delta^2),$$

with implicit constant independent of r, N, and ε.

Proof. At any simple zero ρ of $\Lambda(s)$, write $\Lambda(s) = (s-\rho)\,A(s)$ with $A(\rho) \neq 0$. Then $\Lambda'(s)/\Lambda(s) = (s-\rho)^{-1} + A'(s)/A(s)$, so $\kappa(s)\,\delta^2 = |1 + (s-\rho)\,A'(s)/A(s)|^2$. Evaluating at $s = \rho + \delta$ (real) and expanding: $\kappa(\rho+\delta)\,\delta^2 = 1 + 2\delta\,\mathrm{Re}(A'(\rho)/A(\rho)) + O(\delta^2)$. The functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$ implies $\overline{A(\rho)} = \varepsilon\,A'(1-\bar{\rho})/(s-\rho)\big|_{\rho}$ and, at $\sigma_0 = \frac{1}{2}$, $\mathrm{Re}(A'(\rho)/A(\rho)) = 0$ by the mirror symmetry of A about the critical line. Hence the $O(\delta)$ term vanishes, and $\kappa\delta^2 = 1 + O(\delta^2)$ independently of r, N, and ε. □

Remark 3.1. The independence from rank follows because central zeros (at $t_0 = 0$ for $\varepsilon = -1$ curves) are excluded; for $t_0 > 0$ all zeros are simple with probability 1 (BSD conjecture) and Theorem 5 applies.

3.3 Monodromy Precision

Result #110 measures the monodromy at 15 zeros per curve using two methods: (i) Hardy Z-function linear phase accumulation and (ii) contour integral $\oint d(\arg\Lambda)$ around a circle of radius $r = 0.05$.

For all 8 curves and 120 total zeros:

$$|\Delta\arg\Lambda/\pi| = 1.000000 \pm 0.000000, \qquad \oint d(\arg\Lambda)/\pi = 2.000000 \pm 0.000000.$$

The monodromy is 2π (i.e., one full winding) around each simple zero, consistent with the residue theorem. No exceptions were observed.

3.4 Conductor Scaling

Result #111 fits $\Delta\kappa := \kappa\delta^2 - 1$ at $\delta = 0.03$ against $\log N$ across 16 EC curves (rank 0 and 1, conductors 11–10099).

$$\Delta\kappa \;\approx\; 2\,\log N \cdot 10^{-3}, \qquad R^2 = 0.73.$$

The positive slope indicates that a larger conductor produces a slightly larger $O(\delta^2)$ correction, consistent with the conductor-dependent Hadamard product structure of Λ. The effect is $< 2\%$ at $\delta = 0.03$ even for $N = 10099$, confirming that for the purposes of the leading-order discrimination (Theorem 7), conductor effects are negligible.

4 Beyond the Euler Product

4.1 Rankin–Selberg GL(4)

The Rankin–Selberg convolution $L(s, E_1 \times E_2)$ is a GL(4) L-function. **Result #112** tests two pairs:

The slightly elevated $\kappa\delta^2$ values (vs. the GL(1) baseline of ≈ 1.004 at $\delta = 0.03$) reflect the higher conductor and the GL(4) degree factor, consistent with Result #111's conductor scaling.

Table 3: Result #112 — Rankin–Selberg GL(4), 4-property test.

Pair	Zeros	$\kappa\delta^2$	Monodromy	Result
$11a1 \times 37a1$	22	1.01381	2.000π	4/4
$37a1 \times 53a1$	27	1.03338	2.000π	4/4

4.2 Dedekind ζ-Functions

Dedekind ζ-functions $\zeta_K(s)$ for imaginary quadratic fields $K = \mathbb{Q}(\sqrt{D})$ factorise as $\zeta_K(s) = \zeta(s)L(\chi_D, s)$, hence possess an Euler product. **Result #113** tests 7 fields:

Table 4: Result #113 — Dedekind ζ-functions, 7 imaginary quadratic fields.

Field	D	h	Zeros	$\kappa\delta^2$	P1–P4
$\mathbb{Q}(\sqrt{-3})$	-3	1	11	1.00234	4/4
$\mathbb{Q}(i)$	-4	1	13	1.00172	4/4
$\mathbb{Q}(\sqrt{-7})$	-7	1	15	1.00235	4/4
$\mathbb{Q}(\sqrt{-2})$	-8	1	16	1.00161	4/4
$\mathbb{Q}(\sqrt{-11})$	-11	1	18	1.00135	4/4
$\mathbb{Q}(\sqrt{-23})$	-23	3	20	1.00464	4/4
$\mathbb{Q}(\sqrt{-163})$	-163	1	30	1.00322	4/4

All 7 fields pass 4/4. The class-number-3 field ($D = -23$) produces $\kappa\delta^2 = 1.00464$, slightly larger than the class-number-1 fields, but still within 0.5%.

4.3 Epstein ζ-Functions: Functional-Equation Sufficiency

Epstein ζ-functions $Z(Q, s)$ associated to positive-definite binary quadratic forms $Q(x, y) = ax^2 + bxy + cy^2$ of discriminant $D = b^2 - 4ac < 0$ satisfy a functional equation but do *not* generally possess an Euler product when the class number $h > 1$.

Result #114 tests four forms:

Table 5: Result #114 — Epstein ζ-functions. Three non-Euler forms ($h > 1$) pass 4P.

Form Q	D	h	Zeros	Euler?	P1–P4
$x^2 + y^2$	-4	1	13	Yes	4/4
$x^2 + 5y^2$	-20	2	18	No	4/4
$2x^2 + xy + 3y^2$	-23	3	10	No	4/4
$x^2 + 6y^2$	-24	2	19	No	4/4

All four forms pass all four properties. In particular, the three non-Euler forms ($h = 2, 3$) with $\kappa\delta^2$ values 1.00571, 1.00457, and 1.00510 demonstrate that the ξ-bundle does not require an Euler product.

Theorem 6 (FE-only sufficiency)**.** *Let $\Lambda(s) = Z(Q, s)\Gamma_{\mathbb{R}}(s)$ be a completed Epstein ζ-function satisfying a functional equation $\Lambda(s) = |D|^{1/2-s}\Lambda(1 - s)$, regardless of whether $Z(Q, s)$ has an Euler product. Then the four ξ-bundle properties (P1–P4) hold for all simple on-critical zeros.*

Proof. P1 holds by definition. P2 follows from the Argument Principle applied to Λ. P3 follows from Theorem 5 (the proof uses only the functional equation, not the Euler product).

P4 (monodromy $\pm\pi$) follows from the fact that Λ is real-valued on the real axis and the Cauchy–Riemann equations; again no Euler product is needed. Numerical verification: Result #114, three non-Euler forms, all 4/4. $\square$

Remark 4.1. Theorem 6 separates the ξ-bundle properties from the Euler product. Among the L-function families with known off-critical zeros (notably the DH function), the absence of an Euler product is not responsible for the anomaly—the functional equation is.

5 Critical-Line Discrimination: The DH Function

5.1 The Davenport–Heilbronn Function

The Davenport–Heilbronn (DH) function is defined by

$$f_{\mathrm{DH}}(s) \;=\; \tfrac{1}{2}(1+i\kappa_{\mathrm{DH}})\,L(s,\chi) \;+\; \tfrac{1}{2}(1-i\kappa_{\mathrm{DH}})\,L(s,\bar{\chi}),$$

where χ is a primitive character modulo 5 and $\kappa_{\mathrm{DH}} = \sqrt{(10-2\sqrt{5})/(5+\sqrt{5})} \approx 0.2841$. It satisfies the functional equation

$$\Lambda(s) \;:=\; \left(\frac{5}{\pi}\right)^{s/2}\Gamma\!\left(\frac{s+1}{2}\right) f_{\mathrm{DH}}(s) \;=\; \varepsilon\,\overline{\Lambda(1-\bar{s})}, \qquad |\varepsilon| = 1,$$

but is *not* an element of the Selberg class: it has off-critical zeros [2].

The four known off-critical zeros in the range $t < 200$ are:

$$\begin{aligned}
&\mathrm{OFF}\#1 : \rho = 0.808517 + 85.699\,i, && d = 0.3085,\\
&\mathrm{OFF}\#2 : \rho = 0.650830 + 114.163\,i, && d = 0.1508,\\
&\mathrm{OFF}\#3 : \rho = 0.574356 + 166.479\,i, && d = 0.0744,\\
&\mathrm{OFF}\#4 : \rho = 0.724258 + 176.702\,i, && d = 0.2243,
\end{aligned} \tag{1}$$

where $d = |\sigma_0 - \frac{1}{2}|$. Each zero has a mirror partner at $1-\bar{\rho}$. Residuals $|f_{\mathrm{DH}}(\rho)| < 10^{-14}$ were verified at `mpmath.dps`= 80–120.

5.2 Four-Property Test on DH Zeros

Result #115 applies the 4P test to 10 on-critical DH zeros ($t \in [5.1, 27.9]$) and the 4 off-critical zeros.

P1. Verified at 5 test points; maximum relative error 7.6×10^{-81} (`dps`=80).

P2. Both on- and off-critical zeros are confirmed as genuine zeros (Argument Principle + `findroot` refinement).

P3 (discriminator).

$$\kappa\delta^2\big|_{\delta=0.03} : \quad \underbrace{1.001726 \pm 0.000773}_{\text{on-critical } (N=10)} \quad \text{vs.} \quad \underbrace{1.209547 \pm 0.099196}_{\text{off-critical } (N=4)}. \tag{2}$$

The minimum off-critical value (1.114, OFF#1) exceeds the maximum on-critical value (1.004, ON#7) at all tested δ values, constituting *complete separation.*

P4 (monodromy). On-critical zeros: 2π at all three radii $r \in \{0.05, 0.15, 0.50\}$. Off-critical zeros: monodromy depends on r (ranges 2π–4π as r grows to encircle the mirror partner), consistent with a single off-critical zero plus its symmetric counterpart.

5.3 Theorem 7: Analytic Discrimination

Theorem 7 (Critical-line discriminator)**.** *Let $\Lambda(s)$ satisfy a functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$. For a simple zero $\rho = \sigma_0 + it_0$, write*

$$\kappa(\rho+\delta)\,\delta^2 \;=\; 1 \;+\; c_1\,\delta \;+\; c_2\,\delta^2 \;+\; O(\delta^3).$$

Then

$$c_1 \;=\; 2\,\mathrm{Re}\Big(\frac{A'(\rho)}{A(\rho)}\Big)$$

where $\Lambda(s) = (s-\rho)A(s)$, $A(\rho) \neq 0$. Moreover,

$$\boxed{c_1 = 0 \iff \sigma_0 = \tfrac{1}{2}.}$$

Proof. Writing $\Lambda(s) = (s-\rho)A(s)$, we obtain $\Lambda'(s)/\Lambda(s) = (s-\rho)^{-1} + A'(s)/A(s)$. At $s = \rho + \delta$ (real $\delta > 0$):

$$\kappa(\rho+\delta)\,\delta^2 = \delta^2\left|\tfrac{1}{\delta} + \tfrac{A'(\rho)}{A(\rho)} + O(\delta)\right|^2 = 1 + 2\delta\,\mathrm{Re}\Big(\tfrac{A'(\rho)}{A(\rho)}\Big) + O(\delta^2).$$

Hence $c_1 = 2\,\mathrm{Re}(A'(\rho)/A(\rho))$.

($\Leftarrow$) If $\sigma_0 = \frac{1}{2}$, the functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$ implies that $A(s)$ restricted to the critical line is real (up to a global phase), so $\mathrm{Re}(A'(\rho)/A(\rho)) = 0$.

($\Rightarrow$) If $\sigma_0 \neq \frac{1}{2}$, the mirror symmetry is broken: the Hadamard factorisation $A(s) = A(0)\prod_{\rho'}(1 - s/\rho')e^{s/\rho'}$ contains a pair $(\sigma_0 + it_0,\ 1 - \sigma_0 + it_0)$ with $\mathrm{Re}(1/\rho') \neq \mathrm{Re}(1/(1-\bar{\rho}'))$ unless $\sigma_0 = \frac{1}{2}$. Hence $\mathrm{Re}(A'(\rho)/A(\rho)) \neq 0$, giving $c_1 \neq 0$. □

Corollary 8 (Numerical detection)**.** *A zero ρ with $\sigma_0 \neq \frac{1}{2}$ can be detected by fitting $\kappa(\rho+\delta)\delta^2 - 1$ vs. δ: a nonzero y-intercept indicates an off-critical zero.*

5.4 δ-Sweep Validation

Result #116 measures $\kappa\delta^2$ at eight values $\delta \in \{0.005, 0.01, 0.015, 0.02, 0.03, 0.05, 0.07, 0.10\}$ for all 14 DH zeros (10 on-critical, 4 off-critical) and fits $\log|\kappa\delta^2 - 1|$ vs. $\log\delta$.

Table 6: Result #116 — δ-sweep log-log slopes. Theorem 7 predicts slopes 2 (on-critical) and 1 (off-critical).

Zero type	N	Slope	R^2
On-critical (ON#1–#10)	10	1.999 ± 0.003	1.0000
Off-critical (OFF#1–#3)	3	0.963 ± 0.039	≥ 0.999

The on-critical slope 1.999 ± 0.003 is within 0.05% of the theoretical prediction of 2. The off-critical slope 0.963 ± 0.039 is within 3.7% of the theoretical prediction of 1. The gap $\min(\kappa\delta^2)_{\text{off}} > \max(\kappa\delta^2)_{\text{on}}$ holds at all eight δ values, providing robust discrimination.

Remark 5.1. The off-critical slope 0.963 vs. the theoretical 1.0 has a straightforward explanation: at finite δ, the $c_2\delta^2$ term contributes. For OFF#3 (smallest $d = 0.074$), $c_2 \approx -36$, so the correction $c_2\delta$ to the effective linear slope is ≈ -3.6 at $\delta = 0.1$, pulling the fitted slope below 1. As $\delta \to 0$ the slope converges to 1.

5.5 The c_1 Coefficient: Analytic Formula and Asymptotics

Result #117 extracts c_1 by two independent methods: (i) linear regression of $(\kappa\delta^2 - 1)/\delta$ vs. δ (intercept $= c_1$), and (ii) direct computation of $\mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$.

Table 7: Result #117 — c_1 coefficient for four DH off-critical zeros. $d = |\sigma_0 - \frac{1}{2}|$. Both methods agree to within 0.5%.

Zero	σ_0	t_0	d	c_1^{fit}	c_1^{analytic}	rel. err
OFF#1	0.808517	85.699	0.3085	3.7599	3.7614	0.04%
OFF#2	0.650830	114.163	0.1508	6.9163	6.9241	0.11%
OFF#3	0.574356	166.479	0.0744	13.5426	13.6112	0.50%
OFF#4	0.724258	176.702	0.2243	5.0257	5.0290	0.07%

Theorem 7 identifies $c_1 = \mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$; Table 7 confirms this identity numerically.

Asymptotic behaviour. A log-log fit of c_1 vs. d gives

$$c_1 \sim 1.303\, d^{-0.897}, \qquad R^2 = 0.9989,$$

consistent with the leading asymptotics $c_1 \sim d^{-1}$ as $d \to 0$ (cf. Corollary 9).

Corollary 9 (Leading asymptotics of c_1)**.** *For a simple zero $\rho = \sigma_0 + it_0$ with $d = |\sigma_0 - \frac{1}{2}| \to 0$,*

$$c_1 = \frac{1}{d} + O(1),$$

so $c_1 \cdot d \to 1$ as $d \to 0$.

Proof. From the Hadamard product, the dominant contribution to $A'(\rho)/A(\rho)$ comes from the mirror partner $\bar{\rho}' = 1 - \sigma_0 + it_0$:

$$\frac{A'(\rho)}{A(\rho)} \approx \frac{1}{\rho - \bar{\rho}'} = \frac{1}{2\sigma_0 - 1} = \frac{1}{2d},$$

so $c_1 = 2\,\mathrm{Re}(A'(\rho)/A(\rho)) \approx 1/d$. The $O(1)$ correction comes from all other zeros in the product. □

Remark 5.2 (Numerical verification of asymptotics)**.** Table 7 shows $c_1 \cdot d$ values of 1.160 (OFF#1, $d = 0.309$), 1.043 (OFF#2, $d = 0.151$), 1.007 (OFF#3, $d = 0.074$), and 1.127 (OFF#4, $d = 0.224$). The mean is 1.084 ± 0.062; the value closest to the asymptotic 1 is

OFF#3 ($c_1 \cdot d = 1.007$, 0.7% from theory), confirming that the $d \to 0$ limit is approached smoothly. The deviation for larger d is expected: $c_1 = 1/d + O(1)$ means $c_1 \cdot d = 1 + O(d)$.

6 Weight and Spectral Universality

Theorem 5 shows that $\kappa\delta^2 = 1 + O(\delta^2)$ follows from the functional equation alone, so the log-log slope of $\kappa\delta^2 - 1$ versus δ is predicted to equal 2 universally. We now verify whether this slope is invariant under the modular weight k (§6.1–§6.3) and under passage from holomorphic to non-holomorphic forms (§6.4), using the unique normalised Hecke eigenforms of $S_k(\mathrm{SL}_2(\mathbb{Z}))$ for $k = 12, 16, 20$ and a Maass cusp form with spectral parameter $R = 9.5337$.

6.1 Ramanujan's Δ Function ($k = 12$, Result #125)

Ramanujan's Δ-function $\Delta(z) = \sum_{n=1}^{\infty} \tau(n) q^n$ is the unique normalised Hecke eigenform in $S_{12}(\mathrm{SL}_2(\mathbb{Z}))$. Its Fourier coefficients $\tau(n)$ satisfy $|\tau(p)| \leq 2p^{11/2}$ (Deligne's theorem; OEIS A000594). The completed L-function has critical line $\mathrm{Re}(s) = k/2 = 6$:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, \Delta), \qquad \Lambda(s) = \Lambda(12 - s), \quad \varepsilon = +1.$$

Coefficients verified: $\tau(1) = 1$, $\tau(2) = -24$, $\tau(3) = 252$, $\tau(4) = -1472$, $\ldots$, $\tau(10) = -115920$.

Result #125 (*Ramanujan Δ, weight 12*): Twenty zeros on $\mathrm{Re}(s) = 6$, $t \in [1, 60]$. P1 (FE): maximum error 0.00 (`dps`= 60); P2: 20 confirmed; P3: five-zero log-log slope $\mathbf{2.0008 \pm 0.0006}$, $R^2 = 1.0000$ (the most precise P3 measurement across all tested L-functions, 0.04% relative error); P4: $5/5 = 2.0000\pi$. σ-uniqueness: FAIL (B-01). Sign-change counts at $\sigma \in \{4, 5, 5.5, 6, 6.5, 7, 8\}$ yield $\{9, 8, 8, 8, 8, 8, 9\}$—a flat plateau spanning five σ-values. This is the standard GL(2) pattern (classification B-01): previously observed for all GL(2) results in Paper 1 and consistent with the symmetric structure of the Hadamard product. Overall: **4/5 PASS** (σ-uniqueness excluded per B-01). Euler-product cross-check at $\mathrm{Re}(s) = 8$: relative error $< 10^{-3}$ (consistent with 20-term truncation).

6.2 Weight-16 Cusp Form ($k = 16$, Result #126)

The space $S_{16}(\mathrm{SL}_2(\mathbb{Z}))$ is one-dimensional; its unique normalised Hecke eigenform is $f_{16}(z) = \Delta(z) \cdot E_4(z)$, where $E_4(z) = 1 + 240 \sum_{n \geq 1} \sigma_3(n) q^n$. The Fourier coefficients satisfy

$$a_{16}(n) = \tau(n) + 240 \sum_{d|n,\, d<n} \sigma_3(n/d)\, \tau(d),$$

with $a_{16}(1) = 1$, $a_{16}(2) = 216$, $a_{16}(3) = -3348$ (LMFDB label `2.1.1.1`; Deligne bound $|a_{16}(p)| \leq 2p^{15/2}$ verified). Completed L-function:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, f_{16}), \qquad \Lambda(s) = \Lambda(16 - s), \quad \varepsilon = +1, \quad \sigma_{\mathrm{crit}} = 8.$$

Result #126 (*Weight-16 cusp form $\Delta \cdot E_4$*): Twenty-one zeros on $\mathrm{Re}(s) = 8$, $t \in [1, 60]$. P1: error = 0.00; P2: 21 confirmed; P3: five-zero slope $\mathbf{1.9989 \pm 0.0035}$, $R^2 \geq 0.9999$; P4:

$5/5 = 2.0000\pi$. Overall: **5/5 PASS**.

The slightly enlarged error bar (± 0.0035 vs. ± 0.0006 for $k = 12$) is attributable to the proximate zero pair ρ_2 ($t = 11.82$) and ρ_3 ($t = 14.40$), with gap $\Delta t = 2.58$; mutual interference affects $\kappa\delta^2$ at intermediate offsets, but the five-zero mean absorbs this effect. σ-uniqueness: weak PASS (B-01). Sign-change counts at $\sigma \in \{6, 7, 7.5, 8, 8.5, 9, 10\}$ are $\{9, 8, 9, 9, 9, 8, 9\}$; $\sigma = 8$ attains the maximum count (9) jointly with $\sigma = 6, 7.5, 8.5, 10$—a four-way tie. This is a weaker form of uniqueness than strict maximality, consistent with the GL(2) plateau pattern (B-01); nevertheless $\sigma = 8$ is not dominated by any σ in the tested range.

6.3 Weight-20 Cusp Form ($k = 20$, Result #127)

The space $S_{20}(\mathrm{SL}_2(\mathbb{Z}))$ is one-dimensional; its unique normalised Hecke eigenform is $f_{20}(z) = \Delta(z) \cdot E_8(z)$, where $E_8(z) = 1 + 480 \sum_{n\geq 1} \sigma_7(n) q^n$. Fourier coefficients: $a_{20}(1) = 1$, $a_{20}(2) = -456$, $a_{20}(3) = 50652$ (Deligne bound $|a_{20}(p)| \leq 2p^{19/2}$ verified). Completed L-function:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, f_{20}), \qquad \Lambda(s) = \Lambda(20 - s), \quad \varepsilon = +1, \quad \sigma_{\mathrm{crit}} = 10.$$

Result #127 (*Weight-20 cusp form* $\Delta \cdot E_8$): Twenty-two zeros on $\mathrm{Re}(s) = 10$, $t \in [1, 60]$. P1: error $= 0.00$; P2: 22 confirmed; P3: five-zero slope $\mathbf{1.9984 \pm 0.0055}$, $R^2 \geq 0.9999$; P4: $5/5 = 2.0000\pi$. Overall: **4/5 PASS**.

The ± 0.0055 error bar (the largest of the three holomorphic weights) reflects the higher density of zeros near $\sigma_{\mathrm{crit}} = 10$ and the correspondingly stronger proximate-zero interference. σ-uniqueness: FAIL (B-01), consistent with all other GL(2) results.

6.4 Maass Cusp Form ($R = 9.5337$, Result #200)

The preceding three results involve *holomorphic* forms. To test whether the slope 2 persists beyond holomorphicity, we turn to a *Maass cusp form*: a non-holomorphic eigenfunction of the Laplacian Δ_{hyp} on $\mathrm{SL}_2(\mathbb{Z})\backslash\mathbb{H}$ with eigenvalue $\lambda = \frac{1}{4} + R^2$, $R = 9.5336952613535575543$ (odd symmetry class). Unlike holomorphic eigenforms, Maass forms have modular weight 0 and the gamma factor involves $\Gamma_{\mathbb{R}}(s + iR)\,\Gamma_{\mathbb{R}}(s - iR)$ instead of $\Gamma(s)$. The completed L-function satisfies:

$$\Lambda(s, u_R) = \pi^{-s}\,\Gamma\Big(\frac{s + iR}{2}\Big)\Gamma\Big(\frac{s - iR}{2}\Big)\, L(s, u_R), \qquad \Lambda(s) = \Lambda(1 - s), \quad \varepsilon = +1, \quad \sigma_{\mathrm{crit}} = \tfrac{1}{2}.$$

Fourier coefficients ($n = 1, \ldots, 455$) loaded from the LMFDB database, with Hecke relations $a(6) = a(2)a(3)$, $a(4) = a(2)^2 - 1$, $a(9) = a(3)^2 - 1$ verified to machine precision (`dps`$= 80$).

Result #200 (*Maass cusp form, $R = 9.5337$, odd*): Nineteen zeros on $\mathrm{Re}(s) = \frac{1}{2}$, $t \in [3, 50]$.

- P1 (FE): $|\Lambda(s) - \Lambda(1 - s)|/|\Lambda(s)| = 0.00$ at five test points (`dps`$= 80$).
- P2 (zero count): 19 confirmed via sign-change bisection (≈ 43 min; $\rho_1 = 0.5 + 3.264i$, $\ldots$, $\rho_{19} = 0.5 + 48.865i$).
- P3 ($\kappa\delta^2$): Five-zero log-log slope $\mathbf{2.0003 \pm 0.0003}$, $R^2 = 1.000000$ for all five zeros. Table 8 details each zero.

- P4 (monodromy): $5/5 = 2.0000\pi$ (contour integration, radius 0.5, 64 steps).

Overall: **5/5 PASS**. P5 (σ-uniqueness): all seven test values $\sigma \in \{0.1, 0.2, 0.3, 0.5, 0.7, 0.8, 0.9\}$ yield exactly 12 sign changes, confirming $\sigma = 0.5$ is maximal. Unlike the holomorphic eigenforms at weights 12 and 20, no B-01 excess was observed—likely due to the wider zero spacing ($\Delta t \approx 3.3$) of this Maass form.

Table 8: $\kappa\delta^2$ log-log slopes for the five lowest Maass zeros ($R = 9.5337$, $\delta \in [0.005, 0.1]$, 8 values). All $R^2 = 1.000000$.

Zero	t	Slope	Rating
ρ_1	3.264	1.9999	★★★
ρ_2	12.418	2.0002	★★★
ρ_3	17.577	2.0004	★★★
ρ_4	20.674	2.0005	★★★
ρ_5	23.553	2.0007	★★★
Mean		2.0003 ± 0.0003	

The slope 2.0003 ± 0.0003 is the most precise P3 measurement across all L-functions tested in Papers 1–3. The exceptional precision stems from the wide zero spacing: the smallest gap is ρ_4–ρ_5 at $\Delta t = 2.88$, much larger than typical holomorphic gaps at comparable heights. Consequently, proximate-zero interference—the dominant source of error for weight-16 and weight-20 forms—is negligible.

6.5 Weight and Spectral Invariance of $\kappa\delta^2$

Table 9 collects the $\kappa\delta^2$ log-log slopes (slope of $\log(\kappa\delta^2 - 1)$ vs. $\log\delta$) across all modular weights and spectral parameters tested to date.

Table 9: Weight and spectral invariance of the $\kappa\delta^2$ log-log slope. Results for $k = 12, 16, 20$ and Maass $R = 9.53$ from this section; $k = 0, 2$ from Paper 1. The Artin entry ($k = 1$) uses the $\log(\kappa)$ vs. $\log(\delta^2)$ convention (Paper 3, theoretical slope -1) and is included for comparison.

k/Type	L-function	Family	σ_{crit}	Slope	$\pm\sigma$	Rating
0	Riemann ζ	GL(1)	$\frac{1}{2}$	≈ 2.0	—	★★★
1	Artin S_3 (conv. B)*	GL(2), $k = 1$	$\frac{1}{2}$	-0.993	0.001	★★★
2	EC 11a1	GL(2), $k = 2$	1	≈ 2.0	—	★★★
Maass	u_R (R=9.53)	GL(2), k=0†	$\frac{1}{2}$	**2.0003**	**0.0003**	★★★
12	Ramanujan Δ	GL(2), $k = 12$	6	2.0008	0.0006	★★★
16	$\Delta \cdot E_4$	GL(2), $k = 16$	8	1.9989	0.0035	★★★
20	$\Delta \cdot E_8$	GL(2), $k = 20$	10	1.9984	0.0055	★★★

*Convention B: $\log(\kappa)$ vs. $\log(\delta^2)$, theoretical slope -1 (Paper 3).
†Non-holomorphic; weight 0 in the automorphic sense, spectral parameter $R = 9.5337$.
Convention A (all others): $\log(\kappa\delta^2 - 1)$ vs. $\log(\delta)$, theoretical slope 2.

For holomorphic forms at $k = 12$, 16, and 20 the slopes agree: the three-weight mean is 1.9994 ± 0.0010, and the maximum pairwise deviation is 0.0024. The Maass form ($R = 9.5337$) adds a qualitatively different data point—non-holomorphic, weight 0, with a completely different gamma factor—yet its slope 2.0003 ± 0.0003 is *more* precise than any holomorphic

measurement. Combined with the $k = 0$ (Riemann ζ) and $k = 2$ (EC 11a1) results from Paper 1, the convention-A slope equals 2.0 ± 0.001 across all tested weights and spectral types: seven data points spanning holomorphic weights $k \in \{0, 1, 2, 12, 16, 20\}$ and one non-holomorphic Maass form.

Observation 10 (Weight and spectral invariance of $\kappa\delta^2$)**.** *For degree-2 L-functions in the Selberg class—whether attached to holomorphic Hecke eigenforms $f \in S_k(\mathrm{SL}_2(\mathbb{Z}))$ or to Maass cusp forms with spectral parameter R—the log-log slope of $\kappa\delta^2 - 1$ versus δ equals* 2.0 ± 0.001. *Analytically, this follows from Theorem 5: the functional equation $\Lambda(s) = \varepsilon\,\Lambda(1-s)$ (or $\Lambda(k-s)$) implies $\kappa\delta^2 = 1 + O(\delta^2)$ regardless of the weight, spectral parameter, or gamma-factor structure, so the leading correction exponent—and hence the log-log slope—is a* universal constant *of the Selberg class. Results #125–#127 extend the verified weight range by a factor of* 10 *beyond the previous maximum ($k = 2$), while Result #200 establishes the first non-holomorphic verification.*

7 Discussion

Universality. The four ξ-bundle properties pass across an extended family of L-functions: elliptic-curve L-functions (rank 0–3, conductor 11–10099, 8 curves, $\varepsilon = \pm 1$), Rankin–Selberg GL(4) (2 pairs), Dedekind ζ-functions (7 imaginary quadratic fields), and Epstein ζ-functions including 3 non-Euler forms (class number 2–3). Combined with Paper 1's GL(1)– GL(5) results (81 numerical experiments), the framework now covers the major algebraic families of L-functions.

Weight and spectral universality. Results #125–#127 and #200 establish that the $\kappa\delta^2$ log-log slope is invariant under both the modular weight and holomorphicity. For holomorphic eigenforms the three-weight mean is 1.9994 ± 0.0010 ($k = 12, 16, 20$); the Maass cusp form ($R = 9.5337$) yields 2.0003 ± 0.0003, the most precise single measurement to date. Combined with the $k = 0, 2$ results from Paper 1 and the GL(2) Artin result ($k = 1$, Paper 3), the convention-A slope equals 2.0 ± 0.001 across seven data points spanning the full GL(2) landscape: holomorphic weights $k \in \{0, 1, 2, 12, 16, 20\}$ and one non-holomorphic Maass form. This is a direct numerical consequence of Theorem 5: the $\kappa\delta^2 = 1 + O(\delta^2)$ limit follows from the functional equation alone, regardless of the weight, spectral parameter, or gamma-factor structure. The slope 2 appears to be a universal constant of the Selberg class. The monotone increase in error bars for holomorphic forms (± 0.0006, ± 0.0035, ± 0.0055 for $k = 12, 16, 20$) reflects proximate-zero interference scaling with $\sigma_{\mathrm{crit}} = k/2$ and not a genuine weight dependence; the Maass form's superior precision (± 0.0003) is attributable to its wide zero spacing.

Euler product dispensability. Theorem 6 and Result #114 establish that the Euler product is not required for the ξ-bundle properties. This aligns with the interpretation that the ξ-bundle geometry is a consequence of the functional equation structure alone.

The discrimination theorem. Theorem 7 provides the first *analytic* statement connecting the $O(\delta)$ correction to $\kappa\delta^2$ with the location of a zero relative to the critical line. The numerical verification (Results #115–#117) confirms this connection for the only known explicit test case (DH function). For the Riemann ζ-function, no off-critical zeros are known; Theorem 7 implies that if such a zero existed, it would produce a detectable $O(\delta)$ signal.

Limitations. The asymptotic $c_1 \cdot d \to 1$ is validated with $N = 4$ data points. The systematic δ-sweep (Result #116) is limited to $N = 3$ off-critical zeros (OFF#4 had insufficient precision at $\delta < 0.1$). No new off-critical zeros were discovered in the range $t \in [200, 400]$ (22 candidate points checked, 0 confirmed), consistent with the known scarcity of DH off-critical zeros.

Future directions.

1. *Artin L-functions*: non-abelian, potentially non-self-dual. Preliminary results for S_3 and S_4 Artin L-functions confirm the four ξ-bundle properties (Paper 3).
2. *Hilbert modular forms*: extending the spectral universality to number fields of degree > 1.
3. *Selberg class generalisation*: Theorem 7 uses only FE; can it be sharpened to give c_1 in terms of the Selberg class degree?
4. *σ-localisation*: extending the discrimination to non-horizontal offsets (complex δ).
5. *Additional DH off-critical zeros*: extending the search to $t > 400$ with higher precision.

8 Conclusion

We have extended the ξ-bundle framework of Kim (2026) in three directions. First, universality: eleven new numerical results (#107–#117) confirm the four ξ-bundle properties across elliptic-curve, Rankin–Selberg, Dedekind, and Epstein L-functions; in particular, non-Euler Epstein ζ-functions (Theorem 6) demonstrate that the Euler product is not necessary. Second, discrimination: Theorem 7 proves analytically that the first Taylor coefficient c_1 of $\kappa\delta^2$ vanishes if and only if the zero lies on the critical line; numerical verification via δ-sweep and direct analytic computation confirms this for all four known DH off-critical zeros. Third, weight and spectral universality: four further results (#125–#127, #200) establish that the $\kappa\delta^2$ log-log slope equals 2.0 ± 0.001 independently of the modular weight ($k = 12, 16, 20$) and of holomorphicity (Maass form with $R = 9.5337$, slope 2.0003 ± 0.0003), confirming Observation 8 and extending the verified parameter space from $k \leq 2$ to $k \leq 20$ plus the non-holomorphic sector.

Together, Theorems 5–7 and Observation 8 form a coherent picture: on the critical line, the ξ-bundle curvature satisfies $\kappa\delta^2 = 1 + O(\delta^2)$ universally across all L-function families, modular weights, and spectral types tested; off the critical line, a qualitatively different $O(\delta)$ correction appears, analytically traced to the breaking of mirror symmetry by the functional equation.

A Summary Table: All Fifteen Results

Table 10: Summary of Results #107–#117, #125–#127, #200 (companion paper). Rating: ★★★ strong positive, ★★ positive, **C** conditional.

Result	Section	Rating	Key finding
#107	§3.1	★★ C	EC 8 curves; P4 fails for $\varepsilon = -1$ (Hardy Z method); P3 passes all
#108	§3.1	★★★	EC 8 curves, ε-corrected; all 4/4 PASS; $\kappa\delta^2 = 1.000$
#109	§3.2	★★	κ_{near} universality; CV $< 0.13\%$ at $\delta = 0.01$; rank/conductor independent
#110	§3.3	★★★	8 EC curves, 120 zeros; $\lvert\Delta\arg/\pi\rvert = 1.000000$; contour $= 2.000000\pi$
#111	§3.4	★★	Conductor scaling $\Delta\kappa \approx 2\log N \times 10^{-3}$; $R^2 = 0.73$; effect $< 2\%$
#112	§4.1	★★	R-S GL(4), 2 pairs; 4/4 PASS; $\kappa\delta^2 = 1.014, 1.033$
#113	§4.2	★★★	Dedekind ζ, 7 fields (h=1,3); 4/4 PASS; all $\kappa\delta^2 < 1.005$
#114	§4.3	★★★	Epstein ζ; 3 non-Euler (h=2,3); 4/4 PASS; FE-only sufficiency (Thm 6)
#115	§5.2	★★★	DH on/off discrimination; 1.002 ± 0.001 vs. 1.210 ± 0.099; complete separation
#116	§5.4	★★★	δ-sweep; slope 1.999 ± 0.003 (on) vs. 0.963 ± 0.039 (off); Thm 7 confirmed
#117	§5.5	★★	$c_1 = \mathrm{Re}(\Lambda''/\Lambda')$ verified ($< 0.5\%$, $N = 4$); $c_1 \cdot d \to 1$
#125	§6.1	★★★	Ramanujan Δ ($k = 12$); 4/5 PASS; $\kappa\delta^2$ slope 2.0008 ± 0.0006; σ-uniq. FAIL (B-01)
#126	§6.2	★★★	Weight-16 cusp form ($\Delta \cdot E_4$); 5/5 PASS; slope 1.9989 ± 0.0035; weight invariance (Obs. 8)

Table 10 (continued)

Result	Section	Rating	Key finding
#127	§6.3	★★★	Weight-20 cusp form ($\Delta \cdot E_8$); 4/5 PASS; slope 1.9984 ± 0.0055; σ-uniq. FAIL (B-01)
#200	§6.4	★★★	Maass cusp form (R=9.5337); 5/5 PASS; slope $\mathbf{2.0003 \pm 0.0003}$; most precise P3 to date

B Numerical Data: Off-Critical Zeros (DH)

Table 11 collects the complete δ-sweep data for the three DH off-critical zeros with sufficient precision (OFF#4 included only at $\delta \geq 0.10$).

Table 11: $\kappa\delta^2$ values from Result #116, DH off-critical zeros. ON-critical ensemble mean (10 zeros) shown for comparison.

Zero	δ=0.005	0.01	0.015	0.02	0.03	0.05	0.07	0.10
ON (mean)	1.000048	1.000192	1.000432	1.000767	1.001726	1.004795	1.009399	1.019182
OFF#1	1.018828	1.037696	1.056606	1.075561	1.113605	1.190280	1.267813	1.385945
OFF#2	1.034427	1.068469	1.102132	1.135420	1.200904	1.327729	1.449538	1.624253
OFF#3	1.067034	1.132039	1.195056	1.256139	1.372769	1.585968	1.776429	2.028926
OFF#4	—	—	—	—	—	—	—	1.506337

C Mirror-Pair Decomposition of c_1

For an off-critical zero $\rho = \sigma_0 + it_0$, the mirror partner is $\tilde{\rho} = 1 - \sigma_0 + it_0$. The leading contribution of the mirror pair to $A'(\rho)/A(\rho)$ is

$$\frac{1}{\rho - \tilde{\rho}} = \frac{1}{2\sigma_0 - 1} = \frac{1}{2d},$$

giving $c_1^{\text{mirror}} = 2\,\text{Re}(1/(2d)) = 1/d$. Table 12 shows the decomposition from Result #117.

Table 12: Mirror-pair vs. background decomposition of c_1 (Result #117).

Zero	d	c_1^{fit}	$c_1^{\text{mirror}} = 1/d$	c_1^{bg}	$c_1^{\text{bg}}/c_1^{\text{mirror}}$
OFF#1	0.3085	3.760	3.241	2.139	1.320
OFF#2	0.1508	6.916	6.630	3.601	1.086
OFF#3	0.0744	13.543	13.449	6.818	1.014
OFF#4	0.2243	5.026	4.459	2.796	1.254

As $d \to 0$, the background-to-mirror ratio approaches 1, indicating that the two contributions are of equal magnitude in the limit and that $c_1 \approx 2 \times c_1^{\text{mirror}} = 2/d$ at small d—which exceeds the asymptotic $1/d$ by a factor of 2 until higher-order corrections reduce

the effective value. The data instead show $c_1 \cdot d \to 1$, suggesting significant cancellation among the background terms.

References

[1] Kim, Y.-H. (2026). *Phase Quantisation from Optical Lattices to the Zeta Landscape: A Unified Framework*. Preprint, arXiv:2026.xxxxx.

[2] Titchmarsh, E. C. (1986). *The Theory of the Riemann Zeta-Function* (2nd ed., revised by D. R. Heath-Brown). Oxford University Press.

[3] Edwards, H. M. (1974). *Riemann's Zeta Function*. Academic Press.

[4] Davenport, H., & Heilbronn, H. (1936). On the zeros of certain Dirichlet series. *Journal of the London Mathematical Society*, 11, 181–185.

[5] Silverman, J. H. (1992). *The Arithmetic of Elliptic Curves* (2nd ed.). Springer.

[6] Iwaniec, H., & Kowalski, E. (2004). *Analytic Number Theory*. American Mathematical Society.

[7] Epstein, P. (1903). Zur Theorie allgemeiner Zetafunktionen. *Mathematische Annalen*, 56, 615–644.

[8] Rubinstein, M. (2001). Numerical computations concerning the Riemann hypothesis. *PhD thesis*, Princeton University.

[9] Cremona, J. E. (2019). *Algorithms for Modular Elliptic Curves* (2nd ed.). Cambridge University Press.

[10] PARI/GP (2023). Version 2.15.4. https://pari.math.u-bordeaux.fr/.

[11] Johansson, F., et al. (2023). *mpmath: a Python library for arbitrary-precision floating-point arithmetic* (version 1.3.0). https://mpmath.org/.