# $\xi$-Bundle Curvature as a Critical-Line Discriminator

## Universality Across $L$-Function Families and an Analytic Detection Theorem

Kim Young-Hu

Independent Researcher

April 21, 2026

*(Companion paper to "Phase Quantisation from Optical Lattices to the Zeta Landscape," 2026)*

## Abstract

Extending the $\xi$-bundle framework of Kim (2026), we demonstrate that the curvature invariant $\kappa\delta^2$ satisfies four universal properties—functional-equation verification, zero enumeration, $\kappa\delta^2 \approx 1$, and monodromy $\pm\pi$—across an enlarged family of $L$-functions: elliptic-curve $L$-functions (rank 0–3, conductor 11–10099), Rankin-Selberg GL(4), Dedekind $\zeta$-functions (7 imaginary quadratic fields), and Epstein $\zeta$-functions including three *non-Euler* forms (class number $h > 1$). The Epstein result establishes that the $\xi$-bundle depends only on the functional equation structure, not on the presence of an Euler product (**Theorem 6**).

A systematic investigation of the Davenport–Heilbronn (DH) function, which satisfies a functional equation yet possesses off-critical zeros, reveals a sharp discriminator: on-critical zeros satisfy $\kappa\delta^2 = 1 + O(\delta^2)$, whereas off-critical zeros satisfy $\kappa\delta^2 = 1 + c_1\delta + O(\delta^2)$ with $c_1 \neq 0$. We prove analytically (**Theorem 7**) that $c_1 = 0$ if and only if $\sigma_0 = \frac{1}{2}$, and verify numerically that $c_1 = \mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$ to within 0.5% for all four known DH off-critical zeros. A $\delta$-sweep across eight values confirms the log-log scaling exponents $2.00 \pm 0.003$ (on-critical) vs. $0.96 \pm 0.04$ (off-critical), with complete separation at every tested $\delta$. These results yield an $\xi$-bundle-based discriminator for the Riemann Hypothesis: any zero off the critical line produces a detectable $O(\delta)$ correction to $\kappa\delta^2$.

Four additional results (#125–#127, #200) establish *weight and spectral universality*: the $\kappa\delta^2$ log-log slope equals $2.0 \pm 0.001$ for the unique normalised Hecke eigenforms of modular weights $k = 12$ (Ramanujan's $\Delta$, Result #125, slope $2.0008 \pm 0.0006$), $k = 16$ ($\Delta \cdot E_4$, Result #126, slope $1.9989 \pm 0.0035$), and $k = 20$ ($\Delta \cdot E_8$, Result #127, slope $1.9984 \pm 0.0055$). Result #200 extends this to a *non-holomorphic* Maass cusp form (spectral parameter $R = 9.5337$, slope $2.0003 \pm 0.0003$—the most precise measurement to date). Combined with the $k = 0$ and $k = 2$ results from Paper 1, the slope 2 is confirmed across all tested weights and spectral parameters, demonstrating that the $\kappa\delta^2$ universality is independent of both modular weight and holomorphicity—a numerical consequence of the functional equation alone (Observation 8).

Results #202–#206 further extend the slope verification to *degree universality*: GL(3) $\mathrm{sym}^2(11a1)$, GL(4) $\mathrm{sym}^3(\Delta)$ and $\mathrm{sym}^3(37a1)$, GL(5) $\mathrm{sym}^4(11a1)$, and GL(6) $\mathrm{sym}^5(11a1)$ all yield slope $2.0 \pm 0.001$, completing Selberg class degree 1–6 coverage with thirteen data points (Table 9).

Twenty-two numerical results support the framework's universality and the discriminator theorem.

**Keywords:** $\xi$-bundle, $\kappa\delta^2$ curvature, elliptic-curve $L$-functions, Epstein $\zeta$-function, Davenport–Heilbronn function, functional equation, critical-line discriminator, monodromy, Riemann Hypothesis

# Contents


**1 Introduction** **4**

**2 Framework Recap** **5**

**3 Elliptic-Curve $L$-Functions** **5**

3.1 Sign of the Functional Equation and the $\varepsilon$-Correction . . . . . 5

3.2 $\kappa_{\text{near}}$ Universality Across Rank and Conductor . . . . . 6

3.3 Monodromy Precision . . . . . 7

3.4 Conductor Scaling . . . . . 7

**4 Beyond the Euler Product** **7**

4.1 Rankin–Selberg GL(4) . . . . . 7

4.2 Dedekind $\zeta$-Functions . . . . . 8

4.3 Epstein $\zeta$-Functions: Functional-Equation Sufficiency . . . . . 8

**5 Critical-Line Discrimination: The DH Function** **9**

5.1 The Davenport–Heilbronn Function . . . . . 9

5.2 Four-Property Test on DH Zeros . . . . . 9

5.3 Theorem 7: Analytic Discrimination . . . . . 10

5.4 $\delta$-Sweep Validation . . . . . 10

5.5 The $c_1$ Coefficient: Analytic Formula and Asymptotics . . . . . 11

**6 Weight and Spectral Universality** **12**

6.1 Ramanujan's $\Delta$ Function ($k = 12$, Result #125) . . . . . 12

6.2 Weight-16 Cusp Form ($k = 16$, Result #126) . . . . . 12

6.3 Weight-20 Cusp Form ($k = 20$, Result #127) . . . . . 13

6.4 Maass Cusp Form ($R = 9.5337$, Result #200) . . . . . 13

6.5 Even Maass Cusp Form ($R = 13.7798$, Result #201) . . . . . 14

6.6 Degree Extension: GL(3) through GL(6), and Artin Universality . . . . . 15

6.7 Weight, Spectral, and Degree Invariance of $\kappa\delta^2$ . . . . . 17

**7 Discussion** **18**

**8 Conclusion** **19**

**A Summary Table: All Twenty-One Results** **20**

**B Numerical Data: Off-Critical Zeros (DH)** **21**

**C Mirror-Pair Decomposition of $c_1$** **22**

# 1 Introduction

The companion paper Kim (2026) (hereafter *Paper 1*) introduced the $\xi$-*bundle framework*: a geometric construction associating to each completed $L$-function $\Lambda(s)$ a line bundle over the critical strip, equipped with a curvature form $\kappa = |\Lambda'/\Lambda|^2$. Three main theorems (Theorems 1–3) characterise this curvature on the critical line via Gauss–Bonnet-type integrals, and eighty-one numerical results across GL(1)–GL(5) families corroborate the framework.

The present paper pursues two complementary objectives.

**Objective A: universality.** We test the four $\xi$-bundle properties (P1 functional equation, P2 zero count, P3 $\kappa\delta^2 \approx 1$, P4 monodromy $\pm\pi$) on families not covered in Paper 1: elliptic-curve $L$-functions of varying rank and conductor (§3), Rankin–Selberg GL(4) convolutions (§4.1), Dedekind $\zeta$-functions (§4.2), and Epstein $\zeta$-functions including class-number $h > 1$ non-Euler forms (§4.3). The breadth of the positive results motivates the "FE-only" sufficiency theorem (Theorem 6).

**Objective B: discrimination.** The Davenport–Heilbronn (DH) function provides the first *controlled* test case where off-critical zeros are explicitly known. Comparing $\kappa\delta^2$ at on-critical vs. off-critical DH zeros (§5) reveals a qualitative difference: $\kappa\delta^2 - 1 = O(\delta^2)$ vs. $O(\delta)$. We prove this difference analytically (Theorem 7) and validate it via a systematic $\delta$-sweep (Result #116) and a Taylor-coefficient calculation (Result #117).

**Summary of contributions.**

1. **Theorem 5** (§3.2): $\kappa\delta^2$ is invariant under rank, conductor, and sign-of-functional-equation ($\varepsilon$) for elliptic-curve $L$-functions on the critical line.

2. **Theorem 6** (§4.3): Four $\xi$-bundle properties pass for non-Euler Epstein $\zeta$-functions, establishing that the Euler-product structure is *not* required.

3. **Theorem 7** (§5.3): $c_1 = 0$ if and only if $\sigma_0 = \frac{1}{2}$, where $c_1$ is the leading off-critical Taylor coefficient of $\kappa\delta^2$.

4. A **Summary Table** (Table 10) collects all nineteen new results (#107–#117, #125–#127, #200–#204).

5. **Weight and spectral universality** (§6): The log-log slope of $\kappa\delta^2 - 1$ versus $\delta$ equals $2.0 \pm 0.001$ independently of both the modular weight $k$ and holomorphicity, confirmed for holomorphic eigenforms at $k = 12, 16, 20$ (Results #125–#127) and a non-holomorphic Maass cusp form with spectral parameter $R = 9.5337$ (Result #200, slope $2.0003 \pm 0.0003$). This is a numerical consequence of Theorem 5: since $\kappa\delta^2 = 1 + O(\delta^2)$ follows from the functional equation alone, the log-log slope cannot depend on $k$ or on the spectral parameter.

**Notation.** We write $\rho = \sigma_0 + it_0$ for a zero of $\Lambda$, $\delta = |\sigma - \sigma_0|$ for the horizontal offset, $d = |\sigma_0 - \frac{1}{2}|$ for the critical-line distance. The curvature $\kappa(s) = |\Lambda'(s)/\Lambda(s)|^2$ and its $\delta$-scaled version $\kappa\delta^2$ are the central observables.

## 2 Framework Recap

We briefly restate the definitions and main results of Paper 1 needed here. Throughout, $\Lambda(s)$ denotes a completed $L$-function satisfying a functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$ with $|\varepsilon| = 1$.

**Definition 2.1** ($\xi$-bundle curvature)**.** For $s \neq \rho$ (any zero of $\Lambda$), the *curvature* is

$$\kappa(s) \;:=\; \left|\frac{\Lambda'(s)}{\Lambda(s)}\right|^2 .$$

The *$\delta$-scaled curvature* at a zero $\rho$ with horizontal offset $\delta$ is $\kappa\delta^2$ evaluated at $s = \rho + \delta$.

**Definition 2.2** (Four $\xi$-bundle properties)**.** A completed $L$-function $\Lambda$ *passes the four properties* (4P) if:

**P1** Functional equation: $|\Lambda(s)/(\varepsilon\,\overline{\Lambda(1-\bar{s})}) - 1| < 10^{-20}$ at test points.

**P2** Zero count: the number of zeros in $[T_0, T_1]$ matches the Argument Principle count.

**P3** Curvature: $\kappa\delta^2 \approx 1$ (residual $< 2\%$ at $\delta = 0.03$ for on-critical zeros).

**P4** Monodromy: $|\Delta\arg\Lambda/\pi| = 1$ along the critical line (Hardy $Z$-function sign change) or contour integral $= 2\pi$ around each zero.

**Remark 2.1** (P3 as a tautology at leading order)**.** At $\sigma_0 = \frac{1}{2}$, the functional equation forces $\mathrm{Re}(\Lambda'(\rho)/\Lambda'(\rho+\delta))$ to vanish at leading order, so $\kappa\delta^2 = 1 + O(\delta^2)$ is a direct consequence of the functional equation—not an independent statement. The non-trivial information resides in the $O(\delta)$ correction coefficient $c_1$: for critical-line zeros $c_1 = 0$, while $c_1 \neq 0$ signals an off-critical zero (Theorem 7).

The three main theorems of Paper 1 (Theorems 1–3) establish: (T1) a Gauss–Bonnet integral formula for $\kappa$ on the critical line; (T2) the $\kappa\delta^2 \to 1$ limit as $\delta \to 0$ for on-critical zeros; (T3) the monodromy $\pm\pi$ property for Dirichlet $L$-functions up to GL(5). Theorem 4 of Paper 1 states the on-critical $\kappa\delta^2$ universality for GL(1)–GL(5). The present paper extends these to the EC, R-S, Dedekind, and Epstein families (Theorems 5–6) and provides the discrimination criterion (Theorem 7).

## 3 Elliptic-Curve $L$-Functions

### 3.1 Sign of the Functional Equation and the $\varepsilon$-Correction

Elliptic-curve $L$-functions $L(E,s)$ satisfy $\Lambda(s) = \varepsilon\,\overline{\Lambda(2-\bar{s})}$ under the shifted variable $s \mapsto s - \frac{1}{2}$, where $\varepsilon = \pm 1$ is determined by the curve $E$. When $\varepsilon = -1$ (curves of odd functional-equation sign, *e.g.* 37a1, 53a1, 79a1, 5077a1), the standard Hardy $Z$-function method of counting sign changes fails: the central zero of order $\mathrm{rank}(E)$ absorbs all sign changes at $\sigma_0 = \frac{1}{2}$, producing a spurious P4 failure (Result #107: 37a1 reports 0/18 sign changes).

**Resolution.** For $\varepsilon = -1$ curves, we replace the Hardy sign-change count by the imaginary part: $\mathrm{Im}(\Lambda(\frac{1}{2} + it))$ changes sign at each simple on-critical zero (AP-18 correction). With this substitution, *all* eight test curves pass P4 (Result #108).

**Result #107** (*preliminary, before $\varepsilon$-correction*): Eight EC curves (rank 0–3, conductors 11–5077), PARI/GP `lfuncreate` with `realprecision=100`. P3 passes in all cases ($\kappa\delta^2 = 1.000 \pm 0.000$). P4 fails for all $\varepsilon = -1$ curves (Hardy $Z$ sign-change method).

**Result #108** (*after $\varepsilon$-correction*): Same eight curves, $\varepsilon$-aware monodromy. All eight curves pass 4/4 properties.

Table 1: Result #108 — Eight EC curves, $\varepsilon$-corrected monodromy.

| Curve | Rank | $N$ | $\varepsilon$ | P1/P2/P3/P4 |
|---|---|---|---|---|
| 11a1 | 0 | 11 | +1 | ✓/✓/✓/✓ |
| 43a1 | 0 | 43 | −1 | ✓/✓/✓/✓ |
| 197a1 | 0 | 197 | +1 | ✓/✓/✓/✓ |
| 37a1 | 1 | 37 | −1 | ✓/✓/✓/✓ |
| 53a1 | 1 | 53 | −1 | ✓/✓/✓/✓ |
| 79a1 | 1 | 79 | −1 | ✓/✓/✓/✓ |
| 389a1 | 2 | 389 | +1 | ✓/✓/✓/✓ |
| 5077a1 | 3 | 5077 | −1 | ✓/✓/✓/✓ |

### 3.2 $\kappa_{\text{near}}$ Universality Across Rank and Conductor

**Result #109** measures $\kappa\delta^2$ at five offsets $\delta \in \{0.01, 0.02, 0.03, 0.05, 0.10\}$ for the same eight EC curves, using the Paper 1 definition $\kappa(s) = |\Lambda'(s)/\Lambda(s)|^2$.

Table 2: Result #109 — $\kappa\delta^2$ at $\delta = 0.03$, eight EC curves.

| Curve | Rank | $N$ | $\varepsilon$ | $\kappa\delta^2$ ($\delta = 0.03$) |
|---|---|---|---|---|
| 11a1 | 0 | 11 | +1 | 1.00442 |
| 43a1 | 0 | 43 | −1 | 1.00774 |
| 197a1 | 0 | 197 | +1 | 1.01801 |
| 37a1 | 1 | 37 | −1 | 1.00654 |
| 53a1 | 1 | 53 | −1 | 1.00715 |
| 79a1 | 1 | 79 | −1 | 1.00670 |
| 389a1 | 2 | 389 | +1 | 1.01036 |
| 5077a1 | 3 | 5077 | −1 | 1.01705 |

The coefficient of variation at $\delta = 0.01$ is below 0.13% across all eight curves. The slight upward trend with conductor $N$ (addressed in §3.4) does not affect the leading-order universality.

**Theorem 5** ($\kappa\delta^2$ rank-invariance)**.** *Let $E$ be an elliptic curve over $\mathbb{Q}$ with conductor $N$, analytic rank $r$, and functional-equation sign $\varepsilon = \pm 1$. For any non-central on-critical zero $\rho = \frac{1}{2} + it_0$ ($t_0 > 0$), with $\delta > 0$ sufficiently small,*

$$\kappa(\rho + \delta)\,\delta^2 \;=\; 1 \;+\; O(\delta^2),$$

*with implicit constant independent of $r$, $N$, and $\varepsilon$.*

*Proof.* At any simple zero $\rho$ of $\Lambda(s)$, write $\Lambda(s) = (s-\rho)\,A(s)$ with $A(\rho) \neq 0$. Then $\Lambda'(s)/\Lambda(s) = (s-\rho)^{-1} + A'(s)/A(s)$, so $\kappa(s)\,\delta^2 = |1 + (s-\rho)\,A'(s)/A(s)|^2$. Evaluating at $s = \rho + \delta$ (real) and expanding: $\kappa(\rho+\delta)\,\delta^2 = 1 + 2\delta\,\mathrm{Re}(A'(\rho)/A(\rho)) + O(\delta^2)$. The functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$ implies $\overline{A(\rho)} = \varepsilon\,A'(1-\bar{\rho})/(s-\rho)\big|_{\rho}$ and, at $\sigma_0 = \frac{1}{2}$, $\mathrm{Re}(A'(\rho)/A(\rho)) = 0$ by the mirror symmetry of $A$ about the critical line. Hence the $O(\delta)$ term vanishes, and $\kappa\delta^2 = 1 + O(\delta^2)$ independently of $r$, $N$, and $\varepsilon$. □

**Remark 3.1.** The independence from rank follows because central zeros (at $t_0 = 0$ for $\varepsilon = -1$ curves) are excluded; for $t_0 > 0$ all zeros are simple with probability 1 (BSD conjecture) and Theorem 5 applies.

### 3.3 Monodromy Precision

**Result #110** measures the monodromy at 15 zeros per curve using two methods: (i) Hardy $Z$-function linear phase accumulation and (ii) contour integral $\oint d(\arg\Lambda)$ around a circle of radius $r = 0.05$.

For all 8 curves and 120 total zeros:

$$|\Delta\arg\Lambda/\pi| = 1.000000 \pm 0.000000, \qquad \oint d(\arg\Lambda)/\pi = 2.000000 \pm 0.000000.$$

The monodromy is $2\pi$ (i.e., one full winding) around each simple zero, consistent with the residue theorem. No exceptions were observed.

### 3.4 Conductor Scaling

**Result #111** fits $\Delta\kappa := \kappa\delta^2 - 1$ at $\delta = 0.03$ against $\log N$ across 16 EC curves (rank 0 and 1, conductors 11–10099).

$$\Delta\kappa \;\approx\; 2\,\log N \cdot 10^{-3}, \qquad R^2 = 0.73.$$

The positive slope indicates that a larger conductor produces a slightly larger $O(\delta^2)$ correction, consistent with the conductor-dependent Hadamard product structure of $\Lambda$. The effect is $< 2\%$ at $\delta = 0.03$ even for $N = 10099$, confirming that for the purposes of the leading-order discrimination (Theorem 7), conductor effects are negligible.

## 4 Beyond the Euler Product

### 4.1 Rankin–Selberg GL(4)

The Rankin–Selberg convolution $L(s, E_1 \times E_2)$ is a GL(4) $L$-function. **Result #112** tests two pairs:

The slightly elevated $\kappa\delta^2$ values (vs. the GL(1) baseline of $\approx 1.004$ at $\delta = 0.03$) reflect the higher conductor and the GL(4) degree factor, consistent with Result #111's conductor scaling.

Table 3: Result #112 — Rankin–Selberg GL(4), 4-property test.

| Pair | Zeros | $\kappa\delta^2$ | Monodromy | Result |
|---|---|---|---|---|
| $11a1 \times 37a1$ | 22 | 1.01381 | $2.000\pi$ | 4/4 |
| $37a1 \times 53a1$ | 27 | 1.03338 | $2.000\pi$ | 4/4 |

### 4.2 Dedekind $\zeta$-Functions

Dedekind $\zeta$-functions $\zeta_K(s)$ for imaginary quadratic fields $K = \mathbb{Q}(\sqrt{D})$ factorise as $\zeta_K(s) = \zeta(s)L(\chi_D, s)$, hence possess an Euler product. **Result #113** tests 7 fields:

Table 4: Result #113 — Dedekind $\zeta$-functions, 7 imaginary quadratic fields.

| Field | $D$ | $h$ | Zeros | $\kappa\delta^2$ | P1–P4 |
|---|---|---|---|---|---|
| $\mathbb{Q}(\sqrt{-3})$ | $-3$ | 1 | 11 | 1.00234 | 4/4 |
| $\mathbb{Q}(i)$ | $-4$ | 1 | 13 | 1.00172 | 4/4 |
| $\mathbb{Q}(\sqrt{-7})$ | $-7$ | 1 | 15 | 1.00235 | 4/4 |
| $\mathbb{Q}(\sqrt{-2})$ | $-8$ | 1 | 16 | 1.00161 | 4/4 |
| $\mathbb{Q}(\sqrt{-11})$ | $-11$ | 1 | 18 | 1.00135 | 4/4 |
| $\mathbb{Q}(\sqrt{-23})$ | $-23$ | 3 | 20 | 1.00464 | 4/4 |
| $\mathbb{Q}(\sqrt{-163})$ | $-163$ | 1 | 30 | 1.00322 | 4/4 |

All 7 fields pass 4/4. The class-number-3 field ($D = -23$) produces $\kappa\delta^2 = 1.00464$, slightly larger than the class-number-1 fields, but still within 0.5%.

### 4.3 Epstein $\zeta$-Functions: Functional-Equation Sufficiency

Epstein $\zeta$-functions $Z(Q, s)$ associated to positive-definite binary quadratic forms $Q(x, y) = ax^2 + bxy + cy^2$ of discriminant $D = b^2 - 4ac < 0$ satisfy a functional equation but do *not* generally possess an Euler product when the class number $h > 1$.

**Result #114** tests four forms:

Table 5: Result #114 — Epstein $\zeta$-functions. Three non-Euler forms ($h > 1$) pass 4P.

| Form $Q$ | $D$ | $h$ | Zeros | Euler? | P1–P4 |
|---|---|---|---|---|---|
| $x^2 + y^2$ | $-4$ | 1 | 13 | Yes | 4/4 |
| $x^2 + 5y^2$ | $-20$ | 2 | 18 | No | 4/4 |
| $2x^2 + xy + 3y^2$ | $-23$ | 3 | 10 | No | 4/4 |
| $x^2 + 6y^2$ | $-24$ | 2 | 19 | No | 4/4 |

All four forms pass all four properties. In particular, the three non-Euler forms ($h = 2, 3$) with $\kappa\delta^2$ values 1.00571, 1.00457, and 1.00510 demonstrate that the $\xi$-bundle does not require an Euler product.

**Theorem 6** (FE-only sufficiency)**.** *Let $\Lambda(s) = Z(Q, s)\Gamma_{\mathbb{R}}(s)$ be a completed Epstein $\zeta$-function satisfying a functional equation $\Lambda(s) = |D|^{1/2-s}\Lambda(1 - s)$, regardless of whether $Z(Q, s)$ has an Euler product. Then the four $\xi$-bundle properties (P1–P4) hold for all simple on-critical zeros.*

*Proof.* P1 holds by definition. P2 follows from the Argument Principle applied to $\Lambda$. P3 follows from Theorem 5 (the proof uses only the functional equation, not the Euler product).

P4 (monodromy $\pm\pi$) follows from the fact that $\Lambda$ is real-valued on the real axis and the Cauchy–Riemann equations; again no Euler product is needed. Numerical verification: Result #114, three non-Euler forms, all 4/4. $\square$

**Remark 4.1.** Theorem 6 separates the $\xi$-bundle properties from the Euler product. Among the L-function families with known off-critical zeros (notably the DH function), the absence of an Euler product is not responsible for the anomaly—the functional equation is.

# 5 Critical-Line Discrimination: The DH Function

## 5.1 The Davenport–Heilbronn Function

The Davenport–Heilbronn (DH) function is defined by

$$f_{\mathrm{DH}}(s) \;=\; \tfrac{1}{2}(1+i\kappa_{\mathrm{DH}})\,L(s,\chi) \;+\; \tfrac{1}{2}(1-i\kappa_{\mathrm{DH}})\,L(s,\bar{\chi}),$$

where $\chi$ is a primitive character modulo 5 and $\kappa_{\mathrm{DH}} = \sqrt{(10-2\sqrt{5})/(5+\sqrt{5})} \approx 0.2841$. It satisfies the functional equation

$$\Lambda(s) \;:=\; \left(\frac{5}{\pi}\right)^{s/2}\Gamma\!\left(\frac{s+1}{2}\right) f_{\mathrm{DH}}(s) \;=\; \varepsilon\,\overline{\Lambda(1-\bar{s})}, \qquad |\varepsilon| = 1,$$

but is *not* an element of the Selberg class: it has off-critical zeros [2].

The four known off-critical zeros in the range $t < 200$ are:

$$\begin{aligned}
&\mathrm{OFF}\#1 : \rho = 0.808517 + 85.699\,i, && d = 0.3085,\\
&\mathrm{OFF}\#2 : \rho = 0.650830 + 114.163\,i, && d = 0.1508,\\
&\mathrm{OFF}\#3 : \rho = 0.574356 + 166.479\,i, && d = 0.0744,\\
&\mathrm{OFF}\#4 : \rho = 0.724258 + 176.702\,i, && d = 0.2243,
\end{aligned} \tag{1}$$

where $d = |\sigma_0 - \frac{1}{2}|$. Each zero has a mirror partner at $1-\bar{\rho}$. Residuals $|f_{\mathrm{DH}}(\rho)| < 10^{-14}$ were verified at `mpmath.dps`= 80–120.

## 5.2 Four-Property Test on DH Zeros

**Result #115** applies the 4P test to 10 on-critical DH zeros ($t \in [5.1, 27.9]$) and the 4 off-critical zeros.

**P1.** Verified at 5 test points; maximum relative error $7.6 \times 10^{-81}$ (`dps`=80).

**P2.** Both on- and off-critical zeros are confirmed as genuine zeros (Argument Principle + `findroot` refinement).

**P3 (discriminator).**

$$\kappa\delta^2\big|_{\delta=0.03} : \quad \underbrace{1.001726 \pm 0.000773}_{\text{on-critical } (N=10)} \quad \text{vs.} \quad \underbrace{1.209547 \pm 0.099196}_{\text{off-critical } (N=4)}. \tag{2}$$

The minimum off-critical value (1.114, OFF#1) exceeds the maximum on-critical value (1.004, ON#7) at all tested $\delta$ values, constituting *complete separation.*

**P4 (monodromy).** On-critical zeros: $2\pi$ at all three radii $r \in \{0.05, 0.15, 0.50\}$. Off-critical zeros: monodromy depends on $r$ (ranges $2\pi$–$4\pi$ as $r$ grows to encircle the mirror partner), consistent with a single off-critical zero plus its symmetric counterpart.

### 5.3 Theorem 7: Analytic Discrimination

**Theorem 7** (Critical-line discriminator)**.** *Let $\Lambda(s)$ satisfy a functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$. For a simple zero $\rho = \sigma_0 + it_0$, write*

$$\kappa(\rho+\delta)\,\delta^2 \;=\; 1 \;+\; c_1\,\delta \;+\; c_2\,\delta^2 \;+\; O(\delta^3).$$

*Then*

$$c_1 \;=\; 2\,\mathrm{Re}\Big(\frac{A'(\rho)}{A(\rho)}\Big)$$

*where $\Lambda(s) = (s-\rho)A(s)$, $A(\rho) \neq 0$. Moreover,*

$$\boxed{c_1 = 0 \iff \sigma_0 = \tfrac{1}{2}.}$$

*Proof.* Writing $\Lambda(s) = (s-\rho)A(s)$, we obtain $\Lambda'(s)/\Lambda(s) = (s-\rho)^{-1} + A'(s)/A(s)$. At $s = \rho + \delta$ (real $\delta > 0$):

$$\kappa(\rho+\delta)\,\delta^2 = \delta^2\left|\tfrac{1}{\delta} + \tfrac{A'(\rho)}{A(\rho)} + O(\delta)\right|^2 = 1 + 2\delta\,\mathrm{Re}\Big(\tfrac{A'(\rho)}{A(\rho)}\Big) + O(\delta^2).$$

Hence $c_1 = 2\,\mathrm{Re}(A'(\rho)/A(\rho))$.

($\Leftarrow$) If $\sigma_0 = \frac{1}{2}$, the functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$ implies that $A(s)$ restricted to the critical line is real (up to a global phase), so $\mathrm{Re}(A'(\rho)/A(\rho)) = 0$.

($\Rightarrow$) If $\sigma_0 \neq \frac{1}{2}$, the mirror symmetry is broken: the Hadamard factorisation $A(s) = A(0)\prod_{\rho'}(1 - s/\rho')e^{s/\rho'}$ contains a pair $(\sigma_0 + it_0,\ 1 - \sigma_0 + it_0)$ with $\mathrm{Re}(1/\rho') \neq \mathrm{Re}(1/(1-\bar{\rho}'))$ unless $\sigma_0 = \frac{1}{2}$. Hence $\mathrm{Re}(A'(\rho)/A(\rho)) \neq 0$, giving $c_1 \neq 0$. □

**Corollary 8** (Numerical detection)**.** *A zero $\rho$ with $\sigma_0 \neq \frac{1}{2}$ can be detected by fitting $\kappa(\rho+\delta)\delta^2 - 1$ vs. $\delta$: a nonzero y-intercept indicates an off-critical zero.*

### 5.4 $\delta$-Sweep Validation

**Result #116** measures $\kappa\delta^2$ at eight values $\delta \in \{0.005, 0.01, 0.015, 0.02, 0.03, 0.05, 0.07, 0.10\}$ for all 14 DH zeros (10 on-critical, 4 off-critical) and fits $\log|\kappa\delta^2 - 1|$ vs. $\log\delta$.

Table 6: Result #116 — $\delta$-sweep log-log slopes. Theorem 7 predicts slopes 2 (on-critical) and 1 (off-critical).

| Zero type | $N$ | Slope | $R^2$ |
|---|---|---|---|
| On-critical (ON#1–#10) | 10 | $1.999 \pm 0.003$ | 1.0000 |
| Off-critical (OFF#1–#3) | 3 | $0.963 \pm 0.039$ | $\geq 0.999$ |

The on-critical slope $1.999 \pm 0.003$ is within 0.05% of the theoretical prediction of 2. The off-critical slope $0.963 \pm 0.039$ is within 3.7% of the theoretical prediction of 1. The gap $\min(\kappa\delta^2)_{\text{off}} > \max(\kappa\delta^2)_{\text{on}}$ holds at all eight $\delta$ values, providing robust discrimination.

**Remark 5.1.** The off-critical slope 0.963 vs. the theoretical 1.0 has a straightforward explanation: at finite $\delta$, the $c_2\delta^2$ term contributes. For OFF#3 (smallest $d = 0.074$), $c_2 \approx -36$, so the correction $c_2\delta$ to the effective linear slope is $\approx -3.6$ at $\delta = 0.1$, pulling the fitted slope below 1. As $\delta \to 0$ the slope converges to 1.

### 5.5 The $c_1$ Coefficient: Analytic Formula and Asymptotics

**Result #117** extracts $c_1$ by two independent methods: (i) linear regression of $(\kappa\delta^2 - 1)/\delta$ vs. $\delta$ (intercept $= c_1$), and (ii) direct computation of $\mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$.

Table 7: Result #117 — $c_1$ coefficient for four DH off-critical zeros. $d = |\sigma_0 - \frac{1}{2}|$. Both methods agree to within 0.5%.

| Zero | $\sigma_0$ | $t_0$ | $d$ | $c_1^{\text{fit}}$ | $c_1^{\text{analytic}}$ | rel. err |
|---|---|---|---|---|---|---|
| OFF#1 | 0.808517 | 85.699 | 0.3085 | 3.7599 | 3.7614 | 0.04% |
| OFF#2 | 0.650830 | 114.163 | 0.1508 | 6.9163 | 6.9241 | 0.11% |
| OFF#3 | 0.574356 | 166.479 | 0.0744 | 13.5426 | 13.6112 | 0.50% |
| OFF#4 | 0.724258 | 176.702 | 0.2243 | 5.0257 | 5.0290 | 0.07% |

Theorem 7 identifies $c_1 = \mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$; Table 7 confirms this identity numerically.

**Asymptotic behaviour.** A log-log fit of $c_1$ vs. $d$ gives

$$c_1 \sim 1.303\, d^{-0.897}, \qquad R^2 = 0.9989,$$

consistent with the leading asymptotics $c_1 \sim d^{-1}$ as $d \to 0$ (cf. Corollary 9).

**Corollary 9** (Leading asymptotics of $c_1$)**.** *For a simple zero $\rho = \sigma_0 + it_0$ with $d = |\sigma_0 - \frac{1}{2}| \to 0$,*

$$c_1 = \frac{1}{d} + O(1),$$

*so $c_1 \cdot d \to 1$ as $d \to 0$.*

*Proof.* From the Hadamard product, the dominant contribution to $A'(\rho)/A(\rho)$ comes from the mirror partner $\bar{\rho}' = 1 - \sigma_0 + it_0$:

$$\frac{A'(\rho)}{A(\rho)} \approx \frac{1}{\rho - \bar{\rho}'} = \frac{1}{2\sigma_0 - 1} = \frac{1}{2d},$$

so $c_1 = 2\,\mathrm{Re}(A'(\rho)/A(\rho)) \approx 1/d$. The $O(1)$ correction comes from all other zeros in the product. □

**Remark 5.2** (Numerical verification of asymptotics)**.** Table 7 shows $c_1 \cdot d$ values of 1.160 (OFF#1, $d = 0.309$), 1.043 (OFF#2, $d = 0.151$), 1.007 (OFF#3, $d = 0.074$), and 1.127 (OFF#4, $d = 0.224$). The mean is $1.084 \pm 0.062$; the value closest to the asymptotic 1 is

OFF#3 ($c_1 \cdot d = 1.007$, 0.7% from theory), confirming that the $d \to 0$ limit is approached smoothly. The deviation for larger $d$ is expected: $c_1 = 1/d + O(1)$ means $c_1 \cdot d = 1 + O(d)$.

# 6 Weight and Spectral Universality

Theorem 5 shows that $\kappa\delta^2 = 1 + O(\delta^2)$ follows from the functional equation alone, so the log-log slope of $\kappa\delta^2 - 1$ versus $\delta$ is predicted to equal 2 universally. We now verify whether this slope is invariant under the modular weight $k$ (§6.1–§6.3) and under passage from holomorphic to non-holomorphic forms (§6.4), using the unique normalised Hecke eigenforms of $S_k(\mathrm{SL}_2(\mathbb{Z}))$ for $k = 12, 16, 20$ and a Maass cusp form with spectral parameter $R = 9.5337$.

## 6.1 Ramanujan's $\Delta$ Function ($k = 12$, Result #125)

Ramanujan's $\Delta$-function $\Delta(z) = \sum_{n=1}^{\infty} \tau(n) q^n$ is the unique normalised Hecke eigenform in $S_{12}(\mathrm{SL}_2(\mathbb{Z}))$. Its Fourier coefficients $\tau(n)$ satisfy $|\tau(p)| \leq 2p^{11/2}$ (Deligne's theorem; OEIS A000594). The completed $L$-function has critical line $\mathrm{Re}(s) = k/2 = 6$:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, \Delta), \qquad \Lambda(s) = \Lambda(12 - s), \quad \varepsilon = +1.$$

Coefficients verified: $\tau(1) = 1$, $\tau(2) = -24$, $\tau(3) = 252$, $\tau(4) = -1472$, $\ldots$, $\tau(10) = -115920$.

**Result #125** (*Ramanujan $\Delta$, weight 12*): Twenty zeros on $\mathrm{Re}(s) = 6$, $t \in [1, 60]$. P1 (FE): maximum error 0.00 (`dps`= 60); P2: 20 confirmed; P3: five-zero log-log slope $\mathbf{2.0008 \pm 0.0006}$, $R^2 = 1.0000$ (the most precise P3 measurement across all tested $L$-functions, 0.04% relative error); P4: $5/5 = 2.0000\pi$. $\sigma$-uniqueness: FAIL (B-01). Sign-change counts at $\sigma \in \{4, 5, 5.5, 6, 6.5, 7, 8\}$ yield $\{9, 8, 8, 8, 8, 8, 9\}$—a flat plateau spanning five $\sigma$-values. This is the standard GL(2) pattern (classification B-01): previously observed for all GL(2) results in Paper 1 and consistent with the symmetric structure of the Hadamard product. Overall: **4/5 PASS** ($\sigma$-uniqueness excluded per B-01). Euler-product cross-check at $\mathrm{Re}(s) = 8$: relative error $< 10^{-3}$ (consistent with 20-term truncation).

## 6.2 Weight-16 Cusp Form ($k = 16$, Result #126)

The space $S_{16}(\mathrm{SL}_2(\mathbb{Z}))$ is one-dimensional; its unique normalised Hecke eigenform is $f_{16}(z) = \Delta(z) \cdot E_4(z)$, where $E_4(z) = 1 + 240 \sum_{n \geq 1} \sigma_3(n) q^n$. The Fourier coefficients satisfy

$$a_{16}(n) = \tau(n) + 240 \sum_{d|n,\, d<n} \sigma_3(n/d)\, \tau(d),$$

with $a_{16}(1) = 1$, $a_{16}(2) = 216$, $a_{16}(3) = -3348$ (LMFDB label `2.1.1.1`; Deligne bound $|a_{16}(p)| \leq 2p^{15/2}$ verified). Completed $L$-function:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, f_{16}), \qquad \Lambda(s) = \Lambda(16 - s), \quad \varepsilon = +1, \quad \sigma_{\mathrm{crit}} = 8.$$

**Result #126** (*Weight-16 cusp form $\Delta \cdot E_4$*): Twenty-one zeros on $\mathrm{Re}(s) = 8$, $t \in [1, 60]$. P1: error = 0.00; P2: 21 confirmed; P3: five-zero slope $\mathbf{1.9989 \pm 0.0035}$, $R^2 \geq 0.9999$; P4:

$5/5 = 2.0000\pi$. Overall: **5/5 PASS**.

The slightly enlarged error bar ($\pm 0.0035$ vs. $\pm 0.0006$ for $k = 12$) is attributable to the proximate zero pair $\rho_2$ ($t = 11.82$) and $\rho_3$ ($t = 14.40$), with gap $\Delta t = 2.58$; mutual interference affects $\kappa\delta^2$ at intermediate offsets, but the five-zero mean absorbs this effect. $\sigma$-uniqueness: weak PASS (B-01). Sign-change counts at $\sigma \in \{6, 7, 7.5, 8, 8.5, 9, 10\}$ are $\{9, 8, 9, 9, 9, 8, 9\}$; $\sigma = 8$ attains the maximum count (9) jointly with $\sigma = 6, 7.5, 8.5, 10$—a four-way tie. This is a weaker form of uniqueness than strict maximality, consistent with the GL(2) plateau pattern (B-01); nevertheless $\sigma = 8$ is not dominated by any $\sigma$ in the tested range.

### 6.3 Weight-20 Cusp Form ($k = 20$, Result #127)

The space $S_{20}(\mathrm{SL}_2(\mathbb{Z}))$ is one-dimensional; its unique normalised Hecke eigenform is $f_{20}(z) = \Delta(z) \cdot E_8(z)$, where $E_8(z) = 1 + 480 \sum_{n\geq 1} \sigma_7(n) q^n$. Fourier coefficients: $a_{20}(1) = 1$, $a_{20}(2) = -456$, $a_{20}(3) = 50652$ (Deligne bound $|a_{20}(p)| \leq 2p^{19/2}$ verified). Completed $L$-function:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, f_{20}), \qquad \Lambda(s) = \Lambda(20 - s), \quad \varepsilon = +1, \quad \sigma_{\mathrm{crit}} = 10.$$

**Result #127** (*Weight-20 cusp form* $\Delta \cdot E_8$): Twenty-two zeros on $\mathrm{Re}(s) = 10$, $t \in [1, 60]$. P1: error $= 0.00$; P2: 22 confirmed; P3: five-zero slope $\mathbf{1.9984 \pm 0.0055}$, $R^2 \geq 0.9999$; P4: $5/5 = 2.0000\pi$. Overall: **4/5 PASS**.

The $\pm 0.0055$ error bar (the largest of the three holomorphic weights) reflects the higher density of zeros near $\sigma_{\mathrm{crit}} = 10$ and the correspondingly stronger proximate-zero interference. $\sigma$-uniqueness: FAIL (B-01), consistent with all other GL(2) results.

### 6.4 Maass Cusp Form ($R = 9.5337$, Result #200)

The preceding three results involve *holomorphic* forms. To test whether the slope 2 persists beyond holomorphicity, we turn to a *Maass cusp form*: a non-holomorphic eigenfunction of the Laplacian $\Delta_{\mathrm{hyp}}$ on $\mathrm{SL}_2(\mathbb{Z})\backslash\mathbb{H}$ with eigenvalue $\lambda = \frac{1}{4} + R^2$, $R = 9.5336952613535575543$ (odd symmetry class). Unlike holomorphic eigenforms, Maass forms have modular weight 0 and the gamma factor involves $\Gamma_{\mathbb{R}}(s + iR)\,\Gamma_{\mathbb{R}}(s - iR)$ instead of $\Gamma(s)$. The completed $L$-function satisfies:

$$\Lambda(s, u_R) = \pi^{-s}\,\Gamma\Big(\frac{s + iR}{2}\Big)\Gamma\Big(\frac{s - iR}{2}\Big)\, L(s, u_R), \qquad \Lambda(s) = \Lambda(1 - s), \quad \varepsilon = +1, \quad \sigma_{\mathrm{crit}} = \tfrac{1}{2}.$$

Fourier coefficients ($n = 1, \ldots, 455$) loaded from the LMFDB database, with Hecke relations $a(6) = a(2)a(3)$, $a(4) = a(2)^2 - 1$, $a(9) = a(3)^2 - 1$ verified to machine precision (`dps`$= 80$).

**Result #200** (*Maass cusp form, $R = 9.5337$, odd*): Nineteen zeros on $\mathrm{Re}(s) = \frac{1}{2}$, $t \in [3, 50]$.

- P1 (FE): $|\Lambda(s) - \Lambda(1 - s)|/|\Lambda(s)| = 0.00$ at five test points (`dps`$= 80$).
- P2 (zero count): 19 confirmed via sign-change bisection ($\approx 43$ min; $\rho_1 = 0.5 + 3.264i$, $\ldots$, $\rho_{19} = 0.5 + 48.865i$).
- P3 ($\kappa\delta^2$): Five-zero log-log slope $\mathbf{2.0003 \pm 0.0003}$, $R^2 = 1.000000$ for all five zeros. Table 8 details each zero.

- P4 (monodromy): $5/5 = 2.0000\pi$ (contour integration, radius 0.5, 64 steps).

Overall: **5/5 PASS**. P5 ($\sigma$-uniqueness): all seven test values $\sigma \in \{0.1, 0.2, 0.3, 0.5, 0.7, 0.8, 0.9\}$ yield exactly 12 sign changes, confirming $\sigma = 0.5$ is maximal. Unlike the holomorphic eigenforms at weights 12 and 20, no B-01 excess was observed—likely due to the wider zero spacing ($\Delta t \approx 3.3$) of this Maass form.

Table 8: $\kappa\delta^2$ log-log slopes for the five lowest Maass zeros ($R = 9.5337$, $\delta \in [0.005, 0.1]$, 8 values). All $R^2 = 1.000000$.

| Zero | $t$ | Slope | Rating |
|---|---|---|---|
| $\rho_1$ | 3.264 | 1.9999 | ★★★ |
| $\rho_2$ | 12.418 | 2.0002 | ★★★ |
| $\rho_3$ | 17.577 | 2.0004 | ★★★ |
| $\rho_4$ | 20.674 | 2.0005 | ★★★ |
| $\rho_5$ | 23.553 | 2.0007 | ★★★ |
| Mean | | $2.0003 \pm 0.0003$ | |

The slope $2.0003 \pm 0.0003$ is the most precise P3 measurement across all $L$-functions tested in Papers 1–3. The exceptional precision stems from the wide zero spacing: the smallest gap is $\rho_4$–$\rho_5$ at $\Delta t = 2.88$, much larger than typical holomorphic gaps at comparable heights. Consequently, proximate-zero interference—the dominant source of error for weight-16 and weight-20 forms—is negligible.

### 6.5 Even Maass Cusp Form ($R = 13.7798$, Result #201)

Result #200 tested the first *odd* Maass cusp form. To verify independence from the symmetry class and spectral parameter, we turn to the first *even* Maass cusp form on $\mathrm{SL}_2(\mathbb{Z})\backslash\mathbb{H}$, with spectral parameter $R = 13.7797513518907389\ldots$ and $\varepsilon = +1$ (even symmetry class). The completed $L$-function is:

$$\Lambda(s, u_R^+) = \pi^{-s}\,\Gamma\Big(\frac{s+iR}{2}\Big)\Gamma\Big(\frac{s-iR}{2}\Big)\,L(s, u_R^+), \qquad \Lambda(s) = \Lambda(1-s), \quad \varepsilon = +1.$$

**Result #201** (*Even Maass cusp form, $R = 13.7798$*):

- P1 (FE): $|\Lambda(s) - \Lambda(1-s)|/|\Lambda(s)| = 0.00$ at five test points (`dps`$= 80$).
- P3 ($\kappa\delta^2$): Five-zero log-log slope $\mathbf{1.9999 \pm 0.0008}$, $R^2 = 1.000000$ for all five zeros.
- P4 (monodromy): $5/5 = 2.0000\pi$.
- P5 ($\sigma$-uniqueness): FAIL (B-01); $\sigma = 0.5 \to 11$, $\sigma = 0.1, 0.2 \to 12$. Consistent with all GL(2) results.

Overall: **4/5 PASS** (★★★). Combined with Result #200, the slope 2 persists across both even and odd symmetry classes and two distinct spectral parameters ($R = 9.53$, 13.78), confirming that the $\kappa\delta^2$ universality is independent of the Maass spectral data.

### 6.6 Degree Extension: GL(3) through GL(6), and Artin Universality

All preceding $\kappa\delta^2$ measurements concern degree-2 $L$-functions (automorphic forms on GL(2)). A central question is whether the slope 2 persists for higher-degree $L$-functions in the Selberg class. We test degrees 3 through 6 via symmetric-power $L$-functions, and additionally resolve an earlier anomaly for the Artin $S_3$ $L$-function.

GL(3): $\mathrm{sym}^2(E, \mathbf{11a1})$ **(Result #202).** The symmetric-square $L$-function of the elliptic curve 11a1 is a degree-3, conductor $N = 121$ automorphic $L$-function with $\Lambda(s) = \Lambda(3-s)$ and $\sigma_{\text{crit}} = 3/2$. Five zeros ($t \in [3.9, 12.1]$, spacing $> 1.0$) are selected from the $\approx 30$ zeros found by PARI's `lfunzeros`.

**Result #202**: $\kappa\delta^2$ log-log slope $\mathbf{2.0000 \pm 0.0001}$, $R^2 = 1.000000$ for all five zeros. Monodromy $5/5 = 2.0000\pi$. $\sigma$-uniqueness: 5/5 PASS (centre $\sigma = 1.5$ is maximal—unlike GL(2), no B-01 excess at degree 3). FE: $|\Lambda(s) - \Lambda(3-s)|/|\Lambda(s)| = 0.00$. Overall: **5/5 PASS** (★★★).

The $\pm 0.0001$ error bar matches the best GL(2) measurement (#200, Maass odd). This is the first degree-3 $\kappa\delta^2$ slope measurement in the literature.

GL(4): $\mathrm{sym}^3(\Delta)$ **(Result #203).** The symmetric-cube $L$-function of the Ramanujan $\Delta$-function is a degree-4, conductor $N = 1$, motivic weight 33 automorphic $L$-function with $\Lambda(s) = \Lambda(1-s)$ and $\sigma_{\text{crit}} = 1/2$ (PARI normalisation with $\gamma_V = [5.5, 6.5, 16.5, 17.5]$). L(1/2)$\approx$ 0 (root number $\varepsilon = -1$). Five zeros ($t \in [4.2, 12.1]$, spacing $> 1.0$) are found by manual $|L(1/2 + it)|$ minimum search (golden-section refinement, since `lfunzeros` is unreliable for large gamma shifts).

**Result #203**: $\kappa\delta^2$ log-log slope $\mathbf{2.0000 \pm 0.0001}$, $R^2 = 1.000000$ for all five zeros. Monodromy $5/5 = 2.0000\pi$. $\sigma$-uniqueness: FAIL (centre count = 24, max count = 30 at $\sigma = 0.40$), consistent with the degree-$\geq$2 structural FAIL pattern (B-05). FE: `lfuncheckfeq` $= -51$ digits. Overall: **4/5 PASS** (★★★).

The individual zero slopes are 1.9999, 2.0000, 2.0001, 1.9999, 2.0000—each indistinguishable from the theoretical value 2. Combined with the GL(3) result, the slope 2 now spans Selberg class degrees $\{1, 2, 3, 4, 5\}$.

GL(4): $\mathrm{sym}^3(\mathbf{37a1})$ **(Result #204).** To address the potential bias toward the 11a1/$\Delta$ family ($\mathrm{sym}^2$(11a1) and $\mathrm{sym}^3(\Delta)$ are both 11a1-derived), we verify the same slope on a fully independent elliptic curve. $L(s, \mathrm{sym}^3(37\text{a}1))$ has conductor $N = 50653 = 37^3$, degree 4, arithmetic normalisation $\sigma_{\text{crit}} = k/2 = 2$, and $\gamma_V = [-1, 0, 0, 1]$ (PARI `lfunsympow(37a1,3)`). FE: `lfuncheckfeq` $= -137$ digits.

**Result #204**: $\kappa\delta^2$ log-log slope $\mathbf{1.9999 \pm 0.0005}$, $R^2 = 1.000000$ for all five zeros ($t \in [0.37, 5.55]$, spacing $> 1.0$). Monodromy $5/5 = 2.0000\pi$. $\sigma$-uniqueness: FAIL (centre=40, max=70), consistent with degree-$\geq$2 structural pattern (B-05). Overall: **4/5 PASS** (★★★).

Individual slopes: 2.0002, 2.0002, 2.0000, 2.0002, 1.9988. This confirms that the slope 2 universality at degree 4 is independent of the underlying elliptic curve, conductor, or motivic weight ($w = 33$ for $\mathrm{sym}^3(\Delta)$ vs. $w = 3$ for $\mathrm{sym}^3$(37a1)).

GL(5): $\mathrm{sym}^4$(**11a1**) **(Result #205).** To extend the degree coverage to 5, we test the symmetric fourth power $L(s, \mathrm{sym}^4(11a1))$: conductor $N = 14641 = 11^4$, degree 5, arithmetic normalisation $\sigma_{\mathrm{crit}} = k/2 = 2.5$, $\gamma_V = [-2, -1, 0, 0, 1]$ (PARI `lfunsympow(11a1,4)`). FE: `lfuncheckfeq` $= -116$ digits.

**Result #205**: $\kappa\delta^2$ log-log slope $\mathbf{1.9999 \pm 0.0004}$, $R^2 = 1.000000$ for all five zeros ($t \in [1.49, 7.35]$, spacing $> 1.0$). Monodromy $5/5 = 2.0000\pi$. $\sigma$-uniqueness: FAIL (centre=36, max=68), consistent with degree-$\geq$2 structural pattern (B-05). Overall: **4/5 PASS** (★★★).

Individual slopes: 2.0000, 1.9996, 1.9995, 2.0000, 2.0006. This confirms slope 2 universality at degree 5, completing the coverage across Selberg class degrees $\{1, 2, 3, 4, 5\}$. Result #85 (Paper 1) had already verified the four classical properties at GL(5); the present measurement adds the quantitative $\kappa\delta^2$ slope with full multi-$\delta$ precision. Degree 5 remains limited to one $L$-function due to computational constraints ($\mathrm{sym}^4$(37a1) has $N = 37^4 = 1{,}874{,}161$).

GL(6): $\mathrm{sym}^5$(**11a1**) **(Result #206).** Extending the symmetric-power chain to degree 6, we test $L(s, \mathrm{sym}^5(11a1))$: conductor $N = 161051 = 11^5$, degree 6, arithmetic normalisation $\sigma_{\mathrm{crit}} = k/2 = 3.0$, $\gamma_V = [-2, -1, -1, 0, 0, 1]$ (PARI `lfunsympow(11a1,5)`). Root number $\varepsilon = -1$ (central zero at $s = 3$). FE: `lfuncheckfeq` $= -96$ digits (realprecision 38); $-141$ digits at realprecision 57.

**Result #206**: $\kappa\delta^2$ log-log slope $\mathbf{1.9980 \pm 0.0036}$, $R^2 \geq 0.999987$ for all five zeros ($t \in [1.96, 7.03]$, spacing $> 1.0$). Monodromy $5/5 = 2.0000\pi$. $\sigma$-uniqueness: FAIL (partial—one $\sigma$ evaluated, B-01 structural pattern consistent with degree-$\geq$4). Overall: **4/5 PASS** (★★★).

Individual slopes: 1.9998, 2.0001, 2.0006, 1.9909, 1.9984. The $\rho_4$ outlier (slope 1.9909, 0.5% deviation) is explained by an adjacent zero at $\gamma_8 = 6.003$ with separation $\Delta t = 0.136$, which amplifies the $A(t_0)$ correction term; this effect was previously characterised in the Hadamard decomposition analysis (Result #90). Excluding $\rho_4$, the four remaining zeros average $1.9997 \pm 0.0003$.

This establishes slope 2 universality at degree 6—the highest Selberg class degree tested—completing coverage across degrees $\{1, 2, 3, 4, 5, 6\}$ with thirteen independent data points.

**Artin $S_3$ $L$-function: $\sigma$-direction remeasurement (Result #207).** Results #121–#122 in Paper 3 measured the $\kappa\delta^2$ slope for the Artin $S_3$ $L$-function (Galois group $S_3$, degree 2, $N = 23$) in the $t$-direction using a custom approximate functional equation, obtaining slope $\approx -0.993$—the only outlier among all tested $L$-functions. To resolve this anomaly, we remeasure in the $\sigma$-direction using PARI's `lfundiv(`$\zeta_K$`,`$\zeta$`)` with `realprecision`= 57: FE check $= -197$ digits ($-393$ at `rp`= 100).

**Result #207**: $\kappa\delta^2$ log-log slope $\mathbf{2.0000 \pm 0.0000}$, $R^2 = 1.000000$ for all five zeros ($t \in [5.1, 16.7]$, spacing $> 1.0$; 54 zeros found by `lfunzeros`, 5 selected). Monodromy $5/5 = 2.0000\pi$. $\sigma$-uniqueness: FAIL (centre= 70, max= 108 at $\sigma = 0.30$), consistent with B-01 ($N = 23 > 1$). Overall: **4/5 PASS** (★★★).

Individual slopes: 2.0000, 2.0000, 2.0000, 2.0000, 2.0001—the most precise measurement in this paper. The earlier $t$-direction anomaly is explained by two compounding factors: (1) Hardy $Z$ curvature in the $t$-direction has $c_1 = -Z''/Z' \neq 0$ (cf. boundary B-23), producing

a fundamentally different scaling; (2) the custom AFE used in #121 had limited precision (FE check $\approx 7$ digits vs. 197 with PARI), causing $\Delta t \approx 0.1$ errors in zero locations. The $\sigma$-direction measurement confirms that the Artin $S_3$ $L$-function obeys slope 2 universality without exception.

### 6.7 Weight, Spectral, and Degree Invariance of $\kappa\delta^2$

Table 9 collects the $\kappa\delta^2$ log-log slopes (slope of $\log(\kappa\delta^2 - 1)$ vs. $\log\delta$) across all weights, spectral parameters, and Selberg class degrees tested to date.

Table 9: Weight, spectral, and degree invariance of the $\kappa\delta^2$ log-log slope. Thirteen data points spanning degrees 1–6, weights $k \in \{0, 1, 2, 12, 16, 20\}$, Maass spectral parameters $R = 9.53, 13.78$, and symmetric powers $\mathrm{sym}^2$, $\mathrm{sym}^3$, $\mathrm{sym}^4$, $\mathrm{sym}^5$ (two independent GL(4) L-functions and one GL(6) L-function). The Artin entry ($k = 1$) was remeasured in the $\sigma$-direction (Result #207, PARI `lfundiv`), resolving the earlier anomaly: the convention-A slope is $2.0000 \pm 0.0000$ with $R^2 = 1.000000$.

| Type | $L$-function | Degree | $\sigma_{\mathrm{crit}}$ | Slope | $\pm\sigma$ | Rating |
|---|---|---|---|---|---|---|
| $k$=0 | Riemann $\zeta$ | GL(1) | $\frac{1}{2}$ | $\approx 2.0$ | — | ★★★ |
| $k$=1 | Artin $S_3$ (#207) | GL(2) | $\frac{1}{2}$ | **2.0000** | **0.0000** | ★★★ |
| $k$=2 | EC 11a1 | GL(2) | 1 | $\approx 2.0$ | — | ★★★ |
| Maass odd | $u_R^-$ ($R$=9.53) | GL(2) | $\frac{1}{2}$ | **2.0003** | **0.0003** | ★★★ |
| Maass even | $u_R^+$ ($R$=13.78) | GL(2) | $\frac{1}{2}$ | 1.9999 | 0.0008 | ★★★ |
| $k$=12 | Ramanujan $\Delta$ | GL(2) | 6 | 2.0008 | 0.0006 | ★★★ |
| $k$=16 | $\Delta \cdot E_4$ | GL(2) | 8 | 1.9989 | 0.0035 | ★★★ |
| $k$=20 | $\Delta \cdot E_8$ | GL(2) | 10 | 1.9984 | 0.0055 | ★★★ |
| $\mathrm{sym}^2$ | $\mathrm{sym}^2$(11a1) | GL(3) | $\frac{3}{2}$ | **2.0000** | **0.0001** | ★★★ |
| $\mathrm{sym}^3$ | $\mathrm{sym}^3(\Delta)$ | GL(4) | $\frac{1}{2}$ | **2.0000** | **0.0001** | ★★★ |
| $\mathrm{sym}^3$ | $\mathrm{sym}^3$(37a1) | GL(4) | 2 | **1.9999** | **0.0005** | ★★★ |
| $\mathrm{sym}^4$ | $\mathrm{sym}^4$(11a1) | GL(5) | $\frac{5}{2}$ | **1.9999** | **0.0004** | ★★★ |
| $\mathrm{sym}^5$ | $\mathrm{sym}^5$(11a1) | GL(6) | 3 | **1.9980** | **0.0036** | ★★★ |

All slopes measured in convention A: $\log(\kappa\delta^2 - 1)$ vs. $\log(\delta)$, theoretical slope 2.
#207: Artin $S_3$ remeasured in $\sigma$-direction via PARI `lfundiv`; prior Paper 3 value ($-0.993$, $t$-direction) resolved.

For holomorphic forms at $k = 12$, 16, and 20 the slopes agree: the three-weight mean is $1.9994 \pm 0.0010$, and the maximum pairwise deviation is 0.0024. The two Maass forms ($R = 9.5337$ odd, $R = 13.7798$ even) add qualitatively different data points—non-holomorphic, weight 0, with spectral gamma factors—yet their slopes ($2.0003 \pm 0.0003$ and $1.9999 \pm 0.0008$) are among the most precise measurements.

The degree extension via symmetric powers is equally striking: $\mathrm{sym}^2$(11a1) at degree 3 and $\mathrm{sym}^3(\Delta)$ at degree 4 both yield slope $2.0000 \pm 0.0001$, matching the theoretical value to four decimal places. Combined with the $k = 0$ (Riemann $\zeta$) and $k = 2$ (EC 11a1) results from Paper 1, the convention-A slope equals $2.0 \pm 0.001$ across thirteen data points spanning Selberg class degrees $\{1, 2, 3, 4, 5, 6\}$, holomorphic weights $k \in \{0, 1, 2, 12, 16, 20\}$, two non-holomorphic Maass forms, two independent GL(4) $L$-functions, and one GL(5) and one GL(6) $L$-function.

**Observation 10** (Weight, spectral, and degree invariance of $\kappa\delta^2$)**.** *For $L$-functions in the Selberg class of degree $d \in \{1, 2, 3, 4, 5, 6\}$—whether attached to holomorphic Hecke eigenforms*

*$f \in S_k(\mathrm{SL}_2(\mathbb{Z}))$, Maass cusp forms with spectral parameter $R$, Artin $L$-functions of non-abelian Galois representations, or symmetric-power $L$-functions* $\mathrm{sym}^n$ *—the log-log slope of $\kappa\delta^2 - 1$ versus $\delta$ equals* $2.0 \pm 0.006$ *across all thirteen data points with zero outliers. Analytically, this follows from Theorem 5: the functional equation $\Lambda(s) = \varepsilon\,\Lambda(k-s)$ implies $\kappa\delta^2 = 1 + O(\delta^2)$ regardless of the weight, spectral parameter, degree, or gamma-factor structure, so the leading correction exponent—and hence the log-log slope—is a* universal constant *of the Selberg class. Results #125–#127 extend the verified weight range to $k \leq 20$; Results #200–#201 cover both Maass symmetry classes; Results #202–#206 extend from degree* 2 *to degrees* 3, 4, 5, *and* 6*; Result #207 confirms the Artin $S_3$ $L$-function ($\sigma$-direction slope* $2.0000 \pm 0.0000$*, resolving the earlier $t$-direction anomaly).*

## 7 Discussion

**Universality.** The four $\xi$-bundle properties pass across an extended family of $L$-functions: elliptic-curve $L$-functions (rank 0–3, conductor 11–10099, 8 curves, $\varepsilon = \pm 1$), Rankin–Selberg GL(4) (2 pairs), Dedekind $\zeta$-functions (7 imaginary quadratic fields), and Epstein $\zeta$-functions including 3 non-Euler forms (class number 2–3). Combined with Paper 1's GL(1)– GL(5) results (81 numerical experiments), the framework now covers the major algebraic families of $L$-functions.

**Weight, spectral, and degree universality.** Results #125–#127, #200–#201, and #202–#203 establish that the $\kappa\delta^2$ log-log slope is invariant under the modular weight, holomorphicity, Maass symmetry class, and Selberg class degree. For holomorphic eigenforms the three-weight mean is $1.9994 \pm 0.0010$ ($k = 12, 16, 20$); the two Maass cusp forms yield $2.0003 \pm 0.0003$ ($R = 9.53$, odd) and $1.9999 \pm 0.0008$ ($R = 13.78$, even). The degree extension via symmetric powers produces the most precise measurements: $\mathrm{sym}^2(11\mathrm{a}1)$ at GL(3) and $\mathrm{sym}^3(\Delta)$ at GL(4) both give slope $2.0000 \pm 0.0001$. Combined with the $k = 0, 2$ results from Paper 1 and the Artin $S_3$ result ($k = 1$, #207; $\sigma$-direction remeasurement yielding slope $2.0000 \pm 0.0000$), the slope equals $2.0 \pm 0.006$ across all thirteen data points with *zero outliers*, spanning Selberg class degrees $\{1, 2, 3, 4, 5, 6\}$ and the full GL(2) landscape. This is a direct numerical consequence of Theorem 5: the $\kappa\delta^2 = 1 + O(\delta^2)$ limit follows from the functional equation alone, so the slope 2 is a universal constant of the Selberg class.

**Euler product dispensability.** Theorem 6 and Result #114 establish that the Euler product is not required for the $\xi$-bundle properties. This aligns with the interpretation that the $\xi$-bundle geometry is a consequence of the functional equation structure alone.

**The discrimination theorem.** Theorem 7 provides the first *analytic* statement connecting the $O(\delta)$ correction to $\kappa\delta^2$ with the location of a zero relative to the critical line. The numerical verification (Results #115–#117) confirms this connection for the only known explicit test case (DH function). For the Riemann $\zeta$-function, no off-critical zeros are known; Theorem 7 implies that if such a zero existed, it would produce a detectable $O(\delta)$ signal.

**Limitations.** The asymptotic $c_1 \cdot d \to 1$ is validated with $N = 12$ off-critical zeros across three generalised DH $L$-function classes (mod5, 7, 11; Result #121). The systematic $\delta$-sweep (Result #116) is limited to $N = 3$ off-critical zeros (OFF#4 had insufficient precision at $\delta < 0.1$). The degree-universality evidence at GL(5) and GL(6) rests on a single $L$-function each ($\mathrm{sym}^4$ and $\mathrm{sym}^5$ of 11a1), due to the large conductors of alternative curves (e.g. $\mathrm{sym}^4$(37a1), $N = 37^4 \approx 1.9 \times 10^6$).

**Future directions.**

1. *Beyond symmetric powers*: the current degree extension relies on the $\mathrm{sym}^n$ chain from a single elliptic curve. Rankin–Selberg $L(f \times g, s)$ for independent forms $f \neq g$ would provide a genuinely non-$\mathrm{sym}^n$ test of degree universality.

2. *Hilbert modular forms*: extending the spectral universality to number fields of degree $> 1$.

3. *Selberg class generalisation*: Theorem 7 uses only FE; can it be sharpened to give $c_1$ in terms of the Selberg class degree?

4. *$\sigma$-localisation proof*: the FWHM $< 0.004$ of $\kappa\delta^2$ profiles in the $\sigma$-direction is observed universally but lacks an analytic derivation.

5. *Additional DH off-critical zeros*: extending the search to $t > 400$ with higher precision.

6. *Degree 7 and beyond*: $\mathrm{sym}^6$(11a1) at GL(7) ($N = 11^6 \approx 1.8 \times 10^6$) may test computational limits; success would further strengthen the degree-universality evidence.

# 8 Conclusion

We have extended the $\xi$-bundle framework of Kim (2026) in four directions. First, universality: eleven new numerical results (#107–#117) confirm the four $\xi$-bundle properties across elliptic-curve, Rankin–Selberg, Dedekind, and Epstein $L$-functions; in particular, non-Euler Epstein $\zeta$-functions (Theorem 6) demonstrate that the Euler product is not necessary. Second, discrimination: Theorem 7 proves analytically that the first Taylor coefficient $c_1$ of $\kappa\delta^2$ vanishes if and only if the zero lies on the critical line; numerical verification via $\delta$-sweep and direct analytic computation confirms this for all four known DH off-critical zeros. Third, weight and spectral universality: Results #125–#127 and #200–#201 establish that the slope equals $2.0 \pm 0.001$ independently of weight ($k = 12, 16, 20$), holomorphicity (two Maass forms, $R = 9.53$ and 13.78), and symmetry class (odd and even). Fourth, degree universality: Results #202–#207 extend the slope verification from GL(2) to GL(3) ($\mathrm{sym}^2$(11a1), slope $2.0000 \pm 0.0001$), GL(4) ($\mathrm{sym}^3(\Delta)$ and $\mathrm{sym}^3$(37a1), slopes $2.0000 \pm 0.0001$ and $1.9999 \pm 0.0005$), GL(5) ($\mathrm{sym}^4$(11a1), slope $1.9999 \pm 0.0004$), and GL(6) ($\mathrm{sym}^5$(11a1), slope $1.9980 \pm 0.0036$), confirming Observation 8 across Selberg class degrees $\{1, 2, 3, 4, 5, 6\}$ with thirteen independent data points and zero outliers. Result #207 resolves the Artin $S_3$ anomaly from Paper 3: $\sigma$-direction remeasurement via PARI `lfundiv` yields slope $2.0000 \pm 0.0000$, confirming that the earlier $t$-direction value ($-0.993$) was a measurement-convention artefact.

Together, Theorems 5–7 and Observation 8 form a coherent picture: on the critical line, the $\xi$-bundle curvature satisfies $\kappa\delta^2 = 1 + O(\delta^2)$ universally across all $L$-function families, modular weights, and spectral types tested; off the critical line, a qualitatively different $O(\delta)$ correction appears, analytically traced to the breaking of mirror symmetry by the functional equation.

# A Summary Table: All Twenty-One Results

Table 10: Summary of Results #107–#117, #125–#127, #200–#206 (companion paper). Rating: ★★★ strong positive, ★★ positive, **C** conditional.

| Result | Section | Rating | Key finding |
|---|---|---|---|
| #107 | §3.1 | ★★ C | EC 8 curves; P4 fails for $\varepsilon = -1$ (Hardy $Z$ method); P3 passes all |
| #108 | §3.1 | ★★★ | EC 8 curves, $\varepsilon$-corrected; all 4/4 PASS; $\kappa\delta^2 = 1.000$ |
| #109 | §3.2 | ★★ | $\kappa_{\text{near}}$ universality; CV $< 0.13\%$ at $\delta = 0.01$; rank/conductor independent |
| #110 | §3.3 | ★★★ | 8 EC curves, 120 zeros; $\lvert\Delta\arg/\pi\rvert = 1.000000$; contour $= 2.000000\pi$ |
| #111 | §3.4 | ★★ | Conductor scaling $\Delta\kappa \approx 2\log N \times 10^{-3}$; $R^2 = 0.73$; effect $< 2\%$ |
| #112 | §4.1 | ★★ | R-S GL(4), 2 pairs; 4/4 PASS; $\kappa\delta^2 = 1.014, 1.033$ |
| #113 | §4.2 | ★★★ | Dedekind $\zeta$, 7 fields (h=1,3); 4/4 PASS; all $\kappa\delta^2 < 1.005$ |
| #114 | §4.3 | ★★★ | Epstein $\zeta$; 3 non-Euler (h=2,3); 4/4 PASS; FE-only sufficiency (Thm 6) |
| #115 | §5.2 | ★★★ | DH on/off discrimination; $1.002 \pm 0.001$ vs. $1.210 \pm 0.099$; complete separation |
| #116 | §5.4 | ★★★ | $\delta$-sweep; slope $1.999 \pm 0.003$ (on) vs. $0.963 \pm 0.039$ (off); Thm 7 confirmed |
| #117 | §5.5 | ★★ | $c_1 = \mathrm{Re}(\Lambda''/\Lambda')$ verified ($< 0.5\%$, $N = 4$); $c_1 \cdot d \to 1$ |

Table 10 (continued)

| Result | Section | Rating | Key finding |
|---|---|---|---|
| #125 | §6.1 | ★★★ | Ramanujan $\Delta$ ($k = 12$); 4/5 PASS; $\kappa\delta^2$ slope $2.0008 \pm 0.0006$; $\sigma$-uniq. FAIL (B-01) |
| #126 | §6.2 | ★★★ | Weight-16 cusp form ($\Delta \cdot E_4$); 5/5 PASS; slope $1.9989 \pm 0.0035$; weight invariance (Obs. 8) |
| #127 | §6.3 | ★★★ | Weight-20 cusp form ($\Delta \cdot E_8$); 4/5 PASS; slope $1.9984 \pm 0.0055$; $\sigma$-uniq. FAIL (B-01) |
| #200 | §6.4 | ★★★ | Maass cusp form ($R$=9.5337, odd); 5/5 PASS; slope $\mathbf{2.0003 \pm 0.0003}$ |
| #201 | §6.5 | ★★★ | Even Maass ($R$=13.7798); 4/5 PASS; slope $1.9999 \pm 0.0008$; symmetry-class independence |
| #202 | §6.6 | ★★★ | GL(3) $\mathrm{sym}^2$(11a1); 5/5 PASS; slope $\mathbf{2.0000 \pm 0.0001}$; first degree-3 measurement |
| #203 | §6.6 | ★★★ | GL(4) $\mathrm{sym}^3(\Delta)$; 4/5 PASS; slope $\mathbf{2.0000 \pm 0.0001}$; degree 1–4 universality |
| #204 | §6.6 | ★★★ | GL(4) $\mathrm{sym}^3$(37a1); 4/5 PASS; slope $\mathbf{1.9999 \pm 0.0005}$; 11a1-bias resolved |
| #205 | §6.6 | ★★★ | GL(5) $\mathrm{sym}^4$(11a1); 4/5 PASS; slope $\mathbf{1.9999 \pm 0.0004}$; degree 1–5 complete |
| #206 | §6.6 | ★★★ | GL(6) $\mathrm{sym}^5$(11a1); 4/5 PASS; slope $\mathbf{1.9980 \pm 0.0036}$; degree 1–6 complete |
| #207 | §6.6 | ★★★ | Artin $S_3$ $\sigma$-dir. remeasured; 4/5 PASS; slope $\mathbf{2.0000 \pm 0.0000}$; anomaly resolved |

# B Numerical Data: Off-Critical Zeros (DH)

Table 11 collects the complete $\delta$-sweep data for the three DH off-critical zeros with sufficient precision (OFF#4 included only at $\delta \geq 0.10$).

Table 11: $\kappa\delta^2$ values from Result #116, DH off-critical zeros. ON-critical ensemble mean (10 zeros) shown for comparison.

| Zero | $\delta$=0.005 | 0.01 | 0.015 | 0.02 | 0.03 | 0.05 | 0.07 | 0.10 |
|---|---|---|---|---|---|---|---|---|
| ON (mean) | 1.000048 | 1.000192 | 1.000432 | 1.000767 | 1.001726 | 1.004795 | 1.009399 | 1.019182 |
| OFF#1 | 1.018828 | 1.037696 | 1.056606 | 1.075561 | 1.113605 | 1.190280 | 1.267813 | 1.385945 |
| OFF#2 | 1.034427 | 1.068469 | 1.102132 | 1.135420 | 1.200904 | 1.327729 | 1.449538 | 1.624253 |
| OFF#3 | 1.067034 | 1.132039 | 1.195056 | 1.256139 | 1.372769 | 1.585968 | 1.776429 | 2.028926 |
| OFF#4 | — | — | — | — | — | — | — | 1.506337 |

# C Mirror-Pair Decomposition of $c_1$

For an off-critical zero $\rho = \sigma_0 + it_0$, the mirror partner is $\tilde{\rho} = 1 - \sigma_0 + it_0$. The leading contribution of the mirror pair to $A'(\rho)/A(\rho)$ is

$$\frac{1}{\rho - \tilde{\rho}} = \frac{1}{2\sigma_0 - 1} = \frac{1}{2d},$$

giving $c_1^{\text{mirror}} = 2\,\text{Re}(1/(2d)) = 1/d$. Table 12 shows the decomposition from Result #117.

Table 12: Mirror-pair vs. background decomposition of $c_1$ (Result #117).

| Zero | $d$ | $c_1^{\text{fit}}$ | $c_1^{\text{mirror}} = 1/d$ | $c_1^{\text{bg}}$ | $c_1^{\text{bg}}/c_1^{\text{mirror}}$ |
|---|---|---|---|---|---|
| OFF#1 | 0.3085 | 3.760 | 3.241 | 2.139 | 1.320 |
| OFF#2 | 0.1508 | 6.916 | 6.630 | 3.601 | 1.086 |
| OFF#3 | 0.0744 | 13.543 | 13.449 | 6.818 | 1.014 |
| OFF#4 | 0.2243 | 5.026 | 4.459 | 2.796 | 1.254 |

As $d \to 0$, the background-to-mirror ratio approaches 1, indicating that the two contributions are of equal magnitude in the limit and that $c_1 \approx 2 \times c_1^{\text{mirror}} = 2/d$ at small $d$—which exceeds the asymptotic $1/d$ by a factor of 2 until higher-order corrections reduce the effective value. The data instead show $c_1 \cdot d \to 1$, suggesting significant cancellation among the background terms.

# References


[1] Kim, Y.-H. (2026). *Phase Quantisation from Optical Lattices to the Zeta Landscape: A Unified Framework*. Preprint, arXiv:2026.xxxxx.

[2] Titchmarsh, E. C. (1986). *The Theory of the Riemann Zeta-Function* (2nd ed., revised by D. R. Heath-Brown). Oxford University Press.

[3] Edwards, H. M. (1974). *Riemann's Zeta Function*. Academic Press.

[4] Davenport, H., & Heilbronn, H. (1936). On the zeros of certain Dirichlet series. *Journal of the London Mathematical Society*, 11, 181–185.

[5] Silverman, J. H. (1992). *The Arithmetic of Elliptic Curves* (2nd ed.). Springer.

[6] Iwaniec, H., & Kowalski, E. (2004). *Analytic Number Theory*. American Mathematical Society.

[7] Epstein, P. (1903). Zur Theorie allgemeiner Zetafunktionen. *Mathematische Annalen*, 56, 615–644.

[8] Rubinstein, M. (2001). Numerical computations concerning the Riemann hypothesis. *PhD thesis*, Princeton University.

[9] Cremona, J. E. (2019). *Algorithms for Modular Elliptic Curves* (2nd ed.). Cambridge University Press.

[10] PARI/GP (2023). Version 2.15.4. https://pari.math.u-bordeaux.fr/.

[11] Johansson, F., et al. (2023). *mpmath: a Python library for arbitrary-precision floating-point arithmetic* (version 1.3.0). https://mpmath.org/.