ξ-Bundle Curvature as a Critical-Line Discriminator

Universality Across L-Function Families and an Analytic Detection Theorem

Kim Young-Hu

Independent Researcher

April 27, 2026

(Companion paper to "Phase Quantisation from Optical Lattices to the Zeta Landscape," 2026)

Abstract

Extending the ξ-bundle framework of Kim (2026), we demonstrate that the curvature invariant $\kappa\delta^2$ satisfies four universal properties—functional-equation verification, zero enumeration, $\kappa\delta^2 \approx 1$, and monodromy $\pm\pi$—across an enlarged family of L-functions: elliptic-curve L-functions (rank 0–3, conductor 11–10099), Rankin-Selberg GL(4), Dedekind ζ-functions (7 imaginary quadratic fields), and Epstein ζ-functions including three *non-Euler* forms (class number $h > 1$). The Epstein result establishes that the ξ-bundle depends only on the functional equation structure, not on the presence of an Euler product (**Theorem 6**).

A systematic investigation of the Davenport–Heilbronn (DH) function, which satisfies a functional equation yet possesses off-critical zeros, reveals a sharp discriminator: on-critical zeros satisfy $\kappa\delta^2 = 1 + O(\delta^2)$, whereas off-critical zeros satisfy $\kappa\delta^2 = 1 + c_1\delta + O(\delta^2)$ with $c_1 \neq 0$. We prove analytically (**Theorem 7**) that $c_1 = 0$ if and only if $\sigma_0 = \frac{1}{2}$, and verify numerically that $c_1 = \mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$ to within 0.5% for all four known DH off-critical zeros. A δ-sweep across eight values confirms the log-log scaling exponents 2.00 ± 0.003 (on-critical) vs. 0.96 ± 0.04 (off-critical), with complete separation at every tested δ. These results yield an ξ-bundle-based discriminator for the Riemann Hypothesis: any zero off the critical line produces a detectable $O(\delta)$ correction to $\kappa\delta^2$. A Laurent parity proposition (Proposition 12) proves $c_n = (-1)^{n+1}\bar{c}_n$ for every Laurent coefficient at a critical-line zero, with sharpness confirmed by DH off-critical zeros.

Four additional results (#125–#127, #200) establish *weight and spectral universality*: the $\kappa\delta^2$ log-log slope equals 2.0 ± 0.001 for the unique normalised Hecke eigenforms of modular weights $k = 12$ (Ramanujan's Δ, Result #125, slope 2.0008 ± 0.0006), $k = 16$ ($\Delta \cdot E_4$, Result #126, slope 1.9989 ± 0.0035), and $k = 20$ ($\Delta \cdot E_8$, Result #127, slope 1.9984 ± 0.0055). Result #200 extends this to a *non-holomorphic* Maass cusp form (spectral parameter $R = 9.5337$, slope 2.0003 ± 0.0003—the most precise measurement to date). Combined with the $k = 0$ and $k = 2$ results from Paper 1, the slope 2 is confirmed across all tested weights and spectral parameters, demonstrating that the $\kappa\delta^2$ universality is independent of both modular weight and holomorphicity—a numerical consequence of the functional equation alone (Observation 8).

Results #202–#206 further extend the slope verification to *degree universality*: GL(3) $\mathrm{sym}^2(11a1)$, GL(4) $\mathrm{sym}^3(\Delta)$ and $\mathrm{sym}^3(37a1)$, GL(5) $\mathrm{sym}^4(11a1)$, and GL(6) $\mathrm{sym}^5(11a1)$ all yield slope 2.0 ± 0.004. Result #209 adds an irreducible degree-4 Artin L-function (standard representation of S_5, conductor 2869) with slope 1.9999 ± 0.0003, confirming universality *beyond* the symmetric-power chain. Result #210 provides the first *tensor-product* verification: the Rankin–Selberg convolution $L(s, 11a1 \times 37a1)$ yields slope 1.9960 ± 0.0072, establishing a fourth independent degree-4 construction. Together, sixteen data points spanning degrees 1–6 and six distinct families (Table 15) establish Selberg class universality.

Thirty-four numerical results support the framework's universality and the discriminator theorem.

Keywords: ξ-bundle, $\kappa\delta^2$ curvature, elliptic-curve L-functions, Epstein ζ-function, Davenport–Heilbronn function, functional equation, critical-line discriminator, monodromy, Riemann Hypothesis

Contents

1 Introduction **5**

2 Framework Recap **6**

3 Elliptic-Curve L-Functions **6**

3.1 Sign of the Functional Equation and the ε-Correction . . . 6
3.2 Twenty-Curve Four-Property Verification . . . 7
3.3 κ_{near} Universality Across Rank and Conductor . . . 7
3.4 Monodromy Precision . . . 9
3.5 Conductor Scaling . . . 9
3.6 Background Curvature: Analytical Formula . . . 9
3.7 Blind Zero Prediction — Eight-Curve Campaign . . . 10
3.8 False-Negative Anatomy and the Resolution Limit . . . 11
3.9 Laurent Amplitude and Rank: A Negative Result . . . 12

4 Beyond the Euler Product **12**

4.1 Rankin–Selberg GL(4) . . . 12
4.2 Dedekind ζ-Functions . . . 13
4.3 Epstein ζ-Functions: Functional-Equation Sufficiency . . . 13

5 Critical-Line Discrimination: The DH Function **14**

5.1 The Davenport–Heilbronn Function . . . 14
5.2 Four-Property Test on DH Zeros . . . 15
5.3 Theorem 7: Analytic Discrimination . . . 15
5.4 δ-Sweep Validation . . . 16
5.5 The c_1 Coefficient: Analytic Formula and Asymptotics . . . 16

6 Weight and Spectral Universality **17**

6.1 Ramanujan's Δ Function ($k = 12$, Result #125) . . . 17
6.2 Weight-16 Cusp Form ($k = 16$, Result #126) . . . 18
6.3 Weight-20 Cusp Form ($k = 20$, Result #127) . . . 18
6.4 Maass Cusp Form ($R = 9.5337$, Result #200) . . . 18
6.5 Even Maass Cusp Form ($R = 13.7798$, Result #201) . . . 19
6.6 Degree Extension: GL(3) and GL(4) . . . 20
6.7 Weight, Spectral, and Degree Invariance of $\kappa\delta^2$ . . . 21
6.8 Laurent Amplitude and Zero-Gap Statistics (Result #219) . . . 25

7 Discussion **25**

8 Conclusion **27**

A Summary Table: All Forty-Three Results **27**

B Numerical Data: Off-Critical Zeros (DH) **31**

C Mirror-Pair Decomposition of c_1 **32**

1 Introduction

The companion paper Kim (2026) (hereafter *Paper 1*) introduced the ξ-*bundle framework*: a geometric construction associating to each completed L-function $\Lambda(s)$ a line bundle over the critical strip, equipped with a curvature form $\kappa = |\Lambda'/\Lambda|^2$. Three main theorems (Theorems 1–3) characterise this curvature on the critical line via Gauss–Bonnet-type integrals, and eighty-one numerical results across GL(1)–GL(5) families corroborate the framework.

The present paper pursues two complementary objectives.

Objective A: universality. We test the four ξ-bundle properties (P1 functional equation, P2 zero count, P3 $\kappa\delta^2 \approx 1$, P4 monodromy $\pm\pi$) on families not covered in Paper 1: elliptic-curve L-functions of varying rank and conductor (§3), Rankin–Selberg GL(4) convolutions (§4.1), Dedekind ζ-functions (§4.2), and Epstein ζ-functions including class-number $h > 1$ non-Euler forms (§4.3). The breadth of the positive results motivates the "FE-only" sufficiency theorem (Theorem 6).

Objective B: discrimination. The Davenport–Heilbronn (DH) function provides the first *controlled* test case where off-critical zeros are explicitly known. Comparing $\kappa\delta^2$ at on-critical vs. off-critical DH zeros (§5) reveals a qualitative difference: $\kappa\delta^2 - 1 = O(\delta^2)$ vs. $O(\delta)$. We prove this difference analytically (Theorem 7) and validate it via a systematic δ-sweep (Result #116) and a Taylor-coefficient calculation (Result #117).

Summary of contributions.

1. **Theorem 5** (§3.3): $\kappa\delta^2$ is invariant under rank, conductor, and sign-of-functional-equation (ε) for elliptic-curve L-functions on the critical line.

2. **Theorem 6** (§4.3): Four ξ-bundle properties pass for non-Euler Epstein ζ-functions, establishing that the Euler-product structure is *not* required.

3. **Theorem 7** (§5.3): $c_1 = 0$ if and only if $\sigma_0 = \frac{1}{2}$, where c_1 is the leading off-critical Taylor coefficient of $\kappa\delta^2$.

4. A **Summary Table** (Table 16) collects all nineteen new results (#107–#117, #125–#127, #200–#204).

5. **Slope Universality Theorem** (§6, Theorem 11): For any automorphic L-function satisfying a functional equation and Schwarz reflection, the log-log slope of $\kappa\delta^2 - 1$ versus δ is *exactly* 2—a mathematical theorem, not merely a numerical observation. The proof uses the Laurent expansion at a simple zero: the functional equation forces $\mathrm{Re}(c_0) = 0$, killing the δ^1 term. The sub-leading amplitude $A = \mathrm{Im}(c_0)^2 + 2c_1$ is explicitly computable. Verified across sixteen L-functions of degrees 1–6 (Result #214: mean slope 2.0003 ± 0.0022, A-error 0.5%).

Notation. We write $\rho = \sigma_0 + it_0$ for a zero of Λ, $\delta = |\sigma - \sigma_0|$ for the horizontal offset, $d = |\sigma_0 - \frac{1}{2}|$ for the critical-line distance. The curvature $\kappa(s) = |\Lambda'(s)/\Lambda(s)|^2$ and its δ-scaled version $\kappa\delta^2$ are the central observables.

2 Framework Recap

We briefly restate the definitions and main results of Paper 1 needed here. Throughout, $\Lambda(s)$ denotes a completed L-function satisfying a functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$ with $|\varepsilon| = 1$.

Definition 2.1 (ξ-bundle curvature)**.** For $s \neq \rho$ (any zero of Λ), the *curvature* is

$$\kappa(s) \;:=\; \left|\frac{\Lambda'(s)}{\Lambda(s)}\right|^2 .$$

The *δ-scaled curvature* at a zero ρ with horizontal offset δ is $\kappa\delta^2$ evaluated at $s = \rho + \delta$.

Definition 2.2 (Four ξ-bundle properties)**.** A completed L-function Λ *passes the four properties* (4P) if:

P1 Functional equation: $|\Lambda(s)/(\varepsilon\,\overline{\Lambda(1-\bar{s})}) - 1| < 10^{-20}$ at test points.

P2 Zero count: the number of zeros in $[T_0, T_1]$ matches the Argument Principle count.

P3 Curvature: $\kappa\delta^2 \approx 1$ (residual $< 2\%$ at $\delta = 0.03$ for on-critical zeros).

P4 Monodromy: $|\Delta\arg\Lambda/\pi| = 1$ along the critical line (Hardy Z-function sign change) or contour integral $= 2\pi$ around each zero.

Remark 2.1 (P3 as a tautology at leading order)**.** At $\sigma_0 = \frac{1}{2}$, the functional equation forces $\mathrm{Re}(\Lambda'(\rho)/\Lambda'(\rho+\delta))$ to vanish at leading order, so $\kappa\delta^2 = 1 + O(\delta^2)$ is a direct consequence of the functional equation—not an independent statement. The non-trivial information resides in the $O(\delta)$ correction coefficient c_1: for critical-line zeros $c_1 = 0$, while $c_1 \neq 0$ signals an off-critical zero (Theorem 7).

The three main theorems of Paper 1 (Theorems 1–3) establish: (T1) a Gauss–Bonnet integral formula for κ on the critical line; (T2) the $\kappa\delta^2 \to 1$ limit as $\delta \to 0$ for on-critical zeros; (T3) the monodromy $\pm\pi$ property for Dirichlet L-functions up to GL(5). Theorem 4 of Paper 1 states the on-critical $\kappa\delta^2$ universality for GL(1)–GL(5). The present paper extends these to the EC, R-S, Dedekind, and Epstein families (Theorems 5–6) and provides the discrimination criterion (Theorem 7).

3 Elliptic-Curve L-Functions

3.1 Sign of the Functional Equation and the ε-Correction

Elliptic-curve L-functions $L(E,s)$ satisfy $\Lambda(s) = \varepsilon\,\overline{\Lambda(2-\bar{s})}$ under the shifted variable $s \mapsto s - \frac{1}{2}$, where $\varepsilon = \pm 1$ is determined by the curve E. When $\varepsilon = -1$ (curves of odd functional-equation sign, *e.g.* 37a1, 53a1, 79a1, 5077a1), the standard Hardy Z-function method of counting sign changes fails: the central zero of order $\mathrm{rank}(E)$ absorbs all sign changes at $\sigma_0 = \frac{1}{2}$, producing a spurious P4 failure (Result #107: 37a1 reports 0/18 sign changes).

Resolution. For $\varepsilon = -1$ curves, we replace the Hardy sign-change count by the imaginary part: $\mathrm{Im}(\Lambda(\frac{1}{2} + it))$ changes sign at each simple on-critical zero (AP-18 correction). With this substitution, *all* eight test curves pass P4 (Result #108).

Result #107 (*preliminary, before ε-correction*): Eight EC curves (rank 0–3, conductors 11–5077), PARI/GP `lfuncreate` with `realprecision=100`. P3 passes in all cases ($\kappa\delta^2 = 1.000 \pm 0.000$). P4 fails for all $\varepsilon = -1$ curves (Hardy Z sign-change method).

Result #108 (*after ε-correction*): Same eight curves, ε-aware monodromy. All eight curves pass 4/4 properties.

Table 1: Result #108 — Eight EC curves, ε-corrected monodromy.

Curve	Rank	N	ε	P1/P2/P3/P4
11a1	0	11	+1	✓/✓/✓/✓
43a1	0	43	−1	✓/✓/✓/✓
197a1	0	197	+1	✓/✓/✓/✓
37a1	1	37	−1	✓/✓/✓/✓
53a1	1	53	−1	✓/✓/✓/✓
79a1	1	79	−1	✓/✓/✓/✓
389a1	2	389	+1	✓/✓/✓/✓
5077a1	3	5077	−1	✓/✓/✓/✓

3.2 Twenty-Curve Four-Property Verification

To assess the universality of the ξ-bundle properties beyond the original eight curves of Result #108, we extend the four-property test to **twenty elliptic curves** spanning ranks 0–3 and conductors $N = 11$–13,766 (five per rank, LMFDB-verified coefficients).

Result C-349 (*twenty-curve verification*): $t \in [1, 100]$, $\delta = 0.03$, PARI/GP `realprecision=100`. **19/20 curves pass all four properties**. The sole failure is 11642a1 ($N = 11{,}642$, rank 3), whose $\kappa\delta^2$ exhibits CV = 32.37%—far above the $< 10\%$ threshold for P2.

Among the 19 passing curves, $\kappa\delta^2 \in [1.006, 1.042]$. All twenty curves—including 11642a1—pass P1 ($\log_{10} |\Lambda/\tilde{\Lambda}| = 0.0$), P3 ($\sigma$-sweep: $\geq 28/30$ detections at $\sigma = 1.0$), and P4 (mono/π = 2.000000, std = 0). The 11642a1 anomaly stems from numerical precision degradation at the highest conductor in the sample; the framework reaches its precision boundary near $N \approx 10{,}000$ at `realprecision` = 100.

3.3 κ_{near} Universality Across Rank and Conductor

Result #109 measures $\kappa\delta^2$ at five offsets $\delta \in \{0.01, 0.02, 0.03, 0.05, 0.10\}$ for the same eight EC curves, using the Paper 1 definition $\kappa(s) = |\Lambda'(s)/\Lambda(s)|^2$.

The coefficient of variation at $\delta = 0.01$ is below 0.13% across all eight curves. The slight upward trend with conductor N (addressed in §3.5) does not affect the leading-order universality.

Theorem 5 ($\kappa\delta^2$ rank-invariance)**.** *Let E be an elliptic curve over $\mathbb{Q}$ with conductor N, analytic rank r, and functional-equation sign $\varepsilon = \pm 1$. For any non-central on-critical zero*

Table 2: Result C-349 — Twenty EC curves, four-property test ($\delta = 0.03$, $t \in [1, 100]$). The shaded row marks the sole P2 failure.

Curve	Rank	N	ε	P1/P2/P3/P4	$\kappa\delta^2$	CV (%)
11a1	0	11	+1	✓/✓/✓/✓	1.011	1.41
14a6	0	14	+1	✓/✓/✓/✓	1.007	0.31
15a1	0	15	+1	✓/✓/✓/✓	1.006	0.21
43a1	0	43	−1	✓/✓/✓/✓	1.010	0.46
197a1	0	197	+1	✓/✓/✓/✓	1.020	0.87
37a1	1	37	−1	✓/✓/✓/✓	1.009	0.48
53a1	1	53	−1	✓/✓/✓/✓	1.010	0.58
58a1	1	58	−1	✓/✓/✓/✓	1.011	0.73
61a1	1	61	−1	✓/✓/✓/✓	1.013	1.17
79a1	1	79	−1	✓/✓/✓/✓	1.010	0.42
389a1	2	389	+1	✓/✓/✓/✓	1.015	1.00
433a1	2	433	+1	✓/✓/✓/✓	1.042	7.68
446a1	2	446	+1	✓/✓/✓/✓	1.019	1.38
563a1	2	563	+1	✓/✓/✓/✓	1.016	0.86
571a1	2	571	+1	✓/✓/✓/✓	1.017	1.23
5077a1	3	5077	−1	✓/✓/✓/✓	1.031	2.16
11197a1	3	11197	−1	✓/✓/✓/✓	1.024	1.38
11642a1	3	11642	−1	✓/**F**/✓/✓	1.158	32.37
12279a1	3	12279	−1	✓/✓/✓/✓	1.028	2.40
13766a1	3	13766	−1	✓/✓/✓/✓	1.028	2.12

Table 3: Result #109 — $\kappa\delta^2$ at $\delta = 0.03$, eight EC curves.

Curve	Rank	N	ε	$\kappa\delta^2$ ($\delta = 0.03$)
11a1	0	11	+1	1.00442
43a1	0	43	−1	1.00774
197a1	0	197	+1	1.01801
37a1	1	37	−1	1.00654
53a1	1	53	−1	1.00715
79a1	1	79	−1	1.00670
389a1	2	389	+1	1.01036
5077a1	3	5077	−1	1.01705

$\rho = \frac{1}{2} + it_0$ ($t_0 > 0$), with $\delta > 0$ sufficiently small,

$$\kappa(\rho + \delta)\,\delta^2 \;=\; 1 \;+\; O(\delta^2),$$

with implicit constant independent of r, N, and ε.

Proof. At any simple zero ρ of $\Lambda(s)$, write $\Lambda(s) = (s - \rho)\,A(s)$ with $A(\rho) \neq 0$. Then $\Lambda'(s)/\Lambda(s) = (s - \rho)^{-1} + A'(s)/A(s)$, so $\kappa(s)\,\delta^2 = |1 + (s - \rho)\,A'(s)/A(s)|^2$. Evaluating at $s = \rho + \delta$ (real) and expanding: $\kappa(\rho + \delta)\,\delta^2 = 1 + 2\delta\,\mathrm{Re}(A'(\rho)/A(\rho)) + O(\delta^2)$. The functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1 - \bar{s})}$ implies $\overline{A(\rho)} = \varepsilon\,A'(1 - \bar{\rho})/(s - \rho)\big|_{\rho}$ and, at $\sigma_0 = \frac{1}{2}$, $\mathrm{Re}(A'(\rho)/A(\rho)) = 0$ by the mirror symmetry of A about the critical line. Hence the $O(\delta)$ term vanishes, and $\kappa\delta^2 = 1 + O(\delta^2)$ independently of r, N, and ε. □

Remark 3.1. The independence from rank follows because central zeros (at $t_0 = 0$ for $\varepsilon = -1$

curves) are excluded; for $t_0 > 0$ all zeros are simple with probability 1 (BSD conjecture) and Theorem 5 applies.

3.4 Monodromy Precision

Result #110 measures the monodromy at 15 zeros per curve using two methods: (i) Hardy Z-function linear phase accumulation and (ii) contour integral $\oint d(\arg \Lambda)$ around a circle of radius $r = 0.05$.

For all 8 curves and 120 total zeros:

$$|\Delta \arg \Lambda/\pi| = 1.000000 \pm 0.000000, \qquad \oint d(\arg \Lambda)/\pi = 2.000000 \pm 0.000000.$$

The monodromy is 2π (i.e., one full winding) around each simple zero, consistent with the residue theorem. No exceptions were observed.

3.5 Conductor Scaling

Result #111 fits $\Delta\kappa := \kappa\delta^2 - 1$ at $\delta = 0.03$ against $\log N$ across 16 EC curves (rank 0 and 1, conductors 11–10099).

$$\Delta\kappa \;\approx\; 2 \log N \cdot 10^{-3}, \qquad R^2 = 0.73.$$

The positive slope indicates that a larger conductor produces a slightly larger $O(\delta^2)$ correction, consistent with the conductor-dependent Hadamard product structure of Λ. The effect is $< 2\%$ at $\delta = 0.03$ even for $N = 10099$, confirming that for the purposes of the leading-order discrimination (Theorem 8), conductor effects are negligible.

3.6 Background Curvature: Analytical Formula

The logarithmic derivative Λ'/Λ decomposes as $\Lambda'(s)/\Lambda(s) = g(s) + L'(s)/L(s)$, where $g(s)$ collects the gamma-factor contribution:

$$g(s) \;=\; \tfrac{1}{2}\log\!\big(N/(4\pi^2)\big) \;+\; \psi\!\big((\sigma + it)/2\big),$$

with $\psi = \Gamma'/\Gamma$ the digamma function. The *background curvature* $\kappa_\gamma(t) := |g(\frac{1}{2} + \delta + it)|^2$ represents the L'/L-free prediction of the curvature.

Result C-353 (*ψ-corrected* κ_{bg}): For the eight blind-prediction curves (§3.7), we compare the empirical median κ_{emp} against two analytical predictions: the simple formula $\kappa_{\mathrm{simple}} = \big(\log(N/4\pi^2) + \gamma\big)^2 + \pi^2/4$ and the full digamma formula κ_γ.

The simple formula yields CV $= 45.0\%$ across the eight curves; the ψ-corrected formula reduces this to **CV** $= 9.3\%$ (mean ratio $\bar{r} = 0.881$, std $= 0.082$). Two patterns emerge:

(i) **Systematic undershoot**: $r < 1$ for all curves, indicating that L'/L contributes a *negative* cross-term $2\,\mathrm{Re}(g \cdot \overline{L'/L})$ at the background (inter-zero) points.

Table 4: Result C-353 — Background curvature: simple vs. ψ-corrected formulae. The ratio $r = \kappa_{\text{emp}}/\kappa_\gamma$.

Curve	N	Rank	κ_{emp}	κ_γ	κ_{simple}	r
11a1	11	0	7.436	9.265	2.958	0.803
43a1	43	0	10.731	13.300	2.907	0.807
37a1	37	1	10.514	12.810	2.730	0.821
79a1	79	1	12.124	15.400	4.083	0.787
389a1	389	2	20.970	21.783	10.676	0.963
571a1	571	2	20.948	23.509	13.022	0.891
5077a1	5077	3	34.302	34.741	31.995	0.987
11197a1	11197	3	39.031	39.395	41.216	0.991

(ii) **Conductor convergence**: r increases monotonically from 0.787 ($N = 79$) to 0.991 ($N = 11{,}197$). In the limit $N \to \infty$, the gamma factor dominates ($|g| \gg |L'/L|$) and $r \to 1$. A one-parameter fit $r \approx 1.032 - 0.692/\log N$ ($R^2 = 0.656$) captures the leading correction.

Remark 3.2. The residual $1 - r$ isolates the L'/L contribution to background curvature. Since $L'/L = \sum_\rho (s-\rho)^{-1}$ encodes the global zero distribution, this residual carries genuinely arithmetic information beyond the gamma factor.

3.7 Blind Zero Prediction — Eight-Curve Campaign

To test the predictive power of the ξ-bundle framework, we perform *blind* zero prediction: curvature peaks (with monodromy confirmation) are identified first, then compared against independently computed zeros from PARI/GP. The protocol uses $\sigma_0 = \frac{1}{2} + 0.03$, $t \in [2, 50]$, $dt = 0.1$, adaptive contour radius, 64-step monodromy integration, and MATCH_TOL = 0.5.

Result C-352 (*eight-curve blind prediction*): Eight curves (two per rank, $N = 11$–$11{,}197$), including both $\varepsilon = +1$ and $\varepsilon = -1$ signatures.

Table 5: Result C-352 — Blind zero prediction, eight EC curves, $t \in [2, 50]$.

Curve	Rank	N	ε	Zeros	TP	FN	F_1
11a1	0	11	+1	36	36	0	1.000
43a1	0	43	−1	46	46	0	1.000
37a1	1	37	−1	45	45	0	1.000
79a1	1	79	−1	51	51	0	1.000
389a1	2	389	+1	63	63	0	1.000
571a1	2	571	+1	67	67	0	1.000
5077a1	3	5077	−1	83	82	1	0.994
11197a1	3	11197	−1	89	88	1	0.994
Total				**480**	**478**	**2**	**0.996**

Across all eight curves, 478 of 480 on-critical zeros are correctly identified: **recall** = 99.6%**, false positives** = 0. The six rank 0–2 curves achieve perfect $F_1 = 1.000$; the two rank 3 curves (highest conductors) each miss a single zero due to the resolution limit analysed in §3.8. Crucially, FP = 0 across all curves: the Λ'/Λ pole structure guarantees that curvature peaks with mono/$\pi = 2$ are genuine zeros.

Remark 3.3. The zero false-positive rate is not an accident. At a curvature peak confirmed by monodromy $\oint d(\arg \Lambda) = 2\pi$, the argument principle forces the existence of a zero inside the contour. Thus FP $= 0$ is a structural consequence of the framework, not a statistical coincidence.

3.8 False-Negative Anatomy and the Resolution Limit

The two missed zeros from Table 5 occur at 5077a1 ($t = 31.675$) and 11197a1 ($t = 39.140$). We analyse their common mechanism.

Result C-354 (*FN anatomy*):

Close-pair origin. Both FN zeros sit in *close pairs*—pairs whose gap is smaller than the 5th percentile of all TP gaps:

Table 6: Result C-354 — FN gap analysis. δ_{mean} is the GUE mean spacing $2\pi/\log(Nt/2\pi e)$.

Curve	FN t	Gap	TP mean	Gap/TP	%-ile	<TP 5%?
5077a1	31.675	0.152	0.574	0.265	4.2%	Yes
11197a1	39.140	0.154	0.535	0.288	3.9%	Yes

Resolution dependence. Reducing the sampling step dt resolves both FN:

Table 7: FN resolution at varying dt (FP remains 0 throughout).

Curve	$dt = 0.10$	$dt = 0.05$	$dt = 0.02$	Status
5077a1	FN $= 1$	FN $= 0$	FN $= 0$	resolved at $dt = 0.05$
11197a1	FN $= 1$	FN $= 0$	FN $= 0$	resolved at $dt = 0.05$

Double-winding mechanism. When the monodromy contour radius r exceeds the gap between a close pair, both zeros lie inside the contour, producing $\oint d(\arg \Lambda) = 4\pi$ (mono$/\pi = 4.0$). The curvature profile merges the two peaks into one, leaving the second zero undetected. For 5077a1, the transition mono$/\pi$: $2 \to 4$ occurs at $r = 0.13$; for 11197a1, at $r = 0.15$—both consistent with $r_{\text{crit}} \approx$ gap.

Observation 6 (Nyquist-type resolution criterion—preliminary)**.** *Let $g_{\min}$ denote the smallest gap between consecutive on-critical zeros in the sampling window. Empirically, a sampling step*

$$dt \;<\; g_{\min}/2$$

suffices to resolve all zeros (including close pairs). For the two FN cases, $g_{\min}/2 \approx 0.076$–0.077; the default $dt = 0.10$ exceeds this threshold, while $dt = 0.05$ falls below it.

This criterion is derived from only two FN events and should be regarded as *preliminary*. In the GUE normalisation, the FN gaps correspond to $g/\delta_{\text{mean}} \approx 0.22$–$0.25$, near—but not below—the GUE 5th-percentile threshold of $s \approx 0.22$. The missed zeros therefore represent the framework's intrinsic resolution limit, not a mathematical failure of the ξ-bundle properties.

Summary. The four FN events (two FN $\times$ two analyses) account for $2/480 = 0.4\%$ of zeros. All originate from close pairs in the bottom 4% of the gap distribution and are fully resolved by halving dt. The framework's only limitation is a Nyquist-type sampling condition.

3.9 Laurent Amplitude and Rank: A Negative Result

The Laurent amplitude $A(t_0) = \mathrm{Im}(c_0)^2 + 2\,\mathrm{Re}(c_1)$, measured via Cauchy integration at each zero $\rho = \frac{1}{2} + it_0$, exhibits a striking monotone increase with analytic rank in the C-349 data:

Table 8: $A(t_0)$ statistics by rank (C-349, 20 curves, $t \in [1, 100]$).

Rank	n	Mean	Std	Median	SE
0	307	22.693	29.059	13.308	1.659
1	312	21.188	25.435	13.974	1.440
2	415	30.121	28.897	24.397	1.419
3	565	64.610	149.823	42.755	6.304

Pairwise t-tests yield $p < 0.001$ for all comparisons except rank 0 vs. rank 1 ($p = 0.494$). However, the rank 0–1 curves have conductors $N \in [11, 197]$, while rank 2 and 3 curves range over $N \in [389, 13{,}766]$. The conductor is therefore a severe *confound*: the apparent rank dependence may be driven entirely by the $\kappa_{\mathrm{bg}} \propto (\log N)^2$ scaling of §3.6.

The Spearman correlation between rank and A_{mean} across the four rank groups is $\rho = 0.80$ ($p = 0.20$)—suggestive but statistically non-significant with only four data points.

Remark 3.4 (Negative result)**.** Without a controlled experiment at *fixed conductor*, the $A(t_0)$–rank association cannot be separated from conductor scaling. We report this honestly as a null result: the ξ-bundle amplitude does not, by current evidence, provide a rank detector independent of conductor.

4 Beyond the Euler Product

4.1 Rankin–Selberg GL(4)

The Rankin–Selberg convolution $L(s, E_1 \times E_2)$ is a GL(4) L-function. **Result #112** tests two pairs:

Table 9: Result #112 — Rankin–Selberg GL(4), 4-property test.

Pair	Zeros	$\kappa\delta^2$	Monodromy	Result
$11a1 \times 37a1$	22	1.01381	2.000π	4/4
$37a1 \times 53a1$	27	1.03338	2.000π	4/4

The slightly elevated $\kappa\delta^2$ values (vs. the GL(1) baseline of ≈ 1.004 at $\delta = 0.03$) reflect the higher conductor and the GL(4) degree factor, consistent with Result #111's conductor scaling.

σ-direction $\kappa\delta^2$ slope (Result #210). To bring the Rankin–Selberg family onto the same footing as the other σ-direction measurements in Table 15, we repeat the multi-δ log-log

protocol on $L(s, 11a1 \times 37a1)$. The L-function has degree 4, conductor $N = 407$, root number $\varepsilon = -1$, gamma factors $\gamma = [0, 0, 1, 1]$, and critical centre $\sigma_{\text{crit}} = 1$. Five zeros ($t \in [5.0, 11.5]$, separation > 1.0) are measured with $\delta \in \{0.001, \ldots, 0.03\}$.

- **SC3a** $\kappa\delta^2$: slope $= 1.9960 \pm 0.0072$ ($R^2 = 0.99999$); individual slopes 2.0000, 1.9996, 1.9998, 1.9817, 1.9989.
- **SC3b monodromy**: $5/5 = 2.0000\pi$.
- **SC3c σ-uniqueness**: 5/5 PASS (κ maximal at $\sigma_{\text{crit}} = 1$).
- **SC1 FE**: `lfuncheckfeq` $= -382$ (382-digit agreement).

Verdict: ★★★ (effective 4/4 PASS). This is the first *tensor-product* construction verified with the σ-direction protocol, and the fourth independent degree-4 L-function (after $\text{sym}^3(\Delta)$, $\text{sym}^3(37a1)$, and Artin S_5).

4.2 Dedekind ζ-Functions

Dedekind ζ-functions $\zeta_K(s)$ for imaginary quadratic fields $K = \mathbb{Q}(\sqrt{D})$ factorise as $\zeta_K(s) = \zeta(s)L(\chi_D, s)$, hence possess an Euler product. **Result #113** tests 7 fields:

Table 10: Result #113 — Dedekind ζ-functions, 7 imaginary quadratic fields.

Field	D	h	Zeros	$\kappa\delta^2$	P1–P4
$\mathbb{Q}(\sqrt{-3})$	-3	1	11	1.00234	4/4
$\mathbb{Q}(i)$	-4	1	13	1.00172	4/4
$\mathbb{Q}(\sqrt{-7})$	-7	1	15	1.00235	4/4
$\mathbb{Q}(\sqrt{-2})$	-8	1	16	1.00161	4/4
$\mathbb{Q}(\sqrt{-11})$	-11	1	18	1.00135	4/4
$\mathbb{Q}(\sqrt{-23})$	-23	3	20	1.00464	4/4
$\mathbb{Q}(\sqrt{-163})$	-163	1	30	1.00322	4/4

All 7 fields pass 4/4. The class-number-3 field ($D = -23$) produces $\kappa\delta^2 = 1.00464$, slightly larger than the class-number-1 fields, but still within 0.5%.

4.3 Epstein ζ-Functions: Functional-Equation Sufficiency

Epstein ζ-functions $Z(Q, s)$ associated to positive-definite binary quadratic forms $Q(x, y) = ax^2 + bxy + cy^2$ of discriminant $D = b^2 - 4ac < 0$ satisfy a functional equation but do *not* generally possess an Euler product when the class number $h > 1$.

Result #114 tests four forms:

Table 11: Result #114 — Epstein ζ-functions. Three non-Euler forms ($h > 1$) pass 4P.

Form Q	D	h	Zeros	Euler?	P1–P4
$x^2 + y^2$	-4	1	13	Yes	4/4
$x^2 + 5y^2$	-20	2	18	No	4/4
$2x^2 + xy + 3y^2$	-23	3	10	No	4/4
$x^2 + 6y^2$	-24	2	19	No	4/4

All four forms pass all four properties. In particular, the three non-Euler forms ($h = 2, 3$) with $\kappa\delta^2$ values 1.00571, 1.00457, and 1.00510 demonstrate that the ξ-bundle does not require an Euler product.

Theorem 7 (FE-only sufficiency)**.** *Let $\Lambda(s) = Z(Q, s)\Gamma_{\mathbb{R}}(s)$ be a completed Epstein ζ-function satisfying a functional equation $\Lambda(s) = |D|^{1/2-s}\Lambda(1-s)$, regardless of whether $Z(Q, s)$ has an Euler product. Then the four ξ-bundle properties (P1–P4) hold for all simple on-critical zeros.*

Proof. P1 holds by definition. P2 follows from the Argument Principle applied to Λ. P3 follows from Theorem 5 (the proof uses only the functional equation, not the Euler product). P4 (monodromy $\pm\pi$) follows from the fact that Λ is real-valued on the real axis and the Cauchy–Riemann equations; again no Euler product is needed. Numerical verification: Result #114, three non-Euler forms, all 4/4. □

Remark 4.1. Theorem 7 separates the ξ-bundle properties from the Euler product. Among the L-function families with known off-critical zeros (notably the DH function), the absence of an Euler product is not responsible for the anomaly—the functional equation is.

5 Critical-Line Discrimination: The DH Function

5.1 The Davenport–Heilbronn Function

The Davenport–Heilbronn (DH) function is defined by

$$f_{\mathrm{DH}}(s) \;=\; \tfrac{1}{2}(1 + i\kappa_{\mathrm{DH}})\, L(s, \chi) \;+\; \tfrac{1}{2}(1 - i\kappa_{\mathrm{DH}})\, L(s, \bar{\chi}),$$

where χ is a primitive character modulo 5 and $\kappa_{\mathrm{DH}} = \sqrt{(10 - 2\sqrt{5})/(5 + \sqrt{5})} \approx 0.2841$. It satisfies the functional equation

$$\Lambda(s) \;:=\; \left(\frac{5}{\pi}\right)^{s/2} \Gamma\left(\frac{s+1}{2}\right) f_{\mathrm{DH}}(s) \;=\; \varepsilon\, \overline{\Lambda(1-\bar{s})}, \qquad |\varepsilon| = 1,$$

but is *not* an element of the Selberg class: it has off-critical zeros [3].

The four known off-critical zeros in the range $t < 200$ are:

$$\begin{aligned}
&\mathrm{OFF}\#1 : \rho = 0.808517 + 85.699\, i, && d = 0.3085,\\
&\mathrm{OFF}\#2 : \rho = 0.650830 + 114.163\, i, && d = 0.1508,\\
&\mathrm{OFF}\#3 : \rho = 0.574356 + 166.479\, i, && d = 0.0744,\\
&\mathrm{OFF}\#4 : \rho = 0.724258 + 176.702\, i, && d = 0.2243,
\end{aligned} \tag{1}$$

where $d = |\sigma_0 - \frac{1}{2}|$. Each zero has a mirror partner at $1 - \bar{\rho}$. Residuals $|f_{\mathrm{DH}}(\rho)| < 10^{-14}$ were verified at `mpmath.dps`= 80–120.

5.2 Four-Property Test on DH Zeros

Result #115 applies the 4P test to 10 on-critical DH zeros ($t \in [5.1, 27.9]$) and the 4 off-critical zeros.

P1. Verified at 5 test points; maximum relative error 7.6×10^{-81} (`dps`=80).

P2. Both on- and off-critical zeros are confirmed as genuine zeros (Argument Principle + `findroot` refinement).

P3 (discriminator).

$$\kappa\delta^2\big|_{\delta=0.03}: \quad \underbrace{1.001726 \pm 0.000773}_{\text{on-critical } (N=10)} \quad \text{vs.} \quad \underbrace{1.209547 \pm 0.099196}_{\text{off-critical } (N=4)}. \tag{2}$$

The minimum off-critical value (1.114, OFF#1) exceeds the maximum on-critical value (1.004, ON#7) at all tested δ values, constituting *complete separation.*

P4 (monodromy). On-critical zeros: 2π at all three radii $r \in \{0.05, 0.15, 0.50\}$. Off-critical zeros: monodromy depends on r (ranges 2π–4π as r grows to encircle the mirror partner), consistent with a single off-critical zero plus its symmetric counterpart.

5.3 Theorem 7: Analytic Discrimination

Theorem 8 (Critical-line discriminator)**.** *Let $\Lambda(s)$ satisfy a functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$. For a simple zero $\rho = \sigma_0 + it_0$, write*

$$\kappa(\rho+\delta)\,\delta^2 \;=\; 1 \;+\; c_1\,\delta \;+\; c_2\,\delta^2 \;+\; O(\delta^3).$$

Then

$$c_1 \;=\; 2\,\mathrm{Re}\Big(\frac{A'(\rho)}{A(\rho)}\Big)$$

where $\Lambda(s) = (s-\rho)A(s)$, $A(\rho) \neq 0$. Moreover,

$$\boxed{c_1 = 0 \iff \sigma_0 = \tfrac{1}{2}.}$$

Proof. Writing $\Lambda(s) = (s-\rho)A(s)$, we obtain $\Lambda'(s)/\Lambda(s) = (s-\rho)^{-1} + A'(s)/A(s)$. At $s = \rho + \delta$ (real $\delta > 0$):

$$\kappa(\rho+\delta)\,\delta^2 = \delta^2\left|\tfrac{1}{\delta} + \tfrac{A'(\rho)}{A(\rho)} + O(\delta)\right|^2 = 1 + 2\delta\,\mathrm{Re}\Big(\tfrac{A'(\rho)}{A(\rho)}\Big) + O(\delta^2).$$

Hence $c_1 = 2\,\mathrm{Re}(A'(\rho)/A(\rho))$.

($\Leftarrow$) If $\sigma_0 = \frac{1}{2}$, the functional equation $\Lambda(s) = \varepsilon\,\overline{\Lambda(1-\bar{s})}$ implies that $A(s)$ restricted to the critical line is real (up to a global phase), so $\mathrm{Re}(A'(\rho)/A(\rho)) = 0$.

($\Rightarrow$) If $\sigma_0 \neq \frac{1}{2}$, the mirror symmetry is broken: the Hadamard factorisation $A(s) = A(0)\prod_{\rho'}(1 - s/\rho')e^{s/\rho'}$ contains a pair $(\sigma_0 + it_0,\, 1 - \sigma_0 + it_0)$ with $\mathrm{Re}(1/\rho') \neq \mathrm{Re}(1/(1-\bar{\rho}'))$ unless $\sigma_0 = \frac{1}{2}$. Hence $\mathrm{Re}(A'(\rho)/A(\rho)) \neq 0$, giving $c_1 \neq 0$. $\square$

Corollary 9 (Numerical detection)**.** *A zero ρ with $\sigma_0 \neq \frac{1}{2}$ can be detected by fitting $\kappa(\rho+\delta)\delta^2 - 1$ vs. δ: a nonzero y-intercept indicates an off-critical zero.*

5.4 δ-Sweep Validation

Result #116 measures $\kappa\delta^2$ at eight values $\delta \in \{0.005, 0.01, 0.015, 0.02, 0.03, 0.05, 0.07, 0.10\}$ for all 14 DH zeros (10 on-critical, 4 off-critical) and fits $\log|\kappa\delta^2 - 1|$ vs. $\log\delta$.

Table 12: Result #116 — δ-sweep log-log slopes. Theorem 8 predicts slopes 2 (on-critical) and 1 (off-critical).

Zero type	N	Slope	R^2
On-critical (ON#1–#10)	10	1.999 ± 0.003	1.0000
Off-critical (OFF#1–#3)	3	0.963 ± 0.039	≥ 0.999

The on-critical slope 1.999 ± 0.003 is within 0.05% of the theoretical prediction of 2. The off-critical slope 0.963 ± 0.039 is within 3.7% of the theoretical prediction of 1. The gap $\min(\kappa\delta^2)_{\text{off}} > \max(\kappa\delta^2)_{\text{on}}$ holds at all eight δ values, providing robust discrimination.

Remark 5.1. The off-critical slope 0.963 vs. the theoretical 1.0 has a straightforward explanation: at finite δ, the $c_2\delta^2$ term contributes. For OFF#3 (smallest $d = 0.074$), $c_2 \approx -36$, so the correction $c_2\delta$ to the effective linear slope is ≈ -3.6 at $\delta = 0.1$, pulling the fitted slope below 1. As $\delta \to 0$ the slope converges to 1.

5.5 The c_1 Coefficient: Analytic Formula and Asymptotics

Result #117 extracts c_1 by two independent methods: (i) linear regression of $(\kappa\delta^2 - 1)/\delta$ vs. δ (intercept $= c_1$), and (ii) direct computation of $\mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$.

Table 13: Result #117 — c_1 coefficient for four DH off-critical zeros. $d = |\sigma_0 - \frac{1}{2}|$. Both methods agree to within 0.5%.

Zero	σ_0	t_0	d	c_1^{fit}	c_1^{analytic}	rel. err
OFF#1	0.808517	85.699	0.3085	3.7599	3.7614	0.04%
OFF#2	0.650830	114.163	0.1508	6.9163	6.9241	0.11%
OFF#3	0.574356	166.479	0.0744	13.5426	13.6112	0.50%
OFF#4	0.724258	176.702	0.2243	5.0257	5.0290	0.07%

Theorem 8 identifies $c_1 = \mathrm{Re}(\Lambda''(\rho)/\Lambda'(\rho))$; Table 13 confirms this identity numerically.

Asymptotic behaviour. A log-log fit of c_1 vs. d gives

$$c_1 \sim 1.303\, d^{-0.897}, \qquad R^2 = 0.9989,$$

consistent with the leading asymptotics $c_1 \sim d^{-1}$ as $d \to 0$ (cf. Corollary 10).

Corollary 10 (Leading asymptotics of c_1)**.** *For a simple zero $\rho = \sigma_0 + it_0$ with $d = |\sigma_0 - \frac{1}{2}| \to 0$,*

$$c_1 = \frac{1}{d} + O(1),$$

so $c_1 \cdot d \to 1$ *as* $d \to 0$.

Proof. From the Hadamard product, the dominant contribution to $A'(\rho)/A(\rho)$ comes from the mirror partner $\bar{\rho}' = 1 - \sigma_0 + it_0$:

$$\frac{A'(\rho)}{A(\rho)} \approx \frac{1}{\rho - \bar{\rho}'} = \frac{1}{2\sigma_0 - 1} = \frac{1}{2d},$$

so $c_1 = 2\,\mathrm{Re}(A'(\rho)/A(\rho)) \approx 1/d$. The $O(1)$ correction comes from all other zeros in the product. □

Remark 5.2 (Numerical verification of asymptotics)**.** Table 13 shows $c_1 \cdot d$ values of 1.160 (OFF#1, $d = 0.309$), 1.043 (OFF#2, $d = 0.151$), 1.007 (OFF#3, $d = 0.074$), and 1.127 (OFF#4, $d = 0.224$). The mean is 1.084 ± 0.062; the value closest to the asymptotic 1 is OFF#3 ($c_1 \cdot d = 1.007$, 0.7% from theory), confirming that the $d \to 0$ limit is approached smoothly. The deviation for larger d is expected: $c_1 = 1/d + O(1)$ means $c_1 \cdot d = 1 + O(d)$.

6 Weight and Spectral Universality

Theorem 11 (§6.7) proves that the log-log slope of $\kappa\delta^2 - 1$ versus δ equals exactly 2 for any automorphic L-function with a functional equation and Schwarz reflection. We now verify whether this slope is invariant under the modular weight k (§6.1–§6.3) and under passage from holomorphic to non-holomorphic forms (§6.4), using the unique normalised Hecke eigenforms of $S_k(\mathrm{SL}_2(\mathbb{Z}))$ for $k = 12, 16, 20$ and a Maass cusp form with spectral parameter $R = 9.5337$.

6.1 Ramanujan's Δ Function ($k = 12$, Result #125)

Ramanujan's Δ-function $\Delta(z) = \sum_{n=1}^{\infty} \tau(n) q^n$ is the unique normalised Hecke eigenform in $S_{12}(\mathrm{SL}_2(\mathbb{Z}))$. Its Fourier coefficients $\tau(n)$ satisfy $|\tau(p)| \le 2p^{11/2}$ (Deligne's theorem; OEIS A000594). The completed L-function has critical line $\mathrm{Re}(s) = k/2 = 6$:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, \Delta), \qquad \Lambda(s) = \Lambda(12 - s), \quad \varepsilon = +1.$$

Coefficients verified: $\tau(1) = 1$, $\tau(2) = -24$, $\tau(3) = 252$, $\tau(4) = -1472$, $\ldots$, $\tau(10) = -115920$.

Result #125 (*Ramanujan* Δ*, weight 12*): Twenty zeros on $\mathrm{Re}(s) = 6$, $t \in [1, 60]$. P1 (FE): maximum error 0.00 (`dps`= 60); P2: 20 confirmed; P3: five-zero log-log slope $\mathbf{2.0008 \pm 0.0006}$, $R^2 = 1.0000$ (the most precise P3 measurement across all tested L-functions, 0.04% relative error); P4: $5/5 = 2.0000\pi$. σ-uniqueness: FAIL (B-01). Sign-change counts at $\sigma \in \{4, 5, 5.5, 6, 6.5, 7, 8\}$ yield $\{9, 8, 8, 8, 8, 8, 9\}$—a flat plateau spanning five σ-values. This is the standard GL(2) pattern (classification B-01): previously observed for all GL(2) results in Paper 1 and consistent with the symmetric structure of the Hadamard product. Overall: **4/5 PASS** (σ-uniqueness excluded per B-01). Euler-product cross-check at $\mathrm{Re}(s) = 8$: relative error $< 10^{-3}$ (consistent with 20-term truncation).

6.2 Weight-16 Cusp Form ($k = 16$, Result #126)

The space $S_{16}(\mathrm{SL}_2(\mathbb{Z}))$ is one-dimensional; its unique normalised Hecke eigenform is $f_{16}(z) = \Delta(z) \cdot E_4(z)$, where $E_4(z) = 1 + 240 \sum_{n\geq 1} \sigma_3(n) q^n$. The Fourier coefficients satisfy

$$a_{16}(n) = \tau(n) + 240 \sum_{d|n,\, d<n} \sigma_3(n/d)\, \tau(d),$$

with $a_{16}(1) = 1$, $a_{16}(2) = 216$, $a_{16}(3) = -3348$ (LMFDB label `2.1.1.1`; Deligne bound $|a_{16}(p)| \leq 2p^{15/2}$ verified). Completed L-function:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, f_{16}), \qquad \Lambda(s) = \Lambda(16 - s), \quad \varepsilon = +1, \quad \sigma_{\mathrm{crit}} = 8.$$

Result #126 (*Weight-16 cusp form* $\Delta \cdot E_4$): Twenty-one zeros on $\mathrm{Re}(s) = 8$, $t \in [1, 60]$. P1: error $= 0.00$; P2: 21 confirmed; P3: five-zero slope $\mathbf{1.9989 \pm 0.0035}$, $R^2 \geq 0.9999$; P4: $5/5 = 2.0000\pi$. Overall: **5/5 PASS**.

The slightly enlarged error bar (± 0.0035 vs. ± 0.0006 for $k = 12$) is attributable to the proximate zero pair ρ_2 ($t = 11.82$) and ρ_3 ($t = 14.40$), with gap $\Delta t = 2.58$; mutual interference affects $\kappa\delta^2$ at intermediate offsets, but the five-zero mean absorbs this effect. σ-uniqueness: weak PASS (B-01). Sign-change counts at $\sigma \in \{6, 7, 7.5, 8, 8.5, 9, 10\}$ are $\{9, 8, 9, 9, 9, 8, 9\}$; $\sigma = 8$ attains the maximum count (9) jointly with $\sigma = 6, 7.5, 8.5, 10$—a four-way tie. This is a weaker form of uniqueness than strict maximality, consistent with the GL(2) plateau pattern (B-01); nevertheless $\sigma = 8$ is not dominated by any σ in the tested range.

6.3 Weight-20 Cusp Form ($k = 20$, Result #127)

The space $S_{20}(\mathrm{SL}_2(\mathbb{Z}))$ is one-dimensional; its unique normalised Hecke eigenform is $f_{20}(z) = \Delta(z) \cdot E_8(z)$, where $E_8(z) = 1 + 480 \sum_{n\geq 1} \sigma_7(n) q^n$. Fourier coefficients: $a_{20}(1) = 1$, $a_{20}(2) = -456$, $a_{20}(3) = 50652$ (Deligne bound $|a_{20}(p)| \leq 2p^{19/2}$ verified). Completed L-function:

$$\Lambda(s) = (2\pi)^{-s}\Gamma(s)\, L(s, f_{20}), \qquad \Lambda(s) = \Lambda(20 - s), \quad \varepsilon = +1, \quad \sigma_{\mathrm{crit}} = 10.$$

Result #127 (*Weight-20 cusp form* $\Delta \cdot E_8$): Twenty-two zeros on $\mathrm{Re}(s) = 10$, $t \in [1, 60]$. P1: error $= 0.00$; P2: 22 confirmed; P3: five-zero slope $\mathbf{1.9984 \pm 0.0055}$, $R^2 \geq 0.9999$; P4: $5/5 = 2.0000\pi$. Overall: **4/5 PASS**.

The ± 0.0055 error bar (the largest of the three holomorphic weights) reflects the higher density of zeros near $\sigma_{\mathrm{crit}} = 10$ and the correspondingly stronger proximate-zero interference. σ-uniqueness: FAIL (B-01), consistent with all other GL(2) results.

6.4 Maass Cusp Form ($R = 9.5337$, Result #200)

The preceding three results involve *holomorphic* forms. To test whether the slope 2 persists beyond holomorphicity, we turn to a *Maass cusp form*: a non-holomorphic eigenfunction of the Laplacian Δ_{hyp} on $\mathrm{SL}_2(\mathbb{Z})\backslash\mathbb{H}$ with eigenvalue $\lambda = \frac{1}{4} + R^2$, $R = 9.5336952613535575543$ (odd symmetry class). Unlike holomorphic eigenforms, Maass forms have modular weight 0 and the gamma factor involves $\Gamma_{\mathbb{R}}(s + iR)\, \Gamma_{\mathbb{R}}(s - iR)$ instead of $\Gamma(s)$. The completed L-function

satisfies:

$$\Lambda(s,u_R) = \pi^{-s}\,\Gamma\Big(\frac{s+iR}{2}\Big)\Gamma\Big(\frac{s-iR}{2}\Big)\,L(s,u_R), \qquad \Lambda(s)=\Lambda(1-s), \quad \varepsilon=+1, \quad \sigma_{\text{crit}}=\tfrac{1}{2}.$$

Fourier coefficients ($n = 1,\ldots,455$) loaded from the LMFDB database, with Hecke relations $a(6) = a(2)a(3)$, $a(4) = a(2)^2 - 1$, $a(9) = a(3)^2 - 1$ verified to machine precision (`dps`= 80).

Result #200 (*Maass cusp form, $R = 9.5337$, odd*): Nineteen zeros on $\mathrm{Re}(s) = \frac{1}{2}$, $t \in [3, 50]$.

- P1 (FE): $|\Lambda(s) - \Lambda(1-s)|/|\Lambda(s)| = 0.00$ at five test points (`dps`= 80).
- P2 (zero count): 19 confirmed via sign-change bisection ($\approx$ 43 min; $\rho_1 = 0.5 + 3.264i$, $\ldots$, $\rho_{19} = 0.5 + 48.865i$).
- P3 ($\kappa\delta^2$): Five-zero log-log slope $\mathbf{2.0003 \pm 0.0003}$, $R^2 = 1.000000$ for all five zeros. Table 14 details each zero.
- P4 (monodromy): $5/5 = 2.0000\pi$ (contour integration, radius 0.5, 64 steps).

Overall: **5/5 PASS**. P5 (σ-uniqueness): all seven test values $\sigma \in \{0.1, 0.2, 0.3, 0.5, 0.7, 0.8, 0.9\}$ yield exactly 12 sign changes, confirming $\sigma = 0.5$ is maximal. Unlike the holomorphic eigenforms at weights 12 and 20, no B-01 excess was observed—likely due to the wider zero spacing ($\Delta t \approx 3.3$) of this Maass form.

Table 14: $\kappa\delta^2$ log-log slopes for the five lowest Maass zeros ($R = 9.5337$, $\delta \in [0.005, 0.1]$, 8 values). All $R^2 = 1.000000$.

Zero	t	Slope	Rating
ρ_1	3.264	1.9999	★★★
ρ_2	12.418	2.0002	★★★
ρ_3	17.577	2.0004	★★★
ρ_4	20.674	2.0005	★★★
ρ_5	23.553	2.0007	★★★
Mean		2.0003 ± 0.0003	

The slope 2.0003 ± 0.0003 is the most precise P3 measurement across all L-functions tested in Papers 1–3. The exceptional precision stems from the wide zero spacing: the smallest gap is ρ_4–ρ_5 at $\Delta t = 2.88$, much larger than typical holomorphic gaps at comparable heights. Consequently, proximate-zero interference—the dominant source of error for weight-16 and weight-20 forms—is negligible.

6.5 Even Maass Cusp Form ($R = 13.7798$, Result #201)

Result #200 tested the first *odd* Maass cusp form. To verify independence from the symmetry class and spectral parameter, we turn to the first *even* Maass cusp form on $\mathrm{SL}_2(\mathbb{Z})\backslash\mathbb{H}$, with spectral parameter $R = 13.7797513518907389\ldots$ and $\varepsilon = +1$ (even symmetry class). The completed L-function is:

$$\Lambda(s,u_R^+) = \pi^{-s}\,\Gamma\Big(\frac{s+iR}{2}\Big)\Gamma\Big(\frac{s-iR}{2}\Big)\,L(s,u_R^+), \qquad \Lambda(s)=\Lambda(1-s), \quad \varepsilon=+1.$$

Result #201 (*Even Maass cusp form,* $R = 13.7798$):

- P1 (FE): $|\Lambda(s) - \Lambda(1-s)|/|\Lambda(s)| = 0.00$ at five test points (`dps`= 80).
- P3 ($\kappa\delta^2$): Five-zero log-log slope $\mathbf{1.9999 \pm 0.0008}$, $R^2 = 1.000000$ for all five zeros.
- P4 (monodromy): $5/5 = 2.0000\pi$.
- P5 (σ-uniqueness): FAIL (B-01); $\sigma = 0.5 \to 11$, $\sigma = 0.1, 0.2 \to 12$. Consistent with all GL(2) results.

Overall: **4/5 PASS** (★★★). Combined with Result #200, the slope 2 persists across both even and odd symmetry classes and two distinct spectral parameters ($R = 9.53$, 13.78), confirming that the $\kappa\delta^2$ universality is independent of the Maass spectral data.

6.6 Degree Extension: GL(3) and GL(4)

All preceding $\kappa\delta^2$ measurements concern degree-2 L-functions (automorphic forms on GL(2)). A central question is whether the slope 2 persists for higher-degree L-functions in the Selberg class. We test degree 3 and degree 4 via symmetric-power L-functions.

GL(3)**:** $\mathrm{sym}^2(E, \mathbf{11a1})$ **(Result #202).** The symmetric-square L-function of the elliptic curve 11a1 is a degree-3, conductor $N = 121$ automorphic L-function with $\Lambda(s) = \Lambda(3-s)$ and $\sigma_{\text{crit}} = 3/2$. Five zeros ($t \in [3.9, 12.1]$, spacing > 1.0) are selected from the ≈ 30 zeros found by PARI's `lfunzeros`.

Result #202: $\kappa\delta^2$ log-log slope $\mathbf{2.0000 \pm 0.0001}$, $R^2 = 1.000000$ for all five zeros. Monodromy $5/5 = 2.0000\pi$. σ-uniqueness: 5/5 PASS (centre $\sigma = 1.5$ is maximal—unlike GL(2), no B-01 excess at degree 3). FE: $|\Lambda(s) - \Lambda(3-s)|/|\Lambda(s)| = 0.00$. Overall: **5/5 PASS** (★★★).

The ± 0.0001 error bar matches the best GL(2) measurement (#200, Maass odd). This is the first degree-3 $\kappa\delta^2$ slope measurement in the literature.

GL(4)**:** $\mathrm{sym}^3(\Delta)$ **(Result #203).** The symmetric-cube L-function of the Ramanujan Δ-function is a degree-4, conductor $N = 1$, motivic weight 33 automorphic L-function with $\Lambda(s) = \Lambda(1-s)$ and $\sigma_{\text{crit}} = 1/2$ (PARI normalisation with $\gamma_V = [5.5, 6.5, 16.5, 17.5]$). L(1/2)$\approx$ 0 (root number $\varepsilon = -1$). Five zeros ($t \in [4.2, 12.1]$, spacing > 1.0) are found by manual $|L(1/2 + it)|$ minimum search (golden-section refinement, since `lfunzeros` is unreliable for large gamma shifts).

Result #203: $\kappa\delta^2$ log-log slope $\mathbf{2.0000 \pm 0.0001}$, $R^2 = 1.000000$ for all five zeros. Monodromy $5/5 = 2.0000\pi$. σ-uniqueness: FAIL (centre count = 24, max count = 30 at $\sigma = 0.40$), consistent with the degree-$\geq$2 structural FAIL pattern (B-05). FE: `lfuncheckfeq` $= -51$ digits. Overall: **4/5 PASS** (★★★).

The individual zero slopes are 1.9999, 2.0000, 2.0001, 1.9999, 2.0000—each indistinguishable from the theoretical value 2. Combined with the GL(3) result, the slope 2 now spans Selberg class degrees $\{1, 2, 3, 4, 5\}$.

GL(4): sym^3(**37a1**) **(Result #204).** To address the potential bias toward the 11a1/Δ family (sym^2(11a1) and $\mathrm{sym}^3(\Delta)$ are both 11a1-derived), we verify the same slope on a fully independent elliptic curve. $L(s, \mathrm{sym}^3(37\mathrm{a}1))$ has conductor $N = 50653 = 37^3$, degree 4, arithmetic normalisation $\sigma_{\mathrm{crit}} = k/2 = 2$, and $\gamma_V = [-1, 0, 0, 1]$ (PARI `lfunsympow(37a1,3)`). FE: `lfuncheckfeq` $= -137$ digits.

Result #204: $\kappa\delta^2$ log-log slope $\mathbf{1.9999 \pm 0.0005}$, $R^2 = 1.000000$ for all five zeros ($t \in [0.37, 5.55]$, spacing > 1.0). Monodromy $5/5 = 2.0000\pi$. σ-uniqueness: FAIL (centre=40, max=70), consistent with degree-$\geq$2 structural pattern (B-05). Overall: **4/5 PASS** (★★★).

Individual slopes: 2.0002, 2.0002, 2.0000, 2.0002, 1.9988. This confirms that the slope 2 universality at degree 4 is independent of the underlying elliptic curve, conductor, or motivic weight ($w = 33$ for $\mathrm{sym}^3(\Delta)$ vs. $w = 3$ for $\mathrm{sym}^3(37\mathrm{a}1)$).

[result #204, confirmed 2026-04-21; $\mathrm{sym}^3(37\mathrm{a}1)$: N=50653, k=4, γ_V=$[-1, 0, 0, 1]$; FE= $-$137 digits; slope=1.9999±0.0005; mono 5/5=2π; σ-uniq FAIL (structural); 11-row table; 11a1-bias resolved]

GL(5): sym^4(**11a1**) **(Result #205).** To extend the degree coverage to 5, we test the symmetric fourth power $L(s, \mathrm{sym}^4(11\mathrm{a}1))$: conductor $N = 14641 = 11^4$, degree 5, arithmetic normalisation $\sigma_{\mathrm{crit}} = k/2 = 2.5$, $\gamma_V = [-2, -1, 0, 0, 1]$ (PARI `lfunsympow(11a1,4)`). FE: `lfuncheckfeq` $= -116$ digits.

Result #205: $\kappa\delta^2$ log-log slope $\mathbf{1.9999 \pm 0.0004}$, $R^2 = 1.000000$ for all five zeros ($t \in [1.49, 7.35]$, spacing > 1.0). Monodromy $5/5 = 2.0000\pi$. σ-uniqueness: FAIL (centre=36, max=68), consistent with degree-$\geq$2 structural pattern (B-05). Overall: **4/5 PASS** (★★★).

Individual slopes: 2.0000, 1.9996, 1.9995, 2.0000, 2.0006. This confirms slope 2 universality at degree 5, completing the coverage across Selberg class degrees $\{1, 2, 3, 4, 5\}$. Result #85 (Paper 1) had already verified the four classical properties at GL(5); the present measurement adds the quantitative $\kappa\delta^2$ slope with full multi-δ precision. Degree 5 remains limited to one L-function due to computational constraints ($\mathrm{sym}^4(37\mathrm{a}1)$ has $N = 37^4 = 1{,}874{,}161$).

[result #205, confirmed 2026-04-21; $\mathrm{sym}^4(11\mathrm{a}1)$: N=14641, k=5, γ_V=$[-2, -1, 0, 0, 1]$; FE= $-$ 116 digits; slope=1.9999±0.0004; mono 5/5=2π; σ-uniq FAIL (structural); 12-row table; degree 1–5 complete]

6.7 Weight, Spectral, and Degree Invariance of $\kappa\delta^2$

Table 15 collects the $\kappa\delta^2$ log-log slopes (slope of $\log(\kappa\delta^2 - 1)$ vs. $\log\delta$) across all weights, spectral parameters, and Selberg class degrees tested to date.

For holomorphic forms at $k = 12$, 16, and 20 the slopes agree: the three-weight mean is 1.9994 ± 0.0010, and the maximum pairwise deviation is 0.0024. The two Maass forms ($R = 9.5337$ odd, $R = 13.7798$ even) add qualitatively different data points—non-holomorphic, weight 0, with spectral gamma factors—yet their slopes (2.0003 ± 0.0003 and 1.9999 ± 0.0008) are among the most precise measurements.

The degree extension via symmetric powers is equally striking: $\mathrm{sym}^2(11\mathrm{a}1)$ at degree 3 and $\mathrm{sym}^3(\Delta)$ at degree 4 both yield slope 2.0000 ± 0.0001, matching the theoretical value to four decimal places. The Artin S_5 standard representation ρ_4 (Result #209: irreducible

Table 15: Weight, spectral, degree, and family invariance of the $\kappa\delta^2$ log-log slope. Sixteen data points spanning degrees 1–6, weights $k \in \{0, 1, 2, 12, 16, 20\}$, Maass spectral parameters $R = 9.53, 13.78$, symmetric powers sym^2–sym^5, two Artin L-functions (S_3, S_5), one Rankin–Selberg convolution (11a1 × 37a1), and one Dirichlet L-function (χ_{-7}, conductor 7). All σ-direction measurements use the multi-δ log-log protocol.

Type	L-function	Degree	σ_{crit}	Slope	$\pm\sigma$	Rating
k=0	Riemann ζ	GL(1)	$\frac{1}{2}$	≈ 2.0	—	★★★
Dirichlet	$L(s, \chi_{-7})$ (#213)	GL(1)	$\frac{1}{2}$	**2.0000**	**0.0014**	★★★
k=1	Artin S_3 (#207)	GL(2)	$\frac{1}{2}$	**2.0000**	**0.0000**	★★★
k=2	EC 11a1	GL(2)	1	≈ 2.0	—	★★★
Maass odd	u_R^- (R=9.53)	GL(2)	$\frac{1}{2}$	**2.0003**	**0.0003**	★★★
Maass even	u_R^+ (R=13.78)	GL(2)	$\frac{1}{2}$	1.9999	0.0008	★★★
k=12	Ramanujan Δ	GL(2)	6	2.0008	0.0006	★★★
k=16	$\Delta \cdot E_4$	GL(2)	8	1.9989	0.0035	★★★
k=20	$\Delta \cdot E_8$	GL(2)	10	1.9984	0.0055	★★★
sym^2	sym^2(11a1)	GL(3)	$\frac{3}{2}$	**2.0000**	**0.0001**	★★★
sym^3	sym^3(Δ)	GL(4)	$\frac{1}{2}$	**2.0000**	**0.0001**	★★★
sym^3	sym^3(37a1)	GL(4)	2	**1.9999**	**0.0005**	★★★
Artin S_5	ρ_4 (#209)	GL(4)	$\frac{1}{2}$	**1.9999**	**0.0003**	★★★
R-S	11a1 × 37a1 (#210)	GL(4)	1	**1.9960**	**0.0072**	★★★
sym^4	sym^4(11a1)	GL(5)	$\frac{5}{2}$	**1.9999**	**0.0004**	★★★
sym^5	sym^5(11a1)	GL(6)	3	**1.9980**	**0.0036**	★★★

degree-4, conductor 2869, slope 1.9999 ± 0.0003) confirms universality beyond the symmetric-power chain. The Rankin–Selberg convolution $L(s, 11\text{a}1 \times 37\text{a}1)$ (Result #210: degree 4, $N = 407$, slope 1.9960 ± 0.0072) provides the first tensor-product verification, establishing a fourth independent degree-4 construction. Result #213 adds four quadratic Dirichlet L-functions $L(s, \chi_D)$ with $D \in \{-3, -4, 5, -7\}$ (conductors 3–7), yielding a combined slope of 2.0002 ± 0.0001—the most precise measurement in the table. This confirms that even the simplest non-trivial L-functions (degree 1, $N > 1$) satisfy the universal law, and that conductor alone does not perturb the slope. Combined with the $k = 0$ (Riemann ζ) and $k = 2$ (EC 11a1) results from Paper 1, the slope equals 2.0 ± 0.008 across sixteen data points spanning Selberg class degrees $\{1, 2, 3, 4, 5, 6\}$, holomorphic weights $k \in \{0, 1, 2, 12, 16, 20\}$, two non-holomorphic Maass forms, four Dirichlet L-functions, four independent GL(4) L-functions (two symmetric powers, one Artin, and one Rankin–Selberg), one GL(5) L-function, and one GL(6) L-function.

Theorem 11 (Slope universality)**.** *Let $L(s, \pi)$ be an automorphic L-function with completed function $\Lambda(s, \pi)$ and contragredient $\Lambda(s, \tilde{\pi})$ satisfying:*

(FE) $\Lambda(s, \pi) = \varepsilon\, \Lambda(k - s, \tilde{\pi})$ *for some $k \in \mathbb{R}$, $|\varepsilon| = 1$; and*

(SR) $\Lambda(\bar{s}, \tilde{\pi}) = \overline{\Lambda(s, \pi)}$ (*Schwarz reflection*).

Let $\rho = \frac{k}{2} + i\gamma$ be a simple zero on the critical line. Write the Laurent expansion $\Lambda'/\Lambda(s) = (s - \rho)^{-1} + c_0 + c_1(s - \rho) + \cdots$. Then:

(i) $\mathrm{Re}(c_0) = 0$ (c_0 *is purely imaginary*);

(ii) $\mathrm{Im}(c_1) = 0$ (c_1 *is real*);

(iii) $\kappa(\rho{+}\delta)\,\delta^2 = 1 + A\,\delta^2 + O(\delta^3)$ *with* $A = \mathrm{Im}(c_0)^2 + 2c_1$.

In particular, $\log(\kappa\delta^2 - 1) = \log A + 2\log\delta + O(\delta)$*, so the log-log slope is exactly* 2.

Proof. Write Laurent coefficients c_j for $\Lambda'/\Lambda(s,\pi)$ at ρ and c'_j for $\Lambda'/\Lambda(s,\tilde{\pi})$ at $\bar{\rho} = k - \rho$. From (FE): $\Lambda'/\Lambda(s,\pi) = -\Lambda'/\Lambda(k{-}s,\tilde{\pi})$, so expanding at $s = \rho + \delta$ gives $c'_0 = -c_0$ and $c'_1 = c_1$. From (SR): $\Lambda'/\Lambda(\bar{s},\tilde{\pi}) = \overline{\Lambda'/\Lambda(s,\pi)}$, so expanding at $s = \rho + \delta$ gives $c'_0 = \bar{c}_0$ and $c'_1 = \bar{c}_1$. Combining: $c_0 = -\bar{c}_0$ (so $\mathrm{Re}(c_0) = 0$) and $c_1 = \bar{c}_1$ (so $\mathrm{Im}(c_1) = 0$). Then $\kappa\delta^2 = |1 + c_0\delta + c_1\delta^2|^2 = 1 + A\delta^2 + O(\delta^3)$, since $\mathrm{Re}(c_0) = 0$ kills the δ^1 term. □

Proposition 12 (Laurent coefficient parity—universal)**.** *The FE/SR argument yields* $c_n = (-1)^{n+1}\bar{c}_n$ *for* every *Laurent coefficient* c_n*,* $n \geq 0$*, and for* every *L-function satisfying* $\Lambda(s,\pi) = \varepsilon\,\Lambda(1{-}s,\tilde{\pi})$ *and* $\Lambda(\bar{s},\pi) = \overline{\Lambda(s,\tilde{\pi})}$*. Self-duality* ($\pi = \tilde{\pi}$) *is* ***not required****: the FE relates* $c_n(\pi,\rho)$ *to* $c_n(\tilde{\pi},\bar{\rho})$*, and SR relates* $c_n(\pi,\rho)$ *to* $\bar{c}_n(\tilde{\pi},\bar{\rho})$*; combining gives* $c_n = (-1)^{n+1}\bar{c}_n$ *for any* π*. Consequently,* c_{2k} *is purely imaginary and* c_{2k+1} *is real for all* $k \geq 0$*. Numerical verification at orders* $n = 0$–3*:* 11 *self-dual zeros (*ζ*, 11a1,* χ_{-3}*; Result #C-240) and* 6 *non-self-dual zeros (*$\chi \bmod 5$*,* $\chi \bmod 7$*; Result #C-241) all satisfy* $|\mathrm{Re}(c_{2k})|/|c_{2k}| \leq 3 \times 10^{-7}$ *and* $\mathrm{Im}(c_{2k+1}) = 0$ *to machine precision (*200*-digit arithmetic).* ***Sharpness.*** *The hypothesis* $\sigma = \frac{1}{2}$ *(i.e.* $1{-}\rho = \bar{\rho}$*) is essential: the Davenport–Heilbronn function satisfies both FE and SR, yet its four known off-critical zeros (*$\sigma \neq \frac{1}{2}$*) show* $|\mathrm{Re}(c_0)|/|\mathrm{Im}(c_0)| \sim 2$–$7$*, while five on-critical zeros satisfy* $\sim 10^{-12}$ *(Result #C-242). Thus parity characterises zeros on the critical line.*

Proof. The functional equation maps $c_n(\pi,\rho)$ to $\varepsilon(-1)^n c_n(\tilde{\pi},\bar{\rho})$. The conjugate-centric symmetry maps $c_n(\pi,\rho)$ to $\bar{c}_n(\tilde{\pi},\bar{\rho})$. Combining: $c_n = (-1)^{n+1}\bar{c}_n$. □

Theorem 13 (Off-critical parity protection)**.** *Under the hypotheses of Proposition 12, let* $\rho = (\frac{1}{2} + \delta) + it_0$ *with* $\delta \neq 0$ *be a simple off-critical zero, and let* c_n *denote the Laurent coefficients of* Λ'/Λ *at* ρ*. Then:*

(a) ***Odd-order protection.*** $r(c_{2k+1}) \coloneqq |\mathrm{Im}(c_{2k+1})|/|\mathrm{Re}(c_{2k+1})| = O(|\delta|^{2k+3})$ *as* $\delta \to 0$.

(b) ***Even-order amplification.*** $r(c_{2k}) \coloneqq |\mathrm{Re}(c_{2k})|/|\mathrm{Im}(c_{2k})| = O(|\delta|^{-(2k+1)})$ *as* $\delta \to 0$.

In particular, the parity violation for odd-order coefficients decays as a power law in the off-critical displacement $|\delta|$*, with the exponent growing linearly in the order* k*.*

Proof. By the FE and conjugate-centric symmetry, the zero ρ has a partner $1{-}\bar{\rho} = (\frac{1}{2} - \delta) + it_0$. In the Hadamard product representation, the partner zero at distance 2δ contributes $(-1)^n/(2\delta)^{n+1}$ (a real quantity) to c_n. For odd n, this dominates $\mathrm{Re}(c_n)$ (the parity-preserving component), giving $|\mathrm{Re}(c_{2k+1})| \asymp (2\delta)^{-(2k+2)}$. The parity-violating component $\mathrm{Im}(c_{2k+1})$ arises from the first-order δ-perturbation of on-critical zero contributions in the Hadamard sum; its leading term is $O(\delta)$. The ratio gives $r(c_{2k+1}) = O(\delta/(2\delta)^{-(2k+2)}) = O(\delta^{2k+3})$. For even n, the partner contributes $O((2\delta)^{-(2k+1)})$ to $\mathrm{Re}(c_{2k})$ (the parity-*violating* component), while $|\mathrm{Im}(c_{2k})|$ is dominated by on-critical-zero contributions of order $O(1)$, so $r(c_{2k})$ grows. □

Numerical verification (Result #C-243): Laurent coefficients c_0–c_7 computed at 5 on-critical and 4 off-critical zeros of the Davenport–Heilbronn function (dps = 120, 72 values total). The *partner dominance ratio* $|\mathrm{Re}(c_{2k+1})|/(2\delta)^{-(2k+2)}$ approaches 1 as $\delta \to 0$ (0.98 for c_1 at $\delta = 0.074$; 1.00 for c_3). The δ-exponents $\beta_k := \log r / \log|\delta|$ average 2.91, 4.38, 5.94, 7.74 for $k = 0, 1, 2, 3$ (predicted: $3, 5, 7, 9$); the systematic deficit is attributed to t-dependent prefactors confounded with δ. Even/odd asymmetry ratio: 2.4×10^7.

Corollary 14 (Amplitude formula)**.** *The sub-leading coefficient* $A = \mathrm{Im}(c_0)^2 + 2c_1$ *decomposes into a* local density *contribution* $\mathrm{Im}(c_0)^2$ *(related to nearby zeros via* $c_0 = \sum_{\rho' \neq \rho}(\rho - \rho')^{-1}$*) and a* gamma-factor *contribution* $2c_1$ *(dominated by the digamma terms in* Λ'/Λ*).*

Numerical verification (Result #214): 35 zeros across seven L-functions—four self-dual (ζ, $L(s, \chi_{-3})$, $L(s, \chi_{-7})$, $L(s, \chi_{-11})$) and three non-self-dual (χ mod 5, χ mod 7, χ mod 13 of orders 4, 6, 12)—yield $|\,\mathrm{Re}(c_0)| = 0$ to machine precision (150-digit arithmetic), mean slope 2.0004 ± 0.0023, and mean A-prediction error 0.6% (max 1.7%; Result #216). The amplitude formula is verified at higher degrees (Result #217): 25 zeros across five L-functions spanning GL(1)–GL(4) (ζ, two elliptic curves, sym^2, sym^3) yield A-prediction errors of mean 1.3% (max 4.0% at degree 4). An independent cross-validation (Result #218) adds 20 zeros from four *distinct* families—two Dirichlet L-functions (χ_{-3}, χ_{-7}), the Ramanujan Δ L-function (weight 12, non-EC), and $\mathrm{sym}^2(37\mathrm{a}1)$ (rank-1 curve, independent of 11a1)—all yielding 20/20 PASS with mean error 0.5% (max 2.0%). Together, 45 zeros across nine L-functions confirm Corollary 14 across degrees 1–4, ruling out symmetric-power chain bias. The theorem subsumes the earlier Theorem 5 (which established $\kappa\delta^2 = 1 + O(\delta^2)$ without the explicit A formula) and upgrades the empirical observation of slope 2 in Table 15 from a numerical regularity to a mathematical theorem. Sixteen data points spanning Selberg class degrees $\{1, 2, 3, 4, 5, 6\}$, holomorphic weights $k \in \{0, 1, 2, 12, 16, 20\}$, two Maass forms, four Dirichlet L-functions, four independent GL(4) constructions (symmetric powers, Artin, Rankin–Selberg), and one GL(6) L-function all confirm the predicted slope 2 to within ± 0.008, establishing *construction universality.*

Amplitude scaling: the sub-leading coefficient $A(t_0)$. While the slope is a universal constant ($= 2$), the amplitude $A(t_0) := \kappa(\sigma_{\mathrm{crit}} + \delta,\, t_0) - 1/\delta^2$ at small δ varies across zeros and L-functions. Extracting A at $\delta = 0.01$ from 52 zeros across degrees 1–6 (Result #212) reveals a two-variable dependence:

$$A(d, N) \;\approx\; \alpha\, d^2 \;+\; \beta \log N \;+\; \gamma, \qquad \alpha \approx 0.39,\ \beta \approx 2.19,\ \gamma \approx 0.10, \tag{3}$$

with in-sample $R^2 = 0.983$. The degree-squared term reflects the Hadamard product decomposition $A = B(t_0)^2 + 2H_1(t_0)$, where $H_1 = \sum_{\rho' \neq \rho}(\mathrm{Im}\,\rho' - t_0)^{-2}$ grows with zero density (hence with d), and the $\log N$ term arises from the Stirling approximation of the gamma factors in the completed L-function.

Out-of-sample validation. For the Artin S_5 L-function (Result #209: $d = 4$, $N = 2869$), equation (3) predicts $A \approx 23.9$; the measured mean over five zeros is 24.9 (error 4%). For $L(s, 11\mathrm{a}1 \times 37\mathrm{a}1)$ (Result #210: $d = 4$, $N = 407$), the prediction is 19.6 versus measured 30.7

(high variance due to a close zero pair). The qualitative trend—A growing with both d^2 and $\log N$—is robust; the precise coefficients require further data for stabilization. [result #212, confirmed 2026-04-21; 6 degrees, 52 zeros; out-of-sample Artin S_5 validation 4% error; Observation: $A(d, N) \propto d^2 + \log N$ (tentative coefficients)]

Intrinsic variability of $A(t_0)$. Direct computation of H_1 via the Hadamard sum for 269 zeros of $\zeta(s)$ (Result #215) reveals a coefficient of variation $\mathrm{CV}(H_1) \approx 42\%$, arising from the GUE-type fluctuations in local zero spacing. The normalised ratio $H_1 \cdot \Delta(t_0)^2$ (where $\Delta = 2\pi/\log(t_0/2\pi)$ is the mean spacing) has a stable mean of 4.58 ± 1.95 across $t_0 \in [50, 500]$, but the point-to-point variability precludes a precise universal constant. The $B(t_0)^2$ term (from the Hilbert-type sum $\sum_{\rho' \neq \rho}(\mathrm{Im}\,\rho' - t_0)^{-1}$) contributes ≈24% of A, fluctuating with mean zero and standard deviation ≈1.0. This intrinsic $O(1)$ variability explains why the regression (3) achieves $R^2 = 0.98$ in-sample but remains "tentative" for point predictions. [result #215, confirmed 2026-04-21; 269 zeros of ζ; $H_1 = \sum 1/(\gamma_0 - \gamma_n)^2$ verified vs PARI to 0.7%; CV= 42%; $2H_1$ contributes 76% of A] [result #217, confirmed 2026-04-21; A formula verified degrees 1–4; 25 zeros, 5 L-functions; mean error 1.3%, max 4.0%]

6.8 Laurent Amplitude and Zero-Gap Statistics (Result #219)

The Laurent amplitude $A(\gamma) = \mathrm{Im}(c_0)^2 + 2\,\mathrm{Re}(c_1)$ exhibits a degree-independent negative correlation with GUE-normalised zero gaps ($\rho \in [-0.63, -0.42]$, 12/12 sign-consistent) across *twelve* automorphic L-functions spanning GL(1)–GL(6) (Results #219, #C-261, #C-263–#C-265, #C-268–#C-270, #C-273, #C-275; including $\mathrm{Sym}^5(11a1)$ at GL(6): $\rho = -0.423$, $p = 2.4 \times 10^{-4}$, $n = 71$). The Hadamard lower bound $A \geq 2/g_{\min}^2$ (Proposition 12, Result #C-265) makes $\rho(A, g_{\min}) < 0$ a structural necessity under the GUE pair-correlation conjecture, and $g_{\min}$ is height-stable (8-bin span= 0.163, Kendall τ non-significant; Result #C-270). Complete treatment—including the A_Λ/A_L dichotomy (Result #C-264), the Gamma-factor decomposition (Result #C-268), and all measurements—is given in Kim (2026d) [2].

[Results #219, #C-261, #C-263–#C-265, #C-268–#C-270, #C-273, #C-275; confirmed 2026-04-25; 12/12 GL(1)–GL(6); see Kim (2026d) [2]]

7 Discussion

Universality. The four ξ-bundle properties pass across an extended family of L-functions: elliptic-curve L-functions (rank 0–3, conductor 11–10099, 8 curves, $\varepsilon = \pm 1$), Rankin–Selberg GL(4) (2 pairs), Dedekind ζ-functions (7 imaginary quadratic fields), and Epstein ζ-functions including 3 non-Euler forms (class number 2–3). Combined with Paper 1's GL(1)– GL(5) results (81 numerical experiments), the framework now covers the major algebraic families of L-functions.

Weight, spectral, and degree universality. Results #125–#127, #200–#201, and #202–#203 establish that the $\kappa\delta^2$ log-log slope is invariant under the modular weight, holomorphicity, Maass symmetry class, and Selberg class degree. For holomorphic eigenforms the three-weight mean is 1.9994 ± 0.0010 ($k = 12, 16, 20$); the two Maass cusp forms yield

2.0003 ± 0.0003 ($R = 9.53$, odd) and 1.9999 ± 0.0008 ($R = 13.78$, even). The degree extension via symmetric powers produces the most precise measurements: $\mathrm{sym}^2(11a1)$ at GL(3) and $\mathrm{sym}^3(\Delta)$ at GL(4) both give slope 2.0000 ± 0.0001. Combined with the $k = 0, 2$ results from Paper 1, the Artin result ($k = 1$, Paper 3), and the Rankin–Selberg tensor product (Result #210, slope 1.9960 ± 0.0072), the slope equals 2.0 ± 0.008 across sixteen data points spanning Selberg class degrees $\{1, 2, 3, 4, 5, 6\}$ and six construction families. Theorem 11 proves this analytically: the functional equation and reality condition force $\mathrm{Re}(c_0) = 0$ in the Laurent expansion, killing the δ^1 term and making slope 2 a mathematical necessity for all automorphic L-functions.

A–gap degree-independent universality and mechanism. The Laurent amplitude $A(\gamma)$ correlates negatively with GUE-normalised gaps across *twelve* L-functions spanning GL(1)–GL(6): $\rho \in [-0.63, -0.42]$ with 12/12 sign consistency (Results #219, #C-261, #C-263–#C-265, #C-268–#C-270, #C-273, #C-275). The $g_{\min}$ metric is height-stable (span $= 0.163$, Kendall τ non-significant), and the Hadamard lower bound $A \geq 2/g_{\min}^2$ (Proposition 12) makes the anticorrelation a structural necessity under the GUE pair-correlation conjecture. Full treatment including the A_Λ/A_L dichotomy and Gamma-factor decomposition is given in Kim (2026d) [2].

Euler product dispensability. Theorem 7 and Result #114 establish that the Euler product is not required for the ξ-bundle properties. This aligns with the interpretation that the ξ-bundle geometry is a consequence of the functional equation structure alone.

The discrimination theorem. Theorem 8 provides the first *analytic* statement connecting the $O(\delta)$ correction to $\kappa\delta^2$ with the location of a zero relative to the critical line. The numerical verification (Results #115–#117) confirms this connection for the only known explicit test case (DH function). For the Riemann ζ-function, no off-critical zeros are known; Theorem 8 implies that if such a zero existed, it would produce a detectable $O(\delta)$ signal.

Limitations. The asymptotic $c_1 \cdot d \to 1$ is validated with $N = 4$ data points. The systematic δ-sweep (Result #116) is limited to $N = 3$ off-critical zeros (OFF#4 had insufficient precision at $\delta < 0.1$). No new off-critical zeros were discovered in the range $t \in [200, 400]$ (22 candidate points checked, 0 confirmed), consistent with the known scarcity of DH off-critical zeros.

Future directions.

1. *Hilbert modular forms*: extending the spectral universality to number fields of degree > 1.
2. *Higher-degree extension*: degree 7+ via sym^6 or other automorphic constructions.
3. *Selberg class generalisation*: Theorem 8 uses only FE; can it be sharpened to give c_1 in terms of the Selberg class degree?
4. *σ-localisation*: extending the discrimination to non-horizontal offsets (complex δ).

5. *Additional DH off-critical zeros*: extending the search to $t > 400$ with higher precision.

8 Conclusion

We have extended the ξ-bundle framework of Kim (2026) in four directions. First, universality: eleven new numerical results (#107–#117) confirm the four ξ-bundle properties across elliptic-curve, Rankin–Selberg, Dedekind, and Epstein L-functions; in particular, non-Euler Epstein ζ-functions (Theorem 6) demonstrate that the Euler product is not necessary. Second, discrimination: Theorem 7 proves analytically that the first Taylor coefficient c_1 of $\kappa\delta^2$ vanishes if and only if the zero lies on the critical line; numerical verification via δ-sweep and direct analytic computation confirms this for all four known DH off-critical zeros. Third, weight and spectral universality: Results #125–#127 and #200–#201 establish that the slope equals 2.0 ± 0.001 independently of weight ($k = 12, 16, 20$), holomorphicity (two Maass forms, $R = 9.53$ and 13.78), and symmetry class (odd and even). Fourth, degree and family universality: Results #202–#206 extend the slope verification from GL(2) through GL(6) via symmetric powers; Result #209 adds an irreducible degree-4 Artin L-function (standard representation of S_5, slope 1.9999 ± 0.0003); and Result #210 adds the Rankin–Selberg tensor product $L(s, 11\text{a}1 \times 37\text{a}1)$ (slope 1.9960 ± 0.0072), confirming Observation 8 across Selberg class degrees $\{1, 2, 3, 4, 5, 6\}$ with sixteen independent data points from six distinct families (Dirichlet, symmetric powers, Artin, Rankin–Selberg, holomorphic cusp forms, and Maass forms).

Fifth, the Laurent amplitude $A(\gamma)$ exhibits a degree-independent negative correlation with GUE-normalised zero gaps ($\rho \in [-0.63, -0.42]$, 11/11 sign-consistent) across GL(1)–GL(5) L-functions (Results #219, #C-261, #C-263–#C-265, #C-268–#C-270, #C-273), and Proposition 12 (Result #C-265) identifies the Hadamard mechanism: $A \geq 2/g_{\min}^2$ from nearest-neighbour dominance makes the anticorrelation a structural necessity under the GUE pair-correlation conjecture ($E[A|g]$ monotone decreasing, 7/7 bins, $R^2 = 0.987$). The gap metric $g_{\min}$ is height-stable (Result #C-270, span= 0.163), confirming $g_{\min}$ as the canonical metric; complete treatment in Kim (2026d) [2].

Together, Theorems 5–7 and Observation 8 form a coherent picture: on the critical line, the ξ-bundle curvature satisfies $\kappa\delta^2 = 1 + O(\delta^2)$ universally across all L-function families, modular weights, and spectral types tested; off the critical line, a qualitatively different $O(\delta)$ correction appears, analytically traced to the breaking of mirror symmetry by the functional equation.

A Summary Table: All Forty-Three Results

Table 16: Summary of Results #107–#117, #125–#127, #200–#219, #C-240–#C-243, #C-261, #C-263–#C-265, #C-268–#C-270, #C-273, #C-275 (43 results). Rating: ★★★ strong positive, ★★ positive, **C** conditional.

Result	Section	Rating	Key finding
#107	§3.1	★★ C	EC 8 curves; P4 fails for $\varepsilon = -1$ (Hardy Z method); P3 passes all
#108	§3.1	★★★	EC 8 curves, ε-corrected; all 4/4 PASS; $\kappa\delta^2 = 1.000$
#109	§3.3	★★	κ_{near} universality; CV $< 0.13\%$ at $\delta = 0.01$; rank/conductor independent
#110	§3.4	★★★	8 EC curves, 120 zeros; $\lvert\Delta\arg/\pi\rvert = 1.000000$; contour $= 2.000000\pi$
#111	§3.5	★★	Conductor scaling $\Delta\kappa \approx 2\log N \times 10^{-3}$; $R^2 = 0.73$; effect $< 2\%$
#112	§4.1	★★	R-S GL(4), 2 pairs; 4/4 PASS; $\kappa\delta^2 = 1.014$, 1.033
#113	§4.2	★★★	Dedekind ζ, 7 fields (h=1,3); 4/4 PASS; all $\kappa\delta^2 < 1.005$
#114	§4.3	★★★	Epstein ζ; 3 non-Euler (h=2,3); 4/4 PASS; FE-only sufficiency (Thm 6)
#115	§5.2	★★★	DH on/off discrimination; 1.002 ± 0.001 vs. 1.210 ± 0.099; complete separation
#116	§5.4	★★★	δ-sweep; slope 1.999 ± 0.003 (on) vs. 0.963 ± 0.039 (off); Thm 7 confirmed
#117	§5.5	★★	$c_1 = \mathrm{Re}(\Lambda''/\Lambda')$ verified ($< 0.5\%$, $N = 4$); $c_1 \cdot d \to 1$
#125	§6.1	★★★	Ramanujan Δ ($k = 12$); 4/5 PASS; $\kappa\delta^2$ slope 2.0008 ± 0.0006; σ-uniq. FAIL (B-01)
#126	§6.2	★★★	Weight-16 cusp form ($\Delta \cdot E_4$); 5/5 PASS; slope 1.9989 ± 0.0035; weight invariance (Obs. 8)

Table 16 (continued)

Result	Section	Rating	Key finding
#127	§6.3	★★★	Weight-20 cusp form ($\Delta \cdot E_8$); 4/5 PASS; slope 1.9984 ± 0.0055; σ-uniq. FAIL (B-01)
#200	§6.4	★★★	Maass cusp form (R=9.5337, odd); 5/5 PASS; slope $\mathbf{2.0003 \pm 0.0003}$
#201	§6.5	★★★	Even Maass (R=13.7798); 4/5 PASS; slope 1.9999 ± 0.0008; symmetry-class independence
#202	§6.6	★★★	GL(3) sym^2(11a1); 5/5 PASS; slope $\mathbf{2.0000 \pm 0.0001}$; first degree-3 measurement
#203	§6.6	★★★	GL(4) $\mathrm{sym}^3(\Delta)$; 4/5 PASS; slope $\mathbf{2.0000 \pm 0.0001}$; degree 1–4 universality
#204	§6.6	★★★	GL(4) sym^3(37a1); 4/5 PASS; slope $\mathbf{1.9999 \pm 0.0005}$; 11a1-bias resolved
#205	§6.6	★★★	GL(5) sym^4(11a1); 4/5 PASS; slope $\mathbf{1.9999 \pm 0.0004}$; degree 1–5 complete
#206	§6.6	★★★	GL(6) sym^5(11a1); 4/5 PASS; slope $\mathbf{1.9980 \pm 0.0036}$; degree 1–6 complete
#209	§6.6	★★★	Artin S_5 ρ_4; 4/4 PASS; slope $\mathbf{1.9999 \pm 0.0003}$; independent degree-4 (Paper 3)
#210	§4.1	★★★	R-S 11a1 $\times$ 37a1; eff. 4/4 PASS; slope $\mathbf{1.9960 \pm 0.0072}$; tensor-product construction
#212	§6.7	★★	$A(d, N)$ scaling law; $R^2 = 0.983$; Artin S_5 out-of-sample 4% error; tentative coefficients
#213	§6.7	★★★	Dirichlet $L(s, \chi_D)$, $D \in \{-3, -4, 5, -7\}$; slope $\mathbf{2.0002 \pm 0.0001}$; 6th family (conductor independence)
#215	§6.7	★★	$A(t_0)$ Hadamard decomposition; H_1 verified 0.7%; CV= 42% intrinsic; $2H_1 = 76\%$ of A

Table 16 (continued)

Result	Section	Rating	Key finding
#216	§6.7	★★★★	Non-self-dual ($\chi_5, \chi_7, \chi_{13}$); slope $\mathbf{2.0004 \pm 0.0024}$; FE+SR proof; Thm generalised
#217	§6.7	★★★★	A formula degrees 1–4; 25 zeros, 5 L-functions; mean error 1.3%; Cor. 14 verified
#218	§6.7	★★★★	A formula cross-validation; 20/20 PASS; Δ, χ_{-3}, χ_{-7}, $\mathrm{sym}^2(37a1)$; mean error 0.5%; chain bias ruled out
#C-240	§6.7	★★★	Laurent parity $c_n = (-1)^{n+1}\bar{c}_n$; 11/11 PASS (self-dual); $\lvert\mathrm{Re}(c_2)\rvert/\lvert c_2\rvert \leq 3\times10^{-7}$; Proposition 12
#C-241	§6.7	★★★★	Parity **universal**: non-self-dual χ_5, χ_7 also 6/6 PASS; self-dual not needed (FE+SR suffice); Proposition 12
#C-242	§6.7	★★★★	Parity **sharpness**: DH on-critical 5/5 PASS ($r(c_0)\sim10^{-12}$), off-critical 4/4 FAIL ($r(c_0)\sim4.4$); $\sigma=\frac{1}{2}$ essential; Proposition 12
#C-243	§6.7	★★★★	Parity **protection**: $r(c_{2k+1})=O(\lvert\delta\rvert^{2k+3})$; partner dominance 0.98–1.00; even/odd ratio 2.4×10^7; Theorem 13
#219	§6.8	★★★	$A(\gamma)$–gap correlation; GUE-norm. $\rho=-0.62$ ($p=10^{-22}$); R/L asymmetry 4.4×; pair $\rho=+0.40$ ($p=10^{-9}$); see [2]
#C-261	§6.8	★★★	A–gap Dirichlet universality; χ_3, χ_4, χ_5; $\rho\in[-0.63, -0.55]$; 3/3 sign-consistent; see [2]
#C-263	§6.8	★★★★	A–gap GL(2) universality; E_{11a1} $\rho=-0.57$, E_{37a1} $\rho=-0.55$; 6/6 GL(1)+GL(2) consistent; see [2]
#C-264	§6.8	★★★	GL(3) A–gap; $\mathrm{Sym}^2(11a1)$ $\rho=-0.42$, $\mathrm{Sym}^2(37a1)$ $\rho=-0.49$; A_Λ/A_L dichotomy; see [2]

Table 16 (continued)

Result	Section	Rating	Key finding
#C-265	§6.8	★★★★	Prop 12: Hadamard $A \geq 2/g_{\min}^2$; $E[A\|g]$ monotone 7/7; $\rho=-0.90$; $R^2=0.987$; see [2]
#C-268	§6.8	★★★	Gamma 3-factor decomp.: $\rho(\Delta A, g_{\mathrm{R}})=-0.696$; $\Delta\rho=-0.291$; cross-term dominant; see [2]
#C-269	§6.8	★★★★	GL(4) Sym^3 A–gap; $\rho\in[-0.52,-0.51]$; 10/10 GL(1)–GL(4); see [2]
#C-270	§6.8	★★★★	$g_{\min}$ t-stable: span=0.163; g_{R} decays; canonical metric; B-45/B-46 resolved; see [2]
#C-273	§6.8	★★★★	GL(5) $\mathrm{Sym}^4(11a1)$ A–gap; $\rho(A_L, g_{\min}^{\mathrm{GUE}})=-0.485$ ($p=3.7\times10^{-9}$, $n=132$); 11/11 GL(1)–GL(5); see [2]
#C-275	§6.8	★★★★	GL(6) $\mathrm{Sym}^5(11a1)$ A–gap; $\rho(A_L, g_{\min}^{\mathrm{GUE}})=-0.423$ ($p=2.4\times10^{-4}$, $n=71$); 12/12 GL(1)–GL(6); B-38 resolved; see [2]

B Numerical Data: Off-Critical Zeros (DH)

Table 17 collects the complete δ-sweep data for the three DH off-critical zeros with sufficient precision (OFF#4 included only at $\delta \geq 0.10$).

Table 17: $\kappa\delta^2$ values from Result #116, DH off-critical zeros. ON-critical ensemble mean (10 zeros) shown for comparison.

Zero	δ=0.005	0.01	0.015	0.02	0.03	0.05	0.07	0.10
ON (mean)	1.000048	1.000192	1.000432	1.000767	1.001726	1.004795	1.009399	1.019182
OFF#1	1.018828	1.037696	1.056606	1.075561	1.113605	1.190280	1.267813	1.385945
OFF#2	1.034427	1.068469	1.102132	1.135420	1.200904	1.327729	1.449538	1.624253
OFF#3	1.067034	1.132039	1.195056	1.256139	1.372769	1.585968	1.776429	2.028926
OFF#4	—	—	—	—	—	—	—	1.506337

C Mirror-Pair Decomposition of c_1

For an off-critical zero $\rho = \sigma_0 + it_0$, the mirror partner is $\tilde{\rho} = 1 - \sigma_0 + it_0$. The leading contribution of the mirror pair to $A'(\rho)/A(\rho)$ is

$$\frac{1}{\rho - \tilde{\rho}} = \frac{1}{2\sigma_0 - 1} = \frac{1}{2d},$$

giving $c_1^{\text{mirror}} = 2\,\text{Re}(1/(2d)) = 1/d$. Table 18 shows the decomposition from Result #117.

Table 18: Mirror-pair vs. background decomposition of c_1 (Result #117).

Zero	d	c_1^{fit}	$c_1^{\text{mirror}} = 1/d$	c_1^{bg}	$c_1^{\text{bg}}/c_1^{\text{mirror}}$
OFF#1	0.3085	3.760	3.241	2.139	1.320
OFF#2	0.1508	6.916	6.630	3.601	1.086
OFF#3	0.0744	13.543	13.449	6.818	1.014
OFF#4	0.2243	5.026	4.459	2.796	1.254

As $d \to 0$, the background-to-mirror ratio approaches 1, indicating that the two contributions are of equal magnitude in the limit and that $c_1 \approx 2 \times c_1^{\text{mirror}} = 2/d$ at small d—which exceeds the asymptotic $1/d$ by a factor of 2 until higher-order corrections reduce the effective value. The data instead show $c_1 \cdot d \to 1$, suggesting significant cancellation among the background terms.

References

[1] Kim, Y.-H. (2026). *Phase Quantisation from Optical Lattices to the Zeta Landscape: A Unified Framework*. Preprint, arXiv:2026.xxxxx.

[2] Kim, Y.-H. (2026d). *Amplitude–Gap Anti-Correlation in Automorphic L-Functions: Hadamard Mechanism and Degree-Independent Universality*. Preprint.

[3] Titchmarsh, E. C. (1986). *The Theory of the Riemann Zeta-Function* (2nd ed., revised by D. R. Heath-Brown). Oxford University Press.

[4] Edwards, H. M. (1974). *Riemann's Zeta Function*. Academic Press.

[5] Davenport, H., & Heilbronn, H. (1936). On the zeros of certain Dirichlet series. *Journal of the London Mathematical Society*, 11, 181–185.

[6] Silverman, J. H. (1992). *The Arithmetic of Elliptic Curves* (2nd ed.). Springer.

[7] Iwaniec, H., & Kowalski, E. (2004). *Analytic Number Theory*. American Mathematical Society.

[8] Epstein, P. (1903). Zur Theorie allgemeiner Zetafunktionen. *Mathematische Annalen*, 56, 615–644.

[9] Rubinstein, M. (2001). Numerical computations concerning the Riemann hypothesis. *PhD thesis*, Princeton University.

[10] Cremona, J. E. (2019). *Algorithms for Modular Elliptic Curves* (2nd ed.). Cambridge University Press.

[11] PARI/GP (2023). Version 2.15.4. https://pari.math.u-bordeaux.fr/.

[12] Johansson, F., et al. (2023). *mpmath: a Python library for arbitrary-precision floating-point arithmetic* (version 1.3.0). https://mpmath.org/.