

Formulas

- Bayes' Theorem: If B_1, \dots, B_k form a partition of the sample space S , and A is any event, then for any $i = 1, \dots, k$,

$$\mathbb{P}(B_i | A) = \frac{\mathbb{P}(A | B_i)\mathbb{P}(B_i)}{\sum_{j=1}^k \mathbb{P}(A | B_j)\mathbb{P}(B_j)}.$$

- The following are probability mass functions for distributions we discussed.

1. Bernoulli with parameter p :

$$f(x) = \begin{cases} p & x = 1 \\ 1 - p & x = 0 \\ 0 & \text{otherwise} \end{cases}$$

2. Geometric with parameter p :

$$f(x) = \begin{cases} (1 - p)^{x-1}p & x = 1, 2, \dots \\ 0 & \text{otherwise} \end{cases}$$

3. Binomial with parameters n, p :

$$f(x) = \begin{cases} \binom{n}{x} p^x (1 - p)^{n-x} & x = 0, 1, 2, \dots, n \\ 0 & \text{otherwise} \end{cases}$$

4. Hypergeometric with parameters N_1, N_2, n :

$$f(x) = \begin{cases} \frac{\binom{N_1}{x} \binom{N_2}{n-x}}{\binom{N_1+N_2}{n}} & \text{for } \max\{0, n - N_2\} \leq x \leq \min\{N_1, n\} \\ 0 & \text{otherwise} \end{cases}.$$

5. Negative binomial with parameters r, p :

$$f(x) = \binom{x-1}{r-1} p^r (1-p)^{x-r}, \quad x = r, r+1, \dots$$

6. Poisson with parameter λ :

$$f(x) = e^{-\lambda} \frac{\lambda^x}{x!}, \quad x = 0, 1, 2, \dots$$

7. Trinomial with parameters n, p_A, p_B, p_C :

$$f(x, y) = \frac{n!}{x!y!(n-x-y)!} p_A^x p_B^y p_C^{n-x-y} \text{ for } x, y \geq 0 \text{ and } x + y \leq n.$$

- The following are probability distribution functions for distributions we discussed.

1. Uniform on $[a, b]$:

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{if } x \in [a, b] \\ 0 & \text{otherwise.} \end{cases}$$

2. Exponential with parameter θ :

$$f(x) = \begin{cases} \frac{1}{\theta} e^{-x/\theta} & \text{if } x \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$

3. Gamma with parameters α, θ :

$$f(x) = \begin{cases} \frac{1}{\Gamma(\alpha)\theta^\alpha} x^{\alpha-1} e^{-x/\theta} & \text{if } x \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$

4. Chi-square with r degrees of freedom:

$$f(x) = \begin{cases} \frac{1}{\Gamma(r/2)2^{r/2}} x^{r/2-1} e^{-x/2} & \text{if } x \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$

5. Normal with parameters μ, σ^2 :

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right).$$

- Least squares regression line:

$$y = \mu_Y + \rho \frac{\sigma_Y}{\sigma_X} (x - \mu_X).$$