

## Week 9: Conditional distributions, continuous bivariate distributions

Armenak Petrosyan

Tuesday class

# Conditional pmf

- $X, Y$  be discrete random variables with joint pmf  $f(x, y)$ .


## Definition

Let  $f_Y(y) > 0$ . The **conditional pmf of  $X$  given  $Y = y$**  is the following pmf:

$$g(x|y) = \frac{f(x, y)}{f_Y(y)}, \quad x \in \text{Range}(X)$$

- Notice that

$$g(x|y) = P(X = x|Y = y).$$



- For different values of  $y$ , it defines different pms-s on  $\text{Range}(X)$ .

# Conditional pmf

- $X, Y$  be discrete random variables with joint pmf  $f(x, y)$ .


## Definition

Let  $f_Y(y) > 0$ . The **conditional pmf of  $X$  given  $Y = y$**  is the following pmf:

$$g(x|y) = \frac{f(x, y)}{f_Y(y)}, \quad x \in \text{Range}(X)$$

- Notice that

$$g(x|y) = P(X = x|Y = y).$$



- For different values of  $y$ , it defines different pms-s on  $\text{Range}(X)$ .

# Conditional pmf

- $X, Y$  be discrete random variables with joint pmf  $f(x, y)$ .


## Definition

Let  $f_Y(y) > 0$ . The **conditional pmf of  $X$  given  $Y = y$**  is the following pmf:

$$g(x|y) = \frac{f(x, y)}{f_Y(y)}, \quad x \in \text{Range}(X)$$

- Notice that

$$g(x|y) = P(X = x|Y = y).$$



- For different values of  $y$ , it defines different pms-s on  $\text{Range}(X)$ .

# Conditional pmf

- $X, Y$  be discrete random variables with joint pmf  $f(x, y)$ .


## Definition

Let  $f_Y(y) > 0$ . The **conditional pmf of  $X$  given  $Y = y$**  is the following pmf:

$$g(x|y) = \frac{f(x, y)}{f_Y(y)}, \quad x \in \text{Range}(X)$$

- Notice that

$$g(x|y) = P(X = x|Y = y).$$




- For different values of  $y$ , it defines different pms-s on  $\text{Range}(X)$ .

## Definition

The **conditional pmf of  $Y$  given  $X = x$**  is the following pmf:

$$h(y|x) = \frac{f(x, y)}{f_X(x)}, \quad y \in \text{Range}(Y)$$

assuming  $f_X(x) > 0$ .



► Here


$$h(y|x) = P(Y = y|X = x).$$

## Definition

The **conditional pmf of  $Y$  given  $X = x$**  is the following pmf:

$$h(y|x) = \frac{f(x, y)}{f_X(x)}, \quad y \in \text{Range}(Y)$$

assuming  $f_X(x) > 0$ .



► Here

$$h(y|x) = P(Y = y|X = x).$$



### Problem (4.3-1 in the textbook)

Let  $X$  and  $Y$  have the joint pmf

$$f(x, y) = \frac{x + y}{32} \quad x = 1, 2 \quad y = 1, 2, 3, 4.$$

- (a) Display the joint pmf and the marginal pmfs on a graph.
- (b) Find  $g(x|y)$  and draw a figure depicting the conditional pmfs for  $y = 1, 2, 3$ , and 4.
- (c) Find  $h(y|x)$  and draw a figure depicting the conditional pmfs for  $x = 1, 2$ .
- (d) Find  $P(1 \leq Y \leq 3|X = 1)$ ,  $P(Y \leq 2|X = 2)$ , and  $P(X = 2|Y = 3)$ .
- (e) Find  $E(Y|X = 1)$  and  $\text{Var}(Y|X = 1)$ .

### Problem (4.3-1 in the textbook)

Let  $X$  and  $Y$  have the joint pmf

$$f(x, y) = \frac{x + y}{32} \quad x = 1, 2 \quad y = 1, 2, 3, 4.$$

- (a) *Display the joint pmf and the marginal pmfs on a graph.*
- (b) *Find  $g(x|y)$  and draw a figure depicting the conditional pmfs for  $y = 1, 2, 3$ , and 4.*
- (c) *Find  $h(y|x)$  and draw a figure depicting the conditional pmfs for  $x = 1, 2$ .*
- (d) *Find  $P(1 \leq Y \leq 3|X = 1)$ ,  $P(Y \leq 2|X = 2)$ , and  $P(X = 2|Y = 3)$ .*
- (e) *Find  $E(Y|X = 1)$  and  $\text{Var}(Y|X = 1)$ .*

### Problem (4.3-1 in the textbook)

Let  $X$  and  $Y$  have the joint pmf

$$f(x, y) = \frac{x + y}{32} \quad x = 1, 2 \quad y = 1, 2, 3, 4.$$

- (a) Display the joint pmf and the marginal pmfs on a graph.
- (b) Find  $g(x|y)$  and draw a figure depicting the conditional pmfs for  $y = 1, 2, 3$ , and 4.
- (c) Find  $h(y|x)$  and draw a figure depicting the conditional pmfs for  $x = 1, 2$ .
- (d) Find  $P(1 \leq Y \leq 3|X = 1)$ ,  $P(Y \leq 2|X = 2)$ , and  $P(X = 2|Y = 3)$ .
- (e) Find  $E(Y|X = 1)$  and  $\text{Var}(Y|X = 1)$ .

### Problem (4.3-1 in the textbook)

Let  $X$  and  $Y$  have the joint pmf

$$f(x, y) = \frac{x + y}{32} \quad x = 1, 2 \quad y = 1, 2, 3, 4.$$

- (a) Display the joint pmf and the marginal pmfs on a graph.
- (b) Find  $g(x|y)$  and draw a figure depicting the conditional pmfs for  $y = 1, 2, 3$ , and 4.
- (c) Find  $h(y|x)$  and draw a figure depicting the conditional pmfs for  $x = 1, 2$ .
- (d) Find  $P(1 \leq Y \leq 3|X = 1)$ ,  $P(Y \leq 2|X = 2)$ , and  $P(X = 2|Y = 3)$ .
- (e) Find  $E(Y|X = 1)$  and  $\text{Var}(Y|X = 1)$ .

### Problem (4.3-1 in the textbook)

Let  $X$  and  $Y$  have the joint pmf

$$f(x, y) = \frac{x + y}{32} \quad x = 1, 2 \quad y = 1, 2, 3, 4.$$

- (a) Display the joint pmf and the marginal pmfs on a graph.
- (b) Find  $g(x|y)$  and draw a figure depicting the conditional pmfs for  $y = 1, 2, 3$ , and 4.
- (c) Find  $h(y|x)$  and draw a figure depicting the conditional pmfs for  $x = 1, 2$ .
- (d) Find  $P(1 \leq Y \leq 3|X = 1)$ ,  $P(Y \leq 2|X = 2)$ , and  $P(X = 2|Y = 3)$ .
- (e) Find  $E(Y|X = 1)$  and  $\text{Var}(Y|X = 1)$ .

### Problem (4.3-1 in the textbook)


Let  $X$  and  $Y$  have the joint pmf

$$f(x, y) = \frac{x + y}{32} \quad x = 1, 2 \quad y = 1, 2, 3, 4.$$


- (a) Display the joint pmf and the marginal pmfs on a graph.
- (b) Find  $g(x|y)$  and draw a figure depicting the conditional pmfs for  $y = 1, 2, 3$ , and 4.
- (c) Find  $h(y|x)$  and draw a figure depicting the conditional pmfs for  $x = 1, 2$ .
- (d) Find  $P(1 \leq Y \leq 3|X = 1)$ ,  $P(Y \leq 2|X = 2)$ , and  $P(X = 2|Y = 3)$ .
- (e) Find  $E(Y|X = 1)$  and  $\text{Var}(Y|X = 1)$ .

## Solution


(a) *Joint pmf and marginal pmfs*



(b) *Conditional pmf of  $X$*




(c) *Conditional pmf of  $Y$*




## Solution


(a) *Joint pmf and marginal pmfs*



(b) *Conditional pmf of X*




(c) *Conditional pmf of Y*






## Solution


(a) *Joint pmf and marginal pmfs*



(b) *Conditional pmf of X*



(c) *Conditional pmf of Y*



## Solution

(d) From the graph in (c)

$$P(1 \leq Y \leq 3|X = 1) = \frac{2}{14} + \frac{3}{14} + \frac{4}{14} \boxed{= \frac{9}{14}}.$$

$$P(Y \leq 2|X = 2) = \frac{3}{18} + \frac{4}{18} \boxed{= \frac{7}{18}}.$$

From the graph in (b)

$$P(X = 2|Y = 3) \boxed{= \frac{5}{9}}.$$

(e)

$$E(Y|X = 1) = 1 \cdot \frac{2}{14} + 2 \cdot \frac{3}{14} + 3 \cdot \frac{4}{14} + 4 \cdot \frac{5}{14} \boxed{= \frac{40}{14}}.$$

$$E(Y^2|X = 1) = 1^2 \cdot \frac{2}{14} + 2^2 \cdot \frac{3}{14} + 3^2 \cdot \frac{4}{14} + 4^2 \cdot \frac{5}{14} = \frac{130}{14}.$$

And, finally,

$$\text{Var}(Y|X = 1) = \frac{130}{14} - \left(\frac{40}{14}\right)^2 \boxed{= \frac{220}{196}}.$$

## Solution

(d) From the graph in (c)

$$P(1 \leq Y \leq 3|X = 1) = \frac{2}{14} + \frac{3}{14} + \frac{4}{14} \boxed{= \frac{9}{14}}.$$

$$P(Y \leq 2|X = 2) = \frac{3}{18} + \frac{4}{18} \boxed{= \frac{7}{18}}.$$

From the graph in (b)

$$P(X = 2|Y = 3) \boxed{= \frac{5}{9}}.$$

(e)

$$E(Y|X = 1) = 1 \cdot \frac{2}{14} + 2 \cdot \frac{3}{14} + 3 \cdot \frac{4}{14} + 4 \cdot \frac{5}{14} \boxed{= \frac{40}{14}}.$$

$$E(Y^2|X = 1) = 1^2 \cdot \frac{2}{14} + 2^2 \cdot \frac{3}{14} + 3^2 \cdot \frac{4}{14} + 4^2 \cdot \frac{5}{14} = \frac{130}{14}.$$

And, finally,

$$\text{Var}(Y|X = 1) = \frac{130}{14} - \left(\frac{40}{14}\right)^2 \boxed{= \frac{220}{196}}.$$

## Solution

(d) From the graph in (c)

$$P(1 \leq Y \leq 3|X = 1) = \frac{2}{14} + \frac{3}{14} + \frac{4}{14} \boxed{= \frac{9}{14}}.$$

$$P(Y \leq 2|X = 2) = \frac{3}{18} + \frac{4}{18} \boxed{= \frac{7}{18}}.$$

From the graph in (b)

$$P(X = 2|Y = 3) \boxed{= \frac{5}{9}}.$$

(e)

$$E(Y|X = 1) = 1 \cdot \frac{2}{14} + 2 \cdot \frac{3}{14} + 3 \cdot \frac{4}{14} + 4 \cdot \frac{5}{14} \boxed{= \frac{40}{14}}.$$

$$E(Y^2|X = 1) = 1^2 \cdot \frac{2}{14} + 2^2 \cdot \frac{3}{14} + 3^2 \cdot \frac{4}{14} + 4^2 \cdot \frac{5}{14} = \frac{130}{14}.$$

And, finally,

$$\text{Var}(Y|X = 1) = \frac{130}{14} - \left(\frac{40}{14}\right)^2 \boxed{= \frac{220}{196}}.$$

## Solution

(d) From the graph in (c)

$$P(1 \leq Y \leq 3|X = 1) = \frac{2}{14} + \frac{3}{14} + \frac{4}{14} \boxed{= \frac{9}{14}}.$$

$$P(Y \leq 2|X = 2) = \frac{3}{18} + \frac{4}{18} \boxed{= \frac{7}{18}}.$$

From the graph in (b)

$$P(X = 2|Y = 3) \boxed{= \frac{5}{9}}.$$

(e)

$$E(Y|X = 1) = 1 \cdot \frac{2}{14} + 2 \cdot \frac{3}{14} + 3 \cdot \frac{4}{14} + 4 \cdot \frac{5}{14} \boxed{= \frac{40}{14}}.$$

$$E(Y^2|X = 1) = 1^2 \cdot \frac{2}{14} + 2^2 \cdot \frac{3}{14} + 3^2 \cdot \frac{4}{14} + 4^2 \cdot \frac{5}{14} = \frac{130}{14}.$$

And, finally,

$$\text{Var}(Y|X = 1) = \frac{130}{14} - \left(\frac{40}{14}\right)^2 \boxed{= \frac{220}{196}}.$$

## Solution

(d) From the graph in (c)

$$P(1 \leq Y \leq 3|X = 1) = \frac{2}{14} + \frac{3}{14} + \frac{4}{14} \boxed{= \frac{9}{14}}.$$

$$P(Y \leq 2|X = 2) = \frac{3}{18} + \frac{4}{18} \boxed{= \frac{7}{18}}.$$

From the graph in (b)

$$P(X = 2|Y = 3) \boxed{= \frac{5}{9}}.$$

(e)

$$E(Y|X = 1) = 1 \cdot \frac{2}{14} + 2 \cdot \frac{3}{14} + 3 \cdot \frac{4}{14} + 4 \cdot \frac{5}{14} \boxed{= \frac{40}{14}}.$$

$$E(Y^2|X = 1) = 1^2 \cdot \frac{2}{14} + 2^2 \cdot \frac{3}{14} + 3^2 \cdot \frac{4}{14} + 4^2 \cdot \frac{5}{14} = \frac{130}{14}.$$

And, finally,

$$\text{Var}(Y|X = 1) = \frac{130}{14} - \left(\frac{40}{14}\right)^2 \boxed{= \frac{220}{196}}.$$

## Solution

(d) From the graph in (c)

$$P(1 \leq Y \leq 3|X = 1) = \frac{2}{14} + \frac{3}{14} + \frac{4}{14} \boxed{= \frac{9}{14}}.$$

$$P(Y \leq 2|X = 2) = \frac{3}{18} + \frac{4}{18} \boxed{= \frac{7}{18}}.$$

From the graph in (b)

$$P(X = 2|Y = 3) \boxed{= \frac{5}{9}}.$$

(e)

$$E(Y|X = 1) = 1 \cdot \frac{2}{14} + 2 \cdot \frac{3}{14} + 3 \cdot \frac{4}{14} + 4 \cdot \frac{5}{14} \boxed{= \frac{40}{14}}.$$

$$E(Y^2|X = 1) = 1^2 \cdot \frac{2}{14} + 2^2 \cdot \frac{3}{14} + 3^2 \cdot \frac{4}{14} + 4^2 \cdot \frac{5}{14} = \frac{130}{14}.$$

And, finally,

$$\text{Var}(Y|X = 1) = \frac{130}{14} - \left(\frac{40}{14}\right)^2 \boxed{= \frac{220}{196}}.$$

### Definition

$$\mu_{Y|x} = E[Y|X = x] = \sum_{y \in \text{Range}(Y)} yh(y|x)$$

is called **conditional mean** of  $Y$  given  $X = x$ .

### Definition

$$\sigma_{Y|x}^2 = E[(Y - \mu_{Y|x})^2|X = x] = \sum_{y \in \text{Range}(Y)} [(y - \mu_{Y|x})^2 h(y|x)]$$

is called **conditional variance** of  $Y$  given  $X = x$ .

► The  $\mu_{X|y}, \sigma_{X|y}$  are defined in a similar way.



### Definition

$$\mu_{Y|x} = E[Y|X = x] = \sum_{y \in \text{Range}(Y)} yh(y|x)$$

is called **conditional mean** of  $Y$  given  $X = x$ .

### Definition

$$\sigma_{Y|x}^2 = E[(Y - \mu_{Y|x})^2|X = x] = \sum_{y \in \text{Range}(Y)} [(y - \mu_{Y|x})^2 h(y|x)]$$

is called **conditional variance** of  $Y$  given  $X = x$ .

► The  $\mu_{X|y}, \sigma_{X|y}$  are defined in a similar way.

### Definition

$$\mu_{Y|x} = E[Y|X = x] = \sum_{y \in \text{Range}(Y)} yh(y|x)$$

is called **conditional mean** of  $Y$  given  $X = x$ .

### Definition

$$\sigma_{Y|x}^2 = E[(Y - \mu_{Y|x})^2|X = x] = \sum_{y \in \text{Range}(Y)} [(y - \mu_{Y|x})^2 h(y|x)]$$

is called **conditional variance** of  $Y$  given  $X = x$ .

► The  $\mu_{X|y}, \sigma_{X|y}$  are defined in a similar way.

- Typically there is no function  $m : \text{Range}(X) \rightarrow \mathbb{R}$  so that  $y = m(x)$  for any  $(x, y) \in \text{Range}(X, Y)$ .
- For the same  $x$  there can be multiple  $y$ -s with  $(x, y) \in \text{Range}(X, Y)$ .

## Example


Represent the weights of students in class as function of their heights.

- Conditional mean

$$m(x) = E[Y|X = x]$$

is the “best” function representing  $y$  in terms of  $x$ .

- $\sigma_{Y|x}$  measures how much  $Y$  varies from  $m(x)$  when  $X = x$ .



- Typically there is no function  $m : \text{Range}(X) \rightarrow \mathbb{R}$  so that  $y = m(x)$  for any  $(x, y) \in \text{Range}(X, Y)$ .
- For the same  $x$  there can be multiple  $y$ -s with  $(x, y) \in \text{Range}(X, Y)$ .

## Example


Represent the weights of students in class as function of their heights.

- Conditional mean

$$m(x) = E[Y|X = x]$$

is the “best” function representing  $y$  in terms of  $x$ .

- $\sigma_{Y|x}$  measures how much  $Y$  varies from  $m(x)$  when  $X = x$ .



- ▶ Typically there is no function  $m : \text{Range}(X) \rightarrow \mathbb{R}$  so that  $y = m(x)$  for any  $(x, y) \in \text{Range}(X, Y)$ .
- ▶ For the same  $x$  there can be multiple  $y$ -s with  $(x, y) \in \text{Range}(X, Y)$ .

## Example


Represent the weights of students in class as function of their heights.

- ▶ Conditional mean

$$m(x) = E[Y|X = x]$$

is the “best” function representing  $y$  in terms of  $x$ .

- ▶  $\sigma_{Y|x}$  measures how much  $Y$  varies from  $m(x)$  when  $X = x$ .



- Typically there is no function  $m : \text{Range}(X) \rightarrow \mathbb{R}$  so that  $y = m(x)$  for any  $(x, y) \in \text{Range}(X, Y)$ .
- For the same  $x$  there can be multiple  $y$ -s with  $(x, y) \in \text{Range}(X, Y)$ .

## Example


Represent the weights of students in class as function of their heights.

- Conditional mean

$$m(x) = E[Y|X = x]$$

is the “best” function representing  $y$  in terms of  $x$ .

- $\sigma_{Y|x}$  measures how much  $Y$  varies from  $m(x)$  when  $X = x$ .



- Typically there is no function  $m : \text{Range}(X) \rightarrow \mathbb{R}$  so that  $y = m(x)$  for any  $(x, y) \in \text{Range}(X, Y)$ .
- For the same  $x$  there can be multiple  $y$ -s with  $(x, y) \in \text{Range}(X, Y)$ .

## Example


Represent the weights of students in class as function of their heights.

- Conditional mean

$$m(x) = E[Y|X = x]$$

is the “best” function representing  $y$  in terms of  $x$ .

- $\sigma_{Y|x}$  measures how much  $Y$  varies from  $m(x)$  when  $X = x$ .



- Among all functions  $m : \text{Range}(X) \rightarrow \mathbb{R}$ , conditional mean is the one that minimizes

$$m \mapsto E[(Y - m(X))^2] \quad (\text{homework}).$$

- Among all linear functions  $ax + b$ , the least squares line is the function that minimizes

$$E[(Y - aX - b)^2].$$

- Sometimes  $m(x)$  is a linear function: then conditional mean and least squares lines coincide.



- Among all functions  $m : \text{Range}(X) \rightarrow \mathbb{R}$ , conditional mean is the one that minimizes

$$m \mapsto E[(Y - m(X))^2] \quad (\text{homework}).$$

- Among all linear functions  $ax + b$ , the least squares line is the function that minimizes

$$E[(Y - aX - b)^2].$$

- Sometimes  $m(x)$  is a linear function: then conditional mean and least squares lines coincide.

- Among all functions  $m : \text{Range}(X) \rightarrow \mathbb{R}$ , conditional mean is the one that minimizes

$$m \mapsto E[(Y - m(X))^2] \quad (\text{homework}).$$

- Among all linear functions  $ax + b$ , the least squares line is the function that minimizes

$$E[(Y - aX - b)^2].$$

- Sometimes  $m(x)$  is a linear function: then conditional mean and least squares lines coincide.

## Trinomial distribution case

- Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$

- $p_S + p_I + p_F = 1$ .
- $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .



## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case

- ▶ Consider the trinomial distribution with parameters  $(n, P_S, P_I, p_F)$
- ▶  $p_S + p_I + p_F = 1$ .
- ▶  $\text{Range}(X, Y) = \{(x, y) : x + y \leq n\}$ .
- ▶  $f(x, y) = \binom{n}{x} \binom{n-x}{y} p_S^x p_I^y (1 - p_S - p_I)^{n-x-y}$ .
- ▶  $f_X(x) = \binom{n}{x} p_S^x (1 - p_S)^{n-x}$ .

$$\begin{aligned} h(y|x) &= \frac{f(x, y)}{f_X(x)} = \binom{n-x}{y} \frac{p_I^y (1 - p_S - p_I)^{n-x-y}}{(1 - p_S)^{n-x}} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( \frac{1 - p_S - p_I}{1 - p_S} \right)^{n-x-y} \\ &= \binom{n-x}{y} \left( \frac{p_I}{1 - p_S} \right)^y \left( 1 - \frac{p_I}{1 - p_S} \right)^{n-x-y}. \end{aligned}$$

- ▶ Let  $p = \frac{p_I}{1 - p_S}$ , then

$$h(y|x) = \binom{n-x}{y} p^y (1 - p)^{n-x-y}.$$

- ▶ The conditional pmf is a Binomial distribution with parameters  $(n - x, p)$ .

## Trinomial distribution case (cont)

- Using the formula for the mean and variance of the Binomial distribution with parameters  $(n - x, p)$ .

$$\mu_{Y|x} = (n - x)p = (n - x) \frac{p_I}{1 - p_S}$$

$$\sigma_{Y|x}^2 = (n - x)p(1 - p) = (n - x) \frac{p_I p_F}{(1 - p_S)^2}.$$

- In conclusion,  $m(x) = \mu_{Y|x}$  is a linear function of  $x$ .
- Therefore the least squares line of the trinomial distribution with parameters  $(n, p_S, p_I)$  is

$$y = (n - x) \frac{p_I}{1 - p_S}.$$

- The distribution is closest to linear when either  $p_I \approx 0$  or  $p_F \approx 0$  but not both.

## Trinomial distribution case (cont)

- Using the formula for the mean and variance of the Binomial distribution with parameters  $(n - x, p)$ .

$$\mu_{Y|x} = (n - x)p = (n - x) \frac{p_I}{1 - p_S}$$
$$\sigma_{Y|x}^2 = (n - x)p(1 - p) = (n - x) \frac{p_I p_F}{(1 - p_S)^2}.$$

- In conclusion,  $m(x) = \mu_{Y|x}$  is a linear function of  $x$ .
- Therefore the least squares line of the trinomial distribution with parameters  $(n, p_S, p_I)$  is

$$y = (n - x) \frac{p_I}{1 - p_S}.$$

- The distribution is closest to linear when either  $p_I \approx 0$  or  $p_F \approx 0$  but not both.

## Trinomial distribution case (cont)

- Using the formula for the mean and variance of the Binomial distribution with parameters  $(n - x, p)$ .

$$\mu_{Y|x} = (n - x)p = (n - x) \frac{p_I}{1 - p_S}$$
$$\sigma_{Y|x}^2 = (n - x)p(1 - p) = (n - x) \frac{p_I p_F}{(1 - p_S)^2}.$$

- In conclusion,  $m(x) = \mu_{Y|x}$  is a linear function of  $x$ .
- Therefore the least squares line of the trinomial distribution with parameters  $(n, p_S, p_I)$  is

$$y = (n - x) \frac{p_I}{1 - p_S}.$$

- The distribution is closest to linear when either  $p_I \approx 0$  or  $p_F \approx 0$  but not both.

## Trinomial distribution case (cont)

- Using the formula for the mean and variance of the Binomial distribution with parameters  $(n - x, p)$ .

$$\mu_{Y|x} = (n - x)p = (n - x) \frac{p_I}{1 - p_S}$$
$$\sigma_{Y|x}^2 = (n - x)p(1 - p) = (n - x) \frac{p_I p_F}{(1 - p_S)^2}.$$

- In conclusion,  $m(x) = \mu_{Y|x}$  is a linear function of  $x$ .
- Therefore the least squares line of the trinomial distribution with parameters  $(n, p_S, p_I)$  is

$$y = (n - x) \frac{p_I}{1 - p_S}.$$

- The distribution is closest to linear when either  $p_I \approx 0$  or  $p_F \approx 0$  but not both.



## Trinomial distribution case (cont)

- Using the formula for the mean and variance of the Binomial distribution with parameters  $(n - x, p)$ .

$$\mu_{Y|x} = (n - x)p = (n - x) \frac{p_I}{1 - p_S}$$

$$\sigma_{Y|x}^2 = (n - x)p(1 - p) = (n - x) \frac{p_I p_F}{(1 - p_S)^2}.$$

- In conclusion,  $m(x) = \mu_{Y|x}$  is a linear function of  $x$ .
- Therefore the least squares line of the trinomial distribution with parameters  $(n, p_S, p_I)$  is

$$y = (n - x) \frac{p_I}{1 - p_S}.$$

- The distribution is closest to linear when either  $p_I \approx 0$  or  $p_F \approx 0$  but not both.

## Trinomial distribution case (cont)

- Using the formula for the mean and variance of the Binomial distribution with parameters  $(n - x, p)$ .


$$\mu_{Y|x} = (n - x)p = (n - x) \frac{p_I}{1 - p_S}$$

$$\sigma_{Y|x}^2 = (n - x)p(1 - p) = (n - x) \frac{p_I p_F}{(1 - p_S)^2}.$$


- In conclusion,  $m(x) = \mu_{Y|x}$  is a linear function of  $x$ .
- Therefore the least squares line of the trinomial distribution with parameters  $(n, p_S, p_I)$  is

$$y = (n - x) \frac{p_I}{1 - p_S}.$$

- The distribution is closest to linear when either  $p_I \approx 0$  or  $p_F \approx 0$  but not both.



# Affect of conditional variance



► The shaded area is the region between

$$[m(x) - \sigma_{Y|x}, m(x) + \sigma_{Y|x}]$$