

# Line plot basics

your name

2024-10-22

## Math 2265 Chapter 8. Linear Regression

- Work as a group!
  - You will need to replace "ans" or `your_answer` in the source code
  - Update your name in L3
  - Add your group members' name below; students may lose one point if Question 0 is unanswered
  - Make sure you save and `knit` your work (to html or pdf) before submitting it to Canvas
- 

### Goal

- Review linear (line) equations
  - Learn how to define a `function` in R
  - Review how to define a `data frame` (data set) in R
  - Review vector calculations in R
  - Learn how to plot lines from a (linear) data set
- 

**Question 0. Who are your group members? (List their first names should be sufficient)**

**Answer:**

1. `<name_1>`
  2. `<name_2>`
- 

**If you need more time to get used to Markdown, use the Visual mode.**

The icon is located in the upper-left corner next to `source`.

---

## Functions

Roughly speaking, a function is an assignment that for each value of  $x$ , there is exactly one output  $y$ . Often they are denoted by  $f(x) = \text{expression in } x$  or  $y = \text{expression in } x$ .

Here are some examples of functions.

Linear:

1.  $f(x) = x$
2.  $f(x) = -3x + 1$

Quadratic:

1.  $f(x) = x^2$
2.  $f(x) = -(x - 3)^2 + 5$ .

### Example 1: Compute the $y$ values of a given function

We will use  $f(x) = -3x + 1$  as an example but it applies to any function. Below, we compute  $f(1), f(2), f(3)$ .

```
-3*(1) + 1
```

```
## [1] -2
```

```
-3*(2) + 1
```

```
## [1] -5
```

```
-3*(3) + 1
```

```
## [1] -8
```

Indeed, it is often desired and convenient to use vectors.

```
inputs <- c(1,2,3)
```

```
-3*inputs + 1
```

```
## [1] -2 -5 -8
```

### Task 1: Do the same for $f(x) = -(x - 3)^2 + 5$

Compute  $f(1), f(2), f(3)$ , where  $f(x) = -(x - 3)^2 + 5$ :

```
-(1-3)^2 + 5
```

```
## [1] 1
```

```
-(2-3)^2 + 5
```

```
## [1] 4
```

```
-(3-3)^2 + 5
```

```
## [1] 5
```

Use the vector method to compute the same.

```
inputs <- c(1,2,3)
```

```
-(inputs - 3)^2 + 5
```

```
## [1] 1 4 5
```

### Example 2: Define a function

Let's revisit Example 1, where  $f(x) = -3x + 1$ . This time we will define the function  $f(x)$  first and use it to compute the rest.

```
#define f(x) = -3x + 1
my_func <- function(x){
  -3*x + 1 # be sure to put * to denote multiplication
}
```

```
my_func(1); my_func(2); my_func(3)
```

```
## [1] -2
```

```
## [1] -5
```

```
## [1] -8
inputs <- c(1,2,3)
my_func(inputs)
```

```
## [1] -2 -5 -8
```

**Task 2: Do the same for  $f(x) = -(x - 3)^2 + 5$**

```
my_func2 <- function(x) {
  -(x - 3)^2 + 5
}

my_func2(1); my_func2(2); my_func2(3)
```

```
## [1] 1
```

```
## [1] 4
```

```
## [1] 5
```

Using a vector:

```
inputs <- c(1, 2, 3)
my_func2(inputs)
```

```
## [1] 1 4 5
```

**Example 3: Plot the line  $f(x) = 3x - 1$**


We use the function `my_func` defined above to plot the line  $f(x) = 3x - 1$ . First we make a scatter plot, and then we add a line. To use `ggplot2`, we need to make a data frame consisting of  $x$  and  $y$ .

```
x <- c(-10:10) # c(-10:10) creates a vector -10,-9,...,10
y <- my_func(x)
my_dataframe <- data.frame(
  x,
  y
)
str(my_dataframe)
```

```
## 'data.frame':  21 obs. of  2 variables:
## $ x: int  -10 -9 -8 -7 -6 -5 -4 -3 -2 -1 ...
## $ y: num  31 28 25 22 19 16 13 10 7 4 ...
```


First make a scatter plot:

```
ggplot(data=my_dataframe, mapping = aes(x=x, y=y)) +
  geom_point()
```




We can connect the dots by adding `geom_line()`:

```
ggplot(data=my_dataframe, mapping = aes(x=x, y=y)) +  
  geom_point() +  
  geom_line()
```




Indeed, there is a function `curve` which sketches the function  $f(x)$  without needing to define a data frame.

```
curve(my_func, from=-10,to=10)
```




```
curve(-(x-3)^2 + 5, from=-7, to=13)
```




How-  
ever, for the purpose of linear regression, we will stick to the `ggplot2` package. For lines, the `ggplot2` package has the `abline` function which takes the slope and intercept of a line as parameters. Here is an example for  $f(x) = -3x + 1$  whose slope is  $-3$  and  $y$ -intercept is  $1$ .

```
ggplot() +  
  geom_abline(slope = -3, intercept = 1) +  
  xlim(-10,10) +  
  ylim(-32,32)
```



**Question:**


We will create the possum example in the book link.



- data set: possum
- explanatory variable: total\_l
- response variable: head\_l

```
ggplot(data = possum, mapping = aes(x=total_l, y=head_l)) +  
  geom_point()
```






(a) Make a scatter plot

(b) Add the line with slope 0.59 and intercept 41


```
ggplot(data = possum, mapping = aes(x=total_l, y=head_l)) +  
  geom_point() +  
  geom_abline(slope = 0.59, intercept = 41)
```



### (c) Decorations

We have not been concerned aesthetics so far, but with the a few commands we can make it look similar to the one in the book. Change `red` to the matching color and “`x_title`” and “`y_title`” by “Total Length (cm)” and “Head Length (mm)”, respectively.

```
ggplot(data = possum, mapping = aes(x=total_l, y=head_l)) +
  geom_point(color='red') +
  geom_abline(slope = 0.59, intercept = 41) +
  xlab("Total Length (cm)") +
  ylab("Head Length (mm)")
```



Share your work and help your group members before uploading your work to Canvas