

Parton Distribution Functions

arXiv:hep-lat/9609018

Youngwan Kim

Seoul National University

SNUCMS arXiv Seminar

February 19, 2021

Outline

1. Introduction to PDFs

- Intuition to PDFs

- Factorization

- Significance of PDFs

2. Technical Definition of PDFs

- Technical Definition of PDFs

- Gauge Invariance

- Renormalization

Intuition to PDFs




Figure: Hadron A + Hadron B \rightarrow 2 partons

Consider a collision of two hadrons A and B...

- parton a from hadron A with ξ_A fraction of its momentum
 - ▶ $f_{a/A}(\xi_A) d\xi_A$
- parton b from hadron B with ξ_B fraction of its momentum
 - ▶ $f_{b/B}(\xi_B) d\xi_B$
- $d\hat{\sigma}$ parton level cross section

One can describe $d\sigma$ as a product of these three factors...

Factorization

Hadron Scattering Cross Section

$$\frac{d\sigma}{dp_T} \sim \sum_{a,b} \int d\xi_A f_{a/A}(\xi_A, \mu) \int d\xi_B f_{b/B}(\xi_B, \mu) \frac{d\hat{\sigma}}{dp_T}$$

Parton Level Cross Section

$$\frac{d\hat{\sigma}}{dp_T} \sim \sum_N \left(\frac{\alpha_S(\mu)}{\pi} \right)^N H_N(\xi_A, \xi_B, p_T; a, b, \mu)$$

- The coefficients H_N are calculable via pQCD ; regard factorization as an established theorem of QCD

Another parameter μ ? : there is more to the equation than just a model

- Has dimension of mass \rightarrow renormalization of $\alpha_S(\mu)$ and $f_{a/A}(\xi_A)$
- $d\hat{\sigma}/dp_T$ is calculated straightforwardly at Born level
 - ▶ Divergences appear starting from NLO and beyond, calculation gets tricky
 - ▶ Such divergences are removed, and in their place dependence of the scale μ appears.

Significance of PDFs

Knowledge of PDFs is necessary for the description of hard processes,

Parton-Parton Cross Section (ex. LHC, Fermilab)

$$d\sigma \sim \sum_{a,b} \int d\xi_A f_{a/A}(\xi_A, \mu) \int d\xi_B f_{b/B}(\xi_B, \mu) d\hat{\sigma}$$

DIS Cross Section (ex. HERA)

$$d\sigma \sim \sum_a \int d\xi_A f_{a/A}(\xi_A, \mu) d\hat{\sigma}$$

Obvious as no prediction is available without any knowledge of PDFs.

Significance of PDFs

For both expressions of $d\sigma$ it shows that in high energy, short distance collisions...

- a hard scattering probes the system quickly
- strong binding forces act slowly

→ One needs to know probabilities to find partons in a **fast moving** hadron as seen by an approximately instantaneous probe

→ PDFs should provide a **relativistic** view of partons

Technical Definition of PDFs

- Coordinates : $x^\pm = (x^0 \pm x^3)/\sqrt{2}$, $P^\pm = (P^0 \pm P^3)/\sqrt{2}$, $v = (v^+, v^-, \mathbf{v}_T)$




Figure: partons in a fast moving proton

- ▶ x^+ can be interpreted as time
- ▶ partons in a proton with a small P^- , big P^+ and $\vec{P}_T = 0$, move roughly parallel to the x^+ axis.
- ▶ PDFs are to be defined via **null plane field theory**, a field theory quantized on planes of equal x^+ .
- ▶ $P^\mu x_\mu = P^+ x^- + P^- x^+ - \mathbf{P}_T \cdot \mathbf{x}_T$

Technical Definition of PDFs

We want to define $f_{i/A}(\xi, \mu)$, the probability density for ...

- finding a parton of flavor i (quark, antiquark, gluon ...)
- which carries a fraction ξ of P^+
- in the hadron A with momentum P .

→ Approach by defining a **density operator** using creation, annihilation operators of null plane field theory

Technical Definition of PDFs

Consider $b(\xi P^+, \mathbf{k}_T, s, c; i)$ which annihilates (b^\dagger creates)

- a quark with flavor i , having helicity s and color c
- carrying a fraction ξ of P^+ of the proton with transverse momentum \mathbf{k}_T
- $b(\xi P^+, \mathbf{k}_T, s, c; i) |0\rangle = 0$ for the vacuum state $|0\rangle$
- normalized by the following anticommutation relation :

Anticommutation relation of b, b^\dagger

$$\{b(\xi' P^+, \mathbf{k}'_T, s', c'; i), b(\xi P^+, \mathbf{k}_T, s, c; i)\} = (2\pi)^3 2\xi P^+ \delta(\xi' P^+ - \xi P^+) \delta(\mathbf{k}'_T - \mathbf{k}_T) \delta_{s',s} \delta_{c',c}$$

Technical Definition of PDFs

With b and b^\dagger , one could define $\rho(\xi P^+, \mathbf{k}_T; i)$ which **counts the number of quarks** in a region of ξ and \mathbf{k}_T

Definition of ρ

$$\begin{aligned}\rho(\xi P^+, \mathbf{k}_T; i) \\ = \frac{1}{(2\pi)^3 2\xi} \sum_{s,c} b^\dagger(\xi P^+, \mathbf{k}_T, s, c; i) b(\xi P^+, \mathbf{k}_T, s, c; i)\end{aligned}$$

Technical Definition of PDFs

If $|\Psi\rangle$ is obtained by applying b to the vacuum, ρ counts the number $N(\mathcal{V})$ of quarks in the momentum-space volume \mathcal{V} when integrated,

$$\int_{\mathcal{V}} d\xi d\mathbf{k}_T \rho(\xi P^+, \mathbf{k}_T; i) |\Psi\rangle = N(\mathcal{V}) |\Psi\rangle$$

Here we can take a matrix element of ρ in a proton state to define $f_{i/A}(\xi)$

Preliminary definition of $f_{i/A}(\xi)$ †

$$f_{i/A}(\xi) \langle P' | P \rangle = \int d\mathbf{k}_T \langle P' | \rho(\xi P^+, \mathbf{k}_T; i) | P \rangle$$

Technical Definition of PDFs

In order to make this construction more useful, we relate b, b^\dagger with the quark field operator $\psi_i(x)$ quantized on the null plane as ...

$$P_{dy}\psi_{i,c}(x^+ = 0, x^-, \mathbf{x}_T) = \frac{1}{(2\pi)^3} \int_0^\infty \frac{dk^+}{2k^+} \int d\mathbf{k}_T \\ \times \sum_s \left\{ P_{dy}u(k, s) e^{-ik \cdot x} b(k^+, \mathbf{k}_T, s, c; i) + P_{dy}v(k, s) e^{+ik \cdot x} d^\dagger(k^+, \mathbf{k}_T, s, c; i) \right\}$$

- The **two components** projected by $P_{dy} = \frac{1}{2}\gamma^-\gamma^+$ are the independent dynamical fields representing quarks.
- $u(k, s), v(k, s)$: usual spinor solutions of the free Dirac eq, normalized to $\bar{u}(k, s)\gamma^+u(k, s) = 2k^+$ (same with $v(k, s)$)
- $d^\dagger(k^+, \mathbf{k}_T, s, c; i)$: antiquark creation operator (analogous to b^\dagger)

Technical Definition of PDFs

Then plugging in the above expression into the preliminary definition of $f_{i/A}(\xi)$,

Preliminary definition of $f_{i/A}(\xi)$ II

$$f_{i/A}(\xi) \langle P' | P \rangle = \frac{P^+}{2\pi} \int dy^- e^{-i\xi P^+ y^-} \int dx^- d\mathbf{x}_T \\ \times \langle P' | \bar{\psi}_i(0, x^- + y^-, \mathbf{x}_T) \gamma^+ \psi_i(0, x^-, \mathbf{x}_T) | P \rangle$$

Technical Definition of PDFs

and implementing the following identities :

- Translation invariance :

$$\begin{aligned} & \langle P' | \bar{\psi}_i(0, x^- + y^-, \mathbf{x}_T) \gamma^+ \psi_i(0, x^-, \mathbf{x}_T) | P \rangle \\ &= \exp \left[i(P' - P) \cdot x^- - (\mathbf{P}'_T - \mathbf{P}_T) \cdot \mathbf{x}_T \right] \times \langle P' | \bar{\psi}_i(0, y^-, \mathbf{0}) \gamma^+ \psi_i(\mathbf{0}) | P \rangle \end{aligned}$$

- Proton state normalization : $\langle P' | P \rangle = (2\pi)^3 2P^+ \delta(P'^+ - P^+) \delta(\mathbf{P}'_T - \mathbf{P}_T)$

Preliminary definition of $f_{i/A}(\xi)$

$$f_{i/A}(\xi) = \frac{1}{4\pi} \int dy^- e^{-i\xi P^+ y^-} \langle P | \bar{\psi}_i(0, y^-, \mathbf{0}) \gamma^+ \psi_i(0) | P \rangle$$

Technical Definition of PDFs

Preliminary definition of $f_{i/A}(\xi)$

$$f_{i/A}(\xi) = \frac{1}{4\pi} \int dy^- e^{-i\xi P^+ y^-} \langle P | \bar{\psi}_i(0, y^-, \mathbf{0}) \gamma^+ \psi_i(0) | P \rangle$$

Things to notice from $f_{i/A}(\xi)$...

- It is an expectation value in the hadron state of a certain **operator**.
- This operator is not local but **bilocal** : $(0, y^-, \mathbf{0})$ and $(0, 0, \mathbf{0})$ are light-like separated.
- Integrate over y^- with the right factor so that we annihilate a quark with ξP^+ momentum.

→ Gluons ...? Gauge invariance ...? Renormalization ...?

Gauge Invariance

The $f_{i/A}(\xi)$ above relies on the gluon potential A^μ being in the lightlike axial gauge $A^+ = 0$, so it would be nice to ...

- make the **operator** gauge invariant
- match the previous definition in $A^+ = 0$ gauge

→ Solution : insert a Wilson line factor \mathcal{O} into the operator

Wilson line factor

$$\mathcal{O}(y^-, 0) = \mathcal{P} \exp \left[ig \int_0^{y^-} dz^- A^+(0, z^-, \mathbf{0})_a t_a \right]$$

$t_a : SU(3)_c$ generator matrix in the **3** representation

\mathcal{P} : path ordering with more positive y^- values to the left

Gauge Invariance

Then the revised gauge invariant $f_{i/A}(\xi)$ is defined as :

Gauge invariant definition of $f_{i/A}(\xi)$

$$f_{i/A}(\xi) = \frac{1}{4\pi} \int dy^- e^{-i\xi P^+ y^-} \langle P | \bar{\psi}_i(0, y^-, \mathbf{0}) \gamma^+ \mathcal{O}(y^-, 0) \psi_i(0) | P \rangle$$

one can immediately see that the gauge $A^+ = 0$ corresponds to $\mathcal{O} = 1$, being consistent with the previous definition of $f_{i/A}(\xi)$

Gauge Invariance

Checking gauge invariance under an unitary transformation $U(x)$:

$$\begin{aligned}\psi_i(0) &\rightarrow U(0)\psi_i(0) \\ \bar{\psi}_i(0, y^-, \mathbf{0}) &\rightarrow \bar{\psi}_i(0, y^-, \mathbf{0})U(0, y^-, \mathbf{0})^{-1} \\ \mathcal{O}(y^-, 0) &\rightarrow U(0, y^-, \mathbf{0})\mathcal{O}(y^-, 0)U(0)^{-1}\end{aligned}$$

Thus including $\mathcal{O}(y^-, 0)$, the **operator** maintains invariant under a change of gauge.

Besides gauge invariance what does \mathcal{O} offer? Is there any physical significance ...? A mere technique?

Gauge Invariance




Figure: Effect of the \mathcal{O} operator

- $\psi_i(0)$: absorbs a quark line from the wave function
- $\bar{\psi}_i(0, y^-, \mathbf{0})$: creates a quark line that goes into the conjugate wave function
- $\mathcal{O}(y^-, 0)$: contains gluon fields that create and absorb gluons
 - ▶ not just destroying a quark at position 0, leaving its color nowhere to go
 - ▶ but rather scatter it, so that it moves to infinity along a fixed light-like line in the minus direction, carrying its color with it.
 - ▶ then its color comes back to $(0, y^-, \mathbf{0})$ to provide the color for the quark that we create

→ How in detail?

Gauge Invariance

Using the following identity satisfied by any ordered exponential :

$$\mathcal{P} \exp \left[ig \int_0^\eta d\lambda n \cdot A(\lambda n^\mu) \right] = \left[\mathcal{P} \exp \left[ig \int_0^\infty d\lambda n \cdot A((\lambda + \eta)n^\mu) \right] \right]^\dagger \mathcal{P} \exp \left[ig \int_0^\infty d\lambda n \cdot A(\lambda n^\mu) \right]$$

substituting $A^\mu = A_a^\mu t_a$, $\eta = y^-$ and $n = (n^+, n^-, \mathbf{n}_T) = (0, 1, \mathbf{0})$ which leads to $n \cdot A = n^+ A^- + n^- A^+ - \mathbf{n}_T \cdot \mathbf{A}_T = A^+$ and $\lambda n^\mu = z^-(0, 1, \mathbf{0}) = (0, z^-, \mathbf{0})$, gives

$$\mathcal{P} \exp \left[ig \int_0^{y^-} dz^- A^+(0, z^-, \mathbf{0})_a t_a \right] = \left[\mathcal{P} \exp \left[ig \int_{y^-}^\infty dz^- A^+(0, z^-, \mathbf{0})_a t_a \right] \right]^\dagger \mathcal{P} \exp \left[ig \int_0^\infty dz^- A^+(0, z^-, \mathbf{0})_a t_a \right]$$

Gauge Invariance

Which shows that $\mathcal{O}(y^-, 0) = \mathcal{O}(y^-, \infty)\mathcal{O}(\infty, 0)$ and using this expression, one could insert a complete set $|n\rangle \langle n|$ and write

$$f_{i/A}(\xi) = \frac{1}{4\pi} \int dy^- e^{-i\xi P^+ y^-} \sum_n \langle P | \bar{\Psi}(0, y^-, \mathbf{0}) | n \rangle \gamma^+ \langle n | \Psi(0) | P \rangle$$


$$\Psi(x) = \psi(x) \mathcal{P} \exp \left[ig \int_0^\infty dz^- (x^+, x^- + z^-, \mathbf{x}_T)_a t_a \right]$$

From this one can extract Feynman rules via expanding the ordered exponentials and expressing them in momentum space :

$$\mathcal{P} \left[ig \int_0^\infty dz^- A^+(0, z^-, \mathbf{0})_a t_a \right] =$$

$$1 + \mathcal{P} \sum_{n=1}^\infty \prod_{i=1}^n \int \frac{d^4 q_i}{(2\pi)^4} g \tilde{A}^+(q_i^\mu) \frac{1}{\sum_{j=1}^i q_j^- + i\epsilon}$$

Gauge Invariance



- One can read off the Feynman rules from the expansion.
- The **denominators** are represented by double lines, which we shall refer to **eikonal** lines
- They attach to the gluon propagator via a vertex proportional to $-ign^\mu$

Renormalization

The function $f_{i/A}(\xi)$ is **unrenormalized**, being defined using bare fields, a bare coupling, bare parton masses ... and so on

- would like to introduce the \overline{MS} scheme
- perform integration in $4 - 2\epsilon$ dimensions, including a factor that keeps the dimension constant

Renormalized, gauge invariant definition of $f_{i/A}(\xi, \mu)$

$$f_{i/A}(\xi, \mu) = \frac{1}{4\pi} \int dy^- e^{-i\xi P^+ y^-} \langle P | \bar{\psi}_i(0, y^-, \mathbf{0}) \gamma^+ \mathcal{O}(y^-, 0) \psi_i(0) | P \rangle_{\overline{MS}}$$

Antiquarks and Gluons

Will simply state results ...

- Antiquarks : utilize charge conjugation, $\bar{\mathbf{3}}$ representation generators for \mathcal{O}

$$f_{i/A}(\xi, \mu) = \frac{1}{4\pi} \int dy^- e^{-i\xi P^+ y^-} \langle P | \text{Tr} \left[\gamma^+ \psi_i(0, y^-, \mathbf{0}) \mathcal{O}(y^-, 0) \bar{\psi}_i(0) \right] | P \rangle_{\overline{MS}}$$

- Gluons : $A^+ = 0$ gauge, $\mathbf{8}$ representation generators for \mathcal{O}

$$f_{g/A}(\xi, \mu) = \frac{1}{2\pi x P^+} \int dy^- e^{-i\xi P^+ y^-} \langle P | F_a^+(0, y^-, \mathbf{0})^\nu \mathcal{O}_{ab} F_b^+(0)^\nu | P \rangle_{\overline{MS}}$$

Renormalization Group

- A change in the scale μ affects $f_{a/A}(\xi, \mu)$, which comes from the change in the amount of UV divergence that renormalization is removing.
- By examining the structure of UV divergences, one finds $f_{a/A}(\xi, \mu)$ obeys DGLAP evolution equations,


$$\mu^2 \frac{d}{d\mu^2} f_{a/A}(\xi, \mu) = \sum_b \int_{\xi}^1 \frac{dz}{z} P_{ab}(z, \alpha_S(\mu^2)) f_{b/A}(\xi/z, \mu)$$

- ▶ The evolution kernels P_{ab} are perturbatively calculable as expansions in powers of $\alpha_S(\mu)$
- ▶ Exact evolution kernels have been known up to NNLO (2004),

Renormalization Group

Interpretation? The parameter μ corresponds to the physical resolving power of the probe.

- Parton splitting is not well observed at low resolving power
- At higher μ which is enough to resolve mother parton into its daughters, observing a splitting



- ▶ One can find parton distributions at μ when it is already known at $\mu_0 < \mu$, using the RG equation.
- ▶ Gluon distribution at $\mu = 10, 100 \text{ GeV}$ using the CTEQ3M PDF set
- ▶ With greater resolution, smaller momentum fraction due to parton splitting.

Summary

- PDFs can be constructed as a hadron matrix element of the number density operator for quarks.
- PDFs should be renormalized due to divergences from NLO and beyond.
- Renormalization scale corresponds to the physical resolving power of the probe.

Thank You