



Search for HNLs Using τ Decays at *BABAR*

arXiv:2207.09575v1 [hep-ex] ; BABAR-PUB-22/002

Youngwan Kim
arXiv Presentation [8th Dec. 2023]



Introduction

Motivation

- Searching for HNLs
 - Obvious motivations ... SM neutrino sector anomalies
 - ν -MSM as a potential extension : offers HNLs
- Several mass scales available
 - GeV to TeV scale search by HNL decaying channels
 - sub-GeV scale region could be probed by decay modes of hadrons



Introduction

Current Bounds

- HNL Search from Decay Signatures
 - $|U_{eN}|^2$ and $|U_{\mu N}|^2$ limits from pions and kaons leptonic decay
 - E949 (BNL ; $K^+ \rightarrow \mu^+ N_\mu$) , PIENU (TRIUMF ; $\pi^\pm \rightarrow e^\pm N_e$) ... etc
 - $|U_{\tau N}|^2$ limits from ...
 - ArgoNeuT (Fermilab ; $N \rightarrow \nu e^+ e^-$) @ 280 ~ 970 MeV , direct search
 - NOMAD (CERN ; $D_S \rightarrow \tau N_\tau \rightarrow \tau \nu_\tau e^+ e^-$) @ 10 ~ 290 MeV

BABAR Experiment

Collider and Detector Introduction



- PEP-II Accelerator (SLAC)
 - Asymmetric e^+e^- B-factory with at least 4.5 GeV energy difference
 - COM energy optimized for $\Upsilon(4S)$ production , $\Gamma(\Upsilon(4S) \rightarrow B\bar{B}) \sim 96\%$
 - Energy difference to produce a moving COM frame that allows a detector to time-order the decay of the two B mesons produced in the formation of the $\Upsilon(4S)$ resonant state

Resonance	Run	Month/Year
$\Upsilon(4S)$	Run 1	10/1999 – 10/2000
	Run 2	02/2001 – 06/2002
	Run 3	12/2002 – 06/2003
	Run 4	09/2003 – 07/2004
	Run 5	04/2005 – 08/2006
	Run 6	01/2007 – 09/2007
$\Upsilon(3S)$	Run 7	12/2007 – 02/2008
$\Upsilon(2S)$	Run 7	02/2008 – 03/2008




BABAR Experiment

Collider and Detector Introduction

- PEP-II Accelerator (SLAC)

- Asymmetric e^+e^- B-factory with at least 4.5 GeV




produced in the formation of the $\Upsilon(4S)$ resonant state

BABAR Experiment

Collider and Detector Introduction



- Silicon Vertex Tracker (SVT) - provides precise position information on charged tracks, and is also the sole tracking device for very low-energy particles.
- Drift Chamber (DCH) - provides the main momentum measurements for charged particles and helps in particle identification through dE/dx measurements.
- Detector of Internally Reflected Cerenkov radiation (DIRC) - provides charged hadron identification.
- Electromagnetic Calorimeter (EMC) - provides particle identification for electrons, neutrals, electromagnetic particles, and hadrons.
- Solenoid (not a subdetector) - provides 1.5 T magnetic field for needed for charge and momentum measurements.
- Instrumented Flux Return (IFR) - provides muon and neutral hadron identification.




Detector Resolution

$$\frac{\sigma_E}{E} = \frac{(2.32 \pm 0.30)\%}{\sqrt[4]{E}} \oplus (1.82 \pm 0.12)\%$$

$$\frac{\sigma_{p_T}}{p_T} = (0.13 \pm 0.01)\% \cdot p_T + (0.45 \pm 0.03)\%$$

$$\sigma_{\theta,\phi} = \left(\frac{3.87 \pm 0.07}{\sqrt{E}} \pm 0.04 \right) \text{mrad}$$






Search Strategy

HNL Signatures in τ Decays

- Kinematics
 - HNL as a τ decay product differs the kinematics of the visible byproducts.
 - Recoiling effect from the HNL to the hadrons
 - Will be dependent by the HNL mass hypotheses
 - Consider a hadronic τ decay as a 2-body process which consists of the hadronic system + HNL

$$\tau^\pm \rightarrow h^\pm(E_h, \vec{p}_h) + N(E_N, \vec{p}_N)$$





Search Strategy

HNL Signatures in τ Decays

- Phase Space


- For a given value of m_h the range of E_h is determined as ...

$$E_\tau - \sqrt{m_N^2 + q_+^2} < E_h < E_\tau - \sqrt{m_N^2 + q_-^2}$$

- And m_h is allowed between ...

$$3m_\pi < m_h < m_\tau - m_N$$

Phys. Rev. D 91, 053006 (2015)





Search Strategy

HNL Signatures in τ Decays

- Phase Space


- For a given value of m_h , the range of E_h is determined as ...
(Shape of interest)

$E_\tau -$ Presence of an HNL in a τ decay introduces a *curved crescent-shaped endpoint* in the m_h - E_h phase space

- And m_h is allowed between ...

$$3m_\pi < m_h < m_\tau - m_N$$

[Phys. Rev. D 91, 053006 \(2015\)](#)






Search Strategy

HNL Signatures in τ Decays

- Topology based initial selection
 - **1-3 topology** signature
 - “tag-side” with **1-prong leptonic τ**
 - Branching ratio $\sim 35\%$
 - Suppression of hadronic bkg events
 - “signal-side” with **3-prong hadronic τ**
 - 4 well-reconstructed charged particles
 - Total charge should be 0
 - Spatially well-separated tag-signal hemisphere





Samples

Data & MC Simulation


- Data
 - Corresponding to integrated luminosity of 424/fb
 - $\sigma(e^+e^- \rightarrow \tau^+\tau^-) = (0.919 \pm 0.003) \text{ nb}$, $\sim 4 \times 10^8 \tau$ pairs produced
- Simulation
 - Produced via KK2F generator + TAUOLA decay simulation
 - Background
 - τ backgrounds : $\tau^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp \nu$ (SM) + other decay modes that could be misidentified as 3-prong topology
 - Non- τ backgrounds : $B^0 \bar{B}^0$, $B^+ B^-$, $q\bar{q}$, $\mu^+ \mu^-$
 - Signal (26 = 13*2)
 - 100 MeV incremental HNL mass step between 100 ~ 1300 MeV (13)
 - Both charges for tau considered (2)

Samples

Data & MC Simulation



- Data







$m_4 = 1200 \text{ MeV}/c^2$ $m_4 = 1000 \text{ MeV}/c^2$ $m_4 = 800 \text{ MeV}/c^2$ $m_4 = 600 \text{ MeV}/c^2$ $m_4 = 400 \text{ MeV}/c^2$ $m_4 = 200 \text{ MeV}/c^2$
 $m_4 = 1200 \text{ MeV}/c^2$ $m_4 = 1000 \text{ MeV}/c^2$ $m_4 = 800 \text{ MeV}/c^2$ $m_4 = 600 \text{ MeV}/c^2$ $m_4 = 400 \text{ MeV}/c^2$ $m_4 = 200 \text{ MeV}/c^2$

- Non- τ backgrounds : $B^0\bar{B}^0, B^+B^-, q\bar{q}, \mu^+\mu^-$

- Signal

- 100 MeV incremental HNL mass step between 100 ~ 1300 MeV

- Both charges for tau considered





Event Selection

Data & MC Simulation


- Thrust of event > 0.85
 - suppress spherically symmetric shaped hadronic background events
- Reject events with no missing momentum
 - Require missing momentum $> 0.9\% \sqrt{s}$
- Track cuts to ensure PID performance
 - Should be within the acceptance of the DIRC and EMC ($-0.76 < \cos\theta < 0.9$)
 - $pT > 250$ MeV , which is the minimum to reach the DIRC
- Events containing conversion daughter tracks are removed
- EMC energy deposit cuts
 - To reduce backgrounds from events containing neutral particles
 - Apply specific selection criteria to discriminate against reconstructed energy deposits in the EMC not associated with a charged track



Event Selection

Data & MC Simulation

- Thrust of event > 0.85
 - suppress spherically symmetric shaped hadronic background events
- Reject events with no missing momentum
 - Require missing momentum $> 0.9\% \sqrt{s}$
- Track cuts to ensure PID performance
 - Should be within the acceptance of the DIRC and EMC (i.e., $pT > 250$ MeV, which is the minimum to reach the DIRC resolution)
- Events containing conversion daughter tracks are removed
- EMC energy deposit cuts



$$T = \max_{|\hat{n}|=1} \left[\frac{\sum_i |\vec{p}_i \cdot \hat{n}|}{\sum_i |\vec{p}_i|} \right]$$



Distributions






Electron Tag

τ^+

Bkg. Type	Electron Tag MC Yield	[%]
$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \bar{\nu}_\tau$	894864	70
$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \pi^0 \bar{\nu}_\tau$	332008	26
$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- 2\pi^0 \bar{\nu}_\tau$	34050	2.7
$\tau^+ \rightarrow \pi^- \pi^+ K^+ \bar{\nu}_\tau$	3391	0.27
$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- 3\pi^0 \bar{\nu}_\tau$	1541	0.12
$\tau^+ \rightarrow \pi^+ \pi^0 \bar{\nu}_\tau$	498	0.039
$\tau^+ \rightarrow \pi^+ 2\pi^0 \bar{\nu}_\tau$	252	0.02
$\tau^+ \rightarrow 2K^+ \pi^- \bar{\nu}_\tau (\rightarrow K^+ K^- \pi^+ \bar{\nu}_\tau)$	207	0.016
$e^+ e^- \rightarrow Y(4S) \rightarrow c\bar{c}$	8031	0.63
$e^+ e^- \rightarrow Y(4S) \rightarrow u\bar{u}, s\bar{s}, d\bar{d}$	542	0.043
$e^+ e^- \rightarrow Y(4S) \rightarrow B^0 \bar{B}^0$	108	0.009
$e^+ e^- \rightarrow Y(4S) \rightarrow B^+ B^-$	100	0.008
$e^+ e^- \rightarrow \mu^+ \mu^-$	0	0
Total MC	1278339	-
Data	1265698	-

τ^-

Bkg. Type	Electron Tag MC Yield	[%]
$\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau$	900069	70
$\tau^- \rightarrow \pi^- \pi^- \pi^+ \pi^0 \nu_\tau$	334565	26
$\tau^- \rightarrow \pi^- \pi^- \pi^+ 2\pi^0 \nu_\tau$	34255	2.7
$\tau^- \rightarrow \pi^+ \pi^- K^- \nu_\tau$	3567	0.27
$\tau^- \rightarrow \pi^- \pi^- \pi^+ 3\pi^0 \nu_\tau$	1535	0.12
$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	476	0.039
$\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau$	240	0.02
$\tau^- \rightarrow 2K^- \pi^+ \bar{\nu}_\tau (\rightarrow K^- K^+ \pi^- \bar{\nu}_\tau)$	202	0.016
$e^+ e^- \rightarrow Y(4S) \rightarrow c\bar{c}$	8031	0.63
$e^+ e^- \rightarrow Y(4S) \rightarrow u\bar{u}, s\bar{s}, d\bar{d}$	495	0.043
$e^+ e^- \rightarrow Y(4S) \rightarrow B^0 \bar{B}^0$	126	0.009
$e^+ e^- \rightarrow Y(4S) \rightarrow B^+ B^-$	93	0.008
$e^+ e^- \rightarrow \mu^+ \mu^-$	0	0
Total MC	1283654	-
Data	1273291	-





Distributions










Muon Tag

τ^+

Bkg. Type	Muon Tag MC Yield	[%]
$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \bar{\nu}_\tau$	810586	71
$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \pi^0 \bar{\nu}_\tau$	278830	24
$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- 2\pi^0 \bar{\nu}_\tau$	28841	2.5
$\tau^+ \rightarrow \pi^- \pi^+ K^+ \bar{\nu}_\tau$	3101	0.27
$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- 3\pi^0 \bar{\nu}_\tau$	821	0.07
$\tau^+ \rightarrow \pi^+ \pi^0 \bar{\nu}_\tau$	207	0.017
$\tau^+ \rightarrow \pi^+ 2\pi^0 \bar{\nu}_\tau$	92	0.27
$\tau^+ \rightarrow 2K^+ \pi^- \bar{\nu}_\tau (\rightarrow K^+ K^- \pi^+ \bar{\nu}_\tau)$	146	0.013
$e^+ e^- \rightarrow Y(4S) \rightarrow c\bar{c}$	6512	0.55
$e^+ e^- \rightarrow Y(4S) \rightarrow u\bar{u}, s\bar{s}, d\bar{d}$	13898	1.19
$e^+ e^- \rightarrow Y(4S) \rightarrow B^0 \bar{B}^0$	99	0.0084
$e^+ e^- \rightarrow Y(4S) \rightarrow B^+ B^-$	89	0.0076
$e^+ e^- \rightarrow \mu^+ \mu^-$	15	0.0013
Total MC	1143237	-
Data	1137521	-

τ^-

Bkg. Type	Muon Tag MC Yield	[%]
$\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau$	817342	70
$\tau^- \rightarrow \pi^- \pi^- \pi^+ \pi^0 \nu_\tau$	281613	25
$\tau^- \rightarrow \pi^- \pi^- \pi^+ 2\pi^0 \nu_\tau$	29287	2.5
$\tau^- \rightarrow \pi^+ \pi^- K^- \nu_\tau$	3228	0.27
$\tau^- \rightarrow \pi^- \pi^- \pi^+ 3\pi^0 \nu_\tau$	795	0.07
$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	217	0.019
$\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau$	92	0.08
$\tau^- \rightarrow 2K^- \pi^+ \bar{\nu}_\tau (\rightarrow K^- K^+ \pi^- \bar{\nu}_\tau)$	152	0.013
$e^+ e^- \rightarrow Y(4S) \rightarrow c\bar{c}$	6837	0.58
$e^+ e^- \rightarrow Y(4S) \rightarrow u\bar{u}, s\bar{s}, d\bar{d}$	16602	1.42
$e^+ e^- \rightarrow Y(4S) \rightarrow B^0 \bar{B}^0$	98	0.0083
$e^+ e^- \rightarrow Y(4S) \rightarrow B^+ B^-$	103	0.0088
$e^+ e^- \rightarrow \mu^+ \mu^-$	10	0.0009
Total MC	1155920	-
Data	1150350	-




Limit Extraction

Binned Likelihood Approach



Electron Tag
Templates





Systematics

- Normalization Uncertainties
 - Luminosity, cross-section, decay branching fractions, PID efficiency ...
- Shape Uncertainties
 - Non- τ backgrounds
 - $a_1(1260)$ resonance modeling
 - $\pm 1\sigma$ variation for m_{a_1} (~ 40 MeV)
 - Other τ decay channel modeling
 - $\tau \rightarrow 3\pi^\pm + \pi^0$, $\tau \rightarrow 3\pi^\pm + 2\pi^0$

Uncertainty	Yield Change (\pm)
Luminosity	0.44%
$\sigma(ee \rightarrow \tau\tau)$	0.31%
Branching Fractions (1 prong)	e: 0.22% μ : 0.22%
Branching Fractions (3 prong)	3π : 0.57%
PID Efficiency	e: 2% μ : 1% π : 3%
Bhabha Contamination	0.2%
$q\bar{q}$ Contamination (data)	0.1%
Tracking Efficiency	negligible
Detector Modeling	negligible
Beam Energy	negligible
Tau Mass	negligible

Systematics



- Normalization Uncertainties

m_{a_1} [MeV/c ²]	$m_4 = 0$ MeV/c ²	$m_4 = 500$ MeV/c ²	$m_4 = 700$ MeV/c ²	$m_4 = 1000$ MeV/c ²	
1190	Mean m_h/m_τ	-2.09%	-0.87%	-0.15%	-0.08%
	Mean E_h/E_τ	-1.14%	-0.56%	-0.02%	0%
	RMS m_h/m_τ	-1.85%	-1.51%	-1.04%	-0.57%
	RMS E_h/E_τ	+1.96%	+2.42%	-0.21%	-0.17%
1270	Mean m_h/m_τ	+2.10%	+0.87%	+0.14%	+0.07%
	Mean E_h/E_τ	+1.15%	+0.45%	+0.04%	0%
	RMS m_h/m_τ	+1.87%	+1.57%	+1.23%	+0.61%
	RMS E_h/E_τ	-2.06%	-1.72%	+0.81%	+0.14%


- Non- τ backgrounds

- $a_1(1260)$ resonance modeling

- $\pm 1\sigma$ variation for m_{a_1} (~40 MeV)


- Other τ decay channel modeling

- $\tau \rightarrow 3\pi^\pm + \pi^0$, $\tau \rightarrow 3\pi^\pm + 2\pi^0$






Results



Mass [MeV/ c^2]	No Sys.	With Sys.
100	1.58×10^{-2}	2.31×10^{-2}
200	1.33×10^{-2}	1.95×10^{-2}
300	6.91×10^{-3}	9.67×10^{-3}
400	1.57×10^{-3}	2.14×10^{-3}
500	4.65×10^{-4}	5.85×10^{-4}
600	5.06×10^{-4}	6.22×10^{-4}
700	3.82×10^{-4}	4.85×10^{-4}
800	3.12×10^{-4}	3.85×10^{-4}
900	4.70×10^{-5}	5.38×10^{-5}
1000	8.34×10^{-5}	9.11×10^{-5}
1100	4.49×10^{-5}	4.78×10^{-5}
1200	4.70×10^{-6}	5.04×10^{-6}
1300	3.85×10^{-5}	4.09×10^{-5}



Results



Mass [MeV/ c^2]	No Sys.	With Sys.
100	1.58×10^{-2}	2.31×10^{-2}
200	1.33×10^{-2}	1.95×10^{-2}
300	6.91×10^{-3}	9.67×10^{-3}
400	1.57×10^{-3}	2.14×10^{-3}
500	4.65×10^{-4}	5.85×10^{-4}
600	5.06×10^{-4}	6.22×10^{-4}
700	3.82×10^{-4}	4.85×10^{-4}
800	3.12×10^{-4}	3.85×10^{-4}
900	4.70×10^{-5}	5.38×10^{-5}
1000	8.34×10^{-5}	9.11×10^{-5}
1100	4.49×10^{-5}	4.78×10^{-5}
1200	4.70×10^{-6}	5.04×10^{-6}
1300	3.85×10^{-5}	4.09×10^{-5}

Limits provided here are competitive with the current projections for experiments in the 5 – 10 year time frame in this mass range including those from Belle-II, FASER and NA62