

1. The probability of a leap year selected at random contain 53 Sunday is:							
	(a) 53/360	6 (b) 1/7	(c) 2/7	(d) 53/365			
2 .				A marble is drawn at			
rar	ndom. The pr	obability of dra	awing a black ba	all is :			
	(a) 3/5	(b) 2/5	(c) 0/5	(d) 1/5			
3.	3. The probability that it will rain tomorrow is 0.85. What is the						
		it will not rain					
-				(d) none of these			
4.	* *	, ,	* * *	cted from the numbers			
(1,	2, 3,,1	15) is a multiple	e of 4?				
•			(c) 2/15	(d) 1/3			
5 .	What are th	e total outcome	es when we thre	ow three coins?			
	(a) 4	(b) 5	(c) 8	(d) 7			
				cted at random from the			
	-	,35) is :					
	(a) 12/35	(b) 11/3	5 (c) 13/3	35 (d) none of these			
			of an event and				
		-	c) 0 (d) no				
	1 ,	* *		ose the correct answer			
	(a) 0.15	(b) 2/7	(c) 7/5	(d) none of these.			
	9. If three coins are tossed simultaneously, than the probability of						
		two heads, is:	•				
			(c) ½	(d) 1/8			
10	* *	• •	* *	tters of the word			
				ne letter chosen has:			
•		(b) 7/13	-	(d) none of these.			
		() .	()				
11	A dice is the	rown. Find the	probability of g	etting an even number.			
	2/3		(C) 5/6	_			
` '		、 /					
12. Two coins are thrown at the same time. Find the probability of getting both heads.							
_		/4 (C) 1/2	(D) 0				

13. Two dice are thrown simultaneously. The probability of getting a

1

sum of 9 is:



(D) 4/9

14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
• •	(B) 27/50	(C) 1/4	(D) 29/100			
15. A bag contains 5 red balls and some blue balls . If the probability of drawing a blue ball is double that of a red ball, then the number of blue balls in a bag is:							
(A) 5	(B) 10	(C) 15	(D) 20				
taken out at	16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is:						
(A) 143/150	(B) 147	/150 (C)) 1/25	(D) 1/50			
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100							
18. What is (A) 1/7		y of getting 53 (C) 2/7	_	n a leap year? 7/366			
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) $1/26$ (B) $3/26$ (C) $7/52$ (D) $1/13$							
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) $1/6$ (B) $1/2$ (C) $7/12$ (D) $5/12$							
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$							

(B) 3/10 (C) 1/9

(A) 1/10



22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:						
(A) 364/365	(B) 31/36	65 (C) 1/ 3	365 ((D) 1/133225		
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$						
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7						
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$						
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$			
27. Runs scored by batsman in 5 one day matches are 50, 70, 82,						
93, and 20. The a) 25.79				5.69		
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
a) 13, 15				d) 13, 16		
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is						
a) $^{1}/_{2}$	b) $^{1}/_{3}$	c)	±	d) $^{1}/_{6}$		
30. X is a varia		and 3. The va c) 27		·		
	•	•	·	105		
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?						

a) 3	b) 4	c) 5	d) 7				
32.Out of the following values, which one is not possible in probability? a) $P(x) = 1$ b) $\sum x P(x) = 3$							
C) P(X) = 0.3	c) $P(x) = 0.5$ d) $P(x) = -0.5$						
33.If E(x) = a) 2	2 and E(z) = b) 6	4, then E(z - 2 c) 0	•	ufficient data			
34. The covariance of two independent random variable is							
a) 1	b) 0	c) - 1	d) Und	defined			
35.If $\Sigma P(x) = k^2 - 8$ then, the value of k is? a) 0 b) 1 c) 3 d) Insufficient data							
36.If P(x) = a) 1	0.5 and x = 4 b) 0.5	c) 4	d) 2				
37.In a discrete probability distribution, the sum of all probabilities is always?							
a) 0	b) Infinite	c) 1	d) Und	efined			
38.If the probability of hitting the target is 0.4, find mean and variance.							
a) 0.4, 0.24	b) 0.6	, 0.24	c) 0.4, 0.16	d) 0.6, 0.16			
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6							

40. Find the mean of tossing 8 coins.
a) 2
b) 4
c) 8
d) 1
41. What is the mean and variance for standard normal distribution?



d) 3, 2/3

a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0 c) Mean is 0 and variance is ∞ d) Mean is ∞ and variance is 0							
42.\	/ariance	of a rand	lom varia	ıble X is g	iven by		•
					- (E(X))2		d) (E(X))2
43.Mean of a random variable X is given by a) E(X) b) E(X2) c) E(X2) - (E(X))2						d) (E(X))2	
44.Mean of a constant 'a' is a) 0							
45.Variance of a constant 'a' is . a) 0							
46.Find the mean and variance of X?							
г.			T				
	X	0	1	2	3	4	

3/9

2/9

c) 2, 2/3

d) 3.5

47. Find the expectation of a random variable X?

2/9

b) 3, 4/3

1/9

f(x)

a) 2, 4/3

x 0 1 2 3 f(x) 1/6 2/6 2/6 1/6 a) 0.5 b) 1.5 c) 2.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



a) np

b) npq

c) np2q

d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

a)
$$P(X = x) = nCx px qx$$

b)
$$P(X = x) = nCx px q(n-x)$$

c)
$$P(X = x) = xCn qx p(n-x) d$$

$$P(x = x) = xCn pn qx$$

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np}

- b) \sqrt{pq} c) (np)2 d) \sqrt{npq}