




Contents lists available at ScienceDirect

Clinical Nutrition ESPEN

journal homepage: <http://www.clinicalnutritionespen.com>

Corrigendum to “Vinegar (acetic acid) intake on glucose metabolism: A narrative review” [Clin Nutr ESPEN 32 (2019) 1–7]

Heitor O. Santos ^{a,*}, Wilson M.A.M. de Moraes ^b, Guilherme A.R. da Silva ^c,
Jonato Prestes ^b, Brad J. Schoenfeld ^d

^a School of Medicine, Federal University of Uberlandia (UFU), Uberlandia, Minas Gerais, Brazil

^b Graduation Program on Physical Education, Catholic University of Brasilia (UCB), Brasília, Federal District, Brazil

^c Federal University of the State of Rio de Janeiro (UNIRIO), Brazil

^d Department of Health Sciences, CUNY Lehman College, Bronx, NY, USA

The authors regret some mismatches in the use of spoon measures.

Given that 1 teaspoon of vinegar equals ~5 mL, while 1 tablespoon equals ~15 mL, a *corrigendum* is required in four places in this paper, as described below.

- 1 In highlights, consider “10–30 mL/d (~2–6 teaspoons) of vinegar is the more studied dose.” instead of “10–30 mL/d (~2–6 tablespoons) of vinegar is the more studied dose.”
- 2 In the abstract, consider “When evaluating the current body of literature, daily vinegar intake in amounts of ~10–30 mL (~2–6 teaspoons) appear to improve the glycemic response to carbohydrate-rich meals; ” instead of “When evaluating the current body of literature, daily vinegar intake in amounts of ~10–30 mL (~2–6 tablespoons) appear to improve the glycemic response to carbohydrate-rich meals; ”.
- 3 In topic 6, consider “Converting to homemade measures, 20 g of vinegar corresponds to approximately 4 teaspoons [26].” instead of “Converting to homemade measures, 20 g of vinegar corresponds to approximately 4 tablespoons [26].”
- 4 In the conclusion section, consider “Taken together, consumption of 10–30 mL (~2–6 teaspoons) of vinegar as a dressing or diluted in beverages appears to be adequate to acutely alter glycemic profile.” instead of “Taken together, consumption of 10–30 mL (~2–6 tablespoons) of vinegar as a dressing or diluted in beverages appears to be adequate to acutely alter glycemic profile.”

The authors would like to apologize for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.clnesp.2019.05.008>.

* Corresponding author. School of Medicine, Federal University of Uberlandia, Av. Para, no. 1720 Bloco 2U Campus Umuarama, 38400-902, Uberlandia, Minas Gerais, Brazil.
E-mail address: heitoroliveirasantos@gmail.com (H.O. Santos).

<https://doi.org/10.1016/j.clnesp.2024.09.013>

2405-4577/© 2024 European Society for Clinical Nutrition and Metabolism. Published by Elsevier Ltd. All rights are reserved, including those for text and data mining, AI training, and similar technologies.