

**Concordia University**  
Department of Mathematics and Statistics

<b>Course</b>	<b>Number</b>	<b>Section</b>	
MATH	204/3	CA	
<b>Examination</b>	<b>Date</b>	<b>Time</b>	<b>Pages</b>
Mid-term	July 2020	1 hour 15 mins	2
<b>Instructor</b>	<b>Course Examiner</b>		<b>Marks</b>
U. Tiwari	A. Kokotov		20
<b>Special Instructions: Online proctored Closed Book Exam</b>			

**Q 1. [Marks = 5 ]** Use Gauss-Jordan elimination method to solve the system of equations:

$$\begin{cases} x + y + 4z = 2 \\ 2x + 5y + 20z = 10 \\ -x + 2y + 8z = 4 \end{cases} \Rightarrow \begin{cases} x = 0 \\ y = 2 - 4t \\ z = t \end{cases}, -\infty < t < \infty$$

*t is a parameter*

**Q 2. [Marks = 3.5+1.5=5]** For matrix

$$A = \begin{bmatrix} 4 & 2 & 1 \\ -3 & -3 & -1 \\ 3 & 2 & 1 \end{bmatrix} \quad \text{Find } A^{-1} = \begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & -1 \\ -3 & 2 & 6 \end{bmatrix}$$

(a)  $A^{-1}$  if exists

(b) Use  $A^{-1}$  to solve the system

$$\begin{cases} 4x + 2y + z = -3 \\ -3x - 3y - z = -9 \\ 3x + 2y + z = 2 \end{cases} \quad \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -5 \\ 7 \\ 3 \end{pmatrix}$$

Q 3. [Marks = 5] Evaluate the determinant of the matrix

$$A = \begin{bmatrix} 3 & 0 & 1 & 5 \\ 1 & -1 & 2 & 1 \\ 0 & 1 & 0 & -2 \\ 4 & 0 & 3 & 4 \end{bmatrix}$$

$$\det A = 0$$

Q 4. [Marks = 1+2+2] For square matrices  $A$  and  $B$ , prove or disprove each of the following:

(a)  $\det(A^T) = \det A$

See textbook.

(b)  $\det(A + B) = \det A + \det B$

(c)  $(A^{-1})^{-1} = A$

### COPYRIGHT

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.



## Solution

#1 Augmented matrix of the system

$$\begin{bmatrix} 1 & 1 & 4 & 2 \\ 2 & 5 & 20 & 10 \\ -1 & 2 & 8 & 4 \end{bmatrix} \xrightarrow[R_3+R_1]{R_2-2R_1} \begin{bmatrix} 1 & 1 & 4 & 2 \\ 0 & 3 & 12 & 6 \\ 0 & 3 & 12 & 6 \end{bmatrix} \xrightarrow{\frac{1}{3}R_2}$$

$$\begin{bmatrix} 1 & 1 & 4 & 2 \\ 0 & 1 & 4 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1-R_2} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 4 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \Rightarrow \begin{aligned} x &= 0 \\ y &= 2-4t \\ z &= t \end{aligned}$$

Where  $-\infty < t < \infty$ .

$$\#2 \text{ (a)} [A:I] = \begin{bmatrix} 4 & 2 & 1 & 1 & 0 & 0 \\ -3 & -3 & -1 & 0 & 1 & 0 \\ 3 & 2 & 1 & 0 & 0 & 1 \end{bmatrix} \xrightarrow[R_3+R_2]{R_1+R_2}$$

$$\begin{bmatrix} 1 & -1 & 0 & 1 & 1 & 0 \\ -3 & -3 & -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 & 1 & 1 \end{bmatrix} \xrightarrow{R_2+3R_1} \begin{bmatrix} 1 & -1 & 0 & 1 & 1 & 0 \\ 0 & -6 & -1 & 3 & 4 & 0 \\ 0 & -1 & 0 & 0 & 1 & 1 \end{bmatrix} \xrightarrow[R_2 \leftrightarrow R_3]{-R_3}$$

$$\begin{bmatrix} 1 & -1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & -1 & -1 \\ 0 & -6 & -1 & 3 & 4 & 0 \end{bmatrix} \xrightarrow[R_3+6R_2]{R_1+R_2} \begin{bmatrix} 1 & 0 & 0 & 1 & 0 & -1 \\ 0 & 1 & 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & 3 & -2 & -6 \end{bmatrix} \xrightarrow{-R_3}$$

$$\begin{bmatrix} 1 & 0 & 0 & 1 & 0 & -1 \\ 0 & 1 & 0 & 0 & -1 & -1 \\ 0 & 0 & 1 & -3 & 2 & 6 \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & -1 \\ -3 & 2 & 6 \end{bmatrix}$$

(b)

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & -1 \\ -3 & 2 & 6 \end{bmatrix} \begin{bmatrix} -3 \\ -9 \\ 2 \end{bmatrix} = \begin{bmatrix} -5 \\ 7 \\ 3 \end{bmatrix} \checkmark$$

-9-18+11



$$\#3 \quad \left| \begin{array}{cccc} 3 & 0 & 1 & 5 \\ 1 & -1 & 2 & 1 \\ 0 & 1 & 0 & -2 \\ 4 & 0 & 3 & 4 \end{array} \right| \xrightarrow{R_1 \leftrightarrow R_2} \left| \begin{array}{cccc} 1 & -1 & 2 & 1 \\ 3 & 0 & 1 & 5 \\ 0 & 1 & 0 & -2 \\ 4 & 0 & 3 & 4 \end{array} \right| \xrightarrow{\begin{array}{l} R_2 - 3R_1 \\ R_4 - 4R_1 \end{array}}$$

$$= - \left| \begin{array}{cccc} 1 & -1 & 2 & 1 \\ 0 & 3 & -5 & 2 \\ 0 & 1 & 0 & -2 \\ 0 & 4 & -5 & 0 \end{array} \right| \xrightarrow{R_2 \leftrightarrow R_3} + \left| \begin{array}{cccc} 1 & -1 & 2 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 3 & -5 & 2 \\ 0 & 4 & -5 & 0 \end{array} \right|$$

$$\xrightarrow{\begin{array}{l} R_3 - 3R_2 \\ R_4 - 4R_2 \end{array}} \left| \begin{array}{cccc} 1 & -1 & 2 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & -5 & 8 \\ 0 & 0 & -5 & 8 \end{array} \right| \xrightarrow{R_4 - R_3} \left| \begin{array}{cccc} 1 & -1 & 2 & 1 \\ 0 & 1 & 0 & -2 \\ 0 & 0 & -5 & 8 \\ 0 & 0 & 0 & 0 \end{array} \right| = 0$$

$$\#4 \quad (a) \det(A) = \det(A^T):$$

True, since transposing a matrix changes its rows to columns and its columns to rows, the cofactor expansion of  $A$  along any row is the same as cofactor expansion of  $A^T$  along the corresponding column.

$$(b) \det(A+B) \neq \det A + \det B \text{ in general.}$$

$$\text{Example } A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, B = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$\det A = 1 \text{ and } \det B = 0, \det(A+B) = 0$$

(c) See Text book