

CONCORDIA UNIVERSITY
Department of Mathematics & Statistics

Course	Number	Section(s)
Mathematics	204	All except EC
Examination	Date	Pages
Final	April 2018	2
Instructors	Course Examiner	
M. Baker, Y. Bibilo, Q. Zhang	E. Cohen	

Special Instructions:

- ▷ Only approved calculators are allowed.
- ▷ Justify all your answers.
- ▷ All questions have equal value.

MARKS

1. Use the Gauss-Jordan method to find all the solutions of the system:

$$\begin{aligned} -3x_1 + 2x_2 - x_3 + 6x_4 &= -7 \\ 7x_1 - 3x_2 + 2x_3 - 11x_4 &= 14 \\ x_1 &\quad \quad \quad -x_4 = 1 \end{aligned}$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 1+s \\ -1-2s \\ 2-s \\ s \end{pmatrix}$$

2. Find the inverse of the matrix $A = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}$, if it exists.

$$A^{-1} = A^T = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

3. Using Cramer's Rule, find the value of x_2 in the system:

$$\begin{aligned} 3x_2 &\quad \quad \quad - 3x_4 = 1 \\ 2x_1 - x_2 + 3x_3 - 3x_4 &= -2 \\ -2x_1 + 3x_2 + 2x_3 + 2x_4 &= 0 \\ 2x_1 &\quad \quad \quad + 2x_3 + x_4 = -1 \end{aligned}$$

$$x_2 = \frac{-56}{-156} = \frac{14}{39}$$

4. Find the determinant of $A = \begin{pmatrix} 1 & 5 & 3 & -4 \\ -1 & 2 & -3 & 4 \\ 0 & 4 & 2 & -3 \\ 2 & 4 & 2 & -2 \end{pmatrix}$

$$\det A = 0$$

5. a) Let $u = (-3, 1, 2)$, $v = (1, 1, 1)$. Find the orthogonal projection of v on u . $\text{pr}_{\vec{u}} \vec{v} = (0, 0, 0) = \vec{0}$
- b) Let $u_1 = (1, 0, 0)$, $u_2 = (1, 1, 0)$, $u_3 = (1, 1, 1)$. Find c_1, c_2, c_3 such that $c_1 u_1 + c_2 u_2 + c_3 u_3 = (1, 0, 1)$. $(c_1, c_2, c_3) = (1, -1, 1)$
6. a) Find the area of a triangle with vertices $(1, 2, 1)$, $(0, 5, 2)$, $(6, 7, 3)$.
Find a vector orthogonal to the plane of the triangle. $S_{\Delta} = 15/\sqrt{2}$, orth. vect. $(1, 7, -20)$
- b) Find the distance between the point $(2, 4)$ and the line $3x = 2y - 6$. $\text{dist} = 4/\sqrt{13}$
7. a) Let $u = (4, 0, -2)$, $v = (1, 3, 7)$, $w = (3, 3, 5)$.
Are the vectors linearly dependent or independent? independent
- b) Find the parametric equations for the line in \mathbb{R}^3 passing through the point $(1, 2, -7)$ and perpendicular to the plane $2x - 3y + 5z = 4$. $\begin{cases} x = 2t + 1 \\ y = -3t + 2 \\ z = 5t - 7 \end{cases}$
8. Let $A = \begin{bmatrix} 1 & 0 & 2 & 0 & 1 & 4 \\ 0 & 1 & 3 & 0 & 2 & 7 \\ 0 & 0 & 0 & 1 & 8 & 9 \end{bmatrix}$ and $X = \begin{bmatrix} x \\ y \\ z \\ t \\ u \\ v \end{bmatrix}$. Basis: $\begin{pmatrix} -2, -3, 1, 0, 0, 0 \\ -1, -2, 0, -8, 1, 0 \\ -4, -7, 0, -9, 0, 1 \end{pmatrix}$
- Find a basis for the solution space of the homogeneous system $AX = 0$.
9. Find the standard matrices for following operators on \mathbb{R}^2 : $A_{\theta} = \begin{bmatrix} 1/2 & -\sqrt{3}/2 \\ \sqrt{3}/2 & 1/2 \end{bmatrix}$
- a) a rotation counterclockwise of 60° .
- b) a reflection about line $y = -x$. $A_B = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$
10. Let $A = \begin{bmatrix} 1 & 1 & 1 \\ -2 & -2 & -1 \\ 0 & 0 & -1 \end{bmatrix}$. Find a matrix P such that $P^{-1}AP = D$ a diagonal matrix. Eigenvalues $0, -1$ (mult. 2)
 $P = \begin{bmatrix} -1 & -1 & -1 \\ 1 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ for $D = \begin{bmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.