

CONCORDIA UNIVERSITY
Department of Mathematics & Statistics

Course	Number	Sections
Mathematics	205	All
Examination	Date	Pages
Final	December 2015	2
Instructors:	J. Brody, A. Daniels, T. Hughes U. Mgbemena	Course Examiner A. Atoyan and H. Proppe
Special	Only approved calculators are allowed.	
Instructions:	Show all your work for full marks.	

MARKS

[10] **1. a.** Sketch the graph of the function

$$f(x) = \begin{cases} 1 + \sqrt{4 - x^2} & -2 \leq x \leq 2 \\ 3 - x & 2 < x \end{cases}$$

on the interval $[-2, 5]$, and find the definite integral $\int_{-2}^5 f(x) dx$ in terms of area (do **not** antidifferentiate).

b. Use the Fundamental Theorem of Calculus to calculate the derivative of

$$F(x) = \int_{-x^2}^x e^{1-t^2} dt, \text{ and determine whether } F \text{ is increasing or decreasing at } x = 1.$$

[10] **2.** Find the following indefinite integrals:

$$\text{(a)} \int \frac{\cos^3(x)}{\sin^2(x)} dx \qquad \text{(b)} \int (e^x + \ln x) dx$$

[6] **3.** Find $F(x)$ such that $F'(x) = \frac{x^2 + 4}{x^2 - 4}$ and $F(-1) = 0$.

[18] **4.** Evaluate the following definite integrals (give the **exact** answers, do not approximate):

$$\text{(a)} \int_0^{\pi/2} \frac{\cos(x)}{4 + \sin^2(x)} dx \qquad \text{(b)} \int_0^{\pi/4} \sec^4(x) dx \qquad \text{(c)} \int_0^3 x^2 \sqrt{1+x} dx$$

[8] **5.** Evaluate the given improper integral or show that it diverges:

$$\text{(a)} \int_e^{\infty} \frac{dx}{x \ln^3(x)} \qquad \text{(b)} \int_{-1}^1 \frac{x}{x^2 - 1} dx$$

- [16] **6.** **a.** Sketch the curves $y = 6 - x^2$ and $y = 2 - 3x$, and find the area enclosed.
- b.** Find the volume of a solid obtained by rotating the region bounded by the curve $y = \sin(x)$ and the x -axis on the interval $0 \leq x \leq \pi$ about the axis $y = -1$.
- c.** Find the exact average value of $f(x) = \frac{x}{\sqrt{16 + x^2}}$ on the interval $[0, 3]$.
- [6] **7.** Find the limit of the sequence $\{a_n\}$ at $n \rightarrow \infty$ or prove that it does not exist:
- (a) $a_n = \frac{(3^n + 1)^2}{6^n}$ (b) $a_n = \ln(1 + 2n^2) - \ln(30 + 2n^2)$
- [12] **8.** Determine whether the series is divergent or convergent, and if convergent, whether absolutely or conditionally :
- (a) $\sum_{n=1}^{\infty} \frac{(-1)^n \sqrt{1 + n^3}}{n^2}$ (b) $\sum_{n=0}^{\infty} \frac{(-3)^n}{5 + e^n}$ (c) $\sum_{n=2}^{\infty} \frac{1}{n \ln^2(n)}$
- [6] **9.** Find **(a)** the radius of convergence, and **(b)** the interval convergence of the series $\sum_{n=0}^{\infty} \frac{(x + 1)^n}{(n + 1) 2^n}$.
- [8] **10.** **(a)** Derive the Maclaurin series of $f(x) = x^2 \ln(1 + 2x)$ (HINT: start with the series for $\ln(1 + z)$ where $z = 2x$).
- (b)** Use integrability of power series to find the sum $S(x) = \sum_{n=1}^{\infty} n x^{2n-1}$ in the form of an elementary function within the radius of convergence of $S(x)$. (HINT: first find the sum for the antiderivative of $S(x)$, then differentiate).
- [5] **Bonus question.** If f is continuous, prove that

$$\int_0^{\pi/2} f(\cos x) dx = \int_0^{\pi/2} f(\sin x) dx$$

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.