

CONCORDIA UNIVERSITY

Department of Mathematics & Statistics

Course	Number	Section(s)
Mathematics	209	All
Examination	Date	Pages
Final	April 22 2017	3
Instructors	Course Examiner	
ALL	R. Raphael	

Special Instructions

- ▷ Ruled booklets to be used.
- ▷ Approved calculators allowed.

MARKS

[6] 1. (a) Find the following limits

(i) $\lim_{x \rightarrow 1} \frac{2x^5 + 7x - 1}{x^2 + 5x + 3}$

(ii) $\lim_{x \rightarrow 2} \frac{x^2 + x - 6}{2x^2 - 3x - 2}$

(b) Prove or disprove by giving an example: there exists a function f from the real numbers to the real numbers that is discontinuous at exactly three points.

[4] 2. Find the derivative $f'(x)$ of the functions $f(x)$: (Do not simplify)

(a) $f(x) = 4x^5 - 9x^2 + x - 22$

(b) $f(x) = \frac{x^{-7}}{8} + \frac{1}{\sqrt[2]{x}}$

[10] 3. Find $\frac{dy}{dx}$ (do not simplify):

(a) $y = \frac{7 - x^3}{e^{3x}}$

(b) $y = \ln(3x^4 + 7)$

(c) $y = (4x - 5)^3(3x^2 + 4)$

(d) $y = (5 + x^3 \ln x)^3$

[8] 4. Let $f(x) = 4x^4 - x^2 - 7$

(a) Find the slope of the tangent line to the curve when $x = 1$

(b) Find the equation of the tangent line to the curve when $x = 1$

[13] 5. Let $f(x) = x^4 - 2x^3$

Find

(a) the critical and inflection points of $f(x)$

(b) the intervals where $f(x)$ is increasing and where it is decreasing

(c) the intervals on which $f(x)$ is concave up and on which it is concave down

(d) use the above to sketch the graph

[9] 6. If the cost of a seminar is \$400 per person 1000 people attend. For every \$5 dollar reduction in cost 20 more people will attend the seminar. How much should be charged for the seminar to maximize revenue?

[6] 7. Find the absolute extrema of the function $f(x) = x^3 - 12x$ on the interval $[-5, 5]$.

[4] 8. A country has Lorenz curve $f(x) = x$. Find its Gini index. What can you conclude from the Gini index?

[10] 9. Find the equation(s) of the tangent line(s) to the graph of $y - xy^2 + x^2 + 1 = 0$ at the point(s) with $x = 1$.

[10] 10. Compute these antiderivatives:

(a) $\int (5x^7 - 4x^3 - 9) dx$

(b) $\int \frac{e^{-2x}}{4 + e^{-2x}} dx$

(c) $\int \frac{x^2}{\sqrt{x-5}} dx$

[10] 11. Evaluate the integrals:

(a) $\int_0^1 (x^4 - 5) dx$

(b) $\int_6^{10} \frac{2}{x-4} dx$

(c) $\int_4^7 \sqrt{x-2} dx$

[10] 12. Find the area bounded by the graphs of $f(x) = 5 - x^2$ and $g(x) = 2 - 2x$.

copyright

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.

Math 209 April 2017 FINAL EXAM

1a) (i) $\lim_{x \rightarrow 1} \frac{2x^5 + 7x - 1}{x^2 + 5x + 3} = \frac{2(1)^5 + 7(1) - 1}{(1)^2 + 5(1) + 3} = \frac{8}{9}$

(ii) $\lim_{x \rightarrow 2} \frac{x^2 + x - 6}{2x^2 - 3x - 2} = \frac{2^2 + 2 - 6}{2(2)^2 - 3(2) - 2} = \frac{0}{0}$

$\lim_{x \rightarrow 2} \frac{(x+3)(x-2)}{(2x+1)(x-2)} = \frac{2+3}{2(2)+1} = \frac{5}{5} = 1$

b) Need function that is undefined at 3 values of x

$f(x) = \frac{1}{(x-1)(x-2)(x-3)}$ is discont. (DNE) when $x=1$
 $x=2$
 $x=3$

2. a) $f'(x) = 20x^4 - 18x + 1$

b) $f(x) = \frac{1}{8}x^{-7} + x^{-\frac{1}{2}}$

$\Rightarrow f'(x) = \frac{1}{8}(-7)x^{-8} + (-\frac{1}{2})x^{-\frac{3}{2}}$

3. a) $y' = \frac{e^{3x} \frac{d}{dx}(7-x^3) - (7-x^3) \frac{d}{dx} e^{3x}}{(e^{3x})^2}$

$y' = \frac{e^{3x}(-3x^2) - (7-x^3)e^{3x}(3)}{(e^{3x})^2}$

b) $y' = \frac{1}{3x^4+7} \frac{d}{dx}(3x^4+7)$

$y' = \frac{1}{3x^4+7} (12x^3)$

c) $y' = (4x-5)^3 \frac{d}{dx}(3x^2+4) + (3x^2+4) \frac{d}{dx}(4x-5)^3$
 $= (4x-5)^3(6x) + (3x^2+4) 3(4x-5)^2 \frac{d}{dx}(4x-5)$
 $= (4x-5)^3(6x) + (3x^2+4) 3(4x-5)^2(4)$

d) $y' = 3(5+x^3 \ln x)^2 \frac{d}{dx}(5+x^3 \ln x)$


$y' = 3(5+x^3 \ln x)^2 (x^3 \frac{d}{dx} \ln x + \ln x \frac{d}{dx} x^3)$

$y' = 3(5+x^3 \ln x)^2 (x^3 \cdot \frac{1}{x} + \ln x \cdot 3x^2)$

4. a) $f'(x) = 4(4)x^3 - 2x$

$f'(1) = 16(1)^3 - 2(1) = 14$

b)



Step 1 when $x=1$, $f(1) = 4(1)^4 - (1)^2 = 7 =$
 \Rightarrow pt is $(1, -4)$

Step 2 Let $P(x, y)$ be any pt on line

$m_{AP} = \frac{y - (-4)}{x - 1}$

$14 = \frac{y+4}{x-1} \Rightarrow y = 14x - 18$

5. Step 1 $f'(x) = 4x^3 - 6x^2$

$f'(x) = 0$	$f'(x) = \frac{1}{6}$
$2x^2(x-3) = 0$	No x
$2x^2 = 0 \quad \quad 2x-3=0$	
$x=0 \quad \quad x=3/2$	

$f''(x) = 12x^2 - 12x$

$f''(x) = 0$	$f''(x) = \frac{1}{6}$
$12x(x-1) = 0$	No x
$12x = 0 \quad \quad x-1=0$	
$x=0 \quad \quad x=1$	

Step 2


x	$-\infty < x < 0$	$x=0$	$0 < x < 1$	$x=1$	$1 < x < 3/2$	$3/2$	$3/2 < x < \infty$
$f'(x)$	$f'(-1) = -$	0	$f'(1/2) = -$	-	-	0	$f'(2) = +$
$f''(x)$	$f''(-1) = +$	0	$f''(1/2) = -$	0	+	+	$f''(2) = +$
$f(x)$	Dec, CU.	Inf. pt (0,0)	Dec CD.	Dec Inf. pt (1,-1)	Dec cu	CU Loc min (3/2, 27/16)	CU Inf CU

Step 3

Signs: use

$f(x) = 2x^3(x-3)$

$f''(x) = 4x(3x-4)$
 $= 12x(x-1)$



6.

Step 1

$R = x * p$
Rev. (No. persons) (price/person)

Step 2

too many variable

$R = (1000 + 20r)(400 - 5r)$

Step 3

① $R' = (1000 + 20r)(-5)$
 $+ (400 - 5r)(20)$

$R' = -5000 - 100r + 8000 - 100r$

$R' = -200r + 3000$

② $R' = 0$

$R' = \frac{1}{6}$

$-200r + 3000 = 0$

$r = \frac{-3000}{-200}$

$r = 15$

No r

No. person	Cost/person
1000	400\$
1000 + 1(20)	400 - 1(5)
1000 + 2(20)	400 - 2(5)
1000 + 3(20)	400 - 3(5)
1000 + r(20)	400 - r(5)

where r is No. of \$ dec.

③ test

$R'' = - \Rightarrow R$ is cu for all r

$\Rightarrow r = 15$ produces local MAX

Step 4

Cost/person $400 - 5(15) = 325\$$

Note: No. person $1000 + 20(15) = 1300$ persons

7. Step 1 $f'(x) = 3x^2 - 12$

$f'(x) = 0$

$f'(x) = 6$

Step 2 $3x^2 - 12 = 0$

$3x^2 = 12$

$x^2 = 4$

$x = \pm 2$

No x

Step 3


test

$f(-2) = -16$

$f(2) = 16$

$f(-5) = -65$ MIN

$f(5) = 65$ MAX



8 Gini Index $= 2 \int_0^1 [x - f(x)] dx$
 $= 2 \int_0^1 [x - x] dx$
 $= 0$

This is perfect income distribution (Equality)

ie 10% of population 10% inc

20% ..

20% ..

;

x% ..

x% ..

9. $\frac{d}{dx} y - \frac{d}{dx} x y^2 + \frac{d}{dx} x^2 + \frac{d}{dx} 1 = \frac{d}{dx} 0$

$\frac{dy}{dx} - (x \frac{d}{dx} y^2 + y^2 \frac{d}{dx} x) + 2x \frac{d}{dx} x + 0 = 0$

$\frac{dy}{dx} - x(2y) \frac{dy}{dx} + y^2 + 2x = 0$

$\frac{dy}{dx} [1 - 2xy] = -y^2 - 2x$

$\frac{dy}{dx} = \frac{-y^2 - 2x}{1 - 2xy}$

$\frac{dy}{dx} \bigg|_{\substack{x=1 \\ y=-1}} = \frac{(-1)^2 - 2(1)}{1 - 2(1)(-1)}$
 $= -\frac{1}{3}$

$\frac{dy}{dx} \bigg|_{\substack{x=1 \\ y=2}} = \frac{(2)^2 - 2(1)}{1 - 2(1)(2)}$
 $(1, 2) = -\frac{2}{3}$

$B(1, 2)$
 $Q(x, y)$

$m_{BQ} = \frac{y-2}{x-1}$

$-\frac{2}{3} = \frac{y-2}{x-1}$

$3(y-2) = -2(x-1)$

$y = -\frac{2}{3}x + \frac{8}{3}$

$m_{AP} = \frac{y+1}{x-1}$

$-\frac{1}{3} = \frac{y+1}{x-1}$

$3(y+1) = -(x-1) \Rightarrow y = -\frac{1}{3}x - \frac{2}{3}$

when $x=1$

$\Rightarrow y - 1y^2 + 1 + 1 = 0$

$-y^2 + y + 2 = 0$

$y^2 - y - 2 = 0$

$(y+1)(y-2) = 0$

$y+1=0$

$y=-1$

$y-2=0$

$y=2$

$P(1, -1)$

$P(1, 2)$

$$10. \quad a) \quad 5 \int x^7 dx - 4 \int x^3 dx - 9 \int dx$$

$$\frac{5x^8}{8} - \frac{4x^4}{4} - 9x + C$$

$$b) \quad \int \frac{e^{-2x}}{4+e^{-2x}} dx$$

$$= \int \frac{-\frac{1}{2} du}{u}$$

$$= -\frac{1}{2} \int \frac{1}{u} du$$

$$= -\frac{1}{2} \ln|u| + C$$

$$= -\frac{1}{2} \ln|4+e^{-2x}| + C$$

$$c) \quad \int \frac{x^2}{\sqrt{x-5}} dx$$

$$\int \frac{(u+5)^2}{\sqrt{u}} du$$

$$\int \frac{u^2 + 10u + 25}{\sqrt{u}} du$$

$$\int (u^{3/2} + 10u^{1/2} + 25u^{-1/2}) du$$

$$\frac{2}{5} u^{5/2} + 10 \left(\frac{2}{3} \right) u^{3/2} + 25 \left(2 \right) u^{1/2} + C$$

$$\frac{2}{5} (x-5)^{5/2} + \frac{20}{3} (x-5)^{3/2} + 50 (x-5)^{1/2} + C$$

Answer $\int \frac{1}{x} dx$ #2

Let $u = 4 + e^{-2x}$

$$\frac{du}{dx} = e^{-2x} (-2)$$

$$du = e^{-2x} (-2) dx$$

$$-\frac{1}{2} du = e^{-2x} dx$$

Get on var. in Den
then we can use Div tech.

Let $u = x-5$

$$du = dx$$

Not $u+5=x$

$$(u+5)^2 = x^2$$

$$11. \quad a) \quad \int x^4 dx - 5 \int dx$$

$$\left[\frac{x^5}{5} - 5x \right] \Big|_0^1 = \left[\frac{1^5}{5} - 5(1) \right] - \left[\frac{0^5}{5} - 5(0) \right] = -\frac{24}{5}$$

$$b) \quad \int \frac{2}{x-4} dx$$

Answer $\int \frac{1}{x} dx$ #2

Let $u = x-4$
 $du = dx$

$$= 2 \int \frac{du}{u}$$

$$= 2 \ln|u|$$

$$= \left[2 \ln|x-4| \right]_6^{10} =$$

$$= \left[2 \ln|10-4| \right] - \left[2 \ln|6-4| \right]$$

$$= 2 \ln 6 - 2 \ln 2$$

$$= 2 (\ln 6 - \ln 2)$$

$$= 2 \ln \frac{6}{2}$$

$$11.c) \int_4^7 \sqrt{x-2} \, dx$$

Get one u under Root

$$\text{let } u = x-2$$

$$du = dx$$

$$\text{Note } u+2 = x$$

$$\int \sqrt{u} \, du$$

$$\int u^{\frac{1}{2}} \, du$$

$$\frac{2}{3} u^{\frac{3}{2}}$$

$$\frac{2}{3} (x-2)^{\frac{3}{2}} \Big|_4^7 = \frac{2}{3} (7-2)^{\frac{3}{2}} - \frac{2}{3} (4-2)^{\frac{3}{2}}$$

12. Step 1 Int. pts

$$2-2x = 5-x^2$$

$$x^2 - 2x - 3 = 0$$

$$(x+1)(x-3) = 0$$

$$x = -1$$

$$x = 3$$

$$y = 2-2x$$

$$y = 2-2(-1)$$


$$y = 4$$

$$y = 2-2x$$

$$y = 2-2(3)$$

$$y = -4$$

Step 2 Rough Sketch



Extract a
parabola
if $x=0$
 $y=5$

$$\text{Area} = \int_{-1}^3 |(5-x^2) - (2-2x)| \, dx$$

$$= \int_{-1}^3 |-x^2 + 2x + 3| \, dx$$

$$= \left| -\frac{x^3}{3} + \frac{2x^2}{2} + 3x \right| \Big|_{-1}^3$$

$$= \left| \left(-\frac{3^3}{3} + 3^2 + 3(3) \right) - \left(-\frac{[-1]^3}{3} + [-1]^2 + 3[-1] \right) \right|$$

$$= \left| -9 + 9 + 9 - \frac{1}{3} - 1 + 3 \right|$$

$$= \left| 1 - \frac{1}{3} \right|$$

$$= \frac{33-1}{3}$$

$$= \frac{32}{3}$$