

Analysis for Revision

2026-07-23

HSV1 HSV2

Data is used as is because not skewed.

Question: Is there a difference between asymptomatic and severe at any dose?

```
## boundary (singular) fit: see help('isSingular')  
## boundary (singular) fit: see help('isSingular')
```

Table 1: Linear mixed models.

	3.7U	11.1U	33.3U	100U
HSV1	-18.32 (CI=-45.10, 8.45, p=0.180)	-14.59 (CI=-45.45, 16.26, p=0.354)	-11.79 (CI=-32.02, 8.45, p=0.254)	-0.56 (CI=-4.58, 3.46, p=0.785)
HSV2	-0.60 (CI=-16.02,14.82, p=0.939)	-5.93 (CI=-20.45, 8.60, p=0.424)W	3.78 (CI=-11.36,18.92, p=0.625)W	-0.52 (CI=-6.91, 5.88, p=0.874)

Table 2: Unpaired t test pvals.

	3.7U	11.1U	33.3U	100U
HSV1	0.240	0.404	0.330	0.794
HSV2	0.942	0.447	0.619	0.879

CCL5

Data is log transformed as it is skewed.

```
## boundary (singular) fit: see help('isSingular')  
## boundary (singular) fit: see help('isSingular')  
## boundary (singular) fit: see help('isSingular')  
## boundary (singular) fit: see help('isSingular')
```

Table 3: Linear mixed models.

	0U	0.1U	1U	10U	100U
CCL5	0.00 (CI=-0.33,0.33, p=1.000)W	0.15 (CI=-0.18,0.47, p=0.369)W	0.03 (CI=-0.33,0.38, p=0.881)W	-0.37 (CI=-0.69,-0.05, p=0.023)*W	-0.21 (CI=-0.95,0.54, p=0.587)

Table 4: Unpaired t test pvals.

	0U	0.1U	1U	10U	100U
CCL5	0.090	0.151	0.836	0.047	0.608

HSV2-186

Data is not transformed as it is percentage.

Table 5: Linear mixed models.

	3.7U	11.1U	33.3U	100U
HSV2-186	0.56 (CI=-23.24,24.35, p=0.963)	-8.16 (CI=-35.15,18.83, p=0.553)	14.76 (CI=-25.47,54.98, p=0.472)	8.56 (CI=-17.65,34.78, p=0.522)

Table 6: Unpaired t test pvals.

	3.7U	11.1U	33.3U	100U
HSV2-186	0.876	0.577	0.510	0.562