

Analysis

2026-08-12

Donor 8 is filtered out through a single line in doOneRow.

HSV1 HSV2

Data is used as is because not skewed.

Question: Is there a difference between asymptomatic and severe at any dose?

```
## boundary (singular) fit: see help('isSingular')
## boundary (singular) fit: see help('isSingular')
## boundary (singular) fit: see help('isSingular')
```

Table 1: Linear mixed models.

	3.7U	11.1U	33.3U	100U
HSV1	-7.20 (CI=-27.56,13.15, p=0.488)W	-1.06 (CI=-19.70,17.59, p=0.911)	-2.36 (CI=-12.15, 7.42, p=0.636)	0.72 (CI=-2.99, 4.44, p=0.703)
HSV2	0.31 (CI=-17.81,18.44, p=0.973)	-2.93 (CI=-19.01,13.16, p=0.722)W	6.53 (CI=-10.03,23.08, p=0.440)W	0.70 (CI=-6.33, 7.72, p=0.846)

Table 2: Unpaired t test pvals.

	3.7U	11.1U	33.3U	100U
HSV1	0.455	0.914	0.704	0.706
HSV2	0.975	0.711	0.433	0.851

CCL5

Data is log transformed as it is skewed.

Donor 8 is filtered out

```
## boundary (singular) fit: see help('isSingular')
## boundary (singular) fit: see help('isSingular')
## boundary (singular) fit: see help('isSingular')
## boundary (singular) fit: see help('isSingular')
```

Table 3: Linear mixed models.

	0U	0.1U	1U	10U	100U
CCL5	0.00 (CI=-0.38,0.38, p=1.000)W	0.15 (CI=-0.23,0.52, p=0.444)W	0.04 (CI=-0.36,0.45, p=0.833)W	-0.42 (CI=-0.77,-0.07, p=0.018)*W	-0.01 (CI=-0.75,0.74, p=0.985)

Table 4: Unpaired t test pvals.

	0U	0.1U	1U	10U	100U
CCL5	0.054	0.230	0.768	0.032	0.985

HSV2-186

Data is not transformed as it is percentage.

Table 5: Linear mixed models.

	3.7U	11.1U	33.3U	100U
HSV2-186	3.93 (CI=-22.22,30.08, p=0.768)	-3.93 (CI=-34.32,26.46, p=0.800)	18.36 (CI=-28.50,65.22, p=0.442)	10.57 (CI=-20.10,41.24, p=0.499)

Table 6: Unpaired t test pvals.

	3.7U	11.1U	33.3U	100U
HSV2-186	0.753	0.801	0.438	0.483