

Eexam

Place student sticker here

Note:

- During the attendance check a sticker containing a unique code will be put on this exam.
- This code contains a unique number that associates this exam with your registration number.
- This number is printed both next to the code and to the signature field in the attendance check list.

Machine Learning for Graphs and Sequential Data (Problem sheet)

Graded Exercise: IN2323 / Endterm

Date: Friday 30th July, 2021

Examiner: Prof. Dr. Stephan Günnemann

Time: 11:30 – 12:45

Working instructions

- **DO NOT SUBMIT THIS SHEET! ONLY SUBMIT YOUR PERSONALIZED ANSWER SHEET THAT IS DISTRIBUTED THROUGH TUMEXAM!**
- Make sure that you solve the version of the problem stated on your personalized answer sheet (e.g., Problem 1 (Version B), Problem 2 (Version A), etc.)

Problem 1: Normalizing Flows (Version A)

We use the following normalizing flow model to specify a density $p_2(\mathbf{x})$ over \mathbb{R}^2 . The base density $p_1(\mathbf{z})$ is equal to the Uniform($[0, 2]^2$) distribution, that is

$$p_1(\mathbf{z}) = \begin{cases} \frac{1}{4} & \text{if } z_1 \in [0, 2] \text{ and } z_2 \in [0, 2], \\ 0 & \text{else.} \end{cases}$$

The model is specified by the following forward transformation $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$\mathbf{x} = f(\mathbf{z}) = \mathbf{A}\mathbf{z} = \begin{pmatrix} 1/3 & 0 \\ 0 & 1/4 \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}.$$

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>
3	<input type="checkbox"/>
4	<input type="checkbox"/>
5	<input type="checkbox"/>
6	<input type="checkbox"/>

Compute the density $p_2(\mathbf{x})$ at the point $\mathbf{x}^{(0)} = (1/2, 1/8)^T$. Justify your answer.

Problem 1: Normalizing Flows (Version B)

We use the following normalizing flow model to specify a density $p_2(\mathbf{x})$ over \mathbb{R}^2 . The base density $p_1(\mathbf{z})$ is equal to the $\text{Uniform}([0, 2]^2)$ distribution, that is

$$p_1(\mathbf{z}) = \begin{cases} \frac{1}{4} & \text{if } z_1 \in [0, 2] \text{ and } z_2 \in [0, 2], \\ 0 & \text{else.} \end{cases}$$

The model is specified by the following forward transformation $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$\mathbf{x} = f(\mathbf{z}) = \mathbf{A}\mathbf{z} = \begin{pmatrix} 1/2 & 0 \\ 0 & 1/12 \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}.$$

Compute the density $p_2(\mathbf{x})$ at the point $\mathbf{x}^{(0)} = (1/2, 1/8)^T$. Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6

Problem 1: Normalizing Flows (Version C)

We use the following normalizing flow model to specify a density $p_2(\mathbf{x})$ over \mathbb{R}^2 . The base density $p_1(\mathbf{z})$ is equal to the Uniform($[0, 2]^2$) distribution, that is

$$p_1(\mathbf{z}) = \begin{cases} \frac{1}{4} & \text{if } z_1 \in [0, 2] \text{ and } z_2 \in [0, 2], \\ 0 & \text{else.} \end{cases}$$

The model is specified by the following forward transformation $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$\mathbf{x} = f(\mathbf{z}) = \mathbf{A}\mathbf{z} = \begin{pmatrix} 1/3 & 0 \\ 0 & 1/12 \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}.$$

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>
3	<input type="checkbox"/>
4	<input type="checkbox"/>
5	<input type="checkbox"/>
6	<input type="checkbox"/>

Compute the density $p_2(\mathbf{x})$ at the point $\mathbf{x}^{(0)} = (1/2, 1/8)^T$. Justify your answer.

Problem 1: Normalizing Flows (Version D)

We use the following normalizing flow model to specify a density $p_2(\mathbf{x})$ over \mathbb{R}^2 . The base density $p_1(\mathbf{z})$ is equal to the $\text{Uniform}([0, 2]^2)$ distribution, that is

$$p_1(\mathbf{z}) = \begin{cases} \frac{1}{4} & \text{if } z_1 \in [0, 2] \text{ and } z_2 \in [0, 2], \\ 0 & \text{else.} \end{cases}$$

The model is specified by the following forward transformation $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$\mathbf{x} = f(\mathbf{z}) = \mathbf{A}\mathbf{z} = \begin{pmatrix} 1/2 & 0 \\ 0 & 1/10 \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}.$$

Compute the density $p_2(\mathbf{x})$ at the point $\mathbf{x}^{(0)} = (1/2, 1/8)^T$. Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6

Problem 2: Variational Inference (Version A)

We would like to draw samples from a logistic distribution with reparametrization.

The cumulative distribution function (CDF) of the logistic distribution with scale parameter ϕ is defined as

$$F_x(a) = \Pr(x \leq a) = \frac{1}{1 + \exp(-\phi a)}.$$

We have access to an algorithm that produces samples u from the uniform distribution on $[0, 1]$, that is

$$b(u) = \begin{cases} 1 & \text{if } u \in [0, 1], \\ 0 & \text{else.} \end{cases}$$

0 ☐ Derive a deterministic transformation T_ϕ that converts a sample $u \sim b(u)$ into a sample x from the logistic distribution.

1 ☐

2 ☐

3 ☐

4 ☐

5 ☐

6 ☐

Problem 2: Variational Inference (Version B)

We would like to draw samples from a logistic distribution with reparametrization.

The cumulative distribution function (CDF) of the logistic distribution with scale parameter ϕ is defined as

$$F_x(a) = \Pr(x \leq a) = \frac{1}{1 + \exp(-a + \phi)}.$$

We have access to an algorithm that produces samples u from the uniform distribution on $[0, 1]$, that is

$$b(u) = \begin{cases} 1 & \text{if } u \in [0, 1], \\ 0 & \text{else.} \end{cases}$$

Derive a deterministic transformation T_ϕ that converts a sample $u \sim b(u)$ into a sample x from the logistic distribution.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6

Problem 2: Variational Inference (Version C)

We would like to draw samples from a logistic distribution with reparametrization.

The cumulative distribution function (CDF) of the logistic distribution with scale parameter ϕ is defined as

$$F_x(a) = \Pr(x \leq a) = \frac{1}{1 + \exp(-a/\phi)}.$$

We have access to an algorithm that produces samples u from the uniform distribution on $[0, 1]$, that is

$$b(u) = \begin{cases} 1 & \text{if } u \in [0, 1], \\ 0 & \text{else.} \end{cases}$$

0 ☐ Derive a deterministic transformation T_ϕ that converts a sample $u \sim b(u)$ into a sample x from the logistic distribution.

1 ☐

2 ☐

3 ☐

4 ☐

5 ☐

6 ☐

Problem 2: Variational Inference (Version D)

We would like to draw samples from a logistic distribution with reparametrization.

The cumulative distribution function (CDF) of the logistic distribution with scale parameter ϕ is defined as

$$F_x(a) = \Pr(x \leq a) = \frac{1}{1 + \exp(-a - \phi)}.$$

We have access to an algorithm that produces samples u from the uniform distribution on $[0, 1]$, that is

$$b(u) = \begin{cases} 1 & \text{if } u \in [0, 1], \\ 0 & \text{else.} \end{cases}$$

Derive a deterministic transformation T_ϕ that converts a sample $u \sim b(u)$ into a sample x from the logistic distribution.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6

Problem 3: Deep Generative Models (Version A)

Suppose $f: \mathbb{R}^N \rightarrow \mathbb{R}^M$ is a fully-connected neural network, where $N < M$.

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>

a) Can we use f as the generator in a generative adversarial network? Justify your answer.

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>

b) Can we use f as the transformation to define a normalizing flow model? Justify your answer.

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>

c) Can we use f as the decoder in a variational autoencoder model? Justify your answer.

Problem 3: Deep Generative Models (Version B)

Suppose $f: \mathbb{R}^N \rightarrow \mathbb{R}^M$ is a fully-connected neural network, where $N < M$.

a) Can we use f as the decoder in a variational autoencoder model? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

b) Can we use f as the transformation to define a normalizing flow model? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

c) Can we use f as the generator in a generative adversarial network? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

Problem 3: Deep Generative Models (Version C)

Suppose $f: \mathbb{R}^N \rightarrow \mathbb{R}^M$ is a fully-connected neural network, where $N < M$.

0 ☐
1 ☐
2 ☐ a) Can we use f as the transformation to define a normalizing flow model? Justify your answer.

0 ☐
1 ☐
2 ☐ b) Can we use f as the generator in a generative adversarial network? Justify your answer.

0 ☐
1 ☐
2 ☐ c) Can we use f as the decoder in a variational autoencoder model? Justify your answer.

Problem 3: Deep Generative Models (Version D)

Suppose $f: \mathbb{R}^N \rightarrow \mathbb{R}^M$ is a fully-connected neural network, where $N < M$.

a) Can we use f as the generator in a generative adversarial network? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

b) Can we use f as the decoder in a variational autoencoder model? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

c) Can we use f as the transformation to define a normalizing flow model? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

Problem 4: Robustness (Version A)

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>
3	<input type="checkbox"/>
4	<input type="checkbox"/>
5	<input type="checkbox"/>
6	<input type="checkbox"/>
7	<input type="checkbox"/>
8	<input type="checkbox"/>

Assume we are interested in robustness certification for a model classifying data from \mathbb{R}^D . So far, we have only considered norm-bound perturbation models that — for the l_2 norm — can be expressed via the constraints

$$\begin{aligned} \|\tilde{\mathbf{x}} - \mathbf{x}\|_2 &\leq \epsilon, \\ \tilde{\mathbf{x}} &\in \mathbb{R}^D, \end{aligned} \tag{4.1}$$

with adversarial budget $\epsilon \geq 0$, perturbed input $\tilde{\mathbf{x}}$ and unperturbed input $\mathbf{x} \in \mathbb{R}^D$.

Now we want to model *sparse* norm-bound perturbations, in which the adversary can perturb at most $\eta \in \mathbb{N}$ vector entries. For this, extend Equation (4.1) with additional **linear constraints**. You may introduce at most $\mathcal{O}(D)$ constraints and $\mathcal{O}(D)$ variables. You are allowed to use integer-valued variables.

More formally, the modeled perturbation set should be

$$\mathcal{P}(\mathbf{x}) = \left\{ \tilde{\mathbf{x}} \in \mathbb{R}^D \mid \|\tilde{\mathbf{x}} - \mathbf{x}\|_2 \leq \epsilon \wedge \sum_{d=0}^{D-1} \mathbb{1}(\tilde{x}_d \neq x_d) \leq \eta \right\},$$

where $\mathbb{1}(\cdot)$ is the indicator function.

Problem 4: Robustness (Version B)

Assume we are interested in robustness certification for a model classifying data from \mathbb{R}^D . So far, we have only considered norm-bound perturbation models that — for the l_2 norm — can be expressed via the constraints

$$\begin{aligned} \|\tilde{\mathbf{x}} - \mathbf{x}\|_2 &\leq \epsilon, \\ \tilde{\mathbf{x}} &\in \mathbb{R}^D, \end{aligned} \tag{4.1}$$


with adversarial budget $\epsilon \geq 0$, perturbed input $\tilde{\mathbf{x}}$ and unperturbed input $\mathbf{x} \in \mathbb{R}^D$.

Now we want to model *sparse* norm-bound perturbations, in which the adversary can perturb at most $\eta \in \mathbb{N}$ vector entries. For this, extend Equation (4.1) with additional **linear constraints**. You may introduce at most $\mathcal{O}(D)$ constraints and $\mathcal{O}(D)$ variables. You are allowed to use integer-valued variables.

More formally, the modeled perturbation set should be

$$\mathcal{P}(\mathbf{x}) = \left\{ \tilde{\mathbf{x}} \in \mathbb{R}^D \mid \|\tilde{\mathbf{x}} - \mathbf{x}\|_2 \leq \epsilon \wedge \sum_{d=0}^{D-1} \mathbb{1}(\tilde{x}_d \neq x_d) \leq \eta \right\},$$

where $\mathbb{1}(\cdot)$ is the indicator function.



Problem 4: Robustness (Version C)

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>
3	<input type="checkbox"/>
4	<input type="checkbox"/>
5	<input type="checkbox"/>
6	<input type="checkbox"/>
7	<input type="checkbox"/>
8	<input type="checkbox"/>

Assume we are interested in robustness certification for a model classifying data from \mathbb{R}^D . So far, we have only considered norm-bound perturbation models that — for the l_2 norm — can be expressed via the constraints

$$\begin{aligned} \|\tilde{\mathbf{x}} - \mathbf{x}\|_2 &\leq \epsilon, \\ \tilde{\mathbf{x}} &\in \mathbb{R}^D, \end{aligned} \tag{4.1}$$

with adversarial budget $\epsilon \geq 0$, perturbed input $\tilde{\mathbf{x}}$ and unperturbed input $\mathbf{x} \in \mathbb{R}^D$.

Now we want to model *sparse* norm-bound perturbations, in which the adversary can perturb at most $\eta \in \mathbb{N}$ vector entries. For this, extend Equation (4.1) with additional **linear constraints**. You may introduce at most $\mathcal{O}(D)$ constraints and $\mathcal{O}(D)$ variables. You are allowed to use integer-valued variables.

More formally, the modeled perturbation set should be

$$\mathcal{P}(\mathbf{x}) = \left\{ \tilde{\mathbf{x}} \in \mathbb{R}^D \mid \|\tilde{\mathbf{x}} - \mathbf{x}\|_2 \leq \epsilon \wedge \sum_{d=0}^{D-1} \mathbb{1}(\tilde{x}_d \neq x_d) \leq \eta \right\},$$

where $\mathbb{1}(\cdot)$ is the indicator function.

Problem 4: Robustness (Version D)

Assume we are interested in robustness certification for a model classifying data from \mathbb{R}^D . So far, we have only considered norm-bound perturbation models that — for the l_2 norm — can be expressed via the constraints

$$\begin{aligned} \|\tilde{\mathbf{x}} - \mathbf{x}\|_2 &\leq \epsilon, \\ \tilde{\mathbf{x}} &\in \mathbb{R}^D, \end{aligned} \tag{4.1}$$

with adversarial budget $\epsilon \geq 0$, perturbed input $\tilde{\mathbf{x}}$ and unperturbed input $\mathbf{x} \in \mathbb{R}^D$.

Now we want to model *sparse* norm-bound perturbations, in which the adversary can perturb at most $\eta \in \mathbb{N}$ vector entries. For this, extend Equation (4.1) with additional **linear constraints**. You may introduce at most $\mathcal{O}(D)$ constraints and $\mathcal{O}(D)$ variables. You are allowed to use integer-valued variables.

More formally, the modeled perturbation set should be

$$\mathcal{P}(\mathbf{x}) = \left\{ \tilde{\mathbf{x}} \in \mathbb{R}^D \mid \|\tilde{\mathbf{x}} - \mathbf{x}\|_2 \leq \epsilon \wedge \sum_{d=0}^{D-1} \mathbb{1}(\tilde{x}_d \neq x_d) \leq \eta \right\},$$

where $\mathbb{1}(\cdot)$ is the indicator function.



Problem 5: Autoregressive Models (Version A)

Consider a sequential process described by the following autoregressive model.

$$X_t = X_{t-1} + \varepsilon_t$$

The noise variables ε_t are independently $\mathcal{N}(0, 1)$ distributed and we condition the process on $X_0 = 0$.

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>

a) Is this autoregressive process stationary? Justify your answer.

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>

b) Now we define a second process Y_t as following

$$Y_t = X_t - X_{t-1}.$$

Is the process Y_t stationary? Justify your answer.

Problem 5: Autoregressive Models (Version B)

Consider a sequential process described by the following autoregressive model.

$$X_t = X_{t-1} + \varepsilon_t$$

The noise variables ε_t are independently $\mathcal{N}(0, 1)$ distributed and we condition the process on $X_0 = 0$.

a) Is this autoregressive process stationary? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

b) Now we define a second process Y_t as following

$$Y_t = X_t - X_{t-1}.$$

Is the process Y_t stationary? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

Problem 5: Autoregressive Models (Version C)

Consider a sequential process described by the following autoregressive model.

$$X_t = X_{t-1} + \varepsilon_t$$

The noise variables ε_t are independently $\mathcal{N}(0, 1)$ distributed and we condition the process on $X_0 = 0$.

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>

a) Is this autoregressive process stationary? Justify your answer.

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>

b) Now we define a second process Y_t as following

$$Y_t = X_t - X_{t-1}.$$

Is the process Y_t stationary? Justify your answer.

Problem 5: Autoregressive Models (Version D)

Consider a sequential process described by the following autoregressive model.

$$X_t = X_{t-1} + \varepsilon_t$$

The noise variables ε_t are independently $\mathcal{N}(0, 1)$ distributed and we condition the process on $X_0 = 0$.

a) Is this autoregressive process stationary? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

b) Now we define a second process Y_t as following

$$Y_t = X_t - X_{t-1}.$$

Is the process Y_t stationary? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2

Problem 6: Hidden Markov Models (Version A)

Consider a hidden Markov model with 3 states 1, 2, 3. There are 3 possible observations a, b, c. The initial distribution π , transition probabilities \mathbf{A} and emission probabilities \mathbf{B} are

$$\pi = \begin{matrix} & 1 & 2 & 3 \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 2/5 \\ 2/5 \\ 1/5 \end{pmatrix} \end{matrix}$$

$$\mathbf{A} = \begin{matrix} & 1 & 2 & 3 \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 2/5 & 1/5 & 2/5 \\ 2/5 & 2/5 & 1/5 \\ 1/5 & 0 & 4/5 \end{pmatrix} \end{matrix}$$

$$\mathbf{B} = \begin{matrix} & a & b & c \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 3/5 & 1/5 & 1/5 \\ 1/5 & 3/5 & 1/5 \\ 2/5 & 0 & 3/5 \end{pmatrix} \end{matrix},$$

where \mathbf{A}_{ij} specifies the probability of transitioning from state i to state j .

0 ☐ We have observed the sequence $X_{1:3} = [bbc]$. What is the most likely latent state Z_2 given these observations?
 1 ☐ Justify your answer.
 2 ☐
 3 ☐
 4 ☐
 5 ☐
 6 ☐
 7 ☐
 8 ☐

Problem 6: Hidden Markov Models (Version B)

Consider a hidden Markov model with 3 states 1, 2, 3. There are 3 possible observations a, b, c. The initial distribution π , transition probabilities \mathbf{A} and emission probabilities \mathbf{B} are

$$\pi = \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} \begin{pmatrix} 2/5 \\ 1/5 \\ 2/5 \end{pmatrix} \quad \mathbf{A} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 2/5 & 2/5 & 1/5 \\ 1/5 & 4/5 & 0 \\ 2/5 & 1/5 & 2/5 \end{pmatrix} \end{matrix} \quad \mathbf{B} = \begin{matrix} & \begin{matrix} a & b & c \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 1/5 & 1/5 & 3/5 \\ 0 & 3/5 & 2/5 \\ 3/5 & 1/5 & 1/5 \end{pmatrix} \end{matrix},$$

where \mathbf{A}_{ij} specifies the probability of transitioning from state i to state j .

We have observed the sequence $X_{1:3} = [\text{aab}]$. What is the most likely latent state Z_2 given these observations? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6
<input type="checkbox"/>	7
<input type="checkbox"/>	8

Problem 6: Hidden Markov Models (Version C)

Consider a hidden Markov model with 3 states 1, 2, 3. There are 3 possible observations a, b, c. The initial distribution π , transition probabilities \mathbf{A} and emission probabilities \mathbf{B} are

$$\pi = \begin{matrix} & 1 & 2 & 3 \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 1/5 \\ 2/5 \\ 2/5 \end{pmatrix} \end{matrix}$$

$$\mathbf{A} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 4/5 & 0 & 1/5 \\ 1/5 & 2/5 & 2/5 \\ 2/5 & 1/5 & 2/5 \end{pmatrix} \end{matrix}$$

$$\mathbf{B} = \begin{matrix} & \begin{matrix} a & b & c \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 2/5 & 3/5 & 0 \\ 1/5 & 1/5 & 3/5 \\ 3/5 & 1/5 & 1/5 \end{pmatrix} \end{matrix},$$

where \mathbf{A}_{ij} specifies the probability of transitioning from state i to state j .

We have observed the sequence $X_{1:3} = [\text{ccb}]$. What is the most likely latent state Z_2 given these observations? Justify your answer.

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>
3	<input type="checkbox"/>
4	<input type="checkbox"/>
5	<input type="checkbox"/>
6	<input type="checkbox"/>
7	<input type="checkbox"/>
8	<input type="checkbox"/>

Problem 6: Hidden Markov Models (Version D)

Consider a hidden Markov model with 3 states 1, 2, 3. There are 3 possible observations a, b, c. The initial distribution π , transition probabilities \mathbf{A} and emission probabilities \mathbf{B} are

$$\pi = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 2/5 \\ 2/5 \\ 1/5 \end{pmatrix} \end{matrix} \quad \mathbf{A} = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 2/5 & 1/5 & 2/5 \\ 2/5 & 2/5 & 1/5 \\ 1/5 & 0 & 4/5 \end{pmatrix} \end{matrix} \quad \mathbf{B} = \begin{matrix} & \begin{matrix} a & b & c \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 1/5 & 3/5 & 1/5 \\ 1/5 & 1/5 & 3/5 \\ 3/5 & 2/5 & 0 \end{pmatrix} \end{matrix},$$

where \mathbf{A}_{ij} specifies the probability of transitioning from state i to state j .

We have observed the sequence $X_{1:3} = [\text{cca}]$. What is the most likely latent state Z_2 given these observations? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6
<input type="checkbox"/>	7
<input type="checkbox"/>	8

Problem 7: Temporal point processes (Version A)

0 ☐
1 ☐
2 ☐
3 ☐
4 ☐
5 ☐
6 ☐

We simulate a homogeneous Poisson process with intensity μ on the interval $[0, T]$. What is the probability that we observe no events (i.e., the first event t_1 will happen after T)? Justify your answer.

Problem 7: Temporal point processes (Version B)

We simulate a homogeneous Poisson process with intensity μ on the interval $[0, T]$. What is the probability that we observe no events (i.e., the first event t_1 will happen after T)? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6

Problem 7: Temporal point processes (Version C)

0	<input type="checkbox"/>
1	<input type="checkbox"/>
2	<input type="checkbox"/>
3	<input type="checkbox"/>
4	<input type="checkbox"/>
5	<input type="checkbox"/>
6	<input type="checkbox"/>


We simulate a homogeneous Poisson process with intensity μ on the interval $[0, T]$. What is the probability that we observe no events (i.e., the first event t_1 will happen after T)? Justify your answer.

Problem 7: Temporal point processes (Version D)

We simulate a homogeneous Poisson process with intensity μ on the interval $[0, T]$. What is the probability that we observe no events (i.e., the first event t_1 will happen after T)? Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6

Problem 8: Graphs - Clustering (Version A)



In this task, we consider a Stochastic Block Model with 2 communities that can generate *weighted* graphs. We assume that the edge weights are drawn from the Poisson distribution, that is

$$p(A_{ij} = k | z_i, z_j, \boldsymbol{\eta}) = \frac{\eta_{z_i z_j}^k}{k!} \exp(-\eta_{z_i z_j})$$

Suppose we observed the weighted adjacency matrix $\mathbf{A} \in \{0, 1, 2, \dots\}^{N \times N}$ and the cluster indicators $\mathbf{z} \in \{1, 2\}^N$. Derive the maximum likelihood estimate of the parameters $\boldsymbol{\eta} = \begin{pmatrix} \eta_{11} & \eta_{12} \\ \eta_{21} & \eta_{22} \end{pmatrix}$.

Hint: It might be helpful to introduce shorthand notation

$$N_p = \sum_{i=1}^N \mathbb{1}(z_i = p) \quad \text{and} \quad M_{pq} = \sum_{i=1}^N \sum_{j=1}^N A_{ij} \mathbb{1}(z_i = p, z_j = q),$$

where $\mathbb{1}(\cdot)$ is the indicator function.

Problem 8: Graphs - Clustering (Version B)

In this task, we consider a Stochastic Block Model with 2 communities that can generate *weighted* graphs. We assume that the edge weights are drawn from the Poisson distribution, that is


$$p(A_{ij} = k | z_i, z_j, \boldsymbol{\eta}) = \frac{\eta_{z_i z_j}^k}{k!} \exp(-\eta_{z_i z_j})$$

Suppose we observed the weighted adjacency matrix $\mathbf{A} \in \{0, 1, 2, \dots\}^{N \times N}$ and the cluster indicators $\mathbf{z} \in \{1, 2\}^N$. Derive the maximum likelihood estimate of the parameters $\boldsymbol{\eta} = \begin{pmatrix} \eta_{11} & \eta_{12} \\ \eta_{21} & \eta_{22} \end{pmatrix}$.


Hint: It might be helpful to introduce shorthand notation

$$N_p = \sum_{i=1}^N \mathbb{1}(z_i = p) \quad \text{and} \quad M_{pq} = \sum_{i=1}^N \sum_{j=1}^N A_{ij} \mathbb{1}(z_i = p, z_j = q),$$

where $\mathbb{1}(\cdot)$ is the indicator function.



Problem 8: Graphs - Clustering (Version C)



In this task, we consider a Stochastic Block Model with 2 communities that can generate *weighted* graphs. We assume that the edge weights are drawn from the Poisson distribution, that is

$$p(A_{ij} = k | z_i, z_j, \boldsymbol{\eta}) = \frac{\eta_{z_i z_j}^k}{k!} \exp(-\eta_{z_i z_j})$$

Suppose we observed the weighted adjacency matrix $\mathbf{A} \in \{0, 1, 2, \dots\}^{N \times N}$ and the cluster indicators $\mathbf{z} \in \{1, 2\}^N$. Derive the maximum likelihood estimate of the parameters $\boldsymbol{\eta} = \begin{pmatrix} \eta_{11} & \eta_{12} \\ \eta_{21} & \eta_{22} \end{pmatrix}$.

Hint: It might be helpful to introduce shorthand notation

$$N_p = \sum_{i=1}^N \mathbb{1}(z_i = p) \quad \text{and} \quad M_{pq} = \sum_{i=1}^N \sum_{j=1}^N A_{ij} \mathbb{1}(z_i = p, z_j = q),$$

where $\mathbb{1}(\cdot)$ is the indicator function.

Problem 8: Graphs - Clustering (Version D)

In this task, we consider a Stochastic Block Model with 2 communities that can generate *weighted* graphs. We assume that the edge weights are drawn from the Poisson distribution, that is


$$p(A_{ij} = k | z_i, z_j, \boldsymbol{\eta}) = \frac{\eta_{z_i z_j}^k}{k!} \exp(-\eta_{z_i z_j})$$

Suppose we observed the weighted adjacency matrix $\mathbf{A} \in \{0, 1, 2, \dots\}^{N \times N}$ and the cluster indicators $\mathbf{z} \in \{1, 2\}^N$. Derive the maximum likelihood estimate of the parameters $\boldsymbol{\eta} = \begin{pmatrix} \eta_{11} & \eta_{12} \\ \eta_{21} & \eta_{22} \end{pmatrix}$.

Hint: It might be helpful to introduce shorthand notation

$$N_p = \sum_{i=1}^N \mathbb{1}(z_i = p) \quad \text{and} \quad M_{pq} = \sum_{i=1}^N \sum_{j=1}^N A_{ij} \mathbb{1}(z_i = p, z_j = q),$$

where $\mathbb{1}(\cdot)$ is the indicator function.



Problem 9: Graphs - Ranking (Version A)

We consider a directed graph $G = (V, E)$ with adjacency matrix

$$\mathbf{A} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

where A_{ij} indicates if there exists an edge from node i to node j .

- 0 ☐ a) Find the stationary distribution $\pi(\infty)$ associated with the random walk on graph G . Justify your answer.
1 ☐
2 ☐
3 ☐

- 0 ☐ b) Explain why the Markov chain corresponding to the random walk on graph G is not irreducible.
1 ☐

- 0 ☐ c) We want to add a single directed edge (u, v) to the graph G and create a new graph $G' = (V, E \cup \{(u, v)\})$, such
1 ☐ that the Markov chain corresponding to the random walk on graph G' becomes irreducible. What edge (u, v) should
2 ☐ we add such that the PageRank score for node 1 in G' is also maximized? Also, compute the PageRank scores for
3 ☐ all the nodes in G' after adding the chosen edge (u, v) .
4 ☐
5 ☐
6 ☐

Problem 9: Graphs - Ranking (Version B)

We consider a directed graph $G = (V, E)$ with adjacency matrix

$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

where A_{ij} indicates if there exists an edge from node i to node j .

a) Find the stationary distribution $\pi(\infty)$ associated with the random walk on graph G . Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3

b) Explain why the Markov chain corresponding to the random walk on graph G is not irreducible.

<input type="checkbox"/>	0
<input type="checkbox"/>	1

c) We want to add a single directed edge (u, v) to the graph G and create a new graph $G' = (V, E \cup \{(u, v)\})$, such that the Markov chain corresponding to the random walk on graph G' becomes irreducible. What edge (u, v) should we add such that the PageRank score for node 1 in G' is also maximized? Also, compute the PageRank scores for all the nodes in G' after adding the chosen edge (u, v) .

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6

Problem 9: Graphs - Ranking (Version C)

We consider a directed graph $G = (V, E)$ with adjacency matrix

$$\mathbf{A} = \begin{bmatrix} 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

where A_{ij} indicates if there exists an edge from node i to node j .

- 0 ☐ a) Find the stationary distribution $\pi(\infty)$ associated with the random walk on graph G . Justify your answer.
1 ☐
2 ☐
3 ☐

- 0 ☐ b) Explain why the Markov chain corresponding to the random walk on graph G is not irreducible.
1 ☐

- 0 ☐ c) We want to add a single directed edge (u, v) to the graph G and create a new graph $G' = (V, E \cup \{(u, v)\})$, such
1 ☐ that the Markov chain corresponding to the random walk on graph G' becomes irreducible. What edge (u, v) should
2 ☐ we add such that the PageRank score for node 1 in G' is also maximized? Also, compute the PageRank scores for
3 ☐ all the nodes in G' after adding the chosen edge (u, v) .
4 ☐
5 ☐
6 ☐

Problem 9: Graphs - Ranking (Version D)

We consider a directed graph $G = (V, E)$ with adjacency matrix

$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

where A_{ij} indicates if there exists an edge from node i to node j .

a) Find the stationary distribution $\pi(\infty)$ associated with the random walk on graph G . Justify your answer.

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3

b) Explain why the Markov chain corresponding to the random walk on graph G is not irreducible.

<input type="checkbox"/>	0
<input type="checkbox"/>	1


c) We want to add a single directed edge (u, v) to the graph G and create a new graph $G' = (V, E \cup \{(u, v)\})$, such that the Markov chain corresponding to the random walk on graph G' becomes irreducible. What edge (u, v) should we add such that the PageRank score for node 1 in G' is also maximized? Also, compute the PageRank scores for all the nodes in G' after adding the chosen edge (u, v) .

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5
<input type="checkbox"/>	6

Problem 10: Graphs - Semi-Supervised Learning (Version A)

We consider the graph G with the following adjacency matrix \mathbf{A} .

$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



We assume that the vector of labels is $\mathbf{y} = \begin{bmatrix} \hat{\mathbf{y}}_S \\ \mathbf{y}_U \end{bmatrix}$ where $\hat{\mathbf{y}}_S = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ are known labels and $\mathbf{y}_U = \begin{bmatrix} * \\ * \\ * \end{bmatrix}$ are unknown labels.

In other words, $S = \{1, 2\}$ is the set of labeled nodes and $U = \{3, 4, 5\}$ is the set of unlabeled nodes.

0 ☐
1 ☐
2 ☐
3 ☐

a) Find the optimal cost and all the optimal solutions to the following optimization problem:

$$\begin{aligned} \mathbf{y}^* &= \arg \min_{\mathbf{y} \in \{0,1\}^5} \mathbf{y}^T \mathbf{L} \mathbf{y} \\ \text{subject to } y_i &= \hat{y}_{S,i} \text{ for } i \in \{1, 2\} \end{aligned} \quad (10.1)$$

where $\mathbf{L} = \mathbf{D} - \mathbf{A}$ denotes the Laplacian of the graph G , and \mathbf{D} is the degree matrix. Justify your answer.

0 ☐
1 ☐
2 ☐
3 ☐
4 ☐
5 ☐


b) We consider a relaxation of the optimization problem from Equation (10.1) (note that we now optimize over \mathbb{R}^6):

$$\begin{aligned} \mathbf{y}^* &= \arg \min_{\mathbf{y} \in \mathbb{R}^5} \mathbf{y}^T \mathbf{L} \mathbf{y} \\ \text{subject to } y_i &= \hat{y}_{S,i} \text{ for } i \in \{1, 2\} \end{aligned} \quad (10.2)$$

Suppose that the solution \mathbf{y}^* of this optimization problem is given.

We now consider a modified graph G' with adjacency matrix

$$\mathbf{A}' = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



We denote the labels of G' as $\mathbf{y}' = \begin{bmatrix} \hat{\mathbf{y}}'_S \\ \mathbf{y}'_U \end{bmatrix}$, where $\mathbf{y}'_S = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ are known labels and $\mathbf{y}'_U = \begin{bmatrix} * \\ * \\ * \end{bmatrix}$ are unknown labels.

Your task is to explain how the solution \mathbf{y}^* of the optimization problem in Equation 10.2 can be used to solve the following optimization problem:


$$\begin{aligned} \arg \min_{\mathbf{y}' \in \mathbb{R}^6} \mathbf{y}'^T \mathbf{L}' \mathbf{y}' \\ \text{subject to } y'_i &= \hat{y}'_{S,i} \text{ for } i \in \{1, 2, 3\} \end{aligned} \quad (10.3)$$

where $\mathbf{L}' = \mathbf{A}' - \mathbf{D}'$ is the Laplacian of G' . Justify your answer.

Problem 10: Graphs - Semi-Supervised Learning (Version B)

We consider the graph G with the following adjacency matrix A .

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



We assume that the vector of labels is $\mathbf{y} = \begin{bmatrix} \hat{\mathbf{y}}_S \\ \mathbf{y}_U \end{bmatrix}$ where $\hat{\mathbf{y}}_S = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ are known labels and $\mathbf{y}_U = \begin{bmatrix} * \\ * \\ * \end{bmatrix}$ are unknown labels.

In other words, $S = \{1, 2\}$ is the set of labeled nodes and $U = \{3, 4, 5\}$ is the set of unlabeled nodes.

a) Find the optimal cost and all the optimal solutions to the following optimization problem:

$$\begin{aligned} \mathbf{y}^* &= \arg \min_{\mathbf{y} \in \{0,1\}^5} \mathbf{y}^T \mathbf{L} \mathbf{y} \\ \text{subject to } y_i &= \hat{y}_{S,i} \text{ for } i \in \{1, 2\} \end{aligned} \quad (10.1)$$

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3

where $\mathbf{L} = \mathbf{D} - \mathbf{A}$ denotes the Laplacian of the graph G , and \mathbf{D} is the degree matrix. Justify your answer.

b) We consider a relaxation of the optimization problem from Equation (10.1) (note that we now optimize over \mathbb{R}^6):


$$\begin{aligned} \mathbf{y}^* &= \arg \min_{\mathbf{y} \in \mathbb{R}^5} \mathbf{y}^T \mathbf{L} \mathbf{y} \\ \text{subject to } y_i &= \hat{y}_{S,i} \text{ for } i \in \{1, 2\} \end{aligned} \quad (10.2)$$

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5

Suppose that the solution \mathbf{y}^* of this optimization problem is given.

We now consider a modified graph G' with adjacency matrix

$$A' = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



We denote the labels of G' as $\mathbf{y}' = \begin{bmatrix} \hat{\mathbf{y}}'_S \\ \mathbf{y}'_U \end{bmatrix}$, where $\hat{\mathbf{y}}'_S = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ are known labels and $\mathbf{y}'_U = \begin{bmatrix} * \\ * \\ * \end{bmatrix}$ are unknown labels.

Your task is to explain how the solution \mathbf{y}^* of the optimization problem in Equation 10.2 can be used to solve the following optimization problem:


$$\begin{aligned} \arg \min_{\mathbf{y}' \in \mathbb{R}^6} \mathbf{y}'^T \mathbf{L}' \mathbf{y}' \\ \text{subject to } y'_i &= \hat{y}'_{S,i} \text{ for } i \in \{1, 2, 3\} \end{aligned} \quad (10.3)$$

where $\mathbf{L}' = \mathbf{A}' - \mathbf{D}'$ is the Laplacian of G' . Justify your answer.

Problem 10: Graphs - Semi-Supervised Learning (Version C)

We consider the graph G with the following adjacency matrix \mathbf{A} .

$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



We assume that the vector of labels is $\mathbf{y} = \begin{bmatrix} \hat{\mathbf{y}}_S \\ \mathbf{y}_U \end{bmatrix}$ where $\hat{\mathbf{y}}_S = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ are known labels and $\mathbf{y}_U = \begin{bmatrix} * \\ * \\ * \end{bmatrix}$ are unknown labels.

In other words, $S = \{1, 2\}$ is the set of labeled nodes and $U = \{3, 4, 5\}$ is the set of unlabeled nodes.

0 ☐
1 ☐
2 ☐
3 ☐

a) Find the optimal cost and all the optimal solutions to the following optimization problem:

$$\begin{aligned} \mathbf{y}^* &= \arg \min_{\mathbf{y} \in \{0,1\}^5} \mathbf{y}^T \mathbf{L} \mathbf{y} \\ \text{subject to } y_i &= \hat{y}_{S,i} \text{ for } i \in \{1, 2\} \end{aligned} \quad (10.1)$$

where $\mathbf{L} = \mathbf{D} - \mathbf{A}$ denotes the Laplacian of the graph G , and \mathbf{D} is the degree matrix. Justify your answer.

0 ☐
1 ☐
2 ☐
3 ☐
4 ☐
5 ☐


b) We consider a relaxation of the optimization problem from Equation (10.1) (note that we now optimize over \mathbb{R}^5):

$$\begin{aligned} \mathbf{y}^* &= \arg \min_{\mathbf{y} \in \mathbb{R}^5} \mathbf{y}^T \mathbf{L} \mathbf{y} \\ \text{subject to } y_i &= \hat{y}_{S,i} \text{ for } i \in \{1, 2\} \end{aligned} \quad (10.2)$$

Suppose that the solution \mathbf{y}^* of this optimization problem is given.

We now consider a modified graph G' with adjacency matrix

$$\mathbf{A}' = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



We denote the labels of G' as $\mathbf{y}' = \begin{bmatrix} \hat{\mathbf{y}}'_S \\ \mathbf{y}'_U \end{bmatrix}$, where $\mathbf{y}'_S = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ are known labels and $\mathbf{y}'_U = \begin{bmatrix} * \\ * \\ * \end{bmatrix}$ are unknown labels.

Your task is to explain how the solution \mathbf{y}^* of the optimization problem in Equation 10.2 can be used to solve the following optimization problem:


$$\begin{aligned} \arg \min_{\mathbf{y}' \in \mathbb{R}^6} \mathbf{y}'^T \mathbf{L}' \mathbf{y}' \\ \text{subject to } y'_i &= \hat{y}'_{S,i} \text{ for } i \in \{1, 2, 3\} \end{aligned} \quad (10.3)$$

where $\mathbf{L}' = \mathbf{A}' - \mathbf{D}'$ is the Laplacian of G' . Justify your answer.

Problem 10: Graphs - Semi-Supervised Learning (Version D)

We consider the graph G with the following adjacency matrix \mathbf{A} .

$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



We assume that the vector of labels is $\mathbf{y} = \begin{bmatrix} \hat{\mathbf{y}}_S \\ \mathbf{y}_U \end{bmatrix}$ where $\hat{\mathbf{y}}_S = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ are known labels and $\mathbf{y}_U = \begin{bmatrix} * \\ * \\ * \end{bmatrix}$ are unknown labels.

In other words, $S = \{1, 2\}$ is the set of labeled nodes and $U = \{3, 4, 5\}$ is the set of unlabeled nodes.

a) Find the optimal cost and all the optimal solutions to the following optimization problem:

$$\begin{aligned} \mathbf{y}^* &= \arg \min_{\mathbf{y} \in \{0,1\}^5} \mathbf{y}^T \mathbf{L} \mathbf{y} \\ \text{subject to } y_i &= \hat{y}_{S,i} \text{ for } i \in \{1, 2\} \end{aligned} \quad (10.1)$$

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3

where $\mathbf{L} = \mathbf{D} - \mathbf{A}$ denotes the Laplacian of the graph G , and \mathbf{D} is the degree matrix. Justify your answer.

b) We consider a relaxation of the optimization problem from Equation (10.1) (note that we now optimize over \mathbb{R}^6):


$$\begin{aligned} \mathbf{y}^* &= \arg \min_{\mathbf{y} \in \mathbb{R}^5} \mathbf{y}^T \mathbf{L} \mathbf{y} \\ \text{subject to } y_i &= \hat{y}_{S,i} \text{ for } i \in \{1, 2\} \end{aligned} \quad (10.2)$$

<input type="checkbox"/>	0
<input type="checkbox"/>	1
<input type="checkbox"/>	2
<input type="checkbox"/>	3
<input type="checkbox"/>	4
<input type="checkbox"/>	5

Suppose that the solution \mathbf{y}^* of this optimization problem is given.

We now consider a modified graph G' with adjacency matrix

$$\mathbf{A}' = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$



We denote the labels of G' as $\mathbf{y}' = \begin{bmatrix} \hat{\mathbf{y}}'_S \\ \mathbf{y}'_U \end{bmatrix}$, where $\hat{\mathbf{y}}'_S = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ are known labels and $\mathbf{y}'_U = \begin{bmatrix} * \\ * \\ * \end{bmatrix}$ are unknown labels.

Your task is to explain how the solution \mathbf{y}^* of the optimization problem in Equation 10.2 can be used to solve the following optimization problem:

$$\begin{aligned} \arg \min_{\mathbf{y}' \in \mathbb{R}^6} \mathbf{y}'^T \mathbf{L}' \mathbf{y}' \\ \text{subject to } y'_i &= \hat{y}'_{S,i} \text{ for } i \in \{1, 2, 3\} \end{aligned} \quad (10.3)$$

where $\mathbf{L}' = \mathbf{A}' - \mathbf{D}'$ is the Laplacian of G' . Justify your answer.

