



Review

Soil carbon sequestration, greenhouse gas emissions, and water pollution under different tillage practices



Siddhartha Shankar Bhattacharyya^a, Fernanda Figueiredo Granja Dorilêo Leite^b, Casey L. France^c, Adetomi O. Adekoya^d, Gerard H. Ros^e, Wim de Vries^e, Elda M. Melchor-Martínez^f, Hafiz M.N. Iqbal^{f,*}, Roberto Parra-Saldívar^{f,*}

^a Department of Agronomy, Faculty of Agriculture, Bangladesh Agricultural University, Mymensingh 2200, Bangladesh

^b Chemistry Institute, Geochemistry Department (Graduate program of Environmental Geochemistry), Federal Fluminense University, Niterói, RJ, Brazil

^c Pivot Bio, 1714 W 56th St., Kearney, NE 68845, USA

^d Department of Crop Protection and Environmental Biology, University of Ibadan, Ibadan, Nigeria

^e Environmental Systems Analysis Group, Wageningen University and Research, Wageningen, the Netherlands

^f Tecnológico de Monterrey, School of Engineering and Science, Monterrey 64849, Mexico

HIGHLIGHTS

- The soil carbon sequestration (SCS) potential of no-tillage (NT) is high compared to other tillage practices.
- Tillage induced erosion, which decrease the magnitude of SCS, is less for NT and RT compared to CT.
- The Eutrophication potential of no-tillage (NT) is higher than CT due to higher NO₃⁻ leaching.
- The SCS potential of NT is counteracted by substantial nitrous oxide (N₂O) emission.
- Nitrification inhibitors coupled with tillage practices can offset both N₂O emission and nitrate pollution.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 27 October 2021

Received in revised form 20 February 2022

Accepted 23 February 2022

Available online 26 February 2022

Editor: Fang Wang

Keywords:

Soil carbon sequestration

Soil health

Tillage systems

Eutrophication

Soil organic matter

ABSTRACT

Tillage is a common agricultural practice and a critical component of agricultural systems that is frequently employed worldwide in croplands to reduce climatic and soil restrictions while also sustaining various ecosystem services. Tillage can affect a variety of soil-mediated processes, e.g., soil carbon sequestration (SCS) or depletion, greenhouse gas (GHG) (CO₂, CH₄, and N₂O) emission, and water pollution. Several tillage practices are in vogue globally, and they exhibit varied impacts on these processes. Hence, there is a dire need to synthesize, collate and comprehensively present these interlinked phenomena to facilitate future researches. This study deals with the co-benefits and trade-offs produced by several tillage practices on SCS and related soil properties, GHG emissions, and water quality. We hypothesized that improved tillage practices could enable agriculture to contribute to SCS and mitigate GHG emissions and leaching of nutrients and pesticides. Based on our current understanding, we conclude that sustainable soil moisture level and soil temperature management is crucial under different tillage practices to offset leaching loss of soil stored nutrients/pesticides, GHG emissions and ensuring SCS. For instance, higher carbon dioxide (CO₂) and nitrous oxide (N₂O) emissions from conventional tillage (CT) and no-tillage (NT) could be attributed to the fluctuations in soil

* Corresponding authors.

E-mail addresses: hafiz.iqbal@tec.mx (H.M.N. Iqbal), r.parra@tec.mx (R. Parra-Saldívar).

Water pollution
Environmental health
GHG emissions

moisture and temperature regimes. In addition, NT may enhance nitrate (NO_3^-) leaching over CT because of improved soil structure, infiltration capacity, and greater water flux, however, suggesting that the eutrophication potential of NT is high. Our study indicates that the evaluation of the eutrophication potential of different tillage practices is still overlooked. Our study suggests that improving tillage practices in terms of mitigation of N_2O emission and preventing NO_3^- pollution may be sustainable if nitrification inhibitors are applied.

Contents

1.	Introduction	2
2.	Tillage practices and their impacts on soil carbon sequestration and related soil properties	3
2.1.	Tillage practices	3
2.2.	Impacts on soil carbon sequestration	4
2.3.	Impacts on soil structure	5
2.4.	Impacts on soil hydrological parameters.	5
3.	Impacts of tillage practices on soil erosion and water pollution	6
3.1.	Impacts on soil erosion	6
3.2.	Impacts on water pollution	7
3.2.1.	Tillage can translocate accumulated nutrients on soil surface to aquatic bodies	7
3.2.2.	Tillage can release soil stored pesticides/herbicides to aquatic bodies	7
4.	Tillage-induced synergies and trade-offs in greenhouse gas emissions	8
4.1.	Tillage induced GHG emissions	8
4.2.	Tillage induced methane emissions	8
4.3.	Tillage induced nitrous oxide emissions and mitigation	10
5.	Overall conclusions and perspectives	10
	CRedit authorship contribution statement	11
	Declaration of competing interest	11
	Acknowledgements	11
	References	11

1. Introduction

Current agricultural practices have enormously intensified by focusing on enhanced crop yield to feed the world's billions of people (Busari et al., 2015). Such agricultural intensification is putting additional pressure on natural resources, altering natural processes, and causing deterioration of soil, air, and water quality, by soil organic carbon (SOC) depletion, elevated greenhouse gases (GHG) emissions, soil nutrient imbalances, and fertilizer and pesticide export to aquatic bodies (Poelau et al., 2011; Vitousek et al., 2009; Subbarao et al., 2017; Sun et al., 2012). However, these impacts and their magnitude vary with site properties and farming practices (van Groenigen et al., 2011). Tillage is an agricultural management practice that affects soil, air, and water quality by affecting the exchange (sequestration of losses) of greenhouse gases to air and the loss of nutrients and pesticides to water. For example, tillage generally decreases soil carbon sequestration (SCS) due to the decomposition of elevated carbon (C). In addition, it may enhance/lessen carbon dioxide (CO_2) emissions (Ogle et al., 2005; Abdalla et al., 2016). Reduced or no-tillage is thus seen as a means to enhance soil carbon sequestration (SCS) and reduce CO_2 emissions. This effect may be counteracted by enhanced emissions of nitrous oxide (N_2O), as shown in several studies (Mei et al., 2018; Van Kessel et al., 2013), while tillage also affects methane (CH_4) emissions. Reduced tillage is also advocated as a means to reduce losses of accumulated pesticides and nutrients through leaching or surface-runoff (Fig. 1) (Luo et al., 2010; Endale et al., 2017; Nevins et al., 2020; Aziz et al., 2013; Zhou et al., 2017; Raiesi and Kabiri, 2016; Haddaway et al., 2017; Çelik et al., 2021; Shang et al., 2021; Elias et al., 2018). Numerous statements regarding the adverse impacts of tillage and thus the benefits of no-tillage or reduced tillage, however, overlook differences among tillage practices, soil properties, and associated land management. Different kinds of tillage practices are being practiced globally (Porwollik et al., 2019), for instance, conventional tillage (CT), no-tillage (NT), reduced tillage (RT) or minimum tillage (MT), ridge Tillage (RiT), strip-tillage (ST), strategic tillage (STR), and deep tillage (DT) (see Section 2.1 for background on these practices). The effects of these tillage types on soil-air-water quality may vary across regions and

are dependent on specific soil-climatic-management conditions (Chatterjee and Lal, 2009; Feng et al., 2020; Hussein et al., 2019; Powlson et al., 2014).

Below we summarize some impacts of soil-climatic-management conditions on (i) SCS and CO_2 exchange, (ii) N_2O and CH_4 emissions that may counteract the CO_2 emissions, and (iii) nutrient (N and P) and pesticide leaching and runoff.

Soil carbon sequestration (SCS) induced by agricultural practices is an important solution for decreasing the overall emissions to cause a net atmospheric C drawdown (Derpsch et al., 2014; Mehra et al., 2018; Zomer et al., 2017; Li et al., 2020). SCS potential of global agricultural soils accounts for $0.90\text{--}1.85 \text{ Pg C yr}^{-1}$ (Zomer et al., 2017), exemplifying the potential of agricultural soil management (Lal, 2015) to mitigate CO_2 emissions (Krupenikov et al., 2011). SCS and GHG emissions within agricultural settings are associated with adopted tillage methods. (Wulannintyas et al., 2021; Shakoor et al., 2021; Shang et al., 2021). Studies have shown that the effect of NT over CT on SCS might be confined to surface soils (0–10 cm) in developing countries (Bhattacharyya et al., 2021), which is also evident in Europe (Soane et al., 2012), China (Wang et al., 2021; Huang et al., 2016). Contrarily, NT may positively influence SCS in terms of whole soil profile compared to CT in Italy (Badagliacca et al., 2018), Mediterranean regions (Mazzoncini et al., 2016), and Japan (Wulannintyas et al., 2021). These studies imply that the potential of tillage practices on SCS fluctuates widely (Chatterjee and Lal, 2009). These above-mentioned variations produced by NT may arise discrepancies (Wang et al., 2021) and must be considered to have a clear idea about future endeavors. Moreover, soil organic matter (SOM), SOC and its fractions, soil aggregates, soil microbial community, soil hydrological attributes (e.g., water retention, saturated hydraulic conductivity, etc.), and their inter- and intra- relationships define effective and sustainable SCS (Bhattacharyya et al., 2022). Several studies showed that intensive tillage operations could augment changes in the chemical composition of SOM, also on the quantity and quality of soil microbial populations, and lead to a reduction in structural stability, thus reducing the SOC




Fig. 1. Conceptual schematic diagram illustrating tillage practices (NT and CT) impact soil-air-water quality. Note: SCS- soil carbon sequestration, POM- particulate organic matter.

storage capacity of the soil (Acir et al., 2020; Assunção et al., 2019; Sithole et al., 2019; Tuzzin de Moraes et al., 2016; Kladvko, 2001).

Tillage practices also influence N_2O emissions by affecting nitrification and denitrification processes (Li et al., 2015) and CH_4 emission by affecting methanotrophs. N_2O and CH_4 are greenhouse gases with a global warming potential of about 300 and 30 times larger than CO_2 . CH_4 and N_2O both can offset the positive benefits gained by SCS (Guenet et al., 2021; Hemes et al., 2018). Shakoor et al. (2021) observed a substantial amount of N_2O and CH_4 releases were associated with NT soils over CT globally. So, these tillage-induced trade-offs should be well-documented to fetch out future research directions.

Soil tillage affects SCS and GHG emissions and soil and water quality by erosion and runoff by affecting soil structure, reducing soil stability, and making it more prone to erosion. Melland et al. (2017) concluded that tillage, e.g., ST may increase the susceptibility of soils to erosion while resistance to erosion has been reported for CT/NT rotation (Zhang et al., 2018). In addition, although tillage may improve soil water storage capacity and saturated hydraulic conductivity (Schlüter et al., 2018; Zhang et al., 2018), tillage may also facilitate particulate nutrient loss (Melland et al., 2017), e.g., particulate organic C (POC) or particulate organic N (PON).

Several studies have revealed that tillage may also either increase (Elias et al., 2018), decrease (Klik and Rosner, 2020) or not affect (Okada et al., 2019) leaching or runoff of soil stored pesticides/fertilizers to aquatic bodies. When nitrogenous fertilizers are applied, soil immediately adsorbs and immobilizes NH_4^+ (Sørensen and Amato, 2002). In contrast, the negative charge of NO_3^- produced by nitrification, is not easily adsorbed and immobilized by soil, resulting in NO_3^- loss through runoff or leaching (Li et al., 2015), tillage practice may influence these processes. Bertol et al. (2017) revealed that runoff is more prominent under NT than CT. However, NT prevents erosional loss, while Daryanto et al. (2017b) stated that NT failed to control erosion-induced phosphorus (P) loss to the water. In light

of these findings, finding a way to resolve this detrimental phenomenon remain as an enormous challenge.

This study jointly presents the varied impact of tillage practices on soil quality, soil carbon sequestration (SCS), GHG emissions, leaching of C and N, and water pollution (Fig. 2). Previous reviews or meta-analyses considered only the effects of NT, CT, or DT on SCS or only their pathways (Bai et al., 2019; Haddaway et al., 2017; Bandyopadhyay, 2019) or only N_2O emissions (Huang et al., 2018; Feng et al., 2020) or only water pollution (Elias et al., 2018; Daryanto et al., 2017a). Optimizing tillage practices for agronomic and environmental benefits requires insights into the synergies and trade-offs of these practices. We hypothesized that improved tillage practices could enable agriculture to contribute to soil carbon sequestration (SCS) and mitigate GHG emissions and leaching of nutrients and pesticides. Our study provides in-depth global insights into the varied impacts of different tillage systems on soil carbon storage and air and water pollution.

2. Tillage practices and their impacts on soil carbon sequestration and related soil properties

2.1. Tillage practices

Conventional tillage (CT) is described as any tillage activity that utilizes cultivation such as ploughing and harrowing to prepare a seedbed, remove plant residues from the previous crops, or manage weeds. CT improves soil porosity, aeration, temperature and favors good root growth. No-tillage (NT)/zero tillage is the extreme form of conservation tillage practice where seeds are sown in a no-tilled soil without incorporating residues into soils from previous crops (Horowitz et al., 2012). Reduced tillage (RT) or minimum tillage (MT) is another conservation tillage technique that does not turn the soil over, thus avoiding changes in soil structure. In contrast, the soil surface is kept partly covered by 15% or 30% of crop




Fig. 2. Conceptualized schematic representation of the relationship between soil C and N pools, GHG emission, and leaching of C and N. Note: SOC- soil organic carbon, POC- particulate organic carbon, DOC- dissolved organic carbon.

residues from the previous crop. Compared to CT, NT/RT practices tend to increase soil moisture retention and improve water infiltration, but the soil remains prone to wind and water erosion. Ridge tillage (RiT) is a method of seedbed preparation where the topsoil is concentrated in a defined region to raise the seedbed above the natural topography. RiT is considered as a sustainable tillage practice in the black soils of China (Liu et al., 2018). Deep tillage (DT) involves mechanical modification of the soil profile such as subsoiling, deep ploughing, and mixing of soil profiles that can alleviate high subsoil compaction, and help deeper rooting of plants to reach water and nutrients from the subsoil (Tian et al., 2016; Scanlan and Davies, 2019; Schneider et al., 2017). This type of tillage may not be practical for sites without root-restricting soil layers and could even decrease the crop yields in that situation. It is only recommended for areas with a presence of root-restricting soil layers, especially in regions with continual aridity and erratic rainfall (Schneider et al., 2017). Strip-Tillage (ST) is a method of tillage where the soil is disturbed just in the seeding row while leaving the inter-row with residue cover, conserving soil moisture, decreasing penetration resistance, and creating an ideal seedbed for plant emergence and development (Licht and Al-Kaisi, 2005). Strategic tillage (STR) aims to achieve a balance between NT and CT, combining the benefits between both methods while minimizing the negative impacts of NT (Dang et al., 2015). The main objective of this kind of tillage is to manage herbicide-resistant weeds, diseases, and insects that have a below-ground pupal stage.

2.2. Impacts on soil carbon sequestration

Soil organic matter (SOM) is a mixture of materials, including organic particles, humus, coal, and living microbial biomass and the fine roots of plants (Stockmann et al., 2013). SOM improves soil chemical and biological characteristics, increases water and nutrient use efficiency, and promotes

improved agricultural output and sustainability (Obalum et al., 2017; Sarker et al., 2018a, 2018b). It contains three-folds more C than any other source (Schmidt et al., 2011). SOM usually comprises almost half or more than half of soil organic carbon (SOC) of its total composition (Pribyl, 2010). This indicates the dependent dynamism (changes in SOM can induce change in SOC) of SOM on SOC. The SOM stabilization is attributed to the humification process, which favors the selective preservation of more recalcitrant organic structures, the physical protection of soil aggregates, and chemical bonds through organo-mineral complexes (Bottinelli et al., 2017; Sarker et al., 2018a, 2018b; Six et al., 2002).

However, SOC is prone to changes due to different tillage practices. For instance, CT increases O₂ availability and microorganisms' access to soil aggregates, speeding up SOM decomposition, stimulating oxidation of SOC to form CO₂ (Sithole et al., 2019; Six et al., 2000). Especially, CT with harrowing and ploughing makes the SOM more exposed to microorganisms and increases mineralization, which may trigger higher CO₂ emissions from soils (Galdos et al., 2019; Marafon et al., 2020). In addition, in a maize cultivation system, mixing crop residues in deeper soil layers through CT may encourage higher microbial activity, enhancing CO₂ emissions compared to RT (Rutkowska et al., 2018), which corroborates with the findings of Franco-Luesma et al. (2020). They concluded that apart from microbial activities, higher soil temperature and around 60% water-filled pore space (WFPS) may contribute to the higher CO₂ emissions from CT soils. In NT systems, soils are not disturbed, kept covered by crop residues, reduce splash erosion of the topsoil so that SOM may remain proportionally higher in the upper soil than in deeper layers (Corbeels et al., 2016; Lal, 2019). A 23 years study in Brazil revealed that NT increases the particulate organic carbon (POC), the mineral-associated organic matter (MAOM), and humin contents. This enhances the efficacy of SCS in diverse organic matter fractions (Ferreira et al., 2020). NT systems can increase SOC by increasing the mean residence time and decreasing turn-over (Rahmati et al., 2020).

Additionally, in a meta-analysis performed by Haddaway et al. (2017), 351 studies were systematically reviewed and found significantly higher in NT relative to forms of CT in the 0–15 cm soil layer. But CO₂ emission may be higher in NT soils. For instance, Li et al. (2010) compared CT and NT in Chinese-directed seeded paddy fields and found NT substantially augmented CO₂ than CT. CO₂ fluxes might be related to soil temperature (Almaraz et al., 2009; Li et al., 2010). STR may increase CO₂ releases two-folds (Melland et al., 2017). Negassa et al. (2015) observed that RiT emits substantial CO₂ instead of chisel tillage in various topographic areas. The root causes identified were high soil moisture and soil temperature. In ST, fuel consumption and working speed of the tractor remain as the determining factor for GHG emissions. For instance, Šarauskius et al. (2017) reported low working speed and working depth augmentation from 0 to 200 mm produced the highest CO₂ from the system. A long-term experiment conducted in Belgium showed higher CO₂ production from RT than CT. The underlying reason might be the uniform incorporation of residues throughout the soil profile, which improved SOM. Hence, greater microbial respiration led to the substantial production of CO₂ (Lognoul et al., 2017). Almaraz et al. (2009) reported tillage-derived CO₂ fluxes as season-dependent in soybean fields. For instance, during summer CO₂ emissions were 160 g m⁻² d⁻¹ and 68 g m⁻² d⁻¹ under CT and NT, respectively, while N₂O emissions remained significantly higher over CT owing to residue retention in soils. The reasons behind higher CO₂ fluxes may be higher soil temperature (21–28 °C) and WFPS (44–60%) in the summer (Horák et al., 2020).

Table 1 shows some studies that compared the SOC stock or SOC content in different tillage systems in different soils and regions of the world. It illustrates that a comparison of no tillage with conventional tillage are the most common studied tillage systems and the intercomparison shows that SOC stocks or contents are always higher in the no tillage (and also strip and ridge tillage) systems as compared to conventional tillage but the impacts of tillage on SCS is affected by soil-climatic conditions (Abdalla et al., 2013).

2.3. Impacts on soil structure

Soil tillage can have multiple impacts on soil structure (Table 2), and improved soil structure is vital for greater SCS (Guo et al., 2020). Good soil structure ensures preserving SOC for a longer time and prevents being respired as CO₂ in the atmosphere (Ewing et al., 2006). Studies have shown that tillage systems exhibit substantial influence on SOC dynamics, pools, stocks, and in SCS (Alvarez and Steinbach, 2009; He et al., 2011; Van Den Putte et al., 2010; Ramnarine et al., 2018; Jha et al., 2020; Awale et al., 2017; Mehra et al., 2018). Two major pools that regulate SOC dynamics are light fractions (LF) and heavy fractions (HF) (Christensen, 2020; Cerli et al., 2012). The constituents of HF include water-soluble and non-water-soluble elements. A long-term study conducted by Ramnarine et al. (2018) revealed NT exerted a substantial impact on HF, especially on the non-water-soluble fractions and sequestered C, which accounted for 28.9 Mg C ha⁻¹. The sequestered C was confined to 0–10 cm soil depth over CT with increasing soil depth. Thus, an insignificant difference exists between NT and CT in these components. Pinheiro et al. (2015) observed SOC stocking in HF around 12.3 g kg⁻¹ soil under NT. Again, these fractions may vary with soil types. Azam et al. (2020) found HF maintains a strong relationship with TOC while LF does not. Therefore, HF regulates SOC dynamics, and undoubtedly, the SCS potential of HF is high. Again, a correlation can be drawn among SOC, soil aggregation, and tillage operations (Guo et al., 2020). Aggregate stability and macroaggregates formation are the prerequisites for effective SCS; aggregate collapse causes SOC loss. Theoretically, C is firstly captured by macroaggregates and then transferred to microaggregates, hence, macroaggregates enrichment is another important key to obtaining abundant SCS. High SOC content is often found in soils with more stable aggregates (Parvin et al., 2021). The stability of soil aggregates increases with microbial activity, which helps create glues such as glomalin, allowing this aggregation.

Physical soil aggregates are formed when the soil matrix is disrupted through wetting and drying cycles (Rodríguez et al., 2021). Because soil aggregates contain nearly 90% of SOC, they are vital components of SOC protection. For instance, Kan et al. (2020) found that NT increases the proportions of macro-aggregates. Hati et al. (2019) reported better aggregate stabilization through enhanced activities of soil biota (more C enriched macroaggregates) in topsoil under conservative tillage in vertisols of India, although SOC and nutrient stratification was evident. Kravchenko et al. (2011) reported long-term CT affects soil porosity and aggregate formation in alfisols. Aggregates, in general, are vulnerable to tillage; disruption of macro-aggregates causes transformations of soil structure to microaggregates. Macro-aggregates are formed by adhesives that result from activities linked to microbial activity (Kan et al., 2020). Resins and glues produced by microbes in the soil help bind soil particles together to form water-stable aggregates. They function as the main reserves of SOC and could be a useful parameter to monitor SOC. Pinheiro et al. (2015) acknowledged the significant role of sand particles in SCS compared to silt and clay but affirmed these fractions mainly protect and store the sequestered C in soil. Again, tillage operations have a significant effect on soil particles. NT with residue retention increased silt plus clay associated C by 45% and 74% at 0–20 and 20–40 cm depth, respectively over CT (Kumar and Nath, 2019). Decreasing mechanical cultivation allows SOC to be protected by reducing macro- and micro-aggregates turnover. This helps minimize exposure of more stable SOC to microbial activities (Brewer and Gaudin, 2020). Overall, soil aggregates can shield SOC substantially from microbial spells (Schmidt et al., 2011). These studies suggest that microaggregates are another important regulator of SOC dynamics and can be used as a sustainable indicator to identify and calculate the SCS rate.

Apart from these, transition from CT to RT may lead to the formation of soil firmness, but remain unpaid, lacking higher actions by soil microorganisms (Schlüter et al., 2018). Moldboard plow appeared as the best option for conserving soil structure (0–20 cm depth) in a long-term experiment in Danish Sandy loam soils compared to RT. Residue retention can compensate for the adverse effects created by RT (Abdollahi and Munkholm, 2017). DT is utilized in arid regions with fine or medium-textured soil with high bulk density and soil compaction, while NT/DT rotation tillage benefits SOC sequestration levels (Feng et al., 2020). In a semi-arid area on the Huang-Huai-Hai Plain of China, DT reduced soil bulk density at 0–40 cm layer, favoring root growth and increased grain filling rate and maize grain yield (Zhai et al., 2017). With improved soybean yields, the contribution of DT was substantial in a dry silt loam soil having low SOM in Arkansas, USA compared to CT due to the breakage of the compact layer at 7 to 15 cm in the soil sub-surface, which makes the soil more favorable for plant growth (Henry et al., 2018). and this practice has site-specific positive and negative effects on grain yield. Additionally, it is very dependent on environmental conditions and agronomic management factors (Feng et al., 2020).

2.4. Impacts on soil hydrological parameters

Tillage practices may influence soil hydrological parameters largely. Recent researches into the impact of NT activities found that, as opposed to other tillage schemes, NT may not facilitate water retention (Blanco-Canqui et al., 2017; Deuschle et al., 2019), while other authors exposed increased hydraulic parameters (e.g., greater saturated hydraulic conductivity) and water availability with improved infiltration in NT systems (Acharya et al., 2019; de Almeida et al., 2018). Again, Bertolino et al. (2010) revealed soils under CT remained saturated for a longer time after each shower event. In the case of torrents, it encouraged interflows while the NT system continued to drain. This phenomenon was identified as the determining factor for increased erosion rate under CT systems compared to NT.

RiT tends to reduce evapotranspiration and increase soil moisture in the upper soil profile, water availability, water use efficiency, and grain yield than CT, especially in the dry season (low rainfall and N) when the water is a limiting factor for crops (Li et al., 2018). Cross-ridge tillage can be

Table 1
Impact of different tillage systems on SOC storage under different soil types.

Reference (s)	Type of Tillage Practice(s)	Cropping systems	SOC/TOC (g kg ⁻¹)	Total SOC Stock (ton ha ⁻¹)	Soil classification	Region/Country	Observation(s)
Nath et al., 2017	No-Tillage	Wheat- mungbean (<i>Vigna radiata</i> (L.) Wilczek)	–	12.5	Inceptisol, mahauli	New Delhi, India	No tillage with residue retention and nitrogen management increased SOC stock.
Dimassi et al., 2014	Conventional Tillage	1st period: maize/winter wheat rotation; 2nd period: crop residues were removed; 3rd period: winter wheat/barley/sugarbeet/pea; 4th period: catch crop (oats/vetch) after wheat crop	–	11.6	Haplic Luvisol	Northern France	No difference between the tillage systems from the layer of 27.9 cm after 40 years with different crop systems.
	No-Tillage		8.25	45.1			
Rahmati et al., 2020	Shallow Tillage	Rotation of vetch and wheat (one crop per year)	8.5	44.5	Typic calcixerepts (clay loam texture)	Maragheh, East Azerbaijan Province, Iran	Conservation tillage practices can increase SOC by increasing the mean residence time and decreasing turn-over.
	Full inversion Tillage		11.0	44.2			
Jha et al., 2020	No-Tillage	Wheat	9.2	–	Vertisol or Ustic Pellustert	Queensland, Australia	The greatest increases in SOC occurred with no tillage at higher rates of N fertilization.
	Conventional Tillage		7.3	–			
Kumar and Nath, 2019	No-Tillage	Wheat/chickpea (winter) and mungbean (summer) after puddled transplanted rice in the rainy season	–	19.9	Inceptisol (Typic Ustochrept, sandy loam texture)	Uttar Pradesh, India	No tillage significantly differs from conventional tillage in the 0–20 cm layer.
	Conventional Tillage		5.2	–			
Pinheiro et al., 2015	No-Tillage	Rotation of vegetables: tomato, green pepper and beans	3.7	–	Dystrophic Red Latosol (Typic Haplortox)	Southeast Brazil	In tropical soils, no tillage is the main strategy for SOM accumulation, even when low input of vegetables crop residues occurs.
	Conventional Tillage		15.2	17.6			
Ramnarine et al., 2018	No-Tillage	Corn-soybean-wheat rotation	10.8	12.3	Typic Hapludalf (silt loam)	Southern Ontario, Canada	Significant differences observed in 0–10 cm layer, but no significant differences of TOC in 10–30 cm for both tillage treatments.
	Conventional Tillage		24.1	–			
Liu et al., 2018	Conventional Tillage	Maize-soybean rotation	21.0	–	Black soil/ Hapudoll (clay loam texture)	Northeast China	Ridge tillage generated a significantly higher SOC than conventional tillage in the 0–5 cm.
	Conventional Tillage		17.2	–			
Alcántara et al., 2016	Ridge-Tillage	Crop rotation: mayze, wheat, oilseed rape, potato,	26.10	–	Cambisols and Podzols	Northern and central Germany	Deep tillage has a high potential for increasing SCS through the incorporation of SOC-rich topsoil into the subsoil.
	No-Tillage		25.0	–			
Jaskulska and Jaskulski, 2020	Deep Tillage	Winter wheat and barley	–	142	Cambisol	Central and Eastern Europe	The TOC in the 0–20 cm soil layer increased more than 1 g kg ⁻¹ after 6 years of ST.
	Strip-Tillage		11.0	–			
Endale et al., 2017	Conventional Tillage	Cotton and peanut	9.9	–	Ultisols	Southeastern USA	TOC storage was greater for strip tillage than CT, due to reduced erosion and decreased runoff.
	Strip-Tillage		86.00	–			
	Conventional Tillage		41.6	–			

used to replace downslope tillage on hilly regions to intercept the surface runoff volume, increase water infiltration and reduce loss of topsoil. This practice greatly reduces the sediment and N and P losses during heavy rainfall events in hilly regions from China. This is accomplished by shortening the slope length and retaining runoff (Guo et al., 2019). In humid Ethiopian Highlands, DT was found to break up the dense soil hardpan, thus, increasing infiltration rates and soil moisture throughout the soil profile, reducing the bulk density, penetration resistance, and soil loss, having high potential to improve degraded soils in that region. The objective of DT is to improve soil functions including water infiltration capacity and root penetration, optimizing conditions for crop growth (Hussein et al., 2019; Alcántara et al., 2016).

3. Impacts of tillage practices on soil erosion and water pollution

3.1. Impacts on soil erosion

SOC is a primary indicator of soil health (Krupenikov et al., 2011; Manlay et al., 2007), and highly sensitive to erosion losses as it remains concentrated in the proximity of soil surface. Soil hydrology, SCS, soil water content, and soil erosion are interlinked with each other, and tillage practices exhibit a significant impact on them (Bertolino et al., 2010; Moreno-Casasola et al., 2017). For instance, in China's calcareous soils, Ye et al. (2020) calculated that tillage-induced SOC loss might account for avg. 77% of total SOC loss during the initial tillage

Table 2

Role of some selected tillage practices in improving soil structure.

Tillage practices	Role (s)
No Tillage (NT)	<ol style="list-style-type: none"> 1. The inclusion of plant residues and lack of disturbance to soil encourage NT to increase SOM, aggregate-induced SOC, stable aggregates, increased micropores, and smart connectivity, improving soil structural stability and resistance against rainfall events near the soil surface (Wang et al., 2019; Guo et al., 2020; Chenu et al., 2019). 2. NT may induce greater bulk density and bearing capacity of soil compared to CT and RT and alter water pathways beneficial for soil microorganisms (Soane et al., 2012).
Reduced Tillage (RT)	<ol style="list-style-type: none"> 1. RT encourages lower bulk density, increased soil porosity, and air space, higher saturated hydraulic conductivity, higher macroporosity, and pore connectivity (Schlüter et al., 2018). 2. Contrarily, lack of soil disturbance may lead to decreased air pore space, increased bulk density, and penetration resistance in the topsoil (Abdollahi and Munkholm, 2017).
Deep Tillage (DT)	<ol style="list-style-type: none"> 1. DT reduces higher bulk density and soil compaction, improves SOM and favors good root growth (Feng et al., 2020; Zhai et al., 2017; Henry et al., 2018), but might not be a good option for the soils with poor structure (Alcántara et al., 2016).
Strip Tillage (ST)	<ol style="list-style-type: none"> 1. ST creates mechanical resistance, providing a good soil structure that is capable of withstanding mechanical stress (Pöhlitz et al., 2018). 2. It may reduce nutrient mineralization in soils (Laufer and Koch, 2017).
Strategic tillage (STR)	<ol style="list-style-type: none"> 1. STR lowers bulk density, improves aggregate stability and penetration resistance induced by NT (Çelik et al., 2019).

period. A long-term comparative study of CT, NT, and RT revealed that depending on soil texture, annual erosion rates from CT, NT, and RT might be average 20.9, 2.45, and 4.45 t ha⁻¹, respectively (Klik and Rosner, 2020). SOC subjected to splash erosion may undergo the erosion-sedimentation process. Thus, SCS may be reduced by 0.6–1.5 Gt yr⁻¹ (Lal, 2005). Klik and Rosner (2020) observed different potentials of runoff of N and P from CT, NT, and RT practices. They quantified total N losses ranged from 13.3–48.1, 1.6–9.4, and 4.5–18.7 Kg ha⁻¹, respectively, from CT, NT, and RT while total P losses were 6.7–29.4, 0.7–2.4 and 2.1–3.7 Kg ha⁻¹, respectively. Chowaniak et al. (2020) observed that NT soils might encourage 4% higher runoff over CT, whereas NT soils may have 67% lower soil loss. In addition, NT can reduce total organic C (TOC) loss and organic C loss through sediments by 46% and 53%, respectively, over CT. NT may reduce runoff volume and sediment losses by 50% and 50%–95%, respectively, in Fluvi-Calcaric Cambisol soils (Carretta et al., 2021). Over CT, RT and NT may decrease soil loss from silty clay loam soil by 38 and 65%, and from silt loam soil values remain 70–88% and 84–93%, respectively (Klik and Rosner, 2020). Olson et al. (2013) and colleagues reported substantial annual losses of C under different tillage practices in the USA during 24 years. They revealed that soils under NT lost less amounts (1.2 Mgha⁻¹) of C compared to chisel plow (9.9 Mgha⁻¹) and moldboard plow (8.2 Mg ha⁻¹). Novara et al. (2019) estimated that 9.5 ± 1.2 Mg C ha⁻¹ may be lost annually due to tillage from a vineyard. Bogunovic et al. (2018) recorded typical yearly soil loss under NT (0.53 t ha⁻¹), DT (3.11 t ha⁻¹), and CT (13.11 t ha⁻¹) in Croatia. These losses might occur due to soil erosion plus tillage-induced erosion and notably, responsible for decreased SCS (Lal, 2005) and water pollution by sedimentation-erosion processes. Again, water erosion may cause the formation of labile C, increased mineralization-immobilization processes accompanied by elevated CH₄ and N₂, consequently water pollution and air pollution may occur simultaneously (Issaka and Ashraf, 2017). Intense tillage (e.g., CT) induced erosion may increase soil erosion (Lal, 2005; Chalise et al., 2019). These studies suggest that tillage practices may facilitate soil erosion and decrease SCS through SOC losses. Moreover, nutrient (N and P) loss through leaching surface/sub-surface runoff may cause water pollution or water quality degradation by causing eutrophication or hypertrophication.

3.2. Impacts on water pollution

3.2.1. Tillage can translocate accumulated nutrients on soil surface to aquatic bodies

Farming intensification has caused greater agricultural input materials, e.g., N and P fertilizers (Lu and Tian, 2017) and pesticides (Kremer, 2018). Unwarranted use of these materials may lead to net anthropogenic imbalances in soil systems, influencing leaching loss or surface runoff of nutrients/contaminants (Fowler et al., 2013). Tillage disturbs the stored soil contaminants/nutrients, causing them to be lost from the system and into surrounding bodies of water. There is a strong link between tillage practices and water contamination (Chalise et al., 2019). Nitrate (NO₃⁻), for example, is primarily lost by leaching and is one of the most common causes of eutrophic or hypertrophic conditions (Lu and Tian, 2017; Reza et al., 2020; Hanrahan et al., 2019; Spiess et al., 2020; Syswerda et al., 2012). Studies have shown that different tillage practices can control the leachate volume and concentration of NO₃⁻ (Khan et al., 2017a, 2017b; Daryanto et al., 2017b). Additionally, Endale et al. (2019) showed higher sediment losses and nutrient runoff in ultisols under NT compared to CT. Residue retention and soil compaction may enhance runoff volume, increased macropores may contribute to leachate volume, and SOM may contribute to runoff and leachate concentration (Daryanto et al., 2017a, 2017b). These studies indicate that tillage systems can influence nutrient loss trails and contribute substantially to the deterioration of water quality and eutrophication.

Khan et al. (2017a, 2017b) demonstrated that soils under CT, RT and DT tend to show varying capacities regarding leaching loss of NO₃⁻. For example, RT can significantly reduce NO₃⁻ concentration at 0–25 cm soil depth compared to CT and DT, but increased NO₃⁻ concentration 17.34% observed at 25–50 cm depth. That indicates CT and DT's potential to surface water contamination through NO₃⁻ elimination from the soil system, while the deeper movement of NO₃⁻ under NT may prevent ground water contamination. Spiess et al. (2020) revealed that NO₃⁻ leaching rates are regulated by the quantity of soil NO₃⁻ and they quantified leaching loss of 50 kg N ha⁻¹ yr⁻¹ under NT and mouldboard plow practices in a diverse crop rotation system.

Further, Hess et al. (2020) found that NT can create a buffer against leaching loss of N. Daryanto et al. (2017b) showed in a meta-analysis that although leachate NO₃⁻ load and runoff NO₃⁻ concentration from NT was determined by water flux, values were higher than CT. RT has a greater leaching potential of NO₃⁻ in sandy soils, especially after rainfall (Waddell and Weil, 2006). Contrarily, RT reduced surface runoff and leaching loss of N in paddy soils compared to CT (Qin et al., 2020). Ridge coupled with NT can slow down NH₄⁺ oxidation, thereby reducing the leaching loss (Li et al., 2015). Under simulated rainfall conditions in wheat-soybean rotation, Daverede et al. (2003) found the concentration of PO₄²⁻ leachate was 0.40 and 0.24 mgL⁻¹ under NT and chisel plow tillage systems. To find out why such a higher rate of leaching loss, Williams et al. (2018) observed that fertilization has a great impact under NT soils to induce higher leaching. They concluded that instead of broadcasting, injection or fertilizer placement through tillage could reduce the volume associated with water quality degradation. However, recent studies investigating the prevention of leaching and runoff from soils revealed that applying animal manures coupled with nitrapyrim could reduce both N and P leaching (Luo et al., 2017; Giacometti et al., 2020).

3.2.2. Tillage can release soil stored pesticides/herbicides to aquatic bodies

Several studies have shown that applied pesticides can cause malfunction of soil biota and decay of soil structure (Kremer, 2018; Yang et al., 2018), even accumulation of heavy metals in cultivated soils (Soane et al., 2012), and they are major agents of eco-degradation. Soils act as a reservoir for contaminants safeguarding them from leaching and entry to the aquatic systems, most importantly, translocation by plants (Düring et al., 2002). Many studies identified the persistent existence of several heavy metals in topsoils of rice (Abuzaid and Bassouny, 2020; Ali et al., 2018), maize, soybean, chihua squash (Dzul-Caamal et al., 2020), etc., even, sub-surface soils may also remain subjected to extreme, high and moderate levels of contamination

(Abuzaid and Bassouny, 2020). Additionally, several studies explicitly showed that SOM plays a leading role in the adsorption, binding, and storing of applied pesticides in soil (Stevenson, 1972; Hayes, 1970; Hance, 1974; Payá-Pérez et al., 1992). Under such context, it's critical to figure out how different tillage techniques affect surface/subsurface runoff processes, including leaching by altering SOM or other soil qualities, or how they affect pollutant transport to aquatic bodies. Mottes et al. (2020) reported that tillage might induce leaching loss of pesticides through i) SOM dilapidation, ii) altered water pathways within soil systems, and iii) amorphous phyllosilicate (e.g., allophane clay) corrosion. In a 13-year field trial, Babal et al. (2021) revealed that pendimethalin herbicide applied in an alluvial sandy loam soil under NT may be filtrated down to the subsoil compared to CT, although NT increased 69% higher SOC stock in 0–5 cm soil while the leaching rate was observed almost same between CT and NT. A study did not find any effect of NT on surface runoff of herbicides in a soybean-corn rotation. At the same time, CT impacted the loss of herbicide due to the wash off of the intercepted herbicides on the plant residue (Gonzalez, 2018). Contrarily, NT may increase glyphosate herbicide loss by surface runoff compared to CT, due to increased residue cover, reducing sorption (Shipitalo and Owens, 2011). Glyphosate may exhibit significant residual levels in soils under NT, which may adversely impact soil microbiome (da Silva et al., 2021). The herbicides atrazine and metolachlor are also lost in higher quantity through surface runoff from NT compared to mulch tillage (Ghidey et al., 2010). Residual herbicides bound to mineral and organic fractions in the soil can be leached out through surface runoff, producing water pollution and adversely impacting human health (Alletto et al., 2010). Compared to CT, a meta-analysis indicated that NT might have a modest effect in lowering pesticide load in the runoff. Still, it postulated that soil properties, such as acid soil pH, could modify pesticides' physicochemical properties (polarity, pKa) and affect their mobility to water bodies (Elias et al., 2018). In moderately fine and medium-textured soils with less than 2.3% SOM, NT may enhance pesticide concentration and loads in runoff (Elias et al., 2018). Potter et al. (2015) quantified 2.5 folds greater sub-surface loss of pesticide fluometuron under ST, although ST may reduce surface runoff of this pesticide by 10 times compared to CT. In addition, surface runoff volume under CT may be 2 times higher over ST.

Kaur et al. (2016) showed clay particles facilitate adsorption and the SOM handles the desorption of pesticides in soils as observed for the fate of pretilachlor in Punjab soils. The releasing capacity of different soils is variable. For instance, loamy sand soils favor more leaching loss of pesticides (Kaur et al., 2016). Apart from these, the role of soil aggregates cannot be overlooked. Soil aggregates play a pivotal role in capturing and accessibility of the accumulated heavy metal in cultivated soils (Huang et al., 2017). Huang et al. (2017) observed reduced action of metals in the greater soil depth, thus indicating the importance of maintaining CT or other tillage methods except NT because CT or other tillage practices tend to invert the soil profile. Similarly, observations of long-term tiling impact on Eutric Cambisol and Luvisol revealed soils under NT accumulated more pollutants compared to CT (Düring et al., 2002).

These studies indicate that waterbodies are at significant risk of pollution by tillage practices worldwide. NT although facilitates condensation of pollutants in surface soils, organic amendments can help to a great extent to reduce the leaching loss of pollutants from the surface soil. For example, chlordecone is considered a major soil contaminant and threat to human and crop health (Clostre et al., 2014; Mottes et al., 2020). Organic amendments can act as a potential sequestration tool in this context. Clostre et al. (2014) observed seven times decrease in leaching, and only 5% organic usage led to a substantial reduction of contaminant uptake by crops.

4. Tillage-induced synergies and trade-offs in greenhouse gas emissions

4.1. Tillage induced GHG emissions

Regulation of greenhouse gases (GHG) in the atmosphere is an important soil ecosystem service that is affected by tillage practices (Table 3)

(Lal, 2019; Prado et al., 2016). Quantifying CO₂/CH₄ output proportion is crucial because the net benefit from SCS could not be achieved due to the excess releases (Yu et al., 2008). But, contrasting research results make tillage's role in GHG emissions questionable (Table 4). For instance, decreases in GHG emissions in NT systems compared to CT were observed (Li et al., 2013; de Figueiredo et al., 2018; Fanguiero et al., 2017; Lu et al., 2016), while some reported no significant difference or increase on the GHG (Oorts et al., 2007; Plaza-Bonilla et al., 2014; Zhang et al., 2016a, 2016b). A global meta-analysis conducted by Shakoor et al. (2021) and colleagues revealed that NT augmented global CO₂, N₂O, and CH₄ emissions by 7.1%, 12%, 20.8%, respectively, compared to CT. More specifically, a substantial increase in N₂O and CH₄ discharges occurred globally in rainfed and irrigated agricultural areas. Another rigorous meta-analysis of 740 paired observations showed that NT has a considerable potential to minimize GHG and the global warming potential (GWP); however, net benefits require further understanding of the agroecological environment (Huang et al., 2018). Additionally, life-cycle assessment of tillage practices revealed that the impact order of GHG production is CT > NT > RT (Sørensen et al., 2014). The underlying cause was the higher mineralization rate in soils. NT with legumes as cover crops can boost N₂O emission, but the SCS rate and SOC deposition rate can dominantly mitigate the CO₂ amount of N₂O, and NT soils act as a potential sink for GHG (Bayer et al., 2016). NT may reduce GHG indirectly via reduced fossil fuel usage and reduced demand for synthetic fertilizers (Haddaway et al., 2017). A recent study in China suggests the combination of RiT and NT may act as a potential inhibitor of GHG compared to CT and NT (Li et al., 2021a, 2021b).

4.2. Tillage induced methane emissions

Studies have shown that soils can act as a net source or sink of CH₄ depending on soil temperature, moisture, N level, soil microbial community, air-filled porosity, spatial heterogeneity of soil profile, and so on (Gregorich et al., 2005; Bréchet et al., 2021; Díaz et al., 2018; Feng et al., 2021). Overall, drained agricultural soils are a small CH₄ sink, while CH₄ emissions occur in irrigated paddy-rice systems. For example, Shang et al. (2021) evaluated CT and RT in a Chinese double paddy cropping system. They found that CT and RT were responsible for a huge volume of CH₄, yet, CH₄ production was related to the SCS potential of the soils.

In drained agricultural soils, tillage can alter the mechanisms that govern the release or oxidation of CH₄ through its impact on the soil water regime (Feng et al., 2021). Moisture generally substantially influences CH₄ uptake rates (Liu et al., 2019). Hence, structural deterioration, more specifically compaction prevalent in extensively tilled soils, can negatively influence CH₄ consumption (Ball et al., 1999). For example, it was found that CH₄ uptake was less under STR over NT due to the decreased soil bulk density (Melland et al., 2017). In addition, enhanced residue retention associated with NT may also improve CH₄ consumption, as observed in a maize monoculture system in which NT combined with residue retention was around 3 and 1.5 times higher than CT and NT only, respectively (Franco-Luesma et al., 2020). It has been found NT and RT, reducing soil compaction, may increase two-fold CH₄ uptake compared to CT due to the greater soil shear strength, pore connectivity, and niche availability. These properties may also improve the functioning of soil methanotrophic population, which may increase CH₄ oxidation resulting in lower CH₄ emissions (Alskaf et al., 2021). In irrigated and/or flooded soils, the impact of tillage practices differs from drained soils. Franco-Luesma et al. (2020) found that shortly after water application in flooded irrigation, no significant changes were observed in CH₄ fluxes in a maize monocropping system under NT and CT practices. Contrarily, Kim et al. (2016) observed that continuous NT in rice monocropping enhanced soil physical properties and reduced CH₄ emission during the early years of NT establishment but increased CH₄ emission over time. The cause is most likely due to increased labile SOC boosting methanogenic abundance. Overall, however, NT combined with residue retention on the soil surface likely lowers CH₄ emission from rice under different tillage practices. For instance, Zhang et al. (2013) showed that residue retention combined with NT enhanced soil porosity

Table 3

Overview of different tillage techniques on soil carbon sequestration and GHG emissions.

Source	Scale	Type of tillage practices	Duration	Impact on GHG emissions	Impact on SOC stock	Impact on SOC quantity (whole soil profile)	Impact order of tillage practices for SCS
Shakoor et al. (2021)	Global	NT and CT	-	NT = ++ CT = -	NT = ++ CT = +	NT = + CT = +	CT > NT
Wang et al. (2021)	Regional	NTS, RTS, CTS, CT	14 years	NTS = n.e. RTS = n.e. CTS = n.e. CT = n.e.	NTS = ++ RTS = ++ CTS = + CT = +	NTS = ++ RTS = ++ CTS = ++ CT = +	NTS ≈ RTS > CTS > CT
Badagliacca et al. (2018)	Regional	NT, CT	23 years	NT = + CT = -	NT = + CT = -	NT = - CT = +	NT > CT
Huang et al. (2016)	Regional	NT, CT	10 years	NT = n.e. CT = n.e.	NT = - CT = +	NT = - CT = +	CT > NT
Chatterjee and Lal (2009)	Regional	NT, PT	>4 years	NT = n.e. PT = n.e.	NT = ++ PT = +	NT = ++ PT = +	NT > PT
Chen et al. (2021)	Regional	NTS, RTS and CTS	09 years	NTS = + RTS = - CTS = -	NTS = + RTS = + CTS = +	NTS = + RTS = + CTS = -	NTS > RTS > CTS
Li et al. (2021a, 2021b)	Global	NT, RT, CT	44 years	NT = n.e. CT = n.e. RT = n.e.	NT = ++ RT = + CT = -	NT = + RT = + CT = -	NT > RT > CT
Olson et al. (2013)	Regional	NT, CP, MP	24 years	NT = n.e. CP = n.e. MP = n.e.	NT = - CP = - MP = -	NT = - CP = - MP = -	NT > MP > CP
Wang et al. (2019)	Regional	NTS, RTS, CTS, CTO	9 years	NTS = n.e. RTS = n.e. CTS = n.e. CTO = n.e.	NTS = + RTS = + CTS = + CTO = -	NTS = + RTS = + CTS = + CTO = -	NTS > RTS > CTS > CTO
Modak et al. (2020)	Regional	NT, CT	9 years	NT = n.e. CT = n.e.	NT = + CT = -	NT = - CT = -	NT > CT
Du et al. (2017)	Regional	NT, CT	≥ 3 years	NT = n.e. CT = n.e.	NT = + CT = -	NT = + CT = -	NT > CT
Lembaid et al. (2021)	Regional	NT, CT	9 years	NT = n.e. CT = n.e.	NT = + CT = -	NT = ++ CT = +	NT > CT
Wulanningtyas et al. (2021)	Regional	NT, MP, R	3 years	NT = n.e.; MP = n.e.; R = n.e.	NT = + MP = - R = -	NT = + MP = - R = -	NT > MP > R
Gwenzi et al. (2009)	Regional	NT, RT, CT	6 years	NTS = n.e. RTR = n.e. CTR = n.e.	NTS = + RTR = + CTR = -	NTS = + RTR = + CTR = -	RTR > NTS > CTR
Yu et al. (2020)	Regional	NT, CT, RT	14 years	NT = + CT = - RT = +	NT = - CT = + RT = +	NT = - CT = + RT = +	RT > CT > NT
Mazzoncini et al. (2016)	Regional	NT, CT	28 years	NT = n.e. CT = n.e.	NT = + CT = -	NT = + CT = -	NT > CT

Indicators and Abbreviations: 0 = Neutral; + = High; ++ = Extremely high; - = Low; -- = Extremely low; CT = Conventional Tillage; CTR = Conventional Tillage with residue retention; NT = No-tillage/Zero tillage; RT = Reduced/ Minimum Tillage; ST = Subsoiling tillage; SCS = Soil Carbon sequestration, R = Rotary tillage; NTS = No tillage with residue retention; RTS = Rotary Tillage with residue retention; RTR = Reduced Tillage with residue retention; CTS = Plow tillage with residue retention; CTO = Plow tillage with no residue retention; PT = Plow Tillage; CP = Chisel plow; MP = Moldboard plow; ASRT = All straw return tillage; SRT = Shallow rotary treatment; n.e. = Not evaluated.

and raised the oxidation layer. Hence, CH₄ emission appeared to be lower from a Chinese paddy field. Similarly, NT with residue retention increased CH₄ consumption around 3 and 1.5 times higher compared to CT and NT,

respectively (Franco-Luesma et al., 2020). In a bioenergy cropping system, rotary tillage increased CH₄ emission compared to NT if combined with soil ameliorant or residue retention (Liu et al., 2015).

Table 4

Impacts of different forms of tillage practices on the direction (+ is increase, 0 is no effect and - is decrease) of GHG emissions from agricultural lands.

Tillage practice(s)	Type of study	Scale	Studied crop(s)/cropping systems	Soil type(s)	CO ₂	N ₂ O	CH ₄	Reference(s)
STR compared to NT	Primary research	Regional	Controlled-traffic farming regimes	Vertosol, and Sodosol	+	-	-	Melland et al. (2017)
NT compared to CT	Meta-analysis	Global	-	-	+	+	+	Shakoor et al. (2021)
NT compared to CT	Meta-analysis	Global	-	-	-	+	-	Huang et al. (2018)
RT compared to CT	Primary research	Regional	Maize	Cutanic Luvisol	+	+	n.e.	Lognoul et al. (2017)
CT compared to NT	Primary research	Regional	Rice	Hydromorphic paddy soil (silty clay loam)	0	+	+	Ahmad et al. (2009)
RiT compared to CT	Primary research	Regional	Rice	Haplic lixisol	0	0	0	Nyamadzawo et al. (2013)
DT compared to CA	Primary research	Regional	Maize and wheat rotation	Haplic Phaeozem	0	0	0	Dendooven et al. (2012)
ST compared to NT and CT	Primary research	Regional	Corn-soybean rotation	Typic Haplaquolls and Typic Hapludolls	-	n.e.	n.e.	Al-Kaisi and Yin (2005)

Abbreviations: NT = no tillage, CT = conventional tillage; RT = reduced tillage, RiT = ridge tillage, DT = deep tillage, ST = strip tillage, CA = conservation agriculture.

4.3. Tillage induced nitrous oxide emissions and mitigation

Nitrous oxide (N_2O) emissions from agricultural lands have become a major concern since i) the global warming potential of N_2O is around 300 times higher than CO_2 thus, playing a direct role in global climate change, and (ii) it is currently the primary driver of stratospheric ozone layer depletion (Ravishankara et al., 2009; Li et al., 2014). Li et al. (2005) acknowledged that 75–100% obvious benefit from SCS could be balanced from the upsurge in N_2O concentrations (Mei et al., 2018; Guenet et al., 2021). Nitrification and denitrification predominantly are the main routes of N_2O releases from cultivated fields (Braker and Conrad, 2011). Nitrification is the aerobic biological oxidation of ammonium (NH_4^+) to form nitrite (NO_2^-), nitrate (NO_3^-) as the main products, and N_2O as a byproduct accomplished by the activities of auto and heterotrophic bacteria. Denitrification is the anaerobic reduction of NO_3^- to NO_2^- , NO, N_2O , and finally N_2 (Butterbach-Bahl et al., 2013; Braker and Conrad, 2011; Lognoul et al., 2017). Both of these processes are dependent on soil microbial metabolic activities (Butterbach-Bahl et al., 2013; Congreves et al., 2017).

Tillage systems exhibit a role in N_2O emission either by favoring or reducing nitrification and/or denitrification and thus, soil N_2O emissions due to impacts on several factors, including soil texture, soil temperature, water-filled pore space (WFPS), soil mineral nitrogen ($\text{NH}_4^+\text{-N}$, NO_3^-N) concentration, and C:N ratio (Davidson and Swank, 1986; Abalos et al., 2014; Liu et al., 2017; Chen et al., 2013).

Empirical evidence thus shows that the introduction of no-till practices on N_2O emissions can cause both increases and decreases in N_2O emissions (van Kessel et al., 2013), since it affects the biophysical factors that influence N_2O emissions in contrasting manners (Snyder et al., 2009). There is generally an increase in soil moisture under NT, which is caused by higher water infiltration rates and reduced soil evaporation. In addition, no-till practices tend to increase bulk density and thus higher relative soil moisture contents affecting nitrification and denitrification rates and, therefore N_2O emissions (van Kessel et al., 2013). In wet regions, as in the tropical and humid areas, nitrification is thus mostly reduced by no-till practices, whereas it increases in dryer regions. Furthermore, no-till can lower soil temperature exchange between soil and atmosphere through the presence of litter residues, which can reduce N_2O emissions (Enrique et al., 1999). However, we lack a complete insight into the impacts of different tillage systems on the emission of N_2O in dependence on on-site properties (Congreves et al., 2017). Studies showed that CT might release 1.4–6.3 folds higher N_2O over NT in cold soils (Congreves et al., 2017). Zhang et al. (2013) showed rotary tillage had the highest potential of N_2O emissions over CT and NT. Regina and Alakukku (2010) reported in various Northern European Boreal soils that there is indeed a chance of elevated N_2O over the first years of NT compared to CT, which may remain almost consistent in both types. Shakoor et al. (2021) demonstrated in a meta-analysis that NT is responsible for higher N_2O over CT. N_2O releases were higher in irrigated sub-tropical rice-wheat wheat-maize systems under NT over CT (Pandey et al., 2012; Niu et al., 2019). These studies indicate that when tillage intensity is reduced, there remains a chance of higher N_2O emissions. As per our discussion in Section 2.0, it can be concluded that higher N_2O discharges in NT systems might be linked up with increased SOM, NH_4^+ plus NO_3^- contents, and WFPS. Moreover, a robust meta-analysis comprising 323 observations concluded that activity and number of denitrifiers and the abundance of denitrifying genes were proportionally higher in response to NT, however, thus enhancing N_2O production as compared to CT (Wang and Zou, 2020). Badagliacca et al. (2018) found almost similar results in the Mediterranean region. They stated that a substantial amount of N_2O was associated with an abundance of denitrifying genes, bacterial activities, greater bulk density, and WFPS. These studies indicate that soil structure and soil biota also control N_2O releases largely. Contrarily, Plaza-Bonilla et al. (2018) reported that soils under NT had 2.8–3.3 times lower N_2O emitting potential in the rainfed Mediterranean region compared to CT. This was because the unaccounted nitrifying and non-denitrifying populations lacking nitrous oxide reductase might be a potential storage for the produced soil N_2O (Hallin et al., 2018; Philippot

et al., 2011; Jones et al., 2013). These studies suggest that benefits from NT should be evaluated in terms of N_2O production since the global areas under NT are increasing. Apart from CT and NT, STR reduces N_2O emissions in dermosols and sodosols (Melland et al., 2017). RiT has almost similar trend of N_2O emissions in paddy fields compared to CT, and even the differences were not statistically significant for paddy soils (Qin et al., 2020). Ridge combined with NT can substantially lower N_2O emissions than CT in paddy soils (Li et al., 2015).

Besides, studies showed that nitrification is responsible for most N_2O emissions in well-aerated soil with less than 60% WFPS, while denitrification is the major agent of N_2O emissions when soil moisture surpasses 70% WFPS (Pihlatie et al., 2004; Bateman and Baggs, 2005). In a meta-analysis, Chen et al. (2013) showed that crop residue retention might be favorable for SCS (Abalos et al., 2013), but it provides a congenial microenvironment to denitrifiers if WFPS remains 60–90%. Consequently, a substantial volume of N_2O releases occur. But at >90% WFPS, it may act as a denitrification inhibitor. Abalos et al. (2013) found 105% increases in N_2O emissions in Mediterranean burley fields when crop residues are retained, but removal greatly cut off the N_2O level.

Nitrification inhibitors (Liu et al., 2017) such as dicyandiamide (DCD), nitrapyrin, pronitrade, etc., are proven to be the most promising means of abating N_2O pollution from agricultural soils. For example, when nitrogenous fertilizers are broadcasted, nitrapyrin can reduce N_2O emissions from cold semiarid climatic soils by 41% and 32%, respectively, as found in maize crop fields under both CT and RT practices (Borzouei et al., 2021). Additionally, nitrapyrin addition can significantly slow down NH_4^+ oxidation (Hayden et al., 2021; Giacometti et al., 2020). Liu et al. (2017) reported nitrapyrin plus urea addition in calcareous drip fertigated cotton soils might significantly stabilize soil $\text{NH}_4^+\text{-N}$ content and decrease N_2O emissions, but dependent on WFPS percentage. Combined application of nitrapyrin and DMPSA (3,4-dimethylpyrazole succinic acid) substantially reduced N_2O emissions in both gray Luvisol 60% and 56% at 60% and 80% WFPS, respectively (Lin and Hernandez-Ramirez, 2020). Meng et al. (2020) found DCD, nitrapyrin, and N- (n-butyl) thiophosphoric triamide (NBPT) as the potential N_2 fixer in paddy soils. These studies indicate that soil water content has a prominent role in both nitrification and denitrification processes also on the effectiveness of nitrification inhibitors. These substances can improve benefits from SCS by abating soil N_2O releases by increasing the use efficiency of nitrogenous fertilizers under different tillage systems.

5. Overall conclusions and perspectives

This review compares and contrasts the co-benefits and trade-offs produced from the impact of different tillage practices. We also highlighted several important soil ecosystem services (soil carbon sequestration, GHG regulation, water, and air quality) and their inter-and intra- relationships with different tillage practices. We confirm our hypothesis that improved tillage practices can enable agriculture to contribute to soil carbon sequestration (SCS) and mitigate GHG emissions and loss of accumulated nutrients and pesticides. Our significant findings are:

- SOM is very prone to different tillage practices. Therefore, SOM protection is crucial for maximized SCS. Although NT practices tend to enrich SOM in the upper surface of soils, our study indicated that its impact on the whole soil profile is still dubious. Therefore, DT is a good option for improving continuous NT soils and mixing up the SOM across the entire soil profile since NT promotes greater bulk density.
- CT emits more CO_2 than NT because of higher mineralization rates, microbial activity, and soil temperature and moisture. Although NT and RT have a greater SCS than other tillage techniques, higher soil temperatures may cause sequestered carbon to be lost. Higher N_2O emission under NT is related to soil moisture, increased denitrifier activity, and higher NO_3^- levels in soils. Therefore, soil moisture management is crucial to prevent N_2O emissions from NT soils.
- The volume and magnitude of leaching loss of pesticides and nutrients

are relatively low in the case of CT and DT compared to NT, as they can bury them deep inside of soils. For example, we observed that NT enhances NO_3^- leaching over CT because of improved soil structure, infiltration capacity, and greater water flux.

- d) Tillage erosion coupled with splash/wind erosion remains a significant problem. These processes may decrease the rate of SCS and increase the magnitude of nutrient pollution (eutrophication) in water bodies.
- e) Nitrification inhibitors such as nitrapyrin or DCD coupled with nitrogenous fertilizers can potentially reduce N_2O emissions from soils. The advantages of nitrapyrin should be evaluated more in field settings (Luo et al., 2017). Future research on how to improve the effectiveness of nitrapyrin should be expanded. Despite preventing N_2O emissions, these substances are also quite capable of avoiding leaching loss of nutrients (N and P). Our study suggests that improvement of tillage practices in terms of mitigation of N_2O emission and preventing NO_3^- pollution may be achievable if these inhibitors are applied.

Apart from these, we also demonstrated that the accumulation of SOC and associated SCS is highly correlated with adopted tillage systems. All of the tillage practices have the potential to improve SCS, nevertheless, the role of NT over other practices on SCS is still unclear (Chatterjee and Lal, 2009), and also full potential is confined to certain pedo-climatic-regional conditions. Based on our current understanding, it can be concluded that if soil-climatic needs are not taken seriously, then it may produce erroneous estimates. The confounding factor that triggers the uncertainties and variations in data might be heterogeneity in the adopted outlook regarding NT (Meurer et al., 2018). Derpsch et al. (2014) proposed to develop a systematic procedure based on spatio-temporal variations to avoid existing doubts regarding the SCS potential of NT.

Additionally, Xiao et al. (2020) and Blanco-Canqui et al. (2021) identified the depth factor as the main reason for fueling debate regarding NT. Moreover, CT can improve SOC contents, ultimately reallocating C content in the whole soil profile. NT/RT practices improve soil moisture, water infiltration, stable aggregates, increased micropores, POC, MAOM, humin contents, TSN, MBC, TOC, etc. RT soils have a lower bulk density over NT, higher macropore, and improved aeration. RiT is associated with decreased evapotranspiration loss, improved water use efficiency, and greater SCS than CT and NT. Reduced erosion and reduced runoff from ST system yield in improved SCS, TOC, and TON. STR improves aggregate stability, bulk density, soil compactions, and a smart quantity of sequestered C. DT effectively raises SOM contents and improves SCS. We found a dearth of research in finding out the GHG emissions caused by STR, ST, and RiT. It's critical to find a sustainable solution for limiting C losses under high soil temperature using various tillage techniques. In addition, evaluation of eutrophication potential of different tillage practices (e.g., NT) should be taken into consideration. Current research is much focused on CT and NT, their SCS potential evaluation, and GHG producing capabilities. Future research on other forms of tillage practices needs to be undertaken to have a clear insight into other practices.

CRedit authorship contribution statement

Siddhartha Shankar Bhattacharyya: Conceptualization, Data Analysis, Tables, Figures, Writing-Original Draft, Reviewing and Editing. **Fernanda Figueiredo Granja Dorilêo Leite:** Data Analysis, Tables, Writing-Original Draft. **Casey L. France:** Tables, Writing-Original Draft. **Adetomi O. Adekoya:** Tables, Writing-Original Draft. **Gerard H. Ros:** Figures, Tables, Writing-Original Draft. **Wim de Vries:** Data Analysis, Figures, Tables, Writing-Original Draft. **Elda M. Melchor-Martínez:** Data Analysis, Figures, Tables, Writing-Original Draft. **Hafiz M.N. Iqbal:** Conceptualization, Data Analysis, Figures, Tables, Writing-Original Draft, Reviewing and Editing, Supervision. **Roberto Parra-Saldívar:** Conceptualization, Data Analysis, Figures, Tables, Writing-Original Draft, Reviewing and Editing, Supervision.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

Consejo Nacional de Ciencia y Tecnología (CONACYT) Mexico is thankfully acknowledged for partially supporting this work under Sistema Nacional de Investigadores (SNI) program awarded to Elda M. Melchor-Martínez (CVU: 230784), Hafiz M.N. Iqbal (CVU: 735340) and Roberto Parra-Saldívar (CVU: 35753). S. S. Bhattacharyya thankfully acknowledges the Department of Agronomy, Bangladesh Agricultural University, Mymensingh, Bangladesh, for literature facilities. The authors would like to express their admiration to the four anonymous reviewers for their insightful and critical remarks and suggestions, which helped to improve a previous version of the paper.

References

- Abalos, D., Sanz-Cobena, A., Garcia-Torres, L., van Groenigen, J.W., Vallejo, A., 2013. Role of maize Stover incorporation on nitrogen oxide emissions in a non-irrigated Mediterranean barley field. *Plant Soil* 364, 357–371. <https://doi.org/10.1007/s11040-012-1367-4>.
- Abalos, D., Sanchez-Martin, L., Garcia-Torres, L., van Groenigen, J.W., Vallejo, A., 2014. Management of irrigation frequency and nitrogen fertilization to mitigate GHG and NO emissions from drip-fertigated crops. *Sci. Total Environ.* 490, 880–888. <https://doi.org/10.1016/j.scitotenv.2014.05.065>.
- Abdalla, M., Osborne, B., Lanigan, G., Forristal, D., Williams, M., Smith, P., Jones, M.B., 2013. Conservation tillage systems: a review of its consequences for greenhouse gas emissions. *Soil Use Manag.* 29, 199–209. <https://doi.org/10.1111/sum.12030>.
- Abdalla, K., Chivenge, P., Ciaes, P., Chaplot, V., 2016. No-tillage lessens soil CO_2 emissions the most under arid and sandy soil conditions: results from a meta-analysis. *Biogeosciences* 13, 3619–3633. <https://doi.org/10.5194/bg-13-3619-2016>.
- Abdollahi, L., Munkholm, L.J., 2017. Eleven years' effect of conservation practices for temperate Sandy loams: II. Soil pore characteristics. *Soil Sci. Soc. Am. J.* 81, 392–403. <https://doi.org/10.2136/sssaj2016.07.0221>.
- Abuzaid, A.S., Bassouny, M.A., 2020. Total and DTPA-extractable forms of potentially toxic metals in soils of rice fields, North Nile Delta of Egypt. *Environ. Technol. Innov.* 18, 100717. <https://doi.org/10.1016/j.eti.2020.100717>.
- Acharya, B.S., Dodla, S., Gaston, L.A., Darapuni, M., Wang, J.J., Sepat, S., Bohara, H., 2019. Winter cover crops effect on soil moisture and soybean growth and yield under different tillage systems. *Soil Tillage Res.* 195, 104430. <https://doi.org/10.1016/j.still.2019.104430>.
- Acir, N., Günel, H., Çelik, İ., Barut, Z.B., Budak, M., Kılıç, Ş., 2020. Effects of long-term conventional and conservation tillage systems on biochemical soil health indicators in the Mediterranean region. *Arch. Agron. Soil Sci.* <https://doi.org/10.1080/03650340.2020.1855327>.
- Ahmad, S., Li, C., Dai, G., Zhan, M., Wang, J., Pan, S., Cao, C., 2009. Greenhouse gas emission from direct seeding paddy field under different rice tillage systems in Central China. *Soil Tillage Res.* 106, 54–61. <https://doi.org/10.1016/j.still.2009.09.005>.
- Alcántara, V., Don, A., Well, R., Nieder, R., 2016. Deep ploughing increases agricultural soil organic matter stocks. *Glob. Chang. Biol.* 22, 2939–2956. <https://doi.org/10.1111/gcb.13289>.
- Ali, I., Khan, M.J., Khan, M., Deeba, F., Hussain, H., Abbas, M., Khan, M.D., 2018. Impact of pollutants on paddy soil and crop quality. In: MH, AV (Eds.), *Environmental Pollution of Paddy Soils*. Soil Biology. Springer, Cham, pp. 125–137. https://doi.org/10.1007/978-3-319-93671-0_8.
- Al-Kaisi, M.M., Yin, X., 2005. Tillage and crop residue effects on soil carbon and carbon dioxide emission in corn-soybean rotations. *J. Environ. Qual.* 34, 437–445. <https://doi.org/10.2134/jeq2005.0437>.
- Alletto, L., Coquet, Y., Benoit, P., Heddad, D., Barriuso, E., 2010. Tillage management effects on pesticide fate in soils. A review. *Agron. Sustain. Dev.* 30, 367–400. <https://doi.org/10.1051/agro/2009018>.
- Almaraz, J.J., Zhou, X., Mabood, F., Madramootoo, C., Rochette, P., Ma, B.L., Smith, D.L., 2009. Greenhouse gas fluxes associated with soybean production under two tillage systems in southwestern Quebec. *Soil Tillage Res.* 104, 134–139. <https://doi.org/10.1016/j.still.2009.02.003>.
- de Almeida, W.S., Panachuki, E., de Oliveira, P.T.S., da Silva Menezes, R., Sobrinho, T.A., de Carvalho, D.F., 2018. Effect of soil tillage and vegetal cover on soil water infiltration. *Soil Tillage Res.* 175, 130–138. <https://doi.org/10.1016/j.still.2017.07.009>.
- Askaf, K., Mooney, S.J., Sparkes, D.L., Wilson, P., Sjögersten, S., 2021. Short-term impacts of different tillage practices and plant residue retention on soil physical properties and greenhouse gas emissions. *Soil Tillage Res.* 206, 104803. <https://doi.org/10.1016/j.still.2020.104803>.
- Alvarez, R., Steinbach, H.S., 2009. A review of the effects of tillage systems on some soil physical properties, water content, nitrate availability and crops yield in the Argentine pampas. *Soil Tillage Res.* 104, 1–15. <https://doi.org/10.1016/j.still.2009.02.005>.

- Assunção, S.A., Pereira, M.G., Rosset, J.S., Berbara, R.L.L., García, A.C., 2019. Carbon input and the structural quality of soil organic matter as a function of agricultural management in a tropical climate region of Brazil. *Sci. Total Environ.* 658, 901–911. <https://doi.org/10.1016/j.scitotenv.2018.12.271>.
- Awale, R., Emeson, M.A., Machado, S., 2017. Soil organic carbon pools as early indicators for soil organic matter stock changes under different tillage practices in inland Pacific north-west. *Front. Ecol. Evol.* 5, 96. <https://doi.org/10.3389/fevo.2017.00096>.
- Azam, A., Akhtar, M.S., Rukh, S., Mehmood, A., Imran, M., Khan, A., Qayyum, A., Ahmad, W., Gurmani, A.R., 2020. Changes in soil organic carbon fractions across a loess toposequence. *J. Soil Sci. Plant Nutr.* 20, 1193–1202. <https://doi.org/10.1007/s42729-020-00204-2>.
- Aziz, I., Mahmood, T., Islam, K.R., 2013. Effect of long term no-till and conventional tillage practices on soil quality. *Soil Tillage Res.* 131, 28–35. <https://doi.org/10.1016/j.still.2013.03.002>.
- Babal, B., Phogat, V.K., Sharma, M.K., Ahlawat, S., 2021. Impact of long-term conservation agriculture induced changes in soil properties on persistence of pendimethalin under different cropping systems. *Soil Res.* 59, 299–308. <https://doi.org/10.1071/SR201222>.
- Badagliacca, G., Benítez, E., Amato, G., Badalucco, L., Giambalvo, D., Laudicina, V.A., Ruisi, P., 2018. Long-term no-tillage application increases soil organic carbon, nitrous oxide emissions and faba bean (*Vicia faba* L.) yields under rain-fed Mediterranean conditions. *Sci. Total Environ.* 639, 350–359. <https://doi.org/10.1016/j.scitotenv.2018.05.157>.
- Bai, X., Huang, Y., Ren, W., Coyne, M., Jacinthe, P.A., Tao, B., Hui, D., Yang, J., Matocha, C., 2019. Responses of soil carbon sequestration to climate-smart agriculture practices: a meta-analysis. *Glob. Chang. Biol.* 25, 2591–2606. <https://doi.org/10.1111/gcb.14658>.
- Ball, B.C., Scott, A., Parker, J.P., 1999. Field N₂O, CO₂ and CH₄ fluxes in relation to tillage, compaction and soil quality in Scotland. *Soil Tillage Res.* 53, 29–39. [https://doi.org/10.1016/S0167-1987\(99\)00074-4](https://doi.org/10.1016/S0167-1987(99)00074-4).
- Bandyopadhyay, K.K., 2019. Effect of tillage on soil carbon sequestration. In: PG, SM, DM, BM, SR (Eds.), *Carbon Management in Tropical and Sub-tropical Terrestrial Systems*. Springer, Singapore, pp. 213–229. https://doi.org/10.1007/978-981-13-9628-1_13.
- Bateman, E.J., Baggs, E.M., 2005. Contributions of nitrification and denitrification to N₂O emissions from soils at different water-filled pore space. *Biol. Fertil. Soils* 41, 379–388. <https://doi.org/10.1007/s00374-005-0858-3>.
- Bayer, C., Gomes, J., Zanatta, J.A., Vieira, F.C.B., Dieckow, J., 2016. Mitigating greenhouse gas emissions from a subtropical ultisol by using long-term no-tillage in combination with legume cover crops. *Soil Tillage Res.* 161, 86–94. <https://doi.org/10.1016/j.still.2016.03.011>.
- Bertol, L., Luciano, R.V., Bertol, C., Bagio, B., 2017. Nutrient and organic carbon losses, enrichment rate, and cost of water erosion. *Rev. Bras. Cienc. Solo* 41. <https://doi.org/10.1590/1869657rbs20160150>.
- Bertolino, A.V.F.A., Fernandes, N.F., Miranda, J.P.L., Souza, A.P., Lopes, M.R.S., Palmieri, F., 2010. Effects of plough pan development on surface hydrology and on soil physical properties in southeastern Brazilian plateau. *J. Hydrol.* 393, 94–104. <https://doi.org/10.1016/j.jhydrol.2010.07.038>.
- Bhattacharyya, S.S., Leite, F.F.G.D., Adeyemi, M.A., Sarker, A.J., Cambareri, G.S., Faverin, C., Trieri, M.P., Castillo-Zacarias, C., Melchor-Martinez, E.M., Iqbal, H.M.N., Parra-Saldivar, R., 2021. A paradigm shift to CO₂ sequestration to manage global warming – with the emphasis on developing countries. *Sci. Total Environ.* 790, 148169. <https://doi.org/10.1016/j.scitotenv.2021.148169>.
- Bhattacharyya, S.S., Ros, G.H., Furtak, K., Iqbal, H.M.N., Parra-Saldivar, R., 2022. Soil carbon sequestration – an interplay between soil microbial community and soil organic matter dynamics. *Sci. Total Environ.* 815, 152928. <https://doi.org/10.1016/j.scitotenv.2022.152928>.
- Blanco-Canqui, H., Wienhold, B.J., Jin, V.L., Schmer, M.R., Kibet, L.C., 2017. Long-term tillage impact on soil hydraulic properties. *Soil Tillage Res.* 170, 38–42. <https://doi.org/10.1016/j.still.2017.03.001>.
- Blanco-Canqui, H., Shapiro, C., Jasa, P., Iqbal, J., 2021. No-till and carbon stocks: is deep soil sampling necessary? Insights from long-term experiments. *Soil Tillage Res.* 206, 104840. <https://doi.org/10.1016/j.still.2020.104840>.
- Bogunovic, I., Pereira, P., Kisić, I., Sajko, K., Sraka, M., 2018. Tillage management impacts on soil compaction, erosion and crop yield in stagnosols (Croatia). *Catena* 160, 376–384. <https://doi.org/10.1016/j.catena.2017.10.009>.
- Borzouei, A., Mander, U., Teemusk, A., Sanz-Cobena, A., Zaman, M., Kim, D.G., Muller, C., Kelestanie, A.A., Amin, P.S., Moghiseh, E., Dawar, K., Pérez-Castillo, A.G., 2021. Effects of the nitrification inhibitor nitrapyrin and tillage practices on yield-scaled nitrous oxide emission from a maize field in Iran. *Pedosphere* 31, 314–322. [https://doi.org/10.1016/S1002-0160\(20\)60067-4](https://doi.org/10.1016/S1002-0160(20)60067-4).
- Bottinelli, N., Angers, D.A., Hallaire, V., Michot, D., Le Guillou, C., Cluzeau, D., Heddad, D., Menasseri-Aubry, S., 2017. Tillage and fertilization practices affect soil aggregate stability in a humic cambisol of Northwest France. *Soil Tillage Res.* 170, 14–17. <https://doi.org/10.1016/j.still.2017.02.008>.
- Braker, G., Conrad, R., 2011. Diversity, structure, and size of N₂O-producing microbial communities in soils-what matters for their functioning? *Adv. Appl. Microbiol.* 75, 33–70. <https://doi.org/10.1016/B978-0-12-387046-9.00002-5>.
- Bréchet, L.M., Daniel, W., Stahl, C., Burban, B., Goret, J.Y., Salomón, R.L., Janssens, I.A., 2021. Simultaneous tree stem and soil greenhouse gas (CO₂, CH₄, N₂O) flux measurements: a novel design for continuous monitoring towards improving flux estimates and temporal resolution. *New Phytol.* 230, 2487–2500. <https://doi.org/10.1111/nph.17352>.
- Brewer, K.M., Gaudin, A.C.M., 2020. Potential of crop-livestock integration to enhance carbon sequestration and agroecosystem functioning in semi-arid croplands. *Soil Biol. Biochem.* 149, 107936. <https://doi.org/10.1016/j.soilbio.2020.107936>.
- Busari, M.A., Kukal, S.S., Kaur, A., Bhatt, R., Dulazi, A.A., 2015. Conservation tillage impacts on soil, crop and the environment. *Int. Soil Water Conserv. Res.* 3, 119–129. <https://doi.org/10.1016/j.iswcr.2015.05.002>.
- Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., Zechmeister-Boltenstern, S., 2013. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philos. Trans. R. Soc. B Biol. Sci.* 368, 20130122.
- Carretta, L., Tarolli, P., Cardinali, A., Nasta, P., Romano, N., Masin, R., 2021. Evaluation of runoff and soil erosion under conventional tillage and no-till management: a case study in Northeast Italy. *Catena* 197, 104972. <https://doi.org/10.1016/j.catena.2020.104972>.
- Çelik, İ., Günel, H., Acar, M., Acir, N., Bereket Barut, Z., Budak, M., 2019. Strategic tillage may sustain the benefits of long-term no-till in a vertisol under Mediterranean climate. *Soil Tillage Res.* 185, 17–28. <https://doi.org/10.1016/j.still.2018.08.015>.
- Çelik, İ., Günel, H., Acir, N., Barut, Z.B., Budak, M., 2021. Soil quality assessment to compare tillage systems in Cukurova Plain, Turkey. *Soil Tillage Res.* 208, 104892. <https://doi.org/10.1016/j.still.2020.104892>.
- Cerli, C., Celi, L., Kalbitz, K., Guggenberger, G., Kaiser, K., 2012. Separation of light and heavy organic matter fractions in soil - testing for proper density cut-off and dispersion level. *Geoderma* 170, 403–416. <https://doi.org/10.1016/j.geoderma.2011.10.009>.
- Chalise, D., Kumar, L., Kristiansen, P., 2019. Land degradation by soil erosion in Nepal: a review. *Soil Syst.* 3, 1–18. <https://doi.org/10.3390/soilsystems3010012>.
- Chatterjee, A., Lal, R., 2009. On farm assessment of tillage impact on soil carbon and associated soil quality parameters. *Soil Tillage Res.* 104, 270–277. <https://doi.org/10.1016/j.still.2009.03.006>.
- Chen, H., Li, X., Hu, F., Shi, W., 2013. Soil nitrous oxide emissions following crop residue addition: a meta-analysis. *Glob. Chang. Biol.* 19, 2956–2964. <https://doi.org/10.1111/gcb.12274>.
- Chen, Z.D., Zhang, H., Xue, J., Liu, S., Chen, F., 2021. A nine-year study on the effects of tillage on net annual global warming potential in double rice-cropping systems in southern China. *Soil Tillage Res.* 206, 104797. <https://doi.org/10.1016/j.still.2020.104797>.
- Chenu, C., Angers, D.A., Barré, P., Derrien, D., Arrouays, D., Balesdent, J., 2019. Increasing organic stocks in agricultural soils: knowledge gaps and potential innovations. *Soil Tillage Res.* 188, 41–52. <https://doi.org/10.1016/j.still.2018.04.011>.
- Chowaniak, M., Glab, T., Klima, K., Niemiec, M., Zaleski, T., Zuzek, D., 2020. Effect of tillage and crop management on runoff, soil erosion and organic carbon loss. *Soil Use Manag.* 36, 581–593. <https://doi.org/10.1111/sum.12606>.
- Christensen, B.T., 2020. Carbon in primary and secondary organomineral complexes. *Structure and Organic Matter Storage in Agricultural Soils*, pp. 97–165. <https://doi.org/10.1201/9781003075561-6>.
- Closte, F., Woignier, T., Rangon, L., Fernandes, P., Soler, A., Lesueur-Jannoyer, M., 2014. Field validation of chlordecone soil sequestration by organic matter addition. *J. Soils Sediments* 14, 23–33. <https://doi.org/10.1007/s11368-013-0790-3>.
- Congreves, K.A., Brown, S.E., Németh, D.D., Dunfield, K.E., Wagner-Riddle, C., 2017. Differences in field-scale N₂O flux linked to crop residue removal under two tillage systems in cold climates. *GCB Bioenergy* 9, 666–680. <https://doi.org/10.1111/gcbb.12354>.
- Corbeels, M., Marchão, R.L., Neto, M.S., Ferreira, E.G., Madari, B.E., Scopel, E., Brito, O.R., 2016. Evidence of limited carbon sequestration in soils under no-tillage systems in the cerrado of Brazil. *Sci. Rep.* 6, 21450. <https://doi.org/10.1038/srep21450>.
- Dang, Y.P., Seymour, N.P., Walker, S.M., Bell, M.J., Freebairn, D.M., 2015. Strategic tillage in no-till farming systems in Australia's northern grains-growing regions: I. Drivers and implementation. *Soil Tillage Res.* 152, 104–114. <https://doi.org/10.1016/j.still.2015.03.009>.
- Daryanto, S., Wang, L., Jacinthe, P.A., 2017. Impacts of no-tillage management on nitrate loss from corn, soybean and wheat cultivation: a meta-analysis. *Sci. Rep.* 7, 12117. <https://doi.org/10.1038/s41598-017-12383-7>.
- Daryanto, S., Wang, L., Jacinthe, P.A., 2017. Meta-analysis of phosphorus loss from no-till soils. *J. Environ. Qual.* 46, 1028–1037. <https://doi.org/10.2134/jeq2017.03.0121>.
- Daverde, I.C., Kravchenko, A.N., Hoeft, R.G., Nafziger, E.D., Bullock, D.G., Warren, J.J., Gonzini, L.C., 2003. Phosphorus runoff. *J. Environ. Qual.* 32, 1436. <https://doi.org/10.2134/jeq2003.1436>.
- Davidson, E.A., Swank, W.T., 1986. Environmental parameters regulating gaseous nitrogen losses from two forested ecosystems via nitrification and denitrification. *Appl. Environ. Microbiol.* 52, 1287–1292. <https://doi.org/10.1128/aem.52.6.1287-1292.1986>.
- Dendooven, L., Gutiérrez-Oliva, V.F., Patiño-Zúñiga, L., Ramírez-Villanueva, D.A., Verhulst, N., Luna-Guido, M., Marsch, R., Montes-Molina, J., Gutiérrez-Miceli, F.A., Vásquez-Murrieta, S., Govaerts, B., 2012. Greenhouse gas emissions under conservation agriculture compared to traditional cultivation of maize in the central highlands of Mexico. *Sci. Total Environ.* 431, 237–244. <https://doi.org/10.1016/j.scitotenv.2012.05.029>.
- Derpsch, R., Franzluebbers, A.J., Duiker, S.W., Reicosky, D.C., Koeller, K., Friedrich, T., Sturny, W.G., Sá, J.C.M., Weiss, K., 2014. Why do we need to standardize no-tillage research? *Soil Tillage Res.* <https://doi.org/10.1016/j.still.2013.10.002>.
- Deuschle, D., Minella, J.P.G., Londero, A.L., Schneider, F.J.A., Hörbe, T.de A.N., 2019. Erosion and hydrological response in no-tillage subjected to crop rotation intensification in southern Brazil. *Geoderma* 340, 157–163. <https://doi.org/10.1016/j.geoderma.2019.01.010>.
- Díaz, M.A., Bown, H.E., Fuentes, J.P., Martínez, A.M., 2018. Soils act as sinks or sources of CH₄ depending on air-filled porosity in sclerophyllous ecosystems in semiarid Central Chile. *Appl. Soil Ecol.* 130, 13–20. <https://doi.org/10.1016/j.apsoil.2018.05.017>.
- Dimassi, B., Mary, B., Wylleman, R., Labreuche, J.O., Couture, D., Piraux, F., Cohan, J.P., 2014. Long-term effect of contrasted tillage and crop management on soil carbon dynamics during 41 years. *Agric. Ecosyst. Environ.* 188, 134–146. <https://doi.org/10.1016/j.agee.2014.02.014>.
- Du, Z., Angers, D.A., Ren, T., Zhang, Q., Li, G., 2017. The effect of no-till on organic C storage in Chinese soils should not be overemphasized: a meta-analysis. *Agric. Ecosyst. Environ.* 236, 1–11. <https://doi.org/10.1016/j.agee.2016.11.007>.
- Düring, R.A., Hoß, T., Gath, S., 2002. Depth distribution and bioavailability of pollutants in long-term differently tilled soils. *Soil Tillage Res.* 66, 183–195. [https://doi.org/10.1016/S0167-1987\(02\)00026-0](https://doi.org/10.1016/S0167-1987(02)00026-0).
- Dzul-Caamal, R., Vega-López, A., von Osten, J.R., 2020. Distribution of heavy metals in crop soils from an agricultural region of the Yucatan peninsula and biochemical changes in

- earthworm *Eisenia foetida* exposed experimentally. *Environ. Monit. Assess.* 192, 338. <https://doi.org/10.1007/s10661-020-08273-7>.
- Elias, D., Wang, L., Jacinthe, P.A., 2018. A meta-analysis of pesticide loss in runoff under conventional tillage and no-till management. *Environ. Monit. Assess.* 190. <https://doi.org/10.1007/s10661-017-6441-1>.
- Endale, D.M., Potter, T.L., Strickland, T.C., Bosch, D.D., 2017. Sediment-bound total organic carbon and total organic nitrogen losses from conventional and strip tillage cropping systems. *Soil Tillage Res.* 171, 25–34. <https://doi.org/10.1016/j.still.2017.04.004>.
- Endale, D.M., Schomberg, H.H., Truman, C.C., Franklin, D.H., Tazisong, I.A., Jenkins, M.B., Fisher, D.S., Campbell, J.P., 2019. Runoff and nutrient losses from conventional and conservation tillage systems during fixed and variable rate rainfall simulation. *J. Soil Water Conserv.* 74, 594–612. <https://doi.org/10.2489/JSWC.74.6.594>.
- Enrique, G.S., Braud, I., Jean-Louis, T., Michel, V., Pierre, B., Jean-Christophe, C., 1999. Modelling heat and water exchanges of fallow land covered with plant residue mulch. *Agric. For. Meteorol.* 97, 151–169. [https://doi.org/10.1016/S0168-1923\(99\)00081-7](https://doi.org/10.1016/S0168-1923(99)00081-7) WOS:000083648600002.
- Ewing, S.A., Sanderman, J., Baisden, W.T., Wang, Y., Amundson, R., 2006. Role of large-scale soil structure in organic carbon turnover: evidence from California grassland soils. *J. Geophys. Res. Biogeosci.* 111. <https://doi.org/10.1029/2006JG000174>.
- Fangueiro, D., Becerra, D., Albarrán, Á., Peña, D., Sanchez-Llerena, J., Rato-Nunes, J.M., López-Piñero, A., 2017. Effect of tillage and water management on GHG emissions from Mediterranean rice growing ecosystems. *Atmos. Environ.* 150, 303–312. <https://doi.org/10.1016/j.atmosenv.2016.11.020>.
- Feng, Q., An, C., Chen, Z., Wang, Z., 2020. Can deep tillage enhance carbon sequestration in soils? A meta-analysis towards GHG mitigation and sustainable agricultural management. *Renew. Sust. Energ. Rev.* 133, 110293. <https://doi.org/10.1016/j.rser.2020.110293>.
- Feng, J., Yang, T., Li, F., Zhou, X., Xu, C., Fang, F., 2021. Impact of tillage on the spatial distribution of CH₄ and N₂O in the soil profile of late rice fields. *Soil Tillage Res.* 211, 105029. <https://doi.org/10.1016/j.still.2021.105029>.
- Ferreira, C.dos R., Silva Neto, E.C.da, Pereira, M.G., Rosset, J.S., Anjos, L.H.C.dos, Guedes, J. do N., 2020. Dynamics of soil aggregation and organic carbon fractions over 23 years of no-till management. *Soil Tillage Res.* 198, 104533. <https://doi.org/10.1016/j.still.2019.104533>.
- de Figueiredo, C.C., de Oliveira, A.D., dos Santos, I.L., Ferreira, E.A.B., Malaquias, J.V., de Sá, M.A.C., de Carvalho, A.M., dos Santos, J.de D.G., 2018. Relationships between soil organic matter pools and nitrous oxide emissions of agroecosystems in the Brazilian Cerrado. *Sci. Total Environ.* 618, 1572–1582. <https://doi.org/10.1016/j.scitotenv.2017.09.333>.
- Fowler, D., Coyle, M., Skiba, U., Sutton, M.A., Cape, J.N., Reis, S., Sheppard, L.J., Jenkins, A., Grizzetti, B., Galloway, J.N., Vitousek, P., Leach, A., Bouwman, A.F., Butterbach-Bahl, K., Dentener, F., Stevenson, D., Amann, M., Voss, M., 2013. The global nitrogen cycle in the twenty-first century. *Philos. Trans. R. Soc. B Biol. Sci.* 368, 20130164. <https://doi.org/10.1098/rstb.2013.0164>.
- Franco-Luesma, S., Caverio, J., Plaza-Bonilla, D., Cantero-Martínez, C., Arrúe, J.L., Álvaro-Fuentes, J., 2020. Tillage and irrigation system effects on soil carbon dioxide (CO₂) and methane (CH₄) emissions in a maize monoculture under Mediterranean conditions. *Soil Tillage Res.* 196, 104488. <https://doi.org/10.1016/j.still.2019.104488>.
- Galdos, M.V., Pires, L.F., Cooper, H.V., Calonego, J.C., Rosolem, C.A., Mooney, S.J., 2019. Assessing the long-term effects of zero-tillage on the macroporosity of Brazilian soils using X-ray computed tomography. *Geoderma* 337, 1126–1135. <https://doi.org/10.1016/j.geoderma.2018.11.031>.
- Ghidey, F., Baffaut, C., Lerch, R.N., Kitchen, N.R., Sadler, E.J., Sudduth, K.A., 2010. Herbicide transport to surface runoff from a claypan soil: scaling from plots to fields. *J. Soil Water Conserv.* 65, 168–179. <https://doi.org/10.2489/jswc.65.3.168>.
- Giacometti, C., Mazzon, M., Cavani, L., Ciavatta, C., Marzadori, C., 2020. A nitrification inhibitor, nitrpyrin, reduces potential nitrate leaching through soil columns treated with animal slurries and anaerobic digestate. *Agronomy* 10, 865. <https://doi.org/10.3390/agronomy10060865>.
- Gonzalez, J.M., 2018. Runoff and losses of nutrients and herbicides under long-term conservation practices (no-till and crop rotation) in the U.S. Midwest: a variable intensity simulated rainfall approach. *Int. Soil Water Conserv. Res.* 6, 265–274. <https://doi.org/10.1016/j.iswcr.2018.07.005>.
- Gregorich, E.G., Rochette, P., VandenBygaart, A.J., Angers, D.A., 2005. Greenhouse gas contributions of agricultural soils and potential mitigation practices in eastern Canada. *Soil Tillage Res.* 83, 53–72. <https://doi.org/10.1016/j.still.2005.02.009>.
- van Groenigen, K.J., Hastings, A., Forristal, D., Roth, B., Jones, M., Smith, P., 2011. Soil C storage as affected by tillage and straw management: an assessment using field measurements and model predictions. *Agric. Ecosyst. Environ.* 140, 218–225. <https://doi.org/10.1016/j.agee.2010.12.008>.
- Guenet, B., Gabrielle, B., Chenu, C., Arrouays, D., Balesdent, J., Bernoux, M., Bruni, E., Caliman, J.P., Cardinael, R., Chen, S., Ciais, P., Desbois, D., Fouché, J., Frank, S., Henault, C., Lugato, E., Naipal, V., Nesme, T., Obersteiner, M., Pellerin, S., Powlson, D.S., Rasse, D.P., Rees, F., Soussana, J.F., Su, Y., Tian, H., Valin, H., Zhou, F., 2021. Can N₂O emissions offset the benefits from soil organic carbon storage? *Glob. Chang. Biol.* 27, 237–256. <https://doi.org/10.1111/gcb.15342>.
- Guo, S., Zhai, L., Liu, J., Liu, H., Chen, A., Wang, H., Wu, S., Lei, Q., 2019. Cross-ridge tillage decreases nitrogen and phosphorus losses from sloping farmlands in southern hilly regions of China. *Soil Tillage Res.* 191, 48–56. <https://doi.org/10.1016/j.still.2019.03.015>.
- Guo, Y., Fan, R., Zhang, X., Zhang, Y., Wu, D., McLaughlin, N., Zhang, S., Chen, X., Jia, S., Liang, A., 2020. Tillage-induced effects on SOC through changes in aggregate stability and soil pore structure. *Sci. Total Environ.* 703, 134617. <https://doi.org/10.1016/j.scitotenv.2019.134617>.
- Gwenzi, W., Gotosa, J., Chakanetsa, S., Mutema, Z., 2009. Effects of tillage systems on soil organic carbon dynamics, structural stability and crop yields in irrigated wheat (*Triticum aestivum* L.)-cotton (*Gossypium hirsutum* L.) rotation in semi-arid Zimbabwe. *Nutr. Cycl. Agroecosyst.* 83, 211–221. <https://doi.org/10.1007/s10705-008-9211-1>.
- Haddaway, N.R., Hedlund, K., Jackson, L.E., Kätterer, T., Lugato, E., Thomsen, I.K., Jørgensen, H.B., Isberg, P.E., 2017. How does tillage intensity affect soil organic carbon? A systematic review. *Environ. Evid.* 6. <https://doi.org/10.1186/s13750-017-0108-9>.
- Hallin, S., Philippot, L., Löffler, F.E., Sanford, R.A., Jones, C.M., 2018. Genomics and ecology of novel N₂O-reducing microorganisms. *Trends Microbiol.* 26, 43–55. <https://doi.org/10.1016/j.tim.2017.07.003>.
- Hance, R.J., 1974. Soil organic matter and the adsorption and decomposition of the herbicides atrazine and linuron. *Soil Biol. Biochem.* 6, 39–42. [https://doi.org/10.1016/0038-0717\(74\)90009-1](https://doi.org/10.1016/0038-0717(74)90009-1).
- Hanrahan, B.R., King, K.W., Williams, M.R., Duncan, E.W., Pease, L.A., LaBarge, G.A., 2019. Nutrient balances influence hydrologic losses of nitrogen and phosphorus across agricultural fields in northwestern Ohio. *Nutr. Cycl. Agroecosyst.* 113, 231–245. <https://doi.org/10.1007/s10705-019-09981-4>.
- Hati, K.M., Biswas, A.K., Somasundaram, J., Mohanty, M., Singh, R.K., Sinha, N.K., Chaudhary, R.S., 2019. Soil organic carbon dynamics and carbon sequestration under conservation tillage in tropical vertisols. In: PG, SM, DM, BM, SR (Eds.), *Carbon Management in Tropical and Sub-tropical Terrestrial Systems*. Springer, Singapore, pp. 201–212. https://doi.org/10.1007/978-981-13-9628-1_12.
- Hayden, H.L., Phillips, L.A., Marshall, A.J., Condon, J.R., Doran, G.S., Wells, G.S., Mele, P.M., 2021. Nitrpyrin reduced ammonia oxidation with different impacts on the abundance of bacterial and archaeal ammonia oxidisers in four agricultural soils. *Appl. Soil Ecol.* 157, 103759. <https://doi.org/10.1016/j.apsoil.2020.103759>.
- Hayes, M.H., 1970. Adsorption of triazine herbicides on soil organic matter, including a short review on soil organic matter chemistry. *Residue Rev.* 32, 131–174. https://doi.org/10.1007/978-1-4615-8464-3_6.
- He, J., Li, H., Rasaily, R.G., Wang, Q., Cai, G., Su, Y., Qiao, X., Liu, L., 2011. Soil properties and crop yields after 11 years of no tillage farming in wheat-maize cropping system in North China plain. *Soil Tillage Res.* 113, 48–54. <https://doi.org/10.1016/j.still.2011.01.005>.
- Hemes, K.S., Chamberlain, S.D., Eichmann, E., Knox, S.H., Baldocchi, D.D., 2018. A biogeochemical compromise: the high methane cost of sequestering carbon in restored wetlands. *Geophys. Res. Lett.* 45, 6081–6091. <https://doi.org/10.1029/2018GL077747>.
- Henry, C.G., Sartori, G.M.S., Gaspar, J.P., Marchesan, E., Hirsh, S.M., Horton, A.P., Espinoza, L., James, H., 2018. Deep tillage and gypsum amendments on fully, deficit irrigated, and dryland soybean. *Agron. J.* 110, 737–748. <https://doi.org/10.2134/agronj2015.11.0567>.
- Hess, L.J.T., Hinckley, E.L.S., Robertson, G.P., Matson, P.A., 2020. Rainfall intensification increases nitrate leaching from tilled but not no-till cropping systems in the U.S. Midwest. *Agric. Ecosyst. Environ.* 290, 106747. <https://doi.org/10.1016/j.agee.2019.106747>.
- Horák, J., Igaz, D., Aydin, E., Šimanský, V., Buchkina, N., Balashov, E., 2020. Changes in direct CO₂ and N₂O emissions from a loam haplic luvisol under conventional moldboard and reduced tillage during growing season and post-harvest period of red clover. *J. Hydrol. Hydromech.* 68, 271–278. <https://doi.org/10.2478/johb-2020-0023>.
- Horowitz, J., Ebel, R., Ueda, K., 2012. “No-till” farming is a growing practice. *Horizons Crop Prod. Precip. Agric. No-Till Farming* (No. 1476-2016-120976).
- Huang, M., Zhou, X., Cao, F., Zou, Y., 2016. Long-term effect of no-tillage on soil organic carbon and nitrogen in an irrigated rice-based cropping system. *Paddy Water Environ.* 14, 367–371. <https://doi.org/10.1007/s10333-015-0506-y>.
- Huang, B., Li, Z., Li, D., Yuan, Z., Chen, Z., Huang, J., 2017. Distribution characteristics of heavy metal(loid)s in aggregates of different size fractions along contaminated paddy soil profile. *Environ. Sci. Pollut. Res.* 24, 23939–23952. <https://doi.org/10.1007/s11356-017-0012-4>.
- Huang, Y., Ren, W., Wang, L., Hui, D., Grove, J.H., Yang, X., Tao, B., Goff, B., 2018. Greenhouse gas emissions and crop yield in no-tillage systems: a meta-analysis. *Agric. Ecosyst. Environ.* 268, 144–153. <https://doi.org/10.1016/j.agee.2018.09.002>.
- Hussein, M.A., Muche, H., Schmitter, P., Nakawuka, P., Tilahun, S.A., Langan, S., Barron, J., Steenhuis, T.S., 2019. Deep tillage improves degraded soils in the (sub) humid Ethiopian highlands. *Land* 8, 159. <https://doi.org/10.3390/land8110159>.
- Issaka, S., Ashraf, M.A., 2017. Impact of soil erosion and degradation on water quality: a review. *Geol. Ecol. Landscapes* 1, 1–11. <https://doi.org/10.1080/24749508.2017.1301053>.
- Jaskulska, I., Jaskulski, D., 2020. Strip-till one-pass technology in central and eastern Europe: a MZURI pro-till hybrid machine case study. *Agronomy* 10, 925. <https://doi.org/10.3390/agronomy10070925>.
- Jha, P., Hati, K.M., Dalal, R.C., Dang, Y.P., Kopitke, P.M., Menzies, N.W., 2020. Soil carbon and nitrogen dynamics in a vertisol following 50 years of no-tillage, crop stubble retention and nitrogen fertilization. *Geoderma* 358, 113996. <https://doi.org/10.1016/j.geoderma.2019.113996>.
- Jones, C.M., Graf, D.R.H., Bru, D., Philippot, L., Hallin, S., 2013. The unaccounted yet abundant nitrous oxide-reducing microbial community: a potential nitrous oxide sink. *ISME J.* 7, 417–426. <https://doi.org/10.1038/ismej.2012.125>.
- Kan, Z.R., Ma, S.T., Liu, Q.Y., Liu, B.Y., Virk, A.L., Qi, J.Y., Zhao, X., Lal, R., Zhang, H.L., 2020. Carbon sequestration and mineralization in soil aggregates under long-term conservation tillage in the North China plain. *Catena* 188, 104428. <https://doi.org/10.1016/j.catena.2019.104428>.
- Kaur, P., Paawan, Kaur, Pervinder, Singh, K., Kaur, M., 2016. Adsorption and desorption characteristics of pretilachlor in three soils of Punjab. *Water Air Soil Pollut.* 227, 1–10. <https://doi.org/10.1007/s11270-016-3074-x>.
- van Kessel, C., Venterea, R., Six, J., Adviento-Borbe, M.A., Linquist, B., van Groenigen, K.J., 2013. Climate, duration, and N placement determine N₂O emissions in reduced tillage systems: a meta-analysis. *Glob. Chang. Biol.* 19, 33–44. <https://doi.org/10.1111/j.1365-2486.2012.02779.x>.
- Khan, S., Shah, A., Nawaz, M., Khan, M., 2017. Impact of different tillage practices on soil physical properties, nitrate leaching and yield attributes of maize (Zea mays L.). *J. Soil Sci. Plant Nutr.* 17, 240–252. <https://doi.org/10.4067/S0718-95162017005000019>.
- Khan, Z.I., Ahmad, K., Rehman, S., Siddique, S., Bashir, H., Zafar, A., Sohail, M., Ali, S.A., Cazzato, E., De Mastro, G., 2017. Health risk assessment of heavy metals in wheat

- using different water qualities: implication for human health. *Environ. Sci. Pollut. Res.* 24, 947–955. <https://doi.org/10.1007/s11356-016-7865-9>.
- Kim, S.Y., Gutierrez, J., Kim, P.J., 2016. Unexpected stimulation of CH₄ emissions under continuous no-tillage system in mono-rice paddy soils during cultivation. *Geoderma* 267, 34–40. <https://doi.org/10.1016/j.geoderma.2015.12.021>.
- Kladivko, E.J., 2001. Tillage systems and soil ecology. *Soil Tillage Res.* 61, 61–76. [https://doi.org/10.1016/S0167-1987\(01\)00179-9](https://doi.org/10.1016/S0167-1987(01)00179-9).
- Klik, A., Rosner, J., 2020. Long-term experience with conservation tillage practices in Austria: impacts on soil erosion processes. *Soil Tillage Res.* 203, 104669. <https://doi.org/10.1016/j.still.2020.104669>.
- Kravchenko, A.N., Wang, A.N.W., Smucker, A.J.M., Rivers, M.L., 2011. Long-term differences in tillage and land use affect intra-aggregate pore heterogeneity. *Soil Sci. Soc. Am. J.* 75, 1658–1666. <https://doi.org/10.2136/sssaj2011.0096>.
- Kremer, R.J., 2018. Pesticide use and biodiversity in soils, burleigh douds series in agricultural. *Science*, 247–263 <https://doi.org/10.19103/as.2017.0033.29>.
- Krupenikov, I.A., Boincean, B.P., Dent, D., 2011. Humus – Guardian of fertility and global carbon sink. *The Black Earth*. Springer, Dordrecht, pp. 39–50 https://doi.org/10.1007/978-94-007-0159-5_7.
- Kumar, N., Nath, C.P., 2019. Impact of zero-till residue management and crop diversification with legumes on soil aggregation and carbon sequestration. *Soil Tillage Res.* 189, 158–167. <https://doi.org/10.1016/j.still.2019.02.001>.
- Lal, R., 2005. Soil erosion and carbon dynamics. *Soil Tillage Res.* <https://doi.org/10.1016/j.still.2004.09.002>.
- Lal, R., 2015. Restoring soil quality to mitigate soil degradation. *Sustain.* 7, 5875–5895. <https://doi.org/10.3390/su7055875>.
- Lal, R., 2019. Accelerated soil erosion as a source of atmospheric CO₂. *Soil Tillage Res.* 188, 35–40. <https://doi.org/10.1016/j.still.2018.02.001>.
- Laufer, D., Koch, H.J., 2017. Growth and yield formation of sugar beet (*Beta vulgaris* L.) under strip tillage compared to full width tillage on silt loam soil in Central Europe. *Eur. J. Agron.* 82, 182–189. <https://doi.org/10.1016/j.eja.2016.10.017>.
- Lembaid, I., Moussadek, R., Mrabet, R., Douaik, A., Bouhaouss, A., 2021. Modeling the effects of farming management practices on soil organic carbon stock under two tillage practices in a semi-arid region, Morocco. *Heliyon* 7, 5889. <https://doi.org/10.1016/j.heliyon.2020.e05889>.
- Li, C., Frolking, S., Butterbach-Bahl, K., 2005. Carbon sequestration in arable soils is likely to increase nitrous oxide emissions, offsetting reductions in climate radiative forcing. *Clim. Chang.* 72, 321–338. <https://doi.org/10.1007/s10584-005-6791-5>.
- Li, C.F., Kou, Z.K., Yang, J.H., Cai, M.L., Wang, J.P., Cao, C.G., 2010. Soil CO₂ fluxes from direct seeding rice fields under two tillage practices in Central China. *Atmos. Environ.* 44, 2696–2704. <https://doi.org/10.1016/j.atmosenv.2010.04.045>.
- Li, C., Zhang, Z., Guo, L., Cai, M., Cao, C., 2013. Emissions of CH₄ and CO₂ from double rice cropping systems under varying tillage and seeding methods. *Atmos. Environ.* 80, 438–444. <https://doi.org/10.1016/j.atmosenv.2013.08.027>.
- Li, L., Xu, J., Hu, J., Han, J., 2014. Reducing nitrous oxide emissions to mitigate climate change and protect the ozone layer. *Environ. Sci. Technol.* 48, 5290–5297. <https://doi.org/10.1021/es404728s>.
- Li, S., Jiang, X., Wang, X., Wright, A.L., 2015. Tillage effects on soil nitrification and the dynamic changes in nitrifying microorganisms in a subtropical rice-based ecosystem: a long-term field study. *Soil Tillage Res.* 150, 132–138. <https://doi.org/10.1016/j.still.2015.02.005>.
- Li, N., Zhou, C., Sun, X., Jing, J., Tian, X., Wang, L., 2018. Effects of ridge tillage and mulching on water availability, grain yield, and water use efficiency in rain-fed winter wheat under different rainfall and nitrogen conditions. *Soil Tillage Res.* 179, 86–95. <https://doi.org/10.1016/j.still.2018.01.003>.
- Li, Y., Are, K.S., Qin, Z., Huang, Z., Abegunrin, T.P., Houssou, A.A., Guo, H., Gu, M., Wei, L., 2020. Farmland size increase significantly accelerates road surface fill erosion and nutrient losses in southern subtropics of China. *Soil Tillage Res.* 204, 104689. <https://doi.org/10.1016/j.still.2020.104689>.
- Li, S., Hao, Guo, L., Jin, Cao, C., Gui, Li, C., Fang, 2021. Integrated assessment of carbon footprint, energy budget and net ecosystem economic efficiency from rice fields under different tillage modes in central China. *J. Clean. Prod.* 295, 126398. <https://doi.org/10.1016/j.jclepro.2021.126398>.
- Li, Y., Li, Z., Cui, S., Liang, G., Zhang, Q., 2021. Microbial-derived carbon components are critical for enhancing soil organic carbon in no-tillage croplands: a global perspective. *Soil Tillage Res.* 205, 104758. <https://doi.org/10.1016/j.still.2020.104758>.
- Licht, M.A., Al-Kaisi, M., 2005. Strip-tillage effect on seedbed soil temperature and other soil physical properties. *Soil Tillage Res.* 80, 233–249. <https://doi.org/10.1016/j.still.2004.03.017>.
- Lin, S., Hernandez-Ramirez, G., 2020. Nitrous oxide emissions from manured soils as a function of various nitrification inhibitor rates and soil moisture contents. *Sci. Total Environ.* 738, 139669. <https://doi.org/10.1016/j.scitotenv.2020.139669>.
- Liu, S., Zhao, C., Zhang, Y., Hu, Z., Wang, C., Zong, Y., Zhang, L., Zou, J., 2015. Annual net greenhouse gas balance in a halophyte (*Helianthus tuberosus*) bioenergy cropping system under various soil practices in Southeast China. *GCB Bioenergy* 7, 690–703. <https://doi.org/10.1111/gcbb.12185>.
- Liu, T., Liang, Y., Chu, G., 2017. Nitrapyrin addition mitigates nitrous oxide emissions and raises nitrogen use efficiency in plastic-film-mulched drip-fertigated cotton field. *PLoS One* 12, 176305. <https://doi.org/10.1371/journal.pone.0176305>.
- Liu, S., Zhang, X., Liang, A., Zhang, J., Müller, C., Cai, Z., 2018. Ridge tillage is likely better than no tillage for 14-year field experiment in black soils: insights from a 15N-tracing study. *Soil Tillage Res.* 179, 38–46. <https://doi.org/10.1016/j.still.2018.01.011>.
- Liu, L., Estiarte, M., Peñuelas, J., 2019. Soil moisture as the key factor of atmospheric CH₄ uptake in forest soils under environmental change. *Geoderma* 355, 113920. <https://doi.org/10.1016/j.geoderma.2019.113920>.
- Lognoul, M., Theodorakopoulos, N., Hiel, M.P., Regaert, D., Broux, F., Heinesch, B., Bodson, B., Vandenberg, M., Aubinet, M., 2017. Impact of tillage on greenhouse gas emissions by an agricultural crop and dynamics of N₂O fluxes: insights from automated closed chamber measurements. *Soil Tillage Res.* 167, 80–89. <https://doi.org/10.1016/j.still.2016.11.008>.
- Lu, C., Tian, H., 2017. Global nitrogen and phosphorus fertilizer use for agriculture production in the past half century: shifted hot spots and nutrient imbalance. *Earth Syst. Sci. Data* 9, 181–192. <https://doi.org/10.5194/essd-9-181-2017>.
- Lu, Xingli, Lu, Xingneng, Tanveer, S.K., Wen, X., Liao, Y., 2016. Effects of tillage management on soil CO₂ emission and wheat yield under rain-fed conditions. *Soil Res.* 54, 38–48. <https://doi.org/10.1071/SR14300>.
- Luo, Z., Wang, E., Sun, O.J., 2010. Can no-tillage stimulate carbon sequestration in agricultural soils? A meta-analysis of paired experiments. *Agric. Ecosyst. Environ.* 139, 224–231. <https://doi.org/10.1016/j.agee.2010.08.006>.
- Luo, Y., Benke, M.B., Hao, X., 2017. Nutrient uptake and leaching from soil amended with cattle manure and nitrapyrin. *Commun. Soil Sci. Plant Anal.* 48, 1438–1454. <https://doi.org/10.1080/00103624.2017.1373786>.
- Manlay, R.J., Feller, C., Swift, M.J., 2007. Historical evolution of soil organic matter concepts and their relationships with the fertility and sustainability of cropping systems. *Agric. Ecosyst. Environ.* 119, 217–233. <https://doi.org/10.1016/j.agee.2006.07.011>.
- Marafon, G., Barbosa, R.S., de Jesus Lacerda, J.J., Martins, V., Costa, O.S., e Silva, J.de D.F., 2020. C and P pool restoration by a no-tillage system on Brazilian Cerrado Oxisol in Piauí State. *Environ. Monit. Assess.* 192, 254. <https://doi.org/10.1007/s10661-020-8221-6>.
- Mazzoncini, M., Antichi, D., Di Bene, C., Risaliti, R., Petri, M., Bonari, E., 2016. Soil carbon and nitrogen changes after 28 years of no-tillage management under Mediterranean conditions. *Eur. J. Agron.* 77, 156–165. <https://doi.org/10.1016/j.eja.2016.02.011>.
- Mehra, P., Baker, J., Sojka, R.E., Bolan, N., Desbiolles, J., Kirkham, M.B., Ross, C., Gupta, R., 2018. A review of tillage practices and their potential to impact the soil carbon dynamics. *Adv. Agron.*, 185–230 <https://doi.org/10.1016/bs.agron.2018.03.002>.
- Mei, K., Wang, Z., Huang, H., Zhang, C., Shang, X., Dahlgren, R.A., Zhang, M., Xia, F., 2018. Stimulation of N₂O emission by conservation tillage management in agricultural lands: a meta-analysis. *Soil Tillage Res.* 182, 86–93. <https://doi.org/10.1016/j.still.2018.05.006>.
- Melland, A.R., Antille, D.L., Dang, Y.P., 2017. Effects of strategic tillage on short-term erosion, nutrient loss in runoff and greenhouse gas emissions. *Soil Res.* 55, 201–214. <https://doi.org/10.1071/SR16136>.
- Meng, X., Li, Y., Yao, H., Wang, J., Dai, F., Wu, Y., Chapman, S., 2020. Nitrification and urease inhibitors improve rice nitrogen uptake and prevent denitrification in alkaline paddy soil. *Appl. Soil Ecol.* 154, 103665. <https://doi.org/10.1016/j.apsoil.2020.103665>.
- Meurer, K.H.E., Haddaway, N.R., Bolinder, M.A., Kätterer, T., 2018. Tillage intensity affects total SOC stocks in boreo-temperate regions only in the topsoil—a systematic review using an ESM approach. *EarthSci. Rev.* 177, 613–622. <https://doi.org/10.1016/j.earscirev.2017.12.015>.
- Modak, K., Biswas, D.R., Ghosh, A., Pramanik, P., Das, T.K., Das, S., Kumar, S., Krishnan, P., Bhattacharyya, R., 2020. Zero tillage and residue retention impact on soil aggregation and carbon stabilization within aggregates in subtropical India. *Soil Tillage Res.* 202, 104649. <https://doi.org/10.1016/j.still.2020.104649>.
- Moreno-Casasola, P., Hernández, M.E., Campos, A., 2017. Hydrology, soil carbon sequestration and water retention along a coastal wetland gradient in the Alvarado Lagoon System, Veracruz, Mexico. *J. Coast. Res.*, 104–115 <https://doi.org/10.2112/SI77-011.1> 2017-Sprin.
- Mottes, C., Deffontaines, L., Charlier, J.B., Comte, I., Della Rossa, P., Lesueur-Jannoyer, M., Woignier, T., Adele, G., Tailame, A.L., Arnaud, L., Plet, J., Rangon, L., Bricquet, J.P., Cattani, P., 2020. Spatio-temporal variability of water pollution by chlordecone at the watershed scale: what insights for the management of polluted territories? *Environ. Sci. Pollut. Res.* 27, 40999–41013. <https://doi.org/10.1007/s11356-019-06247-y>.
- Nath, C.P., Das, T.K., Rana, K.S., Bhattacharyya, R., Pathak, H., Paul, S., Meena, M.C., Singh, S.B., 2017. Greenhouse gases emission, soil organic carbon and wheat yield as affected by tillage systems and nitrogen management practices. *Arch. Agron. Soil Sci.* 63, 1644–1660. <https://doi.org/10.1080/03650340.2017.1300657>.
- Negassa, W., Price, R.F., Basir, A., Snapp, S.S., Kravchenko, A., 2015. Cover crop and tillage systems effect on soil CO₂ and N₂O fluxes in contrasting topographic positions. *Soil Tillage Res.* 154, 64–74. <https://doi.org/10.1016/j.still.2015.06.015>.
- Nevens, C.J., Lacey, C., Armstrong, S., 2020. The synchrony of cover crop decomposition, enzyme activity, and nitrogen availability in a corn agroecosystem in the Midwest United States. *Soil Tillage Res.* 197, 104518. <https://doi.org/10.1016/j.still.2019.104518>.
- Niu, Y., Cai, Y., Chen, Z., Luo, J., Di, H.J., Yu, H., Zhu, A., Ding, W., 2019. No-tillage did not increase organic carbon storage but stimulated N₂O emissions in an intensively cultivated sandy loam soil: a negative climate effect. *Soil Tillage Res.* 195, 104419. <https://doi.org/10.1016/j.still.2019.104419>.
- Novara, A., Stallone, G., Cerdà, A., Gristina, L., 2019. The effect of shallow tillage on soil erosion in a semi-arid vineyard. *Agronomy* 9, 257. <https://doi.org/10.3390/agronomy9050257>.
- Nyamadzawo, G., Wuta, M., Chirinda, N., Mujuru, L., Smith, J.L., 2013. Greenhouse gas emissions from intermittently flooded (Dambo) Rice under different tillage practices in chiota smallholder farming area of Zimbabwe. *Atmos. Clim. Sci.* 03, 13–20. <https://doi.org/10.4236/acs.2013.34a003>.
- Obalum, S.E., Chibuike, G.U., Peth, S., Ouyang, Y., 2017. Soil organic matter as sole indicator of soil degradation. *Environ. Monit. Assess.* 189, 176. <https://doi.org/10.1007/s10661-017-5881-y>.
- Ogle, S.M., Breidt, F.J., Paustian, K., 2005. Agricultural management impacts on soil organic carbon storage under moist and dry climatic conditions of temperate and tropical regions. *Biogeochemistry* 72, 87–121. <https://doi.org/10.1007/s10533-004-0360-2>.
- Okada, E., Costa, J.L., Bedmar, F., 2019. Glyphosate dissipation in different soils under no-till and Conventional tillage. *Pedosphere* 29, 773–783. [https://doi.org/10.1016/S1002-0160\(17\)60430-2](https://doi.org/10.1016/S1002-0160(17)60430-2).

- Olson, K.R., Ebelhar, S.A., Lang, J.M., 2013. Effects of 24 years of conservation tillage systems on soil organic carbon and soil productivity. *Appl. Environ. Soil Sci.* 2013, 1–10. <https://doi.org/10.1155/2013/617504>.
- Oorts, K., Merckx, R., Gréhan, E., Labreuche, J., Nicolardot, B., 2007. Determinants of annual fluxes of CO₂ and N₂O in long-term no-tillage and conventional tillage systems in northern France. *Soil Tillage Res.* 95, 133–148. <https://doi.org/10.1016/j.still.2006.12.002>.
- Pandey, D., Agrawal, M., Bohra, J.S., 2012. Greenhouse gas emissions from rice crop with different tillage permutations in rice-wheat system. *Agric. Ecosyst. Environ.* 159, 133–144. <https://doi.org/10.1016/j.agee.2012.07.008>.
- Parvin, N., Sandin, M., Larsbo, M., 2021. Seedbed consolidation and surface sealing for soils of different texture and soil organic carbon contents. *Soil Tillage Res.* 206, 104849. <https://doi.org/10.1016/j.still.2020.104849>.
- Payá-Pérez, A.B., Cortés, A., Sala, M.N., Larsen, B., 1992. Organic matter fractions controlling the sorption of atrazine in sandy soils. *Chemosphere* 25, 887–898. [https://doi.org/10.1016/0045-6535\(92\)90080-B](https://doi.org/10.1016/0045-6535(92)90080-B).
- Philippot, L., Andert, J., Jones, C.M., Bru, D., Hallin, S., 2011. Importance of denitrifiers lacking the genes encoding the nitrous oxide reductase for N₂O emissions from soil. *Glob. Chang. Biol.* 17, 1497–1504. <https://doi.org/10.1111/j.1365-2486.2010.02334.x>.
- Pihlatie, M., Syväsalo, E., Simojoki, A., Esala, M., Regina, K., 2004. Contribution of nitrification and denitrification to N₂O production in peat, clay and loamy sand soils under different soil moisture conditions. *Nutr. Cycl. Agroecosyst.* 70, 135–141. <https://doi.org/10.1023/B:FRES.0000048475.81211.3c>.
- Pineiro, E.F.M., de Campos, D.V.B., de Carvalho Balieiro, F., dos Anjos, L.H.C., Pereira, M.G., 2015. Tillage systems effects on soil carbon stock and physical fractions of soil organic matter. *Agric. Syst.* 132, 35–39. <https://doi.org/10.1016/j.agry.2014.08.008>.
- Plaza-Bonilla, D., Álvaro-Fuentes, J., Artie, J.L., Cantero-Martínez, C., 2014. Tillage and nitrogen fertilization effects on nitrous oxide yield-scaled emissions in a rainfed Mediterranean area. *Agric. Ecosyst. Environ.* 189, 43–52. <https://doi.org/10.1016/j.agee.2014.03.023>.
- Plaza-Bonilla, D., Álvaro-Fuentes, J., Bareche, J., Pareja-Sánchez, E., Justes, E., Cantero-Martínez, C., 2018. No-tillage reduces long-term yield-scaled soil nitrous oxide emissions in rainfed Mediterranean agroecosystems: a field and modelling approach. *Agric. Ecosyst. Environ.* 262, 36–47. <https://doi.org/10.1016/j.agee.2018.04.007>.
- Poepel, C., Don, A., Vesterdal, L., Leifeld, J., Van Wesemael, B., Schumacher, J., Gensior, A., 2011. Temporal dynamics of soil organic carbon after land-use change in the temperate zone - carbon response functions as a model approach. *Glob. Chang. Biol.* 17, 2415–2427. <https://doi.org/10.1111/j.1365-2486.2011.02408.x>.
- Pöhlitz, J., Rücknagel, J., Koblenz, B., Schlüter, S., Vogel, H.J., Christen, O., 2018. Computed tomography and soil physical measurements of compaction behaviour under strip tillage, mulch tillage and no tillage. *Soil Tillage Res.* 175, 205–216. <https://doi.org/10.1016/j.still.2017.09.007>.
- Porwollik, V., Rolinski, S., Heinke, J., Müller, C., 2019. Generating a rule-based global gridded tillage dataset. *Earth Syst. Sci. Data* 11, 823–843. <https://doi.org/10.5194/essd-11-823-2019>.
- Potter, T.L., Bosch, D.D., Strickland, T.C., 2015. Tillage impact on herbicide loss by surface runoff and lateral subsurface flow. *Sci. Total Environ.* 530–531, 357–366. <https://doi.org/10.1016/j.scitotenv.2015.05.079>.
- Powlson, D.S., Stirling, C.M., Jat, M.L., Gerard, B.G., Palm, C.A., Sanchez, P.A., Cassman, K.G., 2014. Limited potential of no-till agriculture for climate change mitigation. *Nat. Clim. Chang.* 4, 678–683. <https://doi.org/10.1038/nclimate2292>.
- Prado, R.B., Fidalgo, E.C.C., Monteiro, J.M.G., Schuler, A.E., Vezzani, F.M., Garcia, J.R., de Oliveira, A.P., Viana, J.H.M., Reatto, A., Parron, L.M., Donagemma, G.K., Turetta, A.P.D., Simões, M., Gomes, B.da C.C.P., Mendes, I.de C., Clemente, E.de P., 2016. Current overview and potential applications of the soil ecosystem services approach in Brazil. *Pesqui. Agropecu. Bras.* 51, 1021–1038. <https://doi.org/10.1590/S0100-204X2016000900002>.
- Pribyl, D.W., 2010. A critical review of the conventional SOC to SOM conversion factor. *Geoderma* 156, 75–83. <https://doi.org/10.1016/j.geoderma.2010.02.003>.
- Qin, C., Wright, A.L., Ma, L., He, X., Xie, D., Jiang, X., 2020. Improving nitrogen-use efficiency by using ridge tillage in rice paddy soils. *Soil Use Manag.*, 1–9. <https://doi.org/10.1111/sum.12675>.
- Rahmati, M., Eskandari, I., Kouselou, M., Feiziasl, V., Mahdavinia, G.R., Aliasgharzad, N., McKenzie, B.M., 2020. Changes in soil organic carbon fractions and residence time five years after implementing conventional and conservation tillage practices. *Soil Tillage Res.* 200, 104632. <https://doi.org/10.1016/j.still.2020.104632>.
- Raiesi, F., Kabiri, V., 2016. Identification of soil quality indicators for assessing the effect of different tillage practices through a soil quality index in a semi-arid environment. *Ecol. Indic.* 71, 198–207. <https://doi.org/10.1016/j.ecolind.2016.06.061>.
- Ramnarine, R., Voroney, R.P., Dunfield, K.E., Wagner-Riddle, C., 2018. Characterization of the heavy, hydrolysable and non-hydrolysable fractions of soil organic carbon in conventional and no-tillage soils. *Soil Tillage Res.* 181, 144–151. <https://doi.org/10.1016/j.still.2018.04.010>.
- Ravishankara, A.R., Daniel, J.S., Portmann, R.W., 2009. Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the 21st century. *Science* (80-.) 326, 123–125. <https://doi.org/10.1126/science.1176985>.
- Regina, K., Alakukku, L., 2010. Greenhouse gas fluxes in varying soils types under conventional and no-tillage practices. *Soil Tillage Res.* 109, 144–152. <https://doi.org/10.1016/j.still.2010.05.009>.
- Reza, A., Shim, S., Kim, S., Ahn, S., Won, S., Ra, C., 2020. Rational budgeting approach as a nutrient management tool for mixed crop-swine farms in Korea. *Asian-Australasian J. Anim. Sci.* 33, 1520–1532. <https://doi.org/10.5713/ajas.19.0640>.
- Rodríguez, L., Suárez, J.C., Rodríguez, W., Artunduaga, K.J., Lavelle, P., 2021. Agroforestry systems impact soil macroaggregation and enhance carbon storage in colombian deforested Amazonia. *Geoderma* 384, 114810. <https://doi.org/10.1016/j.geoderma.2020.114810>.
- Rutkowska, B., Szulc, W., Sosulski, T., Skowrońska, M., Szczepaniak, J., 2018. Impact of reduced tillage on CO₂ emission from soil under maize cultivation. *Soil Tillage Res.* 180, 21–28. <https://doi.org/10.1016/j.still.2018.02.012>.
- Šarauskis, E., Vaitauskienė, K., Romanekas, K., Jasinskas, A., Butkus, V., Kriaučiūnienė, Z., 2017. Fuel consumption and CO₂ emission analysis in different strip tillage scenarios. *Energy* 118, 957–968. <https://doi.org/10.1016/j.energy.2016.10.121>.
- Sarker, J.R., Singh, B.P., Dougherty, W.J., Fang, Y., Badgery, W., Hoyle, F.C., Dalal, R.C., Cowie, A.L., 2018. Impact of agricultural management practices on the nutrient supply potential of soil organic matter under long-term farming systems. *Soil Tillage Res.* 175, 71–81. <https://doi.org/10.1016/j.still.2017.08.005>.
- Sarker, T.C., Incerti, G., Spaccini, R., Piccolo, A., Mazzoleni, S., Bonanomi, G., 2018. Linking organic matter chemistry with soil aggregate stability: insight from 13C NMR spectroscopy. *Soil Biol. Biochem.* 117, 175–184. <https://doi.org/10.1016/j.soilbio.2017.11.011>.
- Scanlan, C.A., Davies, S.L., 2019. Soil mixing and redistribution by strategic deep tillage in a sandy soil. *Soil Tillage Res.* 185, 139–145. <https://doi.org/10.1016/j.still.2018.09.008>.
- Schlüter, S., Großmann, C., Diel, J., Wu, G.M., Tischer, S., Deubel, A., Rücknagel, J., 2018. Long-term effects of conventional and reduced tillage on soil structure, soil ecological and soil hydraulic properties. *Geoderma* 332, 10–19. <https://doi.org/10.1016/j.geoderma.2018.07.001>.
- Schmidt, M.W.L., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P., Weiner, S., Trumbore, S.E., 2011. Persistence of soil organic matter as an ecosystem property. *Nature* 478, 49–56. <https://doi.org/10.1038/nature10386>.
- Schneider, F., Don, A., Hennings, I., Schmittmann, O., Seidel, S.J., 2017. The effect of deep tillage on crop yield – what do we really know? *Soil Tillage Res.* <https://doi.org/10.1016/j.still.2017.07.005>.
- Shakoor, A., Shahbaz, M., Farooq, T.H., Sahar, N.E., Shahzad, S.M., Altaf, M.M., Ashraf, M., 2021. A global meta-analysis of greenhouse gases emission and crop yield under no-tillage as compared to conventional tillage. *Sci. Total Environ.* 750, 142299. <https://doi.org/10.1016/j.scitotenv.2020.142299>.
- Shang, Q., Cheng, C., Wang, J., Luo, K., Zeng, Y., Yang, X., 2021. Net global warming potential, greenhouse gas intensity and carbon footprint as affected by different tillage systems from chinese double-cropping paddy fields. *Soil Tillage Res.* 209. <https://doi.org/10.1016/j.still.2021.104947>.
- Shipitalo, M.J., Owens, L.B., 2011. Comparative losses of glyphosate and selected residual herbicides in surface runoff from conservation-tilled watersheds planted with corn or soybean. *J. Environ. Qual.* 40, 1281–1289. <https://doi.org/10.2134/jeq2010.0454>.
- da Silva, K.A., Nicola, V.B., Dudas, R.T., Demetrio, W.C., Cunha, L., Bartz, M.L.C., Brown, G.G., Pasini, A., Kille, P., Ferreira, N.G.C., de Oliveira, C.M.R., Maia, L.dos S., 2021. Pesticides in a case study on no-tillage farming systems and surrounding forest patches in Brazil. *Sci. Rep.* 11, 9839. <https://doi.org/10.1038/s41598-021-88779-3>.
- Sithole, N.J., Magwaza, L.S., Thibaud, G.R., 2019. Long-term impact of no-till conservation agriculture and N-fertilizer on soil aggregate stability, infiltration and distribution of C in different size fractions. *Soil Tillage Res.* 190, 147–156. <https://doi.org/10.1016/j.still.2019.03.004>.
- Six, J., Elliott, E.T., Paustian, K., 2000. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biol. Biochem.* 32, 2099–2103. [https://doi.org/10.1016/S0038-0717\(00\)00179-6](https://doi.org/10.1016/S0038-0717(00)00179-6).
- Six, J., Conant, R.T., Paul, E.A., Paustian, K., 2002. Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant Soil* 241, 155–176. <https://doi.org/10.1023/A:1016125726789>.
- Snyder, C.S., Bruulsema, T.W., Jensen, T.L., Fixen, P.E., 2009. Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agric. Ecosyst. Environ.* 133, 247–266. <https://doi.org/10.1016/j.agee.2009.04.021>.
- Soane, B.D., Ball, B.C., Arvidsson, J., Basch, G., Moreno, F., Roger-Estrade, J., 2012. No-till in northern, western and south-western Europe: a review of problems and opportunities for crop production and the environment. *Soil Tillage Res.* 118, 66–87. <https://doi.org/10.1016/j.still.2011.10.015>.
- Sørensen, P., Amato, M., 2002. Remineralisation and residual effects of N after application of pig slurry to soil. *Eur. J. Agron.* 16, 81–95. [https://doi.org/10.1016/S1161-0301\(01\)00119-8](https://doi.org/10.1016/S1161-0301(01)00119-8).
- Sørensen, C.G., Halberg, N., Oudshoorn, F.W., Petersen, B.M., Dalgaard, R., 2014. Energy inputs and GHG emissions of tillage systems. *Biosyst. Eng.* 120, 2–14. <https://doi.org/10.1016/j.biosystemseng.2014.01.004>.
- Spies, E., Humphrys, C., Richner, W., Schneider, M.K., Piepho, H.P., Chervet, A., Prasuhn, V., 2020. Does no-tillage decrease nitrate leaching compared to ploughing under a long-term crop rotation in Switzerland? *Soil Tillage Res.* 199, 104590. <https://doi.org/10.1016/j.still.2020.104590>.
- Stevenson, F.J., 1972. Organic matter reactions involving herbicides in soil. *J. Environ. Qual.* 1, 333–343. <https://doi.org/10.2134/jeq1972.00472425000100040001x>.
- Stockmann, U., Adams, M.A., Crawford, J.W., Field, D.J., Henakaarchchi, N., Jenkins, M., Minasny, B., McBratney, A.B., Singh, K., Wheeler, I., Abbott, L., Angers, D.A., Baldock, J., Bird, M., Brookes, P.C., Chenu, C., Jastrow, J.D., Lal, R., Lehmann, J., O'Donnell, A.G., Parton, W.J., Whitehead, D., Zimmermann, M., Courcelles, V.de R.de, 2013. The knowns, known unknowns and unknowns of sequestration of soil organic carbon. *Agric. Ecosyst. Environ.* 164, 80–99. <https://doi.org/10.1016/j.agee.2012.10.001> (No. 1476-2016-120976).
- Subbarao, G.V., Arango, J., Masahiro, K., Hooper, A.M., Yoshihashi, T., Ando, Y., Nakahara, K., Deshpande, S., Ortiz-Monasterio, I., Ishitani, M., Peters, M., Chirinda, N., Wollenberg, L., Lata, J.C., Gerard, B., Tobita, S., Rao, I.M., Braun, H.J., Kommerell, V., Tohme, J., Iwanaga, M., 2017. Genetic mitigation strategies to tackle agricultural GHG emissions: the case for biological nitrification inhibition technology. *Plant Sci.* 262, 165–168. <https://doi.org/10.1016/j.plantsci.2017.05.004>.
- Sun, B., Zhang, L., Yang, L., Zhang, F., Norse, D., Zhu, Z., 2012. Agricultural non-point source pollution in China: causes and mitigation measures. *Ambio* 41, 370–379. <https://doi.org/10.1007/s13280-012-0249-6>.
- Syswerda, S.P., Basso, B., Hamilton, S.K., Tausig, J.B., Robertson, G.P., 2012. Long-term nitrate loss along an agricultural intensity gradient in the Upper Midwest USA. *Agric. Ecosyst. Environ.* 149, 10–19. <https://doi.org/10.1016/j.agee.2011.12.007>.

- Tian, S., Ning, T., Wang, Y., Liu, Z., Li, G., Li, Z., Lal, R., 2016. Crop yield and soil carbon responses to tillage method changes in North China. *Soil Tillage Res.* 163, 207–213. <https://doi.org/10.1016/j.still.2016.06.005>.
- Tuzzin de Moraes, M., Debiassi, H., Carlesso, R., Cezar Franchini, J., Rodrigues da Silva, V., Bonini da Luz, F., 2016. Soil physical quality on tillage and cropping systems after two decades in the subtropical region of Brazil. *Soil Tillage Res.* 155, 351–362. <https://doi.org/10.1016/j.still.2015.07.015>.
- Van den Putte, A., Govers, G., Diels, J., Gillijns, K., Demuzere, M., 2010. Assessing the effect of soil tillage on crop growth: a meta-regression analysis on European crop yields under conservation agriculture. *Eur. J. Agron.* 33, 231–241. <https://doi.org/10.1016/j.eja.2010.05.008>.
- Vitousek, P.M., Naylor, R., Crews, T., David, M.B., Drinkwater, L.E., Holland, E., Johnes, P.J., Katzenberger, J., Martinelli, L.A., Matson, P.A., Nziguheba, G., Ojima, D., Palm, C.A., Robertson, G.P., Sanchez, P.A., Townsend, A.R., Zhang, F.S., 2009. Nutrient imbalances in agricultural development. *Science* (80-.) 324, 1519–1520. <https://doi.org/10.1126/science.1170261>.
- Waddell, J.T., Weil, R.R., 2006. Effects of fertilizer placement on solute leaching under ridge tillage and no tillage. *Soil Tillage Res.* 90, 194–204. <https://doi.org/10.1016/j.still.2005.09.002>.
- Wang, J., Zou, J., 2020. No-till increases soil denitrification via its positive effects on the activity and abundance of the denitrifying community. *Soil Biol. Biochem.* 142, 107706. <https://doi.org/10.1016/j.soilbio.2020.107706>.
- Wang, X., Qi, J.Y., Zhang, X.Z., Li, S.S., Latif Virk, A., Zhao, X., Xiao, X.P., Zhang, H.L., 2019. Effects of tillage and residue management on soil aggregates and associated carbon storage in a double paddy cropping system. *Soil Tillage Res.* 194, 104339. <https://doi.org/10.1016/j.still.2019.104339>.
- Wang, X., Jing, Z.H., He, C., Liu, Q.Y., Qi, J.Y., Zhao, X., Xiao, X.P., Zhang, H.L., 2021. Temporal variation of SOC storage and crop yield and its relationship - a fourteen year field trial about tillage practices in a double paddy cropping system, China. *Sci. Total Environ.* 759. <https://doi.org/10.1016/j.scitotenv.2020.143494>.
- Williams, M.R., King, K.W., Duncan, E.W., Pease, L.A., Penn, C.J., 2018. Fertilizer placement and tillage effects on phosphorus concentration in leachate from fine-textured soils. *Soil Tillage Res.* 178, 130–138. <https://doi.org/10.1016/j.still.2017.12.010>.
- Wulannityas, H.S., Gong, Y., Li, P., Sakagami, N., Nishiwaki, J., Komatsuzaki, M., 2021. A cover crop and no-tillage system for enhancing soil health by increasing soil organic matter in soybean cultivation. *Soil Tillage Res.* 205, 104749. <https://doi.org/10.1016/j.still.2020.104749>.
- Xiao, L., Zhou, S., Zhao, R., Greenwood, P., Kuhn, N.J., 2020. Evaluating soil organic carbon stock changes induced by no-tillage based on fixed depth and equivalent soil mass approaches. *Agric. Ecosyst. Environ.* 300, 106982. <https://doi.org/10.1016/j.agee.2020.106982>.
- Yang, X., Bento, C.P.M., Chen, H., Zhang, H., Xue, S., Lwanga, E.H., Zomer, P., Ritsema, C.J., Geissen, V., 2018. Influence of microplastic addition on glyphosate decay and soil microbial activities in Chinese loess soil. *Environ. Pollut.* 242, 338–347. <https://doi.org/10.1016/j.envpol.2018.07.006>.
- Ye, Y., Xiao, S., Liu, S., Zhang, W., Zhao, J., Chen, H., Guggenberger, G., Wang, K., 2020. Tillage induces rapid loss of organic carbon in large macroaggregates of calcareous soils. *Soil Tillage Res.* 199, 104549. <https://doi.org/10.1016/j.still.2019.104549>.
- Yu, K., Faulkner, S.P., Baldwin, M.J., 2008. Effect of hydrological conditions on nitrous oxide, methane, and carbon dioxide dynamics in a bottomland hardwood forest and its implication for soil carbon sequestration. *Glob. Chang. Biol.* 14, 798–812. <https://doi.org/10.1111/j.1365-2486.2008.01545.x>.
- Yu, Z., Lu, C., Hennessy, D.A., Feng, H., Tian, H., 2020. Impacts of tillage practices on soil carbon stocks in the US corn-soybean cropping system during 1998 to 2016. *Environ. Res. Lett.* 15, 14008. <https://doi.org/10.1088/1748-9326/ab6393>.
- Zhai, Lichao, Xu, P., Zhang, Z., Li, S., Xie, R., Zhai, Lifang, Wei, B., 2017. Effects of deep vertical rotary tillage on dry matter accumulation and grain yield of summer maize in the Huang-Huai-Hai plain of China. *Soil Tillage Res.* 170, 167–174. <https://doi.org/10.1016/j.still.2017.03.013>.
- Zhang, H.L., Bai, X.L., Xue, J.F., Chen, Z.D., Tang, H.M., Chen, F., 2013. Emissions of CH₄ and N₂O under different tillage systems from double-cropped Paddy fields in southern China. *PLoS One* 8, 65277. <https://doi.org/10.1371/journal.pone.0065277>.
- Zhang, Z.S., Cao, C.G., Guo, L.J., Li, C.F., 2016. Emissions of CH₄ and CO₂ from paddy fields as affected by tillage practices and crop residues in Central China. *Paddy Water Environ.* 14, 85–92. <https://doi.org/10.1007/s10333-015-0480-4>.
- Zhang, Z.S., Chen, J., Liu, T.Q., Cao, C.G., Li, C.F., 2016. Effects of nitrogen fertilizer sources and tillage practices on greenhouse gas emissions in paddy fields of Central China. *Atmos. Environ.* 144, 274–281. <https://doi.org/10.1016/j.atmosenv.2016.09.003>.
- Zhang, Yujiao, Wang, S., Wang, H., Ning, F., Zhang, Yuanhong, Dong, Z., Wen, P., Wang, R., Wang, X., Li, J., 2018. The effects of rotating conservation tillage with conventional tillage on soil properties and grain yields in winter wheat-spring maize rotations. *Agric. For. Meteorol.* 263, 107–117. <https://doi.org/10.1016/j.agrformet.2018.08.012>.
- Zhou, B., Sun, X., Ding, Z., Ma, W., Zhao, M., 2017. Effect of tillage practice and fertilization on dry matter accumulation and grain yield of summer maize (zea mays l.). *Sci. Agric. Sin.* 50, 2129–2140. <https://doi.org/10.3864/j.issn.0578-1752.2017.11.018>.
- Zomer, R.J., Bossio, D.A., Sommer, R., Verchot, L.V., 2017. Global sequestration potential of increased organic carbon in cropland soils. *Sci. Rep.* 7, 15554. <https://doi.org/10.1038/s41598-017-15794-8>.