

MINIREVIEW – Biotechnology & Synthetic Biology

The impact of environmental factors on carbon dioxide fixation by microalgae

Marcia Morales^{1,*}, León Sánchez² and Sergio Revah¹

¹Departamento de Procesos y Tecnología, Universidad Autónoma Metropolitana-Cuajimalpa, Av. Vasco de Quiroga 4871, colonia Santa Fe Cuajimalpa, CP 05300, Ciudad de México, Mexico and ²Doctorado en Biotecnología, Universidad Autónoma Metropolitana-Iztapalapa, San Rafael Atlixco 186, CP 09340, Ciudad de México, Mexico

*Corresponding author: Departamento de Procesos y Tecnología, Universidad Autónoma Metropolitana-Cuajimalpa, Av. Vasco de Quiroga 4871, colonia Santa Fe Cuajimalpa, CP 05300, Ciudad de México, Mexico. Tel.: +52-55-58146500 (ext. 3872); E-mail: mmorales@correo.cua.uam.mx

One sentence summary: Cultivation conditions have a direct impact on CO₂ fixation by microalgae.

Editor: Michael Sauer

ABSTRACT

Microalgae are among the most productive biological systems for converting sunlight into chemical energy, which is used to capture and transform inorganic carbon into biomass. The efficiency of carbon dioxide capture depends on the cultivation system configuration (photobioreactors or open systems) and can vary according to the state of the algal physiology, the chemical composition of the nutrient medium, and environmental factors such as irradiance, temperature and pH. This mini-review is focused on some of the most important environmental factors determining photosynthetic activity, carbon dioxide biofixation, cell growth rate and biomass productivity by microalgae. These include carbon dioxide and O₂ concentrations, light intensity, cultivation temperature and nutrients. Finally, a review of the operation of microalgal cultivation systems outdoors is presented as an example of the impact of environmental conditions on biomass productivity and carbon dioxide fixation.

Keywords: carbon dioxide biofixation; microalgae; outdoor cultivations; pH; temperature; irradiance

INTRODUCTION

The accumulation of greenhouse gases in the atmosphere, from human activities and industrialization, has been considered as the main driver of climate change and is recognized as being among the major global problems facing mankind (IPCC 2014). While strong steps are being taken to reduce these emissions, carbon dioxide capture is a viable alternative (Mikkelsen, Jørgensen and Krebs 2010). A microalgal contribution to carbon dioxide capture was suggested in the 1960s by Oswald and Golueke (1960) and much attention has been drawn due to the higher solar energy yield and environmental tolerance of microalgae compared with higher plants, and to the possibility of year-round cultivation (Chisti 2007; John et al. 2011). Microalgae

can also contribute to reducing global warming emissions by recycling carbon dioxide into biofuels (Chisti 2007), or through production of value-added products from flue gases. Furthermore, microalgae can also reduce carbon dioxide emissions from wastewater treatment by lowering the energy needed for aeration (Razzak et al. 2013), or during biogas upgrading (Meier et al. 2015). Nevertheless, worldwide carbon dioxide emissions (≈40 Gt year⁻¹, IPCC 2014) are too high compared with 13 600 t (IEA 2017) of commercialized microalgal biomass (around 27 200 t of carbon dioxide). This extremely low contribution of engineered processes emphasizes the need to increase productivity, as well as to optimize and scale up the existing technologies, in order to produce larger amounts of microalgal biomass. Economic

viability of microalgal cultivation requires natural sunlight for photosynthetic carbon fixation, and an increasing number of scientists consider that large-scale outdoor cultivation remains the only viable option for cell and biofuel production at a commercial scale (Singh and Gu 2010; Chen et al. 2011; de Vree et al. 2015; Eustance et al. 2016; Ho et al. 2017). It has been reported that carbon dioxide fixation by microalgae might become economically viable when production costs fall below $\$500 \text{ t}^{-1} \text{ ha}^{-1} \text{ year}^{-1}$ (Bilanovic and Holland 2012).

Efficient microalgal carbon dioxide mitigation depends on microalgal species, light, carbon dioxide source and bioavailability, as well as photobioreactor design and operating conditions (Cheng et al. 2006), and very exhaustive reviews have reported the carbon dioxide removal efficiencies, biomass productivities and carbon dioxide fixation ability of a broad spectrum of microalgae growing at different operational conditions (Farrelly et al. 2013; Singh and Singh 2014; Zhao and Su 2014; Cuellar-Bermudez et al. 2015).

A key parameter influencing biomass productivity is the photosynthetic conversion efficiency, representing the luminous energy transformed to chemical energy, which is at the base of the anabolic reactions leading to biomass production. The maximum theoretical efficiency of solar energy conversion into biomass is around 12% (Weyer et al. 2010; Williams and Laurens 2010; Ooms et al. 2016). This value is calculated on the basis that only 47% of the total solar spectrum is available for photosynthesis and a maximum absorption of only 27% is achieved by the photosynthetic system from the photosynthetic active radiation. Finally, an additional 10% reduction is considered due to photo-transmission losses. However, photosynthetic conversion efficiency values between 1.5%, for open raceway ponds, and 3.8–5% for tubular or flat panel photobioreactors have been reported in outdoor conditions (Norsker et al. 2011; de Vree et al. 2015) and of 6% under laboratory conditions (Norsker et al. 2011). Higher photosynthetic conversion efficiency should allow higher productivities and lower costs (Wijffels, Barbosa and Eppink 2010; Wijffels and Barbosa 2010).

The main problem associated with outdoor algal cultivation is the limited productivity resulting from (i) light attenuation along the path in the cultivation system; (ii) excessive light impinging on the superficial layers causing photoinhibition (Janssen and Jack 2016); (iii) non-optimal temperatures due to climatic or seasonal conditions; (iv) respiratory losses of biomass; (v) salinity variations driven by water evaporation; (vi) inhibitory high oxygen concentration (Cohen, Koren and Arad 1991); and (vii) cultivation failure by bio-contamination with fungal or bacterial species, competing algae, predators, parasites, viruses, etc. In the case of biofuel production, strains should be found with high lipids or carbohydrates for biodiesel or biogas (hydrogen or methane) that are capable of maintaining fast growth rates under the nutrient limitation required to trigger the accumulation of those compounds. While some key parameters, such as pH, may be maintained, others, including temperature and irradiance, cannot be controlled to the same extent as under laboratory conditions (Wolf et al. 2016). Some of the most important environmental factors affecting growth rate, biomass productivity and carbon fixation will be reviewed here.

PHOTOSYNTHESIS AND CARBON DIOXIDE CONCENTRATING MECHANISM

Photosynthesis allows microalgal carbon dioxide fixation and the production of life-sustaining organic molecules. In the light-dependent cycle, photons from sunlight provide energy

to split water into molecular oxygen, and chlorophylls and carotenoids in the photosystems (I and II) are the key light-harvesting molecules that funnel the absorbed energy into the electron transport chain to create the high-energy and reducing molecules, ATP and NADPH, respectively. These are further used in the Calvin–Benson cycle to capture carbon dioxide through ribulose biphosphate carboxylase–oxygenase (Rubisco) enzyme mediation. (Information about the time scale of these processes can be found in Falkowski and Raven (1997) and Luo and Al-Dahhan (2011).) The low Rubisco specificity is compensated through a carbon dioxide concentrating mechanism by enhancing the carbon dioxide concentration up to 1000 times with respect to the carbon dioxide in the liquid medium (Price et al. 2008) and by decreasing photorespiration (Zhao and Su 2014).

When carbon dioxide dissolves in water, it can be found as carbon dioxide, HCO_3^- , CO_3^{2-} and H_2CO_3 . Among these, carbon dioxide and HCO_3^- are the major species used by microalgal cells. Carbonic anhydrases play a key role in the carbon dioxide concentrating mechanism and are found in the periplasm, cytosol and chloroplasts, having different functions, such as maintaining the carbon dioxide and HCO_3^- equilibrium or fostering transport through membranes. Once Rubisco acts, carbon dioxide is transformed into the carbohydrate precursors used for cell metabolism and growth (Zhao and Su 2014).

MICROALGAL GROWTH

The simple structure of microalgae permits fast cell growth with doubling times of 1 or 2 days or less (a few hours) and higher carbon dioxide fixation efficiencies than terrestrial plants (Richmond 2004). Microalgal growth is influenced by both biotic and abiotic factors that include, among others, temperature, pH, carbon dioxide, dissolved O_2 , light intensity and nutrients present in the medium. Additionally, the chemical composition of microalgal biomass also depends on the microalgal species, the growth phase and even on harvest time (Sánchez-Mirón et al. 2003; Juneja, Ceballos and Murthy 2013). Therefore, it is important to know the single and combined effect of the above-mentioned key parameters to improve the productivity of biomass and its products.

Nutrients

The molecular formula of microalgal biomass has been reported as $\text{CO}_{0.48}\text{H}_{1.83}\text{N}_{0.11}\text{P}_{0.01}$ (Chisti 2007), which can help to determine the nutritional requirements for cell growth. Table 1 summarizes the main nutrients constituting microalgal cells. As can be observed, nutrients include nitrogen (N), phosphorus (P), magnesium (Mg), sulfur (S) and trace element. The major cell component is carbon at around 50%, so to produce 1 ton of algal biomass, 1.8 tons of carbon dioxide need to be fixed (Chisti 2007; Alcántara, García-Encina and Muñoz 2013). Carbon dioxide can be supplied from the atmospheric air, industrial gaseous streams (e.g. flue gas) (Pires et al. 2012; Cuellar-Bermudez et al. 2015; Duarte et al. 2016; Fistarol, Farias and Salomon 2016; Thomas, Mechery and Paulose 2016) or even soluble carbonates (e.g. NaHCO_3 and Na_2CO_3) (Abinandan and Shanthakumar 2016; Mokashi et al. 2016; Wang et al. 2008). Atmospheric carbon dioxide concentration is low (around 400 ppm) and cannot sustain an intensive production of microalgal biomass; therefore, streams with higher carbon dioxide content, such as those found in flue gases, are preferred. Nevertheless, certain aspects of flue gases have to be considered, including their temperature and the presence of other gaseous components, such as SO_x

Table 1. Nutrient functions in microalgal cultivation (adapted from Zeng et al. 2011).

Source	Main nutrient	Example of function	Concentration range
Carbon	CO ₂ , HCO ₃ ⁻ , CO ₃ ²⁻ , etc.	Synthesis of carbohydrates, protein, lipids of the whole cell	1–10 g L ⁻¹
Nitrogen	NO ₃ ⁻ , urea, ammonia, N ₂ , etc.	Synthesis of nucleic acids, proteins, pigments such as chlorophyll and phycocyanin	10–2000 mg L ⁻¹
Phosphorous	Hydrophosphate, phosphate, etc.	Part of the backbone of DNA and RNA, ATP, phospholipids	10–500 mg L ⁻¹
Sulfur	Sulfate	Formation of the amino acids methionine and cysteine	1–200 mg L ⁻¹
Inorganic salts	K, Ca, Na, Mg, etc.	Maintenance of cell structure and activity	0.1–100 mg L ⁻¹
Trace elements	Fe, Zn, Mn, Pb, Cd, etc.	Coenzyme factors	0.01–10 mg L ⁻¹
Vitamins	B, C, E, etc.	Aid to cell division	0.01–1000 µg L ⁻¹

Table 2. Common optimum conditions for microalgae growth.

Microalgae	Temperature (°C)	Irradiance (µmol m ⁻² s ⁻¹)	pH	CO ₂ tolerance (%)	Mixing	Reference
<i>Chlorella</i> sp.	27	100	7	40	Aeration	Hanagata et al. (1992), Kim et al. (2012)
<i>Dunaliella</i> sp.	25	100		10	Aeration	Kim et al. (2012)
<i>Dunaliella salina</i> DCCBC2	27	80	8	5	Aeration	Kim et al. (2012)
<i>Haematococcus pluvialis</i>	25–28	90	7	34	Aeration	Fan, Vonshak and Boussiba (1994), Sarada, Tripathi and Ravishankar (2002), Huntley and Redjalje (2007)
<i>Euglena gracilis</i>	27–31	100	–	45	Fermenter	Nakano et al. (1996), Kitaya, Azuma and Kiyota (2005)
<i>Phaeodactylum tricornutum</i>	20.4	10	7.8	15	Aeration	Bitaubé, Caro and Pérez (2008)
<i>Scenedesmus almeriensis</i>	35	200	7–8	–	Aeration	Costache et al. (2013)
<i>Scenedesmus obtusiusculus</i>	35	300	7–8	15	Aeration	Cabello, Morales and Revah (2015)
<i>Spirulina platensis</i>	30	330	9–10	10	Aeration	Kebede and Ahlgren (1996), Oliveira et al. (1999), Richmond (2004), Colla et al. (2007), Kumar et al. (2010)
<i>Spirulina maxima</i>	35	–	9–10	–	Aeration	Oliveira et al. (1999), Richmond (2004)

and NO_x (Yen et al. 2015; Duarte-Santos et al. 2016; Sun, Zwolin-ska and Chmielewski 2016) or metals (Napan et al. 2015; Hess et al. 2017) that could be toxic to microalgae. Nitrogen is the second major component in importance for microalgal growth and may interact with the carbon source. Bilanovic et al. (2009) used the response surface methodology to define carbon dioxide and N concentrations to maximize both biomass production and carbon dioxide fixation capacity and found that nitrogen concentration for freshwater algae should be between 285 and 427 mg_N L⁻¹ for carbon dioxide concentrations from atmospheric to 70%. Phosphorous is necessary to synthesize amino acids, phospholipids and energy carriers and it is generally provided as phosphates. Nutrient limitation (mainly N or P) shifts the cell metabolism from growth to storage of high energy content molecules, such as lipids and carbohydrates (Takagi et al. 2000; Khozin-Goldberg and Cohen 2006; Rodolfi et al. 2009; Xin et al. 2010; Juneja, Ceballos and Murthy 2013), but the specific carbon capture rate is generally lowered. Oligo- and trace elements are also required in low amounts and sodium chloride (NaCl) affects carbon dioxide and O₂ solubility and helps to regulate biomass growth and product formation (Ben Amotz, Tornabene and Thomas 1985; Takagi, Karseno, Yoshida 2006). In conclusion, medium formulation is a necessary step to increase growth rates and productivity and, therefore, carbon dioxide capture, considering both the cell composition and process requirements (Danquah et al. 2010).

EFFECT OF ENVIRONMENTAL FACTORS ON carbon dioxide BIOFIXATION

In this section, the main environmental factors affecting photosynthesis and, consequently, the growth rate, biomass productivity and carbon fixation are analyzed and Table 2 summarizes the optimum values for some species.

Irradiance

Light intensity is essential for photosynthesis, microalgal growth and, therefore, carbon dioxide fixation (Fernández et al. 2012; Costache et al. 2013). During photosynthesis, the products of light reactions, ATP and NADPH, are used to fix carbon dioxide by Rubisco in the Calvin–Benson cycle. Both molecules are provided from light-dependent reactions; therefore, a good light supply is necessary. On the one hand, low light conditions lead to low biomass concentration and carbon dioxide fixation. On the other hand, excess light increases the production of reactive oxygen species that are responsible for abnormal physiological reactions due to oxidative stress (Seo et al. 2017). They damage the essential proteins required for electron transfer during photosynthesis and, consequently, carbon dioxide fixation is reduced.

The empirical relationship between irradiance and photosynthesis is graphically depicted by photosynthesis-irradiance

(P–I) curves (Béchet, Shilton and Guieysse 2013). These curves are species-dependent and need to be experimentally determined under uniform light intensity (Béchet, Shilton and Guieysse 2013). Generally, three regions are identified: (i) a region limited by the rate of capture of photons where the photosynthetic rate is directly proportional to irradiance; (ii) the 'light-saturation' region where photosynthesis is irradiance-independent—here, photosynthesis is limited by the reaction rates achieving maximum value (Béchet, Shilton and Guieysse 2013); and (iii) the photoinhibition region at high irradiances where a decrease in photosynthesis is observed (Camacho-Rubio et al. 2003). However, an important issue relating to P–I curves is that they are determined under uniform light intensity, contrary to the situation found during cultivation in photobioreactors, where dense cultures can be achieved. Consequently, P–I curves cannot be used directly to predict productivity in dense cultures, where the light regime must be known or some valid correlations must be used to predict light attenuation based on depth, biomass concentration and mixing patterns.

Photo-acclimation allows microalgae to respond to light changes, balancing the light reactions, the energy demand for carbon dioxide fixation and other metabolic reactions (Vonshak and Torzillo 2004). The photo-acclimation periods vary according to species (Falkowski and Chen 2003; Nikolaou et al. 2016). In the short term (time scales of seconds to minutes), adaptation includes adjustments in the delivery of light energy to photosystem II through reorganization of photosynthetic complexes (Mooij et al. 2017). Long-term acclimation comprises changes in enzyme activity and gene expression affecting both the structure and the stoichiometry in the photosystems (Cabello, Morales and Revah 2014). Photoinhibition occurs in systems exposed to high irradiances, just above the value where maximum photosynthetic activity occurs (Camacho-Rubio et al. 2003; Pulz and Gross 2004), and involves deactivation of proteins in photosynthetic units (Camacho-Rubio et al. 2003; Fan et al. 2007). This process is time-dependent and a half of the photosynthetic system might be irreversibly damaged after a period of 10–20 min of exposure (Pulz and Gross 2004).

Different light sources are used for microalgal cultivation, including natural sunlight and artificial light. Kommareddy and Anderson (2003) compared different light sources, including incandescent, halogen, fluorescent, AlInGaP II (aluminum indium gallium phosphide with a peak at a wavelength of 643 nm) and light emitting diodes. They found that light emitting diodes and AlInGaP II were the most efficient and economical light sources. Recent advances include illumination with selected wavelengths, dye compounds to lower the energy photons, fluorescent paints and nanoparticles to filter the light (Ramanna, Rawat and Bux 2017). Although many studies are performed indoors with artificial light under controlled conditions, large scale cultivation systems operate outdoors with solar light depending on the circadian cycle, location, season of the year and climatic conditions. Furthermore, a successful cultivation system with high biomass productivity must have high cell concentrations to minimize costs (Molina-Grima et al. 2003); however, this high biomass content produces light gradients in the medium and cells are exposed to different light intensities (Béchet, Shilton and Guieysse 2013).

Light supply may be continuous or with intermittent photoperiods. Cycles of 12:12 or 18:6 dark–light cycles have been reported as adequate to achieve good carbon dioxide fixation (Ramanna, Rawat and Bux 2017); however, Jacob-Lopes et al. (2009) found that during continuous illumination, carbon dioxide fixation by *Aphanotece microscopica* Nägeli microalgae

reached 99.69% of elimination efficiency. Furthermore, algal growth and carbon dioxide fixation are affected not only by light–dark cycle, but also by light intensity. Thawechai et al. (2016) found that the optimal conditions of 60 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ and continuous illumination allowed a carbon dioxide fixation rate of 0.729 $\text{mg L}^{-1} \text{ day}^{-1}$. Contrary to those results, Sun and Duan (2007) found that for the microalga *Tetraselmis tetrathele*, light cycles ranged from 6 to 18 h, whereas the cell density, chlorophyll *a* and protein content reached high levels suggesting the importance of an optimized light cycle to maximize metabolism and algal growth.

Temperature

Temperature influences both physicochemical carbon dioxide availability to the cells and the metabolic processes within the cell (Ördög et al. 2016; Paliwal et al. 2017). Carbon dioxide solubility depends on temperature; it is known that higher temperatures (>20°C) lower the carbon dioxide solubility. On the other hand, increases in temperature decrease Rubisco's affinity for carbon dioxide (Kumar et al. 2011). However, the temperature effect on the reaction metabolic rate is strain dependent. Light and temperature were the most important factors affecting the biomass productivity and carbon dioxide fixation in a long-term experiment in which *Scenedesmus obliquus* was grown in outdoor photobioreactors with N and carbon dioxide maintained at a constant level (Hindersin et al. 2014).

Ras, Steyer and Bernard (2013) performed a review of the effect of temperature on different species, emphasizing the biochemical effect and the adaptation to high temperatures. Microalgae usually grow in a temperature range between 15 and 30°C with optimal values at 20–25°C. It has also been reported that temperatures below 16°C affect particularly mechanisms of carbon fixation (Ras, Steyer and Bernard 2013) and metabolic activity in general. On the other hand, high temperatures lower carbon dioxide solubility (Zhao and Su 2014) and temperatures above 35°C are normally lethal for many species (Briassoulis et al. 2010), decreasing the growth rate and increasing both respiration and photorespiration. Temperature is especially important for treatment of flue gases (temperatures around 120°C) or large-scale outdoor cultivation with circadian variations between 10 and 45°C. In these cases, the use of temperature-tolerant organisms could result in savings related to cooling costs. Some of the reported photosynthetic organisms include *Chlorella pyrenoidosa* (optimum around 39°C and tolerance up to 45°C), *Chloroglossopsis* sp. (isolated from a hot spring in the Yellowstone National Park; grows at 50°C; Ono and Cuello 2007) and *Synechococcus* sp. (Miller and Castenholz 2000; 42–75°C), which exhibits the thermal maximum for photosynthetic life (Brock 1967; Castenholz 1969).

pH

The pH regulates, on the chemical level, the distribution of different dissolved inorganic species (carbon dioxide, HCO_3^- and CO_3^{2-} ; de Moraes and Costa 2007). It is also one of the important factors for algal growth because it affects the activity of different enzymes (Zhang et al. 2016). Its value has a strong influence on cell growth and the optimum and tolerance are species-dependent. The optimum pH is normally in the neutral to slightly alkaline range (6–8.3; Moss 1973; Azov 1982; Olaizola 2003; Pandey, Pathak and Tiwari 2010; Ying, James Gilmour and Zimmerman 2014). However, some organisms bloom in extreme conditions. *Spirulina* grows in highly alkaline conditions (pH 9–11; Pandey, Pathak

and Tiwari 2010; Koru 2012; Sharma et al. 2014). Alkaline conditions, at pH above 11.0, could help to maintain microalgal cultures free of contaminants, such as protozoa and other competing microalgae (Touloupakis et al. 2016), and also promote neutral lipid accumulation in microalgal cells (Zhang et al. 2016). Other microalgae thriving in more acidic conditions include the autotrophic Chlorophyta (*Chlamydomonas acidophila*, *Dunaliella acidophila*), Chrysophyta (*Ochromonas* sp.) and Euglenophyta (*Euglena mutabilis*), filamentous algae identified as *Zygnema circumcarinatum* and *Klebsormidium acidophilum* (Sung et al. 1998; Novis and Harding 2007; Rowe et al. 2007), and *Chlorella protothecoides* var. *acidicola* isolated from mine-impacted water bodies, which grows optimally at pH 2.5 under heterotrophic conditions (Nancucheo and Johnson 2012).

During carbon dioxide fixation and microalgal growth, pH variations occur in relation to carbon dioxide dissolution, equilibrium of the dissolved inorganic carbon species and the uptake of carbon dioxide and nitrate. However, these changes also depend on the carbon dioxide input concentration. When HCO_3^- is used, the carbonic anhydrase converts it to carbon dioxide resulting in the release of the hydroxyl ion (OH^-) and consequently an increase in pH. This situation indicates carbon dioxide limitation affecting the system performance and carbon dioxide supply and allows the control of culture pH at its optimal value. Recently, Duarte-Santos et al. (2016) have used the strategy of pH control with injection of carbon dioxide-rich gases on demand. This allowed them to meet the carbon requirements of microalgal cultures and to optimize biomass productivity and carbon dioxide fixation. That study was performed in a pilot scale- raceway pond, to produce high value-added compounds, such as phycocyanin (Chen et al. 2016). The same strategy was also reported to control the protozoa (ciliates or amoebae) (Ma et al. 2017). High carbon dioxide concentrations, similar to those found in flue gases (10–25%), acidify the cultivation medium when continuously fed (Zhao and Su 2014; Choi et al. 2017). Furthermore, other chemicals, such as SO_2 or NO_x from flue gases, reduce the pH when they dissolve in water (Kao et al. 2014; Cuellar-Bermudez et al. 2015; Yen et al. 2015; Fistarol, Farias and Salomon 2016). Consequently, pH control by addition of neutralizing agents, by designing buffers or by defining strategies of carbon dioxide supply is critical to avoid growth inhibition.

carbon dioxide concentration

carbon dioxide fixation involves mass transfer and solubilization from the gaseous stream to the liquid phase, attaining the ionic equilibrium between carbon dioxide, HCO_3^- and CO_3^{2-} , and carbon assimilation by the microalgal cells. Tolerance and optimal carbon dioxide concentration are strain-dependent (Solovchenko and Khozin-Goldberg 2013) and most microalgae generally grow well at 2% carbon dioxide, but levels above 5% carbon dioxide (Cheng et al. 2006) may inhibit their growth (Yun et al. 1997; Chiu et al. 2008). This effect may be related to the acidification of the stroma in the chloroplast (Solovchenko and Khozin-Goldberg 2013) causing the inactivation of key enzymes of the Calvin-Benson cycle. Singh and Singh (2014) reviewed the influence of carbon dioxide levels on different microalgal species including *Scenedesmus obliquus*, *Botryococcus braunii*, *Chlorella vulgaris* and *Nannochloropsis oculata*. They reported that the highest carbon dioxide concentration was fixed by *S. aquatilis*, *B. braunii*, *C. vulgaris* and *Synechococcus* species. However, according to Singh and Ahluwalia (2013), although the most carbon dioxide-tolerant strains were *Chlorella* sp. T-1 (100% carbon dioxide), *Scenedesmus* sp. (80% carbon dioxide) and *Eu-*

glena gracilis (45%), their maximum biomass productivity was found at carbon dioxide concentrations of 10%, 10–20% and 5%, respectively. Likewise, *Chlorella* sp. KR-1 showed the maximum growth at 10% carbon dioxide and tolerated up to 70% carbon dioxide (Sung et al. 1998). The above-mentioned range is important because some microalgae can grow with 10–15% carbon dioxide, the range normally found in flue gases, which are an important greenhouse gas source of pollution, but may be an excellent carbon dioxide source for microalgae.

Dissolved O_2 concentration

Dissolved oxygen is an indicator of photosynthetic activity by microalgae, and registered values are higher than those found in equilibrium with air. Low dissolved oxygen indicates problems with the microalgal growth and possibly dissolved oxygen consumption by heterotrophic microorganisms. The build-up of O_2 in microalgal cultivation systems is one of the major problems in obtaining high biomass productivity and carbon dioxide fixation rate. Dissolved oxygen may exceed 250% of saturation during daytime (Fernández et al. 2012; Raso et al. 2012; Costache et al. 2013; Bilanovic et al. 2016). High oxygen concentration leads to the oxidative stress of reactive oxygen species and also competes with carbon dioxide for the Rubisco involved in carbon dioxide fixation. The carbon dioxide and O_2 concentrations define carboxylation (photosynthesis) and oxygenation (photorespiration) prevalence. Low carbon dioxide/ O_2 ratios reduce the photosynthetic rate and, therefore, carbon dioxide fixation and favor photorespiration and carbon dioxide release (Costache et al. 2013). To avoid dissolved oxygen accumulation, both design and operational issues must be considered. They include the establishment of maximum tube length in photobioreactors and promotion of oxygen release to the gas phase by degassing the liquid and promoting turbulence of the medium (Grobbelaar 1994; Camacho-Rubio et al. 1999).

METHODOLOGICAL EVALUATION OF THE EFFECT OF ENVIRONMENTAL VARIABLES (carbon dioxide, TEMPERATURE, PH AND IRRADIANCES)

The experimental assessment of the effect of key factors reviewed above is vital to determine the optimal operation conditions to maximize growth rates and carbon dioxide capture. Data on the effects on photosynthetic activity are generally gathered by cell growth or oxygen production rates. Microalgal growth is normally measured in photobioreactors by biomass accumulation during the cultivation time and results from the combination of biological, environmental and mass transfer effects. This approach does not allow the independent evaluation of key variables and depends on the changing operational conditions and photobioreactor configuration. Furthermore, it requires a time scale of days to obtain significant results (Butterwick, Heaney and Talling 2005; Breuer et al. 2013). On the other hand, the photosynthetic activity can be determined at different culture conditions through dynamic oxygen measurements (Bitaubé, Caro and Pérez 2008; Brindley, Ación and Fernández-Sevilla 2010; Fernández et al. 2012; Costache et al. 2013). This approach yields results on a time scale of minutes and is proportional to the growth rate according to the stoichiometry of the overall photosynthetic reaction (Brindley, Ación and Fernández-Sevilla 2010). Oxygen response to changing conditions also provides information about the photo-acclimation phenomena generated in

photosystem II as a response to fluctuations in light intensity (Sato et al. 2002; Avendaño and Shubert 2005). Furthermore, the time needed for determination of the photosynthetic irradiance curves and the associated kinetic parameters is reduced and some models have been developed for predicting O₂ production (Camacho-Rubio et al. 1999; Reboloso et al. 1999; Concas and Pisu 2010; Fernández et al. 2012; Hu et al. 2012; Béchet, Shilton and Guieysse 2013).

Dynamic dissolved O₂ measurements coupled to mass balance and transfer models have been used to evaluate the effect of irradiance, temperature and pH on the photosynthetic activity for the microalga *Scenedesmus obtusiusculus* (Cabello, Morales and Revah 2015). Here, thermodynamic and intrinsic kinetic pa-

rameter data were evaluated in a mini-photobioreactor, where self-shading, mass transfer effects and carbon dioxide limitation were minimized. Results with the optimized conditions were further tested successfully in controlled flat-panel photobioreactors to produce biomass and lipids (Morales, Cabello and Revah 2015).

Dynamic methods were also used to evaluate the photosynthetic activity of the same microalga to different light intensities in an air-lift photobioreactor through step changes under N-replete and N-starved conditions (Cabello, Morales and Revah 2014), as well as the effect of variations of carbon dioxide concentration (Cabello, Morales and Revah 2017). In this case, during the experiments on-line monitoring of pH, carbon dioxide and

Table 3. Common optimum conditions for microalgae growth.

Microalgae	Temperature (°C)	Irradiance (μmol m ⁻² s ⁻¹)	CO ₂ pH tolerance	Photobioreactor conditions	Growth rate (day ⁻¹)	Biomass Productivity (g L ⁻¹ day ⁻¹)	CO ₂ biofixation rate (mg L ⁻¹ day ⁻¹)	Reference	
<i>Chlorella</i> sp.	27	100	7	40%	Conical flask, 8 days continuous aeration, air + CO ₂ ; horizontal bubble column, 11 days, continuous aeration, N ₂ + CO ₂	0.38	0.09	7.2	Hanagata et al. (1992), Kim et al. (2012)
<i>Dunaliella</i> sp.	25	100	8	5%	Horizontal bubble column, 11 days, continuous aeration, N ₂ + CO ₂	0.25	0.12	10.4	Kim et al. (2012)
<i>Dunalliella salina</i> DCCBC2	27	80	8	10%	Horizontal bubble column, 11 days, continuous aeration, N ₂ + CO ₂	0.42	0.13	8.2	Kim et al. (2012)
<i>Haematococcus pluvialis</i>	25–28	90	7	34%	Bubble column, aeration air + CO ₂	1.29			Fan, Vonshak and Boussiba (1994), Sarada, Tripathi and Ravishankar (2002), Huntley and Redjalje (2007)
<i>Euglena gracilis</i>	27–31	100		45%	Photobioreactor, 28 days, flue gas 11% CO ₂ , continuous operation	0.31	0.29	74	Nakano et al. (1996), Kitaya, Azuma and Kiyota (2005)
<i>Phaeodactylum tricornutum</i>	20.4	10	7.8	15%	Erlenmeyer flask, aeration; Roux bottles, 11 days, continuous aeration air + CO ₂		0.15	280 ^a	Bitaubé, Caro and Pérez (2008)
<i>Scenedesmus almeriensis</i>	35	200	7–8	10%	Tubular photobioreactor, outdoor conditions, flue gas (pure CO ₂) on demand, operated continuously	0.34	0.42	790	Costache et al. (2013)
<i>Scenedesmus obtusiusculus</i>	35	300	7–8	10%	Bubble column, 14 days, continuous aeration air + CO ₂	0.34	0.52	970 ^a	Cabello, Morales and Revah (2015), Toledo-Cervantes et al. (2013)
<i>Spirulina platensis</i>	30	330	9–10	10%	Bubble column, 25 days, continuous aeration, greenhouse conditions	0.65	0.15	280 ^a	Kebede and Ahlgren (1996), Oliveira et al. (1999), Richmond (2004), Colla et al. (2007), Kumar et al. (2010)
<i>Spirulina maxima</i>	35		9–10		Container of glass, 15 days, mechanical agitation	0.6	0.15	280 ^a	Oliveira et al. (1999), Richmond (2004)

^aCalculated from the biomass productivity according to the equation: CO₂ fixation rate (P_{CO₂}) = 1.88 × biomass productivity (g L⁻¹ day⁻¹), which is derived from the molecular formula of microalgal biomass, CO_{0.48}H_{1.83}N_{0.11}P_{0.01}.

Table 4. Seasonal growth of *Scenedesmus obtusiusculus* in air-lift extended loop photobioreactor.

Conditions	Productivity (g L ⁻¹ day ⁻¹)	carbon dioxide fixation ^a (g L ⁻¹ day ⁻¹)	Temperature range (°C)	I_{\max} (μmol m ⁻² s ⁻¹)/mol photons	PE ^b (%)
Winter	0.19	0.36	5–31	1536/92.9	4.11
Spring	0.29	0.55	11–47	2035/98.8	5.34
Summer	0.21	0.40	5–26	2055/71.2	4.13
Autumn	0.22	0.42	7–45	2100/88.5	4.21
Optimum laboratory conditions ^c	0.97	1.84	35.2	300/15.6 ^d	4.4

^aCalculated from theoretical value of 1.8 g carbon dioxide g biomass⁻¹. ^bPhotosynthetic efficiency (PE) is calculated as $PE = (X_t V \Delta H_{EN}) / (E A I_t)$ where X_t is the biomass at time t , A is the illuminated area of the reactor, V is the reactor volume, ΔH is the combustion heat of the microalgal biomass with values between 20.6 and 26.7 kJ g⁻¹ for a microalgal biomass with a lipid content of 54.7% and 18.1%, respectively, I_t is the impinging irradiance at the time t , E_N is the coefficient related to the energy required for nitrogen assimilation with values of 1.25 for nitrate assimilation (Williams and Laurens 2010), E is the energy from photosynthetically active radiation (210 KJ mol photons⁻¹) (Dillschneider et al. 2013). ^c Evaluated in a flat panel photobioreactor. ^dContinuous illumination.

dissolved O₂ was performed. The dynamic experiments allowed a rapid analysis of the short-term carbon dioxide consumption rates in response to changes in inlet gas carbon dioxide concentration.

OUTDOOR OPERATION

The impact of environmental conditions on carbon dioxide capture and biomass growth is clearly observed in outdoor cultivation. The microalgal growth rate is drastically lower than that obtained under controlled conditions in the laboratory and many microalgal strains do not grow reliably outdoors as a consequence of diurnal or seasonal fluctuations in irradiance and temperature (Moheimani 2013; Eustance et al. 2016).

Excellent reviews on algal physiology and large-scale outdoor cultures of microalgae have been published, including those on evaluation of the effect of photobioreactor configuration (see Table 3) on microalgal productivity and photosynthetic efficiency (de Vree et al. 2015). However, limited information exists about large-scale production and long-term operation of outdoor cultivation systems for most microalgal species (Van Bergeijk, Salas-Leiton and Cañavate 2010; Chiu, Soong and Chen 2016), and performance comparisons of a strain at optimum conditions in laboratory and outdoor environments are scant. Therefore, here we will take the case of *Scenedesmus obtusiusculus* as an example (Sánchez et al. 2013; Cabello, Morales and Revah 2015).

Scenedesmus obtusiusculus was grown in a 35 L airlift tubular photobioreactor, based on the design proposed by Molina-Grima et al. (2001). It was operated outdoors during a yearlong cultivation in Mexico City on a building roof located at N 19°21'32.466", W 99°4'25.192" in 2013. Table 4 shows results of biomass productivity and carbon dioxide fixation, including data under controlled optimal conditions in a flat panel photobioreactor. As can be seen, the highest biomass productivity, 0.29 g L⁻¹ day⁻¹, and carbon dioxide fixation rate, 0.55 g L⁻¹ day⁻¹, were obtained during the spring when the highest temperatures and irradiances (photon exposition) were recorded. However, these values were lower than those obtained in laboratory controlled conditions, where the biomass productivity was 0.97 g L⁻¹ day⁻¹ and a carbon dioxide fixation rate of 1.84 g L⁻¹ day⁻¹ was achieved (Toledo-Cervantes et al. 2013). This difference could be attributed to the fact that the above-mentioned conditions (temperature, irradiance) change dynamically making it difficult to clearly isolate the impact of any single factor.

Finally, few works have evaluated the effect of outdoor conditions on the carbon dioxide capture, biomass composition

and biofuel potential. Recently, Ho et al. (2017) have found that *Scenedesmus obliquus* CNW-N exhibited the optimal carbon dioxide fixation rate and carbohydrate productivity during the summer with an average irradiance around 200 μmol m⁻² s⁻¹ and temperatures close to 31°C. They recognize the importance of understanding the seasonal changes and conclude that it is possible to combine carbon dioxide fixation and bioethanol production using microalgae grown outdoors.

CONCLUSION

There is a common consensus in the scientific community about the need to address the significant difficulties involved in the large-scale production of microalgae and their products. Furthermore, to exceed the economic break-even point and make microalgal technologies a relevant contributor to carbon dioxide sequestration, the range of products should be extended, production should be maximized under the constraints (the selected strain) and the environmental conditions set in accordance with the cultivation method and the location of the facilities.

ACKNOWLEDGEMENT

The authors thank the Fondo Sectorial CONACYT-SENER-Sustentabilidad Energética, Clúster Biocombustibles Gaseosos project 247006 and CONACYT projects PN2014-247402 and PD-CPN 2015-241 for financial support. L.S. acknowledges CONACYT for a scholarship.

Conflict of interest. None declared.

REFERENCES

- Abinandan S, Shanthakumar S. Evaluation of photosynthetic efficacy and carbon dioxide removal of microalgae grown in an enriched bicarbonate medium. *3 Biotech* 2016;6:9.
- Alcántara C, García-Encina PA, Muñoz R. Evaluation of mass and energy balances in the integrated microalgae growth-anaerobic digestion process. *Chem Eng* 2013;221:238–46.
- Avendaño D, Shubert H. Oxygen evolution and respiration of the cyanobacterium *Synechocystis* sp. *Physiol Plant* 2005;125:381–91.
- Azov Y. Effect of pH on inorganic carbon uptake in algal cultures. *Appl Environ Microbiol* 1982;43:1300–6.
- Béchet Q, Shilton A, Guieysse B. Modeling the effects of light and temperature on algae growth: State of the art and critical

- assessment for productivity prediction during outdoor cultivation. *Biotech Adv* 2013;**31**:1648–63.
- Ben Amotz B, Tornabene TG, Thomas WH. Chemical profile of selected species of microalgae with emphasis on lipids. *J Phycol* 1985;**21**:72–78.
- Bilanovic D, Andargatchew A, Kroeger T et al. Fresh water and marine microalgae sequestering of carbon dioxide at different C and N concentrations – response surface methodology analysis. *Energy Convers Manage* 2009;**50**:262–7.
- Bilanovic D, Holland M. Microalgal carbon dioxide sequestering – Modeling microalgae production costs. *Energy Convers Manage* 2012;**58**:104–9.
- Bilanovic D, Holland M, Starosvetsky J et al. Co-cultivation of microalgae and nitrifiers for higher biomass production and better carbon capture. *Bioresour Technol* 2016;**220**:282–8.
- Bitaubé E, Caro I, Pérez L. Kinetic model for growth of *Phaedolactynum tricornutum* in intensive culture photobioreactor. *Biochem Eng J* 2008;**40**:520–5.
- Breuer G., Lamers PP, Martens DE et al. Effect of light intensity, pH, and temperature on triacylglycerol (TAG) accumulation induced by nitrogen starvation in *Scenedesmus obliquus*. *Bioresour Technol* 2013;**143**:1–9.
- Briassoulis D, Panagakis P, Chionidis M et al. An experimental helical-tubular photobioreactor for continuous production of *Nannochloropsis* sp. *Bioresour Technol* 2010;**101**:6768–77.
- Brindley C, Acien FG, Fernández-Sevilla JM. The oxygen evolution methodology affects photosynthetic rate measurements of microalgae in well-defined light regimes. *Biotechnol Bioeng* 2010;**106**:228–37.
- Brock TD. Micro-organisms adapted to high temperatures. *Nature* 1967;**214**:882–5.
- Butterwick C, Heaney SI, Talling JF. Diversity in the influence of temperature on the growth rates of freshwater algae, and its ecological relevance. *Freshw Biol* 2005;**50**:291–300.
- Cabello J, Morales M, Revah S. Dynamic photosynthetic response of the microalga *Scenedesmus obtusiusculus* to light intensity perturbations. *Chem Eng J* 2014;**252**:104–11.
- Cabello J, Morales M, Revah S. Effect of the temperature, pH and irradiance on the photosynthetic activity by *Scenedesmus obtusiusculus* under nitrogen replete and deplete conditions. *Bioresour Technol* 2015;**181**:128–35.
- Cabello J, Morales M, Revah S. Carbon dioxide consumption of the microalga *Scenedesmus obtusiusculus* under transient inlet carbon dioxide concentration variations. *Sci Total Environ* 2017;**584–585**:1310–6.
- Camacho-Rubio F, Acien -Fernández FG, Sánchez-Pérez JA et al. Prediction of dissolved oxygen and carbon dioxide concentration profiles in tubular photobioreactors for microalgal culture. *Biotechnol Bioeng* 1999;**62**:71–86.
- Camacho-Rubio F, García-Camacho F, Fernández-Sevilla JM et al. A mechanistic model of photosynthesis in microalgae. *Biotechnol Bioeng* 2003;**81**:459–73.
- Castenholz RW. Thermophilic blue-green algae and the thermal environment. *Bacteriol Rev* 1969;**33**:476–504.
- Chen C-Y, Kao P-Ch, Tan ChH et al. Using an innovative pH-stat carbon dioxide feeding strategy to enhance cell growth and C-phycocyanin production from *Spirulina platensis*. *Biochem Eng J* 2016;**112**:78–85.
- Chen C-Y, Yeh K-L, Aisyah R et al. Cultivation, photobioreactor design and harvesting of microalgae for biodiesel production: a critical review. *Bioresour Technol* 2011;**102**:71–81.
- Cheng L, Zhang L, Chen H et al. Carbon dioxide removal from air by microalgae cultured in a membrane-photobioreactor. *Sep Purif Technol* 2006;**50**:324–9.
- Chisti Y. Biodiesel from microalgae. *Biotechnol Adv* 2007;**25**:294–306.
- Chiu P-H, Soong K, Chen CN. Cultivation of two thermotolerant microalgae under tropical conditions: influences of carbon sources and light duration on biomass and lutein productivity in four seasons. *Bioresour Technol* 2016;**212**:190–8.
- Chiu SY, Kao CY, Chen CH et al. Reduction of carbon dioxide by a high-density culture of *Chlorella* sp. in a semicontinuous photobioreactor. *Bioresour Technol* 2008;**99**:3389–96.
- Choi YY, Joun JM, Lee J et al. Development of large-scale and economic pH control system for outdoor cultivation of microalgae *Haematococcus pluvialis* using industrial flue gas. *Bioresour Technol* 2017;**244**:1235–44.
- Cohen E, Koren A, Arad S. A closed system for outdoor cultivation of microalgae. *Biomass Bioenerg* 1991;**1**:83–88.
- Colla LM, Oliveira Reinehr C, Reichert C et al. Production of biomass and nutraceutical compounds by *Spirulina platensis* under different temperature and nitrogen regimes. *Bioresour Technol* 2007;**98**:1489–93.
- Concas A, Pisu M. Novel simulation model of the solar collector of biocoil photobioreactors for carbon dioxide sequestration with microalgae. *Chem Eng J* 2010;**157**:297–303.
- Costache TA, Acien FG, Morales MM et al. Comprehensive model of microalgae photosynthesis rate as a function of culture conditions in photobioreactor. *Appl Biotechnol* 2013;**97**:7627–37.
- Cuellar-Bermudez SP, Garcia-Perez JS, Rittmann BE et al. Photosynthetic bioenergy utilizing carbon dioxide: an approach on flue gases utilization for third generation biofuels. *J Clean Prod* 2015;**98**:53–65.
- Danquah MK, Harun R, Halim R et al. Cultivation medium design via elemental balancing for *Tetraselmis suecica*. *Chem Biochem Eng Q* 2010;**24**:361–9.
- de Morais MG, Costa JA. Carbon dioxide fixation by *Chlorella kessleri*, *C. vulgaris*, *Scenedesmus obliquus* and *Spirulina* sp. cultivated in flasks and vertical tubular photobioreactors. *Biotechnol Lett* 2007;**29**:1349–52.
- de Vree JH, Bosma R, Janssen M et al. Comparison of four outdoor pilot-scale photobioreactors. *Biotechnol Biofuels* 2015;**8**:8–12.
- Dillschneider R, Steinweg C, Rosello-Sastre R et al. Biofuels from microalgae: Photoconversion efficiency during lipid accumulation. *Bioresour Technol* 2013;**142**:647–54.
- Duarte JH, Schneider-Fanka L, Vieira Costa JA. Utilization of simulated flue gas containing carbon dioxide, SO₂, NO and ash for *Chlorella fusca* cultivation. *Bioresour Technol* 2016;**2014**:159–65.
- Duarte-Santos T, Mendoza-Martín JL, Acien Fernández FG et al. Optimization of carbon dioxide supply in raceway reactors: Influence of carbon dioxide molar fraction and gas flow rate. *Bioresour Technol* 2016;**212**:72–81.
- Eustance E, Wray JT, Badvipour S et al. The effects of cultivation depth, areal density, and nutrient level on lipid accumulation of *Scenedesmus acutus* in outdoor raceway ponds. *J Appl Phycol* 2016;**28**:1459–69.
- Falkowski PG, Chen YB. Photoacclimation of light harvesting systems in eukaryotic algae. In: Green BR, Parson WW (eds). *Light-Harvesting Antennas in Photosynthesis. Advances in Photosynthesis and Respiration*, vol 13. Dordrecht: Springer, 2003, 423–447.
- Falkowski PG, Raven JA. *Aquatic Photosynthesis*. Malden, MA: Blackwell Science, 1997.
- Fan L, Vonshak A, Boussiba S. Effect of temperature and irradiance on growth of *Haematococcus pluvialis* (Chlorophyceae). *J Phycol* 1994;**30**:829–33.

- Fan L-H, Zhang Y-T, Cheng L-H et al. Optimization of carbon dioxide fixation by *Chlorella vulgaris* cultivated in a membrane-photobioreactor. *Chem Eng Technol* 2007;**30**:1094–9.
- Farrelly DJ, Everard CD, Fagan CC et al. Carbon sequestration and the role of biological carbon mitigation: a review. *Renew Sust Energ Rev* 2013;**21**:712–27.
- Fernández I, Acién FG, Fernández JM et al. Dynamic model of microalgal production in tubular photobioreactors. *Bioresour Technol* 2012;**126**:172–81.
- Fistarol GO, Farias M, Salomon PS. Viability of using flue gases as carbon source for microalgae cultivation. *Int J Green Technol* 2016;**2**:13–19.
- Grobbelaar JU. Turbulence in mass algal cultures and the role of light/dark fluctuations. *J Appl Phycol* 1994;**6**:331–5.
- Hanagata N, Takeuchi T, Furuju Y et al. Tolerance of microalgae to high carbon dioxide and high temperature. *Phytochemistry* 1992;**31**:3345–8.
- Hess D, Napan K, McNeil BT et al. Quantification of effects of flue gas derived inorganic contaminants on microalgae growth system and end fate of contaminants. *Algal Res* 2017;**25**:68–75.
- Hindersin S, Leupold M, Kerner M et al. Key parameters for outdoor biomass production of *Scenedesmus obliquus* in solar tracked photobioreactors. *J Appl Phycol* 2014;**26**:2315–25.
- Ho Ah-H, Chen Y-D, Chang Ch-Y et al. Feasibility of carbon dioxide mitigation and carbohydrate production by microalga *Scenedesmus obliquus* CNW-N used for bioethanol fermentation under outdoor conditions: effects of seasonal changes. *Biotechnol Biofuels* 2017;**10**:27.
- Hu D, Li M, Zhou R et al. Design and optimization of photo bioreactor for O₂ regulation and control by system dynamics and computer simulation. *Bioresour Technol* 2012;**104**:608–15.
- Huntley M, Redjalje D. carbon dioxide mitigation and renewable oil from photosynthetic microbes: A new appraisal. *Mitig Adapt Strat Glob Chang* 2007;**12**:573–608.
- IEA. State of Technology Review – Algae Bioenergy. IEA Bioenergy, 2017. Retrieved from: <http://www.ieabioenergy.com/wp-content/uploads/2017/02/IEA-Bioenergy-Algae-report-update-Final-template-20170131.pdf>. August 2017.
- IPCC. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, Pachauri RK, Meyer LA (eds)]. Geneva: IPCC, 2014. Retrieved from: https://www.ipcc.ch/pdf/assessment-report/ar5/syr/AR5_SYR_FINAL_All_Topics.pdf. August 2017.
- Jacob-Lopes E, Scoparo CHG, Lacerda LMCF et al. Effect of light cycles (night/day) on carbon dioxide fixation and biomass production by microalgae in photobioreactors. *Chem Eng Process Process Intensif* 2009;**48**:306–10.
- Janssen M, Jack L. Microalgal photosynthesis and growth in mass culture. *Adv Chem Eng* 2016;**48**:185–256.
- John RP, Anisha GS, Nampoothiri KM et al. Micro and microalgal biomass: a renewable source for bioethanol. *Bioresour Technol* 2011;**102**:186–93.
- Juneja A, Ceballos RM, Murthy GS. Effects of environmental factors and nutrient availability on the biochemical composition of algae for biofuels production: A review. *Energies* 2013;**6**:4607–38.
- Kao CY, Chen TY, Chang YB et al. Utilization of carbon dioxide in industrial flue gases for the cultivation of microalga *Chlorella* sp. *Bioresour Technol* 2014;**166**:485–93.
- Kebede E, Ahlgren G. Optimum growth conditions and light utilization efficiency of *Spirulina platensis* (*Arthrospira fusiformis*) (Cyanophyta) from Lake Chitu, Ethiopia. *Hydrobiologia* 1996;**332**:99–109.
- Khozin-Goldberg I, Cohen Z. The effect of phosphate starvation on the lipid and fatty acid composition of the fresh water eustigmatophyte *Monodus subterraneus*. *Phytochemistry* 2006;**67**:696–701.
- Kim W, Park JM, Gim GH et al. Optimization of culture conditions and comparison of biomass productivity of three green algae. *Bioprocess Biosyst Eng* 2012;**35**:19–27.
- Kitaya Y, Azuma H, Kiyota M. Effect of temperature, carbon dioxide/O₂ concentrations and light intensity on cellular multiplication of microalgae, *Euglena gracilis*. *Adv Space Res* 2005;**35**:1584–8.
- Kommareddy AR, Anderson GA. Study of light as a parameter in the growth of algae in a photo-bio reactor (PBR). ASAE Paper No. 034057. Las Vegas, USA: ASAE, 2003.
- Koru E. Earth food *Spirulina* (*Arthrospira*): production and quality standards. In: El-Samragy Y (ed). *Food Additive*. London: In-Tech, 2012.
- Kumar A, Ergas S, Yuan X et al. Enhanced carbon dioxide fixation and biofuels production via microalgae: recent developments and future directions. *Trends Biotechnol* 2011;**28**:371–80.
- Kumar A, Yuan X, Sahu AK et al. A hollow fiber membrane photobioreactor for carbon dioxide sequestration from combustion gas coupled with wastewater treatment: A process engineering approach. *J Chem Technol Biotechnol* 2010;**85**:387–94.
- Luo HP, Al-Dahhan MH. Verification and validation of CFD simulations for local flow dynamics in a draft tube airlift bioreactor. *Chem Eng Sci* 2011;**66**:907–23.
- Ma M, Yuan D, He Y et al. Effective control of *Poteroochromonas malhamensis* in pilot-scale culture of *Chlorella sorokiniana* GT-1 by maintaining carbon dioxide-mediated low culture pH. *Algal Res* 2017;**26**:436–44.
- Meier L, Pérez R, Azócar L et al. Photosynthetic carbon dioxide uptake by microalgae: An attractive tool for biogas upgrading. *Biomass Bioenerg* 2015;**73**:102–9.
- Mikkelsen M, Jørgensen M, Krebs FC. The teraton challenge. A review of fixation and transformation of carbon dioxide. *Energy Environm Sci* 2010;**3**:43–81.
- Miller SR, Castenholz RW. Evolution of thermotolerance in hot spring cyanobacteria of the genus *Synechococcus*. *Appl Environ Microbiol* 2000;**66**:4222–9.
- Moheimani NR. Long-term outdoor growth and lipid productivity of *Tetraselmis suecica*, *Dunaliella tertiolecta* and *Chlorella* sp. (Chlorophyta) in bag photobioreactors. *J Appl Phycol* 2013;**25**:167–76.
- Mokashi K, Shetty V, George SA et al. Sodium bicarbonate as inorganic carbon source for higher biomass and lipid production integrated carbon capture in *Chlorella vulgaris*. *Achieve Life Sci* 2016;**10**:111–7.
- Molina-Grima E, Belarbi EH, Acién-Fernández FG et al. Recovery of microalgal biomass and metabolites: process options and economics. *Biotechnol Adv* 2003;**20**:491–515.
- Molina-Gima E, Fernández J, Acién FG et al. Tubular photobioreactor design for algal cultures. *J Biotechnol* 2001;**92**:113–31.
- Mooij T, Nejad ZR, van Buren L et al. Effect of photoacclimation on microalgae mass culture productivity. *Algal Res* 2017;**22**:56–67.
- Morales M, Cabello J, Revah M. Gas balances and growth in algal cultures. In: Prokop A, Bajpai R, Zappi M (eds). *Algal Biorefineries*. Vol. 2. Cham: Springer International, 2015, 263–314.
- Moss B. The influence of environmental factors on the distribution of freshwater algae: an experimental study: II. The

- role of pH and the carbon dioxide-bicarbonate system. *J Ecol* 1973;**61**:157–177.
- Nakano Y, Miyatake K, Okuno H et al. Growth of photosynthetic algae *Euglena* in high carbon dioxide conditions and its photosynthetic characteristics. *Acta Hort* 1996;**440**:49–54.
- Nancucheo I, Johnson DB. Acidophilic algae isolated from mine-impacted environments and their roles in sustaining heterotrophic acidophiles. *Front Microbiol* 2012;**3**:325.
- Napan K, Teng L, Quinn JC et al. Impact of heavy metals from flue gas integration with microalgae production. *Algal Res* 2015;**8**:83–88.
- Nikolaou A, Hartmann Ph, Sciandra A et al. Dynamic coupling of photoacclimation and photoinhibition in a model of microalgae growth. *J Theor Biol* 2016;**390**:61–72.
- Norsker NH, Barbosa MJ, Vermuë M et al. Microalgal production – A close look at the economics. *Biotechnol Adv* 2011;**29**:24–27.
- Novis P, Harding JS. Extreme acidophiles: fresh-water algae associated with acid mine drainage. In: Seckbach J (ed). *Algae and Cyanobacteria in Extreme Environments*. Heidelberg: Springer, 2007, 443–63.
- Olaizola M. Commercial development of microalgal biotechnology: from the test tube to the marketplace. *Biomol Eng* 2003;**20**:459–66.
- Oliveira MACL, Monteiro D, Robbs MPC et al. Growth and chemical composition of *Spirulina maxima* and *Spirulina platensis* biomass at different temperatures. *Aquacult Int* 1999;**7**:261–75.
- Ooms MD, Dinh CT, Sargent EH et al. Photon management for augmented photosynthesis. *Nat Commun* 2016;**7**:12699.
- Ono E, Cuello JL. Carbon dioxide mitigation using thermophilic cyanobacteria. *Biosyst Eng* 2007;**96**:129–34.
- Ördög V, Stirk WA, Bálint P et al. Effect of temperature and nitrogen concentration on lipid productivity and fatty acid composition in three *Chlorella* strains. *Algal Res* 2016;**16**:141–9.
- Oswald WJ, Golueke CG. Biological transformation of solar energy. *Adv Appl Microbiol* 1960;**2**:223–62.
- Paliwal C, Mitra M, Bhayani K et al. Abiotic stresses as tools for metabolites in microalgae. *Bioresour Technol* 2017;**244**:1216–26.
- Pandey JP, Pathak N, Tiwari A. Standardization of pH and light intensity for the biomass production of *Spirulina platensis*. *J Algal Biomass Utiln* 2010;**1**:93–102.
- Pires JCM, Alvim-Ferraz MCM, Martins FG et al. Carbon dioxide capture from flue gases using microalgae: Engineering aspects and biorefinery concept. *Renew Sust Energy Rev* 2012;**6**:3043–53.
- Price GD, Badger MR, Woodger FJ et al. Advances in understanding the cyanobacterial carbon dioxide-concentrating-mechanism (CCM): functional components, Ci transporters, diversity, genetic regulation and prospects for engineering into plants. *J Exp Bot* 2008;**59**:1441–61.
- Pulz O, Gross W. Valuable products from biotechnology of microalgae. *Appl Microbiol Biotechnol* 2004;**65**:635–48.
- Ramanna L, Rawat I, Bux F. Light enhancement strategies improve microalgal biomass productivity. *Renew Sust Energy Rev* 2017;**80**:765–73.
- Ras M, Steyer JP, Bernard O. Temperature effect on microalgae: a crucial factor for outdoor production. *Rev Environ Sci Biotechnol* 2013;**12**:153–64.
- Raso S, van Genugten B, Vermuë M et al. Effect of oxygen concentration on the growth of *Nannochloropsis* sp. at low light intensity. *J Appl Phycol* 2012;**24**:863–71.
- Razzak SA, Hossain MM, Lucky RA et al. Integrated carbon dioxide capture, wastewater treatment and biofuel production by microalgae culturing – a review. *Renew Sustain Energy Rev* 2013;**27**:622–53.
- Reboloso MM, García JL, Fernández JM et al. Outdoor continuous culture of *Porphyridium cruentum* in a tubular photobioreactor: quantitative analysis of the daily cyclic variation of culture parameters. *J Biotechnol* 1999;**70**:271–88.
- Richmond A. *Handbook Of Microalgal Culture: Biotechnology and Applied Phycology*. Oxford: Blackwell Science, 2004.
- Rodolfi L, Zittelli GC, Bassi N et al. Microalgae for oil: strain selection, induction of lipid synthesis and outdoor mass cultivation in a low-cost photobioreactor. *Biotechnol Bioeng* 2009;**102**:100–12.
- Rowe OF, Sánchez-España J, Hallberg KB et al. Microbial communities and geochemical dynamics in an extremely acidic, metal-rich stream at an abandoned sulfide mine (Huelva, Spain) underpinned by two functional primary production systems. *Environ. Microbiol* 2007;**9**:1761–71.
- Sánchez L, Rojas A, Revah S et al. Outdoor cultivation of the microalga *Scenedesmus obtusiusculus* for biomass production. XV National Congress of Biotechnology and Bioengineering and 12th International Symposium on the Genetics of Industrial Microorganisms. June 23rd–28th, 2013, Cancún, México. 2013.
- Sánchez-Mirón A, Cerón-García MC, Contreras-Gómez A et al. Shear stress tolerance and biochemical characterization of *Phaeodactylum tricornutum* in quasi steady-state continuous culture in outdoor photobioreactors. *Biochem Eng J* 2003;**16**:287–97.
- Sarada R, Tripathi U, Ravishanker GA. Influence of stress on astaxanthin production in *Haematococcus pluvialis* grown under different culture conditions. *Process Biochem* 2002;**37**:623–7.
- Sato A, Kurano N, Senger H et al. Regulation of energy balance in response to changes in carbon dioxide concentrations and light intensities during growth in extremely-high-carbon dioxide-tolerant green microalgae. *Plant Cell Physiol* 2002;**43**:440–51.
- Seo SH, Ha JSh, Yoo Ch et al. Light intensity as major factor to maximize biomass and lipid productivity of *Ettlia* sp. in carbon dioxide-controlled photoautotrophic chemostat. *Bioresour Technol* 2017;**244**:621–8.
- Sharma G, Kumar M, Ali MI et al. Effect of carbon content, salinity and pH on *Spirulina platensis* for phycocyanin, allophycocyanin and phycoerythrin accumulation. *J Microb Biotechnol* 2014;**6**:202–6.
- Singh J, Gu S. Commercialization potential of microalgae for biofuels production. *Renew Sust Energy Rev* 2010;**14**:2596–610.
- Singh SP, Singh P. Effect of carbon dioxide concentration on algal growth: A review. *Renew Sust Energy Rev* 2014;**38**:172–9.
- Singh UB, Ahluwalia AS. Microalgae: a promising tool for carbon sequestration. *Mitig Adapt Strateg Glob Chang* 2013;**18**:73–95.
- Solovchenko A, Khozin-Goldberg I. High-carbon dioxide tolerance in microalgae: possible mechanisms and implications for biotechnology and bioremediation. *Biotechnol Lett* 2013;**35**:1745–52.
- Sun S, Duan S. Response of marine microalgae *Tetraselmis tetraele* to changes of photoperiods. *Ecol Sci* 2007;**26**:293–7.
- Sun Y, Zwolinska E, Chmielewski AG. Abatement technologies for high concentrations of NO_x and SO₂ removal from exhaust gases: A review. *Crit Rev Environ Sci Technol* 2016;**46**:119–42.
- Sung K D, Lee J S, Shin C S et al. Isolation of a new highly carbon dioxide tolerant fresh water microalgae *Chlorella* sp. KR-1. *Korean J Chem Eng* 1998;**15**:449–50.

- Takagi M, Karseno, Yoshida T. Effect of salt concentration on intracellular accumulation of lipids and triacylglyceride in marine microalgae *Dunaliella* cells. *J Biosci Bioeng* 2006; **101**:223–6.
- Takagi M, Watanabe K, Yamaberi K et al. Limited feeding of potassium nitrate for intracellular lipid and triglyceride accumulation of *Nannochloris* sp. UTEX LB1999. *Appl Microbiol Biotechnol* 2000; **54**:112–7.
- Thawechai T, Cheirsilp B, Louhasakul Y et al. Mitigation of carbon dioxide by oleaginous microalgae for lipids and pigments production: Effect of light illumination and carbon dioxide feeding strategies. *Bioresour Technol* 2016; **219**:139–49.
- Thomas DM, Mechery J, Paulose SV. Carbon dioxide capture strategies from flue gas using microalgae: a review. *Environ Sci Pollut Res* 2016; **23**:16926–40.
- Toledo-Cervantes A, Morales M, Novelo E. et al. Carbon dioxide fixation and lipid storage by *Scenedesmus obtusiusculus*. *Biore-sour Technol* 2013; **130**:652–8.
- Touloupakis E, Cicchi B, Benavides AMS et al. Effect of high pH on growth of *Synechocystis* sp. PCC 6803 cultures and their contamination by golden algae (*Poterioochromonas* sp.). *Appl Microbiol Biotechnol* 2016; **100**:1333.
- Van Bergeijk SA, Salas-Leiton E, Cañavate JP. Low and variable productivity and low efficiency of mass cultures of the hap-tophyte (T-iso) in outdoor tubular photobioreactors. *Aquacult Eng* 2010; **43**:14–23.
- Vonshak A, Torzillo G. Environmental stress physiology. In: Richmond A (ed). *Handbook of Microalgal Culture: Biotechnology and Applied Phycology*. Oxford: Blackwell Science, 2004, 57–82.
- Wang B, Li Y, Wu N et al. carbon dioxide bio-mitigation using microalgae. *Appl Microbiol Biotechnol* 2008; **79**:707–18.
- Weyer KM, Bush DR, Darzins A et al. Theoretical maximum algal oil production. *Bioenergy Res* 2010; **3**:204–13.
- Wijffels RH, Barbosa MJ. An outlook on microalgal biofuels. *Science* 2010; **329**:796–9.
- Wijffels RH, Barbosa MJ, Eppink MHM. Microalgae for the production of bulk chemicals and biofuels. *Biofuel Bioprod Bior* 2010; **4**:287–95.
- Williams PJB, Laurens LML. Microalgae as biodiesel and biomass feedstocks: Review and analysis of the biochemistry, energetics, and economics. *Energy Environ Sci* 2010; **3**:554–90.
- Wolf J, Stephens E, Steinbusch S et al. Multifactorial comparison of photobioreactor geometries in parallel microalgae cultivations. *Algal Res* 2016; **15**:187–201.
- Xin L, Hu H Y, Ke G et al. Effects of different nitrogen and phosphorus concentrations on the growth, nutrient uptake, and lipid accumulation of a freshwater microalga *Scenedesmus* sp. *Bioresour Technol* 2010; **101**:5494–500.
- Ying K, Gilmour DJ, Zimmerman WB. Effects of carbon dioxide and pH on growth of the microalga *Dunaliella salina*. *J Microb Biochem Technol* 2014; **6**:167–73.
- Yen HW, Ho SH, Chen CY et al. carbon dioxide, NO_x and SO_x removal from flue gas via microalgae cultivation: A critical review. *Biotechnol J* 2015; **10**:829–39.
- Yun YS, Lee SB, Park JM et al. Carbon dioxide fixation by algal cultivation using wastewater nutrients. *J Chem Tech Biotechnol* 1997; **69**:451–5.
- Zeng X, Danquah MK, Chen XD et al. Microalgae bioengineering: From carbon dioxide fixation to biofuel production. *Renew Sust Energ Rev* 2011; **15**:3252–60.
- Zhang H, Zeng R, Chen D et al. A pivotal role of vacuolar H⁺-ATPase in regulation of lipid production in *Phaeodactylum tricornutum*. *Sci Rep* 2016; **6**:31319.
- Zhao B, Su Y. Process effect of microalgal-carbon dioxide fixation and biomass production: A review. *Renew Sust Energ Rev* 2014; **31**:121–32.