Table 1. Baseline characteristics of the overall domestic cohort.

Characteristic	Overall cohort (n = 445)
Demographic and background characteristics	
Age (years)	52.0 (45.0, 61.0)
Male sex	313 (70.3)
BMI (kg/m²)	22.49 (20.55, 24.44)
Obesity, n (%)	35 (7.9)
Hypertension, n (%)	51 (11.5)
Diabetes mellitus, n (%)	79 (17.8)
Metabolic comorbidities, n (%)	131 (29.4)
Smoking history	185 (41.6)
Alcohol use history	183 (41.1)
HBV serological, virological and treatment-related characteristics	
HBeAg/anti-HBe status	
HBeAg−/anti-HBe+	313 (70.3)
HBeAg+/anti-HBe−	77 (17.3)
HBeAg−/anti-HBe−	36 (8.1)
HBeAg+/anti-HBe+	19 (4.3)
HBV DNA study category (IU/mL)	**Available n = 442**
<2,000	260 (58.8)
2,000–<20,000	47 (10.6)
≥20,000	135 (30.5)
HBV DNA, $\log_{10}$(IU/mL)	2.70 (2.70, 4.97); n = 437
HBsAg, $\log_{10}$(IU/mL)	2.81 (1.94, 3.16)
Prior antiviral therapy	246 (55.3)
Recorded clinical manifestations	
Ascites	194 (43.6)
EGVH	246 (55.3)
SBP	108 (24.3)
HE	27 (6.1)
HRS	5 (1.1)
Infection	181 (40.7)
Laboratory measurements	
Albumin (g/L)	29.00 (26.00, 34.00)
Total bilirubin (μmol/L)	24.70 (15.70, 41.00)
ALT (U/L)	39.00 (27.00, 57.00)
AST (U/L)	44.00 (30.00, 65.00)
ALP (U/L)	92.00 (67.00, 128.00)
GGT (U/L)	38.00 (24.00, 74.00)
INR	1.37 (1.22, 1.52)

Characteristic	Overall cohort (n = 445)
PT (s)	16.40 (15.10, 18.30)
PTA (%)	61.00 (52.00, 71.00)
Platelet count (10^9/L)	63.00 (44.00, 95.00)
WBC (10^9/L)	3.97 (2.70, 5.95)
Hemoglobin (g/L)	92.00 (73.00, 111.00)
Serum creatinine (μmol/L)	67.00 (56.00, 79.00)
Sodium (mmol/L)	139.00 (137.00, 141.00)
Composite scores	
MELD-Na	12.47 (9.99, 15.54)
ALBI	-1.62 (-2.02, -1.22)
FIB-4	5.76 (3.38, 10.53)
Index hospitalization	
Length of index hospital stay (days)	10.0 (7.0, 15.0)
Observed 1-year outcome	
Study-defined readmission	185 (41.6)

Values are presented as median (interquartile range) or n (%). Percentages were calculated using available nonmissing observations; available n is shown where it differs from 445. Clinical manifestations were not mutually exclusive. Metabolic comorbidities were defined as the presence of obesity, hypertension or diabetes mellitus; obesity was defined as BMI ≥28 kg/m². Retained viral-marker coding, source-scale limitations and derived-index definitions are provided in Supplementary Table S3A and Supplementary Note M1. Abbreviations: BMI, body mass index; HBV, hepatitis B virus; DNA, deoxyribonucleic acid; HBeAg, hepatitis B e antigen; anti-HBe, antibody to hepatitis B e antigen; HBsAg, hepatitis B surface antigen; EGVH, esophagogastric variceal hemorrhage; SBP, spontaneous bacterial peritonitis; HE, hepatic encephalopathy; HRS, hepatorenal syndrome; ALT, alanine aminotransferase; AST, aspartate aminotransferase; ALP, alkaline phosphatase; GGT, gamma-glutamyl transferase; INR, international normalized ratio; PT, prothrombin time; PTA, prothrombin activity; WBC, white blood cell count; MELD-Na, Model for End-Stage Liver Disease-sodium; ALBI, albumin-bilirubin score; FIB-4, fibrosis-4 index.