



BITS Pilani
Dubai Campus

Machine Learning

CS F464

Dr. Pranav M. Pawar

- Hidden Markov Model
 - Markov model
 - Examples
 - Hidden Markov model
 - Examples

- Set of states: $\{s_1, s_2, \dots, s_N\}$
- Process moves from one state to another generating a sequence of states : $s_{i1}, s_{i2}, \dots, s_{ik}, \dots$
- Markov chain property: probability of each subsequent state depends only on what was the previous state:

$$P(s_{ik} \mid s_{i1}, s_{i2}, \dots, s_{ik-1}) = P(s_{ik} \mid s_{ik-1})$$


- To define Markov model, the following probabilities have to be specified:
 - transition probabilities $a_{ij} = P(s_i \mid s_j)$
 - and initial probabilities $\pi_i = P(s_i)$

Markov Model Example 1

innovate

achieve

lead



- Two states
'Rain' and 'Dry'.
- Transition probabilities:
 $P(\text{'Rain'}|\text{'Rain'})=0.3$, $P(\text{'Dry'}|\text{'Rain'})=0.7$,
 $P(\text{'Rain'}|\text{'Dry'})=0.2$, $P(\text{'Dry'}|\text{'Dry'})=0.8$
- Initial probabilities:
 $P(\text{'Rain'})=0.4$, $P(\text{'Dry'})=0.6$.

- By Markov chain property, probability of state sequence can be found by the formula:

$$\begin{aligned} P(s_{i1}, s_{i2}, \dots, s_{ik}) &= P(s_{ik} \mid s_{i1}, s_{i2}, \dots, s_{ik-1}) P(s_{i1}, s_{i2}, \dots, s_{ik-1}) \\ &= P(s_{ik} \mid s_{ik-1}) P(s_{i1}, s_{i2}, \dots, s_{ik-1}) = \dots \\ &= P(s_{ik} \mid s_{ik-1}) P(s_{ik-1} \mid s_{ik-2}) \dots P(s_{i2} \mid s_{i1}) P(s_{i1}) \end{aligned}$$

- For example

$$P(\{\text{'Dry'}, \text{'Dry'}, \text{'Rain'}, \text{'Rain'}\})$$

$$= P(\text{'Rain'} \mid \text{'Rain'}) P(\text{'Rain'} \mid \text{'Dry'}) P(\text{'Dry'} \mid \text{'Dry'}) P(\text{'Dry'})$$

$$= 0.3 * 0.2 * 0.8 * 0.6$$

$$\begin{aligned} P(\{\text{Dry}(s_1), \text{Rain}(s_2), \text{Rain}(s_3), \text{Dry}(s_4)\}) &= P(\text{Dry} \mid \text{Rain}) P(\text{Rain} \mid \text{Rain}) \\ P(\text{Rain} \mid \text{Dry}) P(\text{Dry}) &= \end{aligned}$$

Example 2

innovate

achieve

lead

- In analysing switching by Business Class customers between airlines the following data has been obtained by XYZ airways:

Next flight by			
Last flight by		XYZ	Competition
	XYZ	0.85	0.15
	Competition	0.10	0.90

- For example if the last flight by a Business Class customer was by XYZ the probability that their next flight is by XYZ is 0.85. Business Class customers make 2 flights a year on average.
- Currently XYZ have 30% of the Business Class market. What would you forecast XYZ's share of the Business Class market to be after two years?

Example 2

innovate

achieve

lead

- We have the initial system state s_1 given by $s_1 = [0.30, 0.70]$ and the transition matrix P is given by

$$P = \begin{bmatrix} 0.85 & 0.15 \\ 0.10 & 0.90 \end{bmatrix}^2 = \begin{bmatrix} 0.7375 & 0.2625 \\ 0.1750 & 0.8250 \end{bmatrix}$$


- Here we take square because customer make 2 flights a year on a average.
- After one year, $s_2 = s_1 P = [0.34375 \ 0.65625]$
- After two years, $s_3 = s_2 P = [0.368 \ 0.632]$
- So, after two years have elapses XYZ share of business class market is 36.8%

Example 3

create

achieve

lead



- Given that today the weather is ☀️, what's the probability that tomorrow is ☀️ and the day after is ☁️?

Using the Markov assumption and the probabilities in table 1, this translates into:

$$\begin{aligned} P(q_2 = \text{☀️}, q_3 = \text{☁️} | q_1 = \text{☀️}) &= P(q_3 = \text{☁️} | q_2 = \text{☀️}, q_1 = \text{☀️}) \cdot P(q_2 = \text{☀️} | q_1 = \text{☀️}) \\ &= P(q_3 = \text{☁️} | q_2 = \text{☀️}) \cdot P(q_2 = \text{☀️} | q_1 = \text{☀️}) \quad (\text{Markov assumption}) \\ &= 0.05 \cdot 0.8 \\ &= 0.04 \end{aligned}$$

Example 3

innovate

achieve

lead

2. Assume, the weather yesterday was $q_1 = \text{☁}$, and today it is $q_2 = \text{☁}$, what is the probability that tomorrow it will be $q_3 = \text{☀}$?

$$\begin{aligned} P(q_3 = \text{☀} | q_2 = \text{☁}, q_1 = \text{☁}) &= P(q_3 = \text{☀} | q_2 = \text{☁}) \quad (\text{Markov assumption}) \\ &= 0.2. \end{aligned}$$

Hidden Markov Model

innovate

achieve

lead

- Set of states: $\{s_1, s_2, \dots, s_N\}$
- Process moves from one state to another generating a sequence of states : $s_{i1}, s_{i2}, \dots, s_{ik}, \dots$
- Markov chain property: probability of each subsequent state depends only on what was the previous state:

$$P(s_{ik} \mid s_{i1}, s_{i2}, \dots, s_{ik-1}) = P(s_{ik} \mid s_{ik-1})$$


- States are not visible, but each state randomly generates one of M observations (or visible states) $\{v_1, v_2, \dots, v_M\}$
- To define hidden Markov model, the following probabilities have to be specified:
 - matrix of transition probabilities $A=(a_{ij})$, $a_{ij}= P(s_i \mid s_j)$
 - matrix of observation probabilities $B=(b_i(v_m))$, $b_i(v_m) = P(v_m \mid s_i)$
 - vector of initial probabilities $\pi=(\pi_i)$, $\pi_i = P(s_i)$
- Model is represented as, $M=(A, B, \pi)$

Example 1 HMM

innovate

achieve

lead



Example 1 HMM

innovate

achieve

lead

- Two states : 'Low' and 'High' atmospheric pressure.
- Two observations : 'Rain' and 'Dry'.
- Transition probabilities: $P(\text{'Low'}|\text{'Low'})=0.3$,
 $P(\text{'High'}|\text{'Low'})=0.7$, $P(\text{'Low'}|\text{'High'})=0.2$,
 $P(\text{'High'}|\text{'High'})=0.8$
- Observation probabilities : $P(\text{'Rain'}|\text{'Low'})=0.6$,
 $P(\text{'Dry'}|\text{'Low'})=0.4$, $P(\text{'Rain'}|\text{'High'})=0.4$,
 $P(\text{'Dry'}|\text{'High'})=0.3$.
- Initial probabilities: say $P(\text{'Low'})=0.4$, $P(\text{'High'})=0.6$.

Example 1 HMM

innovate

achieve

lead

- Suppose we want to calculate a probability of a sequence of observations in our example, {'Dry','Rain'}.
- Consider all possible hidden state sequences:

$$P(\{\text{'Dry','Rain'}\}) = P(\{\text{'Dry','Rain'}\}, \{\text{'Low','Low'}\}) + P(\{\text{'Dry','Rain'}\}, \{\text{'Low','High'}\}) + P(\{\text{'Dry','Rain'}\}, \{\text{'High','Low'}\}) + P(\{\text{'Dry','Rain'}\}, \{\text{'High','High'}\})$$

$$\begin{aligned}\text{First term} &\Rightarrow P(\{\text{'Dry','Rain'}\}, \{\text{'Low','Low'}\}) = \\ &P(\{\text{'Dry','Rain'}\} \mid \{\text{'Low','Low'}\}) P(\{\text{'Low','Low'}\}) = \\ &P(\text{'Dry'} \mid \text{'Low'})P(\text{'Rain'} \mid \text{'Low'}) P(\text{'Low'})P(\text{'Low'} \mid \text{'Low'}) \\ &= 0.4 * 0.6 * 0.4 * 0.3 = 0.0288\end{aligned}$$

$$\begin{aligned}\text{Second term} &\Rightarrow P(\{\text{'Dry','Rain'}\}, \{\text{'Low','High'}\}) = \\ &P(\{\text{'Dry','Rain'}\} \mid \{\text{'Low','High'}\}) P(\{\text{'Low','High'}\}) = \\ &P(\text{'Dry'} \mid \text{'Low'})P(\text{'Rain'} \mid \text{'High'}) P(\text{'High'})P(\text{'Low'} \mid \text{'High'}) \\ &= 0.4 * 0.4 * 0.6 * 0.2 = 0.0192\end{aligned}$$

Example 1 HMM

innovate

achieve

lead

Third term =>

$$P(\{\text{'Dry'}, \text{'Rain'}\}, \{\text{'High'}, \text{'Low'}\}) =$$

$$P(\{\text{'Dry'}, \text{'Rain'}\} \mid \{\text{'High'}, \text{'Low'}\}) P(\{\text{'High'}, \text{'Low'}\}) =$$

$$P(\text{'Dry'} \mid \text{'High'}) P(\text{'Rain'} \mid \text{'Low'}) P(\text{'Low'}) P(\text{'High'} \mid \text{'Low'}) \\ = 0.3 * 0.6 * 0.4 * 0.7 = 0.0504$$

$$\text{Fourth term} \Rightarrow P(\{\text{'Dry'}, \text{'Rain'}\}, \{\text{'High'}, \text{'High'}\}) =$$

$$P(\{\text{'Dry'}, \text{'Rain'}\} \mid \{\text{'High'}, \text{'High'}\}) P(\{\text{'High'}, \text{'High'}\}) =$$

$$P(\text{'Dry'} \mid \text{'High'}) P(\text{'Rain'} \mid \text{'High'}) P(\text{'High'}) P(\text{'High'} \mid \text{'High'}) \\ = 0.3 * 0.4 * 0.6 * 0.8 = 0.0576$$

Therefore,

$$P(\{\text{'Dry'}, \text{'Rain'}\}) = 0.0288 + 0.0192 + 0.0504 + 0.0576 \\ = 0.156$$







Example 2 HMM


innovate

achieve

lead

Given,

Today's weather	Tomorrow's weather		
			
	0.8	0.05	0.15
	0.2	0.6	0.2
	0.2	0.3	0.5



Observed Probabilities,

Weather	Probability of umbrella
Sunny	0.1
Rainy	0.8
Foggy	0.3

Example 2 HMM

innovate

achieve

lead

- . Suppose the day you were locked in it was sunny. The next day, the caretaker carried an umbrella into the room. You would like to know, what the weather was like on this second day.

First we calculate the likelihood for the second day to be sunny:

$$\begin{aligned} L(q_2 = \text{☀️} | q_1 = \text{☀️}, x_2 = \text{☂️}) &= P(x_2 = \text{☂️} | q_2 = \text{☀️}) \cdot P(q_2 = \text{☀️} | q_1 = \text{☀️}) \\ &= 0.1 \cdot 0.8 = 0.08, \end{aligned}$$

then for the second day to be rainy:

$$\begin{aligned} L(q_2 = \text{☁️} | q_1 = \text{☀️}, x_2 = \text{☂️}) &= P(x_2 = \text{☂️} | q_2 = \text{☁️}) \cdot P(q_2 = \text{☁️} | q_1 = \text{☀️}) \\ &= 0.8 \cdot 0.05 = 0.04, \end{aligned}$$

and finally for the second day to be foggy:

$$\begin{aligned} L(q_2 = \text{🌫️} | q_1 = \text{☀️}, x_2 = \text{☂️}) &= P(x_2 = \text{☂️} | q_2 = \text{🌫️}) \cdot P(q_2 = \text{🌫️} | q_1 = \text{☀️}) \\ &= 0.3 \cdot 0.15 = 0.045. \end{aligned}$$

Thus, although the caretaker did carry an umbrella, it is most likely that on the second day the weather was sunny.



BITS Pilani
Dubai Campus



Thank You!