An Exponential Succinctness Gap between Three-Variable Logic and the Calculus of Relations

Yuya Uezato　Yoshiki Nakamura
CyberAgent, Inc.　Chiba University

STOC 2027 submission　(author video for the PC)

論文 v8 との差分（用語）：

- fold → flip
- notice → (newly) distinguish：同じ色の辺で真偽が食い違う。notice では「気づいてどうなるか」（そのゲートで初めて区別が生まれる）が伝わらない
- reference structure → $\mathcal{M}_0$ ／ read・witness edge → x-y・x-z edge
- color-determined（v8）→ color-invariant。bisimulation-invariant などと同じ「同色の点対で値が同じ」を意図

論文 v8 との差分（内容）：


- Key Lemma「d 個を見分ける項は合成 $\geq d/2$」論文でもこの形で先に述べるとよい？
  （v8：各 fold はどこかの合成で notice される ＋ 合成 1 つで高々 2）
- distinguish を定義として明示（v8 は prediction と notice で定義）
- 省略：ループ（$(\hbar, \hbar)$ を含む）、端点 witness（p.11 で "see the paper"）。Cor 1.3（CoR 等式）、Cor 1.8・1.9（固定サイズ・乱択・ランダム入力）、項サイズ $(2 + o(1))k2^k$
- 名前のみ：Thm 1.4（FO^3 vs FO^2）、Prop 1.5（CoR vs FO^2）、Thm 1.6（ARA(3)）、Prop 1.7（RTC）は p.4 で結果の名前だけ
- 追加：導入の FO^k の例（3-cycle、推移性、closed walk）とモデル検査の表

補足：

- 投稿版の論文が確定したら定理番号を入れる（v8 の番号）：p.4 の定理 Thm 1.1、各項目 Thm 1.4・Prop 1.5・Thm 1.6・Prop 1.7、Key Lemma → Lemma 3.4・3.5、p.11 の "see the paper" → Lemma 3.5

k-variable logic FO^k

FO^k: first-order logic with k variables.
For example, in FO^3: $\varphi(x,y) := \exists z \left(E(x,y) \wedge E(y,z) \wedge E(z,x)\right)$ (triangle detection)



FO^3 also expresses, e.g.,

- transitivity: $\forall x \forall y \forall z \, (E(x,y) \wedge E(y,z) \rightarrow E(x,z))$,
- closed walk of length 4: $\exists x \exists y \left(E(x,y) \wedge \exists z \, (E(y,z) \wedge \exists y \, (E(z,y) \wedge E(y,x)))\right)$.

Model checking: $\mathcal{M} \models \varphi(a,b)$? (given $\mathcal{M}$ with n elements, a formula φ, and $a, b \in \mathcal{M}$)


☞ $O(|\varphi| \cdot n^k)$ for FO^k by table memoization [Var95, PODS]
E.g., $O(n^3)$ for φ above: try all x, y, z

ページ概要：

- FO^k とモデル検査の問題設定
- 素直に計算すると $O(|\varphi| \cdot n^k)$

k-variable logic FO^k

FO^k: first-order logic with k variables.
For example, in FO^3: $\varphi(x,y) := \exists z \left(E(x,y) \wedge E(y,z) \wedge E(z,x)\right)$ (triangle detection)



FO^3 also expresses, e.g.,

- transitivity: $\forall x \forall y \forall z \, (E(x,y) \wedge E(y,z) \to E(x,z))$,
- closed walk of length 4: $\exists x \exists y \left(E(x,y) \wedge \exists z \, (E(y,z) \wedge \exists y \, (E(z,y) \wedge E(y,x)))\right)$.

Model checking: $\mathcal{M} \models \varphi(a,b)$? (given $\mathcal{M}$ with n elements, a formula φ, and $a, b \in \mathcal{M}$)

👉 $O(|\varphi| \cdot n^k)$ for FO^k by table memoization [Var95, PODS]
E.g., $O(n^3)$ for φ above: try all x, y, z

ページ概要：

- FO^k とモデル検査の問題設定
- 素直に計算すると $O(|\varphi| \cdot n^k)$

FO^k model checking: $O(n^k)$ is 'optimal' for $k \geq 4$; the exception is FO^3

We consider binary relation structures with n elements:

syntax	model checking	
FO^k, $k \geq 4$	no $O(n^{k-\varepsilon})$ under Hyperclique Hypothesis	[BS21, ICALP] [NU27, PODS]
FO^3	$2^{O(\lvert\varphi\rvert)} \cdot n^\omega$ no $2^{o(\lvert\varphi\rvert)} \cdot n^{\omega-\varepsilon}$ under 3-OV Hypothesis	[NU27, PODS] [Wil14, CSL-LICS, for PNF] [NU27, PODS]
CoR [Tar41, JSL] Tarski's Calculus of Relations	$O(\lvert\varphi\rvert \cdot n^\omega)$	

Boolean matrix algorithm of FO^3 model checking via $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ [Tar41, JSL; TG87]:

$$\exists z.\ P(x,z) \wedge Q(z,y) \quad \rightsquigarrow \quad (P;Q)(x,y)$$

composition = Boolean matrix multiplication

To obtain the form of $\exists z.\ \cdots \wedge \cdots$, we use DNF translation.
$\rightsquigarrow$ $2^{O(|\varphi|)}$ compositions are used [NJ+22, JLAMP].

ページ概要：

- $k \geq 4$ では n^k が最適。FO^3 は特別に CoR への翻訳で n^ω まで下がる
- ただし翻訳で合成が $2^{O(|\varphi|)}$ 個に増える

FO^k model checking: $O(n^k)$ is 'optimal' for $k \geq 4$; the exception is FO^3

We consider binary relation structures with n elements:

syntax	model checking	
FO^k, $k \geq 4$	no $O(n^{k-\varepsilon})$ under Hyperclique Hypothesis	[BS21, ICALP] [NU27, PODS]
FO^3	$2^{O(\lvert\varphi\rvert)} \cdot n^\omega$ no $2^{o(\lvert\varphi\rvert)} \cdot n^{\omega-\varepsilon}$ under 3-OV Hypothesis	[NU27, PODS] [Wil14, CSL-LICS, for PNF] [NU27, PODS]
CoR [Tar41, JSL] Tarski's Calculus of Relations	$O(\lvert\varphi\rvert \cdot n^\omega)$	

Boolean matrix algorithm of FO^3 model checking via $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ [Tar41, JSL; TG87]:

$$\exists z.\ P(x,z) \wedge Q(z,y) \quad \rightsquigarrow \quad (P;Q)(x,y)$$

composition = Boolean matrix multiplication

To obtain the form of $\exists z.\ - \wedge -$, we use DNF translation.
$\rightsquigarrow$ $2^{O(\lvert\varphi\rvert)}$ compositions are used [Nak22, JLAMP].

ページ概要：

- $k \geq 4$ では n^k が最適。FO^3 は特別に CoR への翻訳で n^ω まで下がる
- ただし翻訳で合成が $2^{O(|\varphi|)}$ 個に増える

FO^k model checking: $O(n^k)$ is 'optimal' for $k \geq 4$; the exception is FO^3

We consider binary relation structures with n elements:

syntax	model checking	
FO^k, $k \geq 4$	no $O(n^{k-\varepsilon})$ under Hyperclique Hypothesis	[BS21, ICALP] [NU27, PODS]
FO^3	$2^{O(\lvert\varphi\rvert)} \cdot n^\omega$ no $2^{o(\lvert\varphi\rvert)} \cdot n^{\omega-\varepsilon}$ under 3-OV Hypothesis	[NU27, PODS] [Wil14, CSL-LICS, for PNF] [NU27, PODS]
CoR [Tar41, JSL] Tarski's Calculus of Relations	$O(\lvert\varphi\rvert \cdot n^\omega)$	

Boolean matrix algorithm of FO^3 model checking via $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ [Tar41, JSL; TG87]:

$$\exists z.\ P(x,z) \wedge Q(z,y) \quad \rightsquigarrow \quad (P;Q)(x,y)$$

composition = Boolean matrix multiplication

To obtain the form of $\exists z.\ - \wedge -$, we use DNF translation.
$\rightsquigarrow$ $2^{O(\lvert\varphi\rvert)}$ compositions are used [Nak22, JLAMP].

ページ概要：

- $k \geq 4$ では n^k が最適。FO^3 は特別に CoR への翻訳で n^ω まで下がる
- ただし翻訳で合成が $2^{O(|\varphi|)}$ 個に増える

FO^k model checking: $O(n^k)$ is 'optimal' for $k \geq 4$; the exception is FO^3

We consider binary relation structures with n elements:

syntax	model checking	
FO^k, $k \geq 4$	no $O(n^{k-\varepsilon})$ under Hyperclique Hypothesis	[BS21, ICALP] [NU27, PODS]
FO^3	$2^{O(\lvert\varphi\rvert)} \cdot n^\omega$ no $2^{o(\lvert\varphi\rvert)} \cdot n^{\omega-\varepsilon}$ under 3-OV Hypothesis	[NU27, PODS] [Wil14, CSL-LICS, for PNF] [NU27, PODS]
CoR [Tar41, JSL] Tarski's Calculus of Relations	$O(\lvert\varphi\rvert \cdot n^\omega)$	

Boolean matrix algorithm of FO^3 model checking via $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ [Tar41, JSL; TG87]:

$$\exists z.\ P(x,z) \wedge Q(z,y) \quad \rightsquigarrow \quad (P;Q)(x,y)$$

composition = Boolean matrix multiplication

To obtain the form of $\exists z.\ - \wedge -$, we use DNF translation.
$\rightsquigarrow$ $2^{O(\lvert\varphi\rvert)}$ compositions are used [Nak22, JLAMP].

ページ概要：

- $k \geq 4$ では n^k が最適。FO^3 は特別に CoR への翻訳で n^ω まで下がる
- ただし翻訳で合成が $2^{O(\lvert\varphi\rvert)}$ 個に増える

Are $2^{O(|\varphi|)}$ compositions unavoidable?

DNF translation is used, for example:

$$\exists z.\ \big(P_1(x,z) \vee Q_1(z,y)\big) \wedge \cdots \wedge \big(P_k(x,z) \vee Q_k(z,y)\big)$$

$$\rightsquigarrow_{\text{(DNF)}}\ \exists z.\ \bigvee_{A \subseteq [k]} \Big(\bigwedge_{i \in A} P_i(x,z) \wedge \bigwedge_{i \notin A} Q_i(z,y) \Big)$$

$$\rightsquigarrow_{\text{(to CoR)}}\ \Big(\bigcup_{A \subseteq [k]} \Big(\bigcap_{i \in A} P_i \Big) ; \Big(\bigcap_{i \notin A} Q_i \Big) \Big)(x,y)$$

(This translation uses 2^k compositions, since the number of subsets $A \subseteq [k]$ is 2^k.)

Problem (Open [FGL+15, Inf. Sci.; Nak22, JLAMP; NU27, PODS])

Can we prove that there is no polynomial-size translation from FO^3 *to* CoR*?*

Partial answer [Nak22, JLAMP]: yes for **existential** CoR [Nak23, LICS] ($\equiv$ existential FO^3), via preservation under **extensions**, which fails for full CoR.

ページ概要：

- 指数個の合成は避けられるか？ は未解決の問題であった

Are $2^{O(|\varphi|)}$ compositions unavoidable?

DNF translation is used, for example:

$$\exists z.\ \big(P_1(x,z) \vee Q_1(z,y)\big) \wedge \cdots \wedge \big(P_k(x,z) \vee Q_k(z,y)\big)$$

$$\rightsquigarrow_{\text{(DNF)}}\ \exists z.\ \bigvee_{A \subseteq [k]} \Big(\bigwedge_{i \in A} P_i(x,z) \wedge \bigwedge_{i \notin A} Q_i(z,y) \Big)$$

$$\rightsquigarrow_{\text{(to CoR)}}\ \Big(\bigcup_{A \subseteq [k]} \big(\bigcap_{i \in A} P_i \big) ; \big(\bigcap_{i \notin A} Q_i \big) \Big)(x,y)$$

(This translation uses 2^k compositions, since the number of subsets $A \subseteq [k]$ is 2^k.)

Problem (Open [FGL+15, Inf. Sci.; Nak22, JLAMP; NU27, PODS])

Can we prove that there is no polynomial-size translation from FO^3 *to* CoR*?*

Partial answer [Nak22, JLAMP]: yes for **existential** CoR [Nak23, LICS] ($\equiv$ existential FO^3), via preservation under **extensions**, which fails for full CoR.

ページ概要：

- 指数個の合成は避けられるか？ は未解決の問題であった

Are $2^{O(|\varphi|)}$ compositions unavoidable?

DNF translation is used, for example:

$$\begin{aligned} & \exists z.\ \big(P_1(x,z) \vee Q_1(z,y)\big) \wedge \cdots \wedge \big(P_k(x,z) \vee Q_k(z,y)\big) \\ \rightsquigarrow_{\text{(DNF)}} \ & \exists z.\ \bigvee_{A \subseteq [k]} \Big(\bigwedge_{i \in A} P_i(x,z) \wedge \bigwedge_{i \notin A} Q_i(z,y) \Big) \\ \rightsquigarrow_{\text{(to CoR)}} \ & \Big(\bigcup_{A \subseteq [k]} \Big(\bigcap_{i \in A} P_i \Big) ; \Big(\bigcap_{i \notin A} Q_i \Big) \Big)(x,y) \end{aligned}$$

(This translation uses 2^k compositions, since the number of subsets $A \subseteq [k]$ is 2^k.)

Problem (Open [FGL+15, Inf. Sci.; Nak22, JLAMP; NU27, PODS])

Can we prove that there is no polynomial-size translation from FO^3 to CoR?

Partial answer [Nak22, JLAMP]: yes for **existential** CoR [Nak23, LICS] ($\equiv$ existential FO^3), via preservation under **extensions**, which fails for full CoR.

ページ概要：

- 指数個の合成は避けられるか？ は未解決の問題であった

Are $2^{O(|\varphi|)}$ compositions unavoidable?

DNF translation is used, for example:

$$\exists z.\ \big(P_1(x,z) \vee Q_1(z,y)\big) \wedge \cdots \wedge \big(P_k(x,z) \vee Q_k(z,y)\big)$$
$$\rightsquigarrow_{\text{(DNF)}}\ \exists z.\ \bigvee_{A \subseteq [k]} \Big(\bigwedge_{i \in A} P_i(x,z) \wedge \bigwedge_{i \notin A} Q_i(z,y) \Big)$$
$$\rightsquigarrow_{\text{(to CoR)}}\ \Big(\bigcup_{A \subseteq [k]} \Big(\bigcap_{i \in A} P_i \Big) ; \Big(\bigcap_{i \notin A} Q_i \Big) \Big)(x,y)$$

(This translation uses 2^k compositions, since the number of subsets $A \subseteq [k]$ is 2^k.)

Problem (Open [FGL+15, Inf. Sci.; Nak22, JLAMP; NU27, PODS])

Can we prove that there is no polynomial-size translation from FO^3 to CoR?

Partial answer [Nak22, JLAMP]: yes for **existential** CoR [Nak23, LICS] ($\equiv$ existential FO^3), via preservation under **extensions**, which fails for full CoR.

ページ概要：

- 指数個の合成は避けられるか？ は未解決の問題であった

Are $2^{O(|\varphi|)}$ compositions unavoidable?

DNF translation is used, for example:

$$\exists z.\ \big(P_1(x,z) \vee Q_1(z,y)\big) \wedge \cdots \wedge \big(P_k(x,z) \vee Q_k(z,y)\big)$$

$$\rightsquigarrow_{\text{(DNF)}}\ \exists z.\ \bigvee_{A \subseteq [k]} \Big(\bigwedge_{i \in A} P_i(x,z) \wedge \bigwedge_{i \notin A} Q_i(z,y) \Big)$$

$$\rightsquigarrow_{\text{(to CoR)}}\ \Big(\bigcup_{A \subseteq [k]} \Big(\bigcap_{i \in A} P_i \Big) ; \Big(\bigcap_{i \notin A} Q_i \Big) \Big)(x,y)$$

(This translation uses 2^k compositions, since the number of subsets $A \subseteq [k]$ is 2^k.)

Problem (Open [FGL+15, Inf. Sci.; Nak22, JLAMP; NU27, PODS])

Can we prove that there is no polynomial-size translation from FO^3 to CoR?

Partial answer [Nak22, JLAMP]: yes for **existential** CoR [Nak23, LICS] ($\equiv$ existential FO^3), via preservation under **extensions**, which fails for full CoR.

ページ概要：

- 指数個の合成は避けられるか？ は未解決の問題であった

Outline

1. Our results

2. Technical overview

3. Conclusion

Our results

The answer is **YES!**

Thm. (This paper)

There is no polynomial-size translation from FO^3 *to* CoR.
More precisely, there are FO^3 formulas $\varphi_1, \varphi_2, \ldots$ with $|\varphi_k| = O(k)$ such that every CoR term equivalent to φ_k needs $\geq 2^{k-1}$ compositions, even with shared subterms.

Also in this paper: exponential lower bounds for

- $\mathrm{FO}^3 \rightsquigarrow \mathrm{FO}^2$, binary (before: FO^k vs. FO^{k-1} for $k \geq 4$ [VS24, LICS])
- $\mathrm{CoR} \rightsquigarrow \mathrm{FO}^2$
- ARA(3) $\rightsquigarrow$ matrix circuits asked by [BGV19, arXiv; BGV22, AMAI]
- $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ with RTC (reflexive-transitive closure)

Our proof technique applies to all of them!

ページ概要：

- $|\varphi_k| = O(k)$ の式に、合成 $\geq 2^{k-1}$ 個が必須
- 同じ手法が他の設定（FO^2、行列言語、RTC 付き CoR など）にも適用できる

Our results

The answer is **YES!**

Thm. (This paper)

There is no polynomial-size translation from FO^3 *to* CoR.
More precisely, there are FO^3 formulas $\varphi_1, \varphi_2, \ldots$ with $|\varphi_k| = O(k)$ such that every CoR term equivalent to φ_k needs $\geq 2^{k-1}$ compositions, even with shared subterms.

Also in this paper: exponential lower bounds for

- $\mathrm{FO}^3 \rightsquigarrow \mathrm{FO}^2$, binary (before: FO^k vs. FO^{k-1} for $k \geq 4$ [VS24, LICS])
- $\mathrm{CoR} \rightsquigarrow \mathrm{FO}^2$
- ARA(3) $\rightsquigarrow$ matrix circuits asked by [BGV19, arXiv; BGV22, AMAI]
- $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ with RTC (reflexive-transitive closure)

Our proof technique applies to all of them!

ページ概要：

- $|\varphi_k| = O(k)$ の式に、合成 $\geq 2^{k-1}$ 個が必須
- 同じ手法が他の設定（FO^2、行列言語、RTC 付き CoR など）にも適用できる

Our results

The answer is **YES!**

Thm. (This paper)

There is no polynomial-size translation from FO^3 *to* CoR.
More precisely, there are FO^3 formulas $\varphi_1, \varphi_2, \ldots$ with $|\varphi_k| = O(k)$ such that every CoR term equivalent to φ_k needs $\geq 2^{k-1}$ compositions, even with shared subterms.

Also in this paper: exponential lower bounds for

- $\mathrm{FO}^3 \rightsquigarrow \mathrm{FO}^2$, binary (before: FO^k vs. FO^{k-1} for $k \geq 4$ [VS24, LICS])
- $\mathrm{CoR} \rightsquigarrow \mathrm{FO}^2$
- ARA(3) $\rightsquigarrow$ matrix circuits asked by [BGV19, arXiv; BGV22, AMAI]
- $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ with RTC (reflexive-transitive closure)

Our proof technique applies to all of them!

ページ概要：

- $|\varphi_k| = O(k)$ の式に、合成 $\geq 2^{k-1}$ 個が必須
- 同じ手法が他の設定（FO^2、行列言語、RTC 付き CoR など）にも適用できる

Our results

The answer is **YES!**

Thm. (This paper)

There is no polynomial-size translation from FO^3 *to* CoR.
More precisely, there are FO^3 formulas $\varphi_1, \varphi_2, \ldots$ with $|\varphi_k| = O(k)$ such that every CoR term equivalent to φ_k needs $\geq 2^{k-1}$ compositions, even with shared subterms.

Also in this paper: exponential lower bounds for

- $\mathrm{FO}^3 \rightsquigarrow \mathrm{FO}^2$, binary (before: FO^k vs. FO^{k-1} for $k \geq 4$ [VS24, LICS])
- $\mathrm{CoR} \rightsquigarrow \mathrm{FO}^2$
- ARA(3) $\rightsquigarrow$ matrix circuits asked by [BGV19, arXiv; BGV22, AMAI]
- $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ with RTC (reflexive-transitive closure)

Our proof technique applies to all of them!

ページ概要：

- $|\varphi_k| = O(k)$ の式に、合成 $\geq 2^{k-1}$ 個が必須
- 同じ手法が他の設定（FO^2、行列言語、RTC 付き CoR など）にも適用できる

Our results

The answer is **YES!**

Thm. (This paper)

There is no polynomial-size translation from FO^3 *to* CoR.
More precisely, there are FO^3 formulas $\varphi_1, \varphi_2, \ldots$ with $|\varphi_k| = O(k)$ such that every CoR term equivalent to φ_k needs $\geq 2^{k-1}$ compositions, even with shared subterms.

Also in this paper: exponential lower bounds for

- $\mathrm{FO}^3 \rightsquigarrow \mathrm{FO}^2$, binary (before: FO^k vs. FO^{k-1} for $k \geq 4$ [VS24, LICS])
- $\mathrm{CoR} \rightsquigarrow \mathrm{FO}^2$
- ARA(3) $\rightsquigarrow$ matrix circuits asked by [BGV19, arXiv; BGV22, AMAI]
- $\mathrm{FO}^3 \rightsquigarrow \mathrm{CoR}$ with RTC (reflexive-transitive closure)


Our proof technique applies to all of them!

ページ概要：

- $|\varphi_k| = O(k)$ の式に、合成 $\geq 2^{k-1}$ 個が必須
- 同じ手法が他の設定（FO^2、行列言語、RTC 付き CoR など）にも適用できる

The hard formula G_k

$$G_k(x,y) \coloneqq \exists z \bigwedge_{i=1}^{k} \left[(P_i(x,y) \wedge Q_i(x,z)) \vee (Q_i(x,y) \wedge P_i(x,z)) \right] \qquad (|G_k| = O(k))$$



$G_k(x,y)$ holds iff there is z such that
for every $i \in [k]$,
the x-y and x-z edges have labels P_i and Q_i, in either order.

CoR: 2^k **compositions** (by taking DNF, as in the example before)


$$G_k \equiv \bigcup_{A \subseteq [k]} \left[\left(\bigcap_{i \in A} P_i \cap \bigcap_{i \notin A} Q_i \right) \cap \left(\left(\bigcap_{i \in A} Q_i \cap \bigcap_{i \notin A} P_i \right) ; \mathbf{1} \right) \right]$$

ページ概要：

- 下界を与える式 G_k
- その DNF が合成 2^k 個であること（2 倍を除いて最適）

The hard formula G_k

$$G_k(x,y) \coloneqq \exists z \bigwedge_{i=1}^{k} \left[(P_i(x,y) \wedge Q_i(x,z)) \vee (Q_i(x,y) \wedge P_i(x,z)) \right] \qquad (|G_k| = O(k))$$



$G_k(x,y)$ holds iff there is z such that
for every $i \in [k]$,
the x-y and x-z edges have labels P_i and Q_i, in either order.

CoR: 2^k **compositions** (by taking DNF, as in the example before)

$$G_k \equiv \bigcup_{A \subseteq [k]} \left[\left(\bigcap_{i \in A} P_i \cap \bigcap_{i \notin A} Q_i \right) \cap \left(\left(\bigcap_{i \in A} Q_i \cap \bigcap_{i \notin A} P_i \right); \mathbf{1} \right) \right]$$

ページ概要：

- 下界を与える式 G_k
- その DNF が合成 2^k 個であること（2 倍を除いて最適）

Outline

1. Our results

2. Technical overview

3. Conclusion

Proof outline (high level)

Goal Lemma Every CoR term equivalent to G_k needs $2^{\Omega(k)}$ compositions.

Approach We construct structures $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ such that

- G_k distinguishes all of $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$:

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
G_k	T	F	F	$\cdots$	F	2^k	$\geq 2^k/2$

- **Key Lemma** If a CoR term t distinguishes d structures among $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$, then t needs $\geq d/2$ compositions.

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\mathcal{M}_3$	$\mathcal{M}_4$	$\mathcal{M}_5$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
t	T	F	T	F	T	T	$\cdots$	T	2	≥ 1
s	T	T	T	T	F	F	$\cdots$	F	2	≥ 1

☞ If $t \equiv G_k$, then t needs $\geq 2^k/2$ compositions.

TASK How can we construct such $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$?

ページ概要：

- 証明の方針：G_k が区別する構造 $\mathcal{M}_0, \mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ を作る
- Key Lemma：d 個を区別するには合成 $\geq d/2$ 個 $\rightsquigarrow$ G_k には合成 $\geq 2^{k-1}$ 個

Proof outline (high level)

Goal Lemma Every CoR term equivalent to G_k needs $2^{\Omega(k)}$ compositions.

Approach We construct structures $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ such that

- G_k distinguishes all of $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$:

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
G_k	T	F	F	$\cdots$	F	2^k	$\geq 2^k/2$

- **Key Lemma** If a CoR term t distinguishes d structures among $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$, then t needs $\geq d/2$ compositions.

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\mathcal{M}_3$	$\mathcal{M}_4$	$\mathcal{M}_5$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
t	T	F	T	F	T	T	$\cdots$	T	2	≥ 1
s	T	F	T	T	F	F	$\cdots$	F	2	≥ 1

☞ If $t \equiv G_k$, then t needs $\geq 2^k/2$ compositions.

TASK How can we construct such $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$?

ページ概要：

- 証明の方針：G_k が区別する構造 $\mathcal{M}_0, \mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ を作る
- Key Lemma：d 個を区別するには合成 $\geq d/2$ 個 $\rightsquigarrow$ G_k には合成 $\geq 2^{k-1}$ 個

Proof outline (high level)

Goal Lemma Every CoR term equivalent to G_k needs $2^{\Omega(k)}$ compositions.

Approach We construct structures $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ such that

- G_k distinguishes all of $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$:

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
G_k	T	F	F	$\cdots$	F	2^k	$\geq 2^k/2$

- **Key Lemma** If a CoR term t distinguishes d structures among $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$, then t needs $\geq d/2$ compositions.

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\mathcal{M}_3$	$\mathcal{M}_4$	$\mathcal{M}_5$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
t	T	F	T	F	T	T	$\cdots$	T	2	≥ 1
t'	F	T	T	T	T	F	$\cdots$	F	4	≥ 2

☞ If $t \equiv G_k$, then t needs $\geq 2^k/2$ compositions.

TASK How can we construct such $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$?

ページ概要：

- 証明の方針：G_k が区別する構造 $\mathcal{M}_0, \mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ を作る
- Key Lemma：d 個を区別するには合成 $\geq d/2$ 個 $\rightsquigarrow$ G_k には合成 $\geq 2^{k-1}$ 個

Proof outline (high level)

Goal Lemma Every CoR term equivalent to G_k needs $2^{\Omega(k)}$ compositions.

Approach We construct structures $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ such that

- G_k distinguishes all of $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$:

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
G_k	T	F	F	$\cdots$	F	2^k	$\geq 2^k/2$

- **Key Lemma** If a CoR term t distinguishes d structures among $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$, then t needs $\geq d/2$ compositions.

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\mathcal{M}_3$	$\mathcal{M}_4$	$\mathcal{M}_5$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
t	T	F	T	F	T	T	$\cdots$	T	2	≥ 1
t'	F	T	T	T	T	F	$\cdots$	F	4	≥ 2

👉 If $t \equiv G_k$, then t needs $\geq 2^k/2$ compositions.

TASK How can we construct such $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$?

ページ概要：

- 証明の方針：G_k が区別する構造 $\mathcal{M}_0, \mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ を作る
- Key Lemma：d 個を区別するには合成 $\geq d/2$ 個 $\rightsquigarrow$ G_k には合成 $\geq 2^{k-1}$ 個

Proof outline (high level)

Goal Lemma Every CoR term equivalent to G_k needs $2^{\Omega(k)}$ compositions.

Approach We construct structures $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ such that

- G_k distinguishes all of $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$:

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
G_k	T	F	F	$\cdots$	F	2^k	$\geq 2^k/2$

- **Key Lemma** If a CoR term t distinguishes d structures among $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$, then t needs $\geq d/2$ compositions.

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\mathcal{M}_3$	$\mathcal{M}_4$	$\mathcal{M}_5$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
t	T	F	T	F	T	T	$\cdots$	T	2	≥ 1
t'	F	T	T	T	T	F	$\cdots$	F	4	≥ 2

👉 If $t \equiv G_k$, then t needs $\geq 2^k/2$ compositions.

TASK How can we construct such $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$?

ページ概要：

- 証明の方針：G_k が区別する構造 $\mathcal{M}_0, \mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ を作る
- Key Lemma：d 個を区別するには合成 $\geq d/2$ 個 $\rightsquigarrow$ G_k には合成 $\geq 2^{k-1}$ 個

Proof outline (high level)

Goal Lemma Every CoR term equivalent to G_k needs $2^{\Omega(k)}$ compositions.

Approach We construct structures $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ such that

- G_k distinguishes all of $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$:

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
G_k	T	F	F	$\cdots$	F	2^k	$\geq 2^k/2$

- **Key Lemma** If a CoR term t distinguishes d structures among $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$, then t needs $\geq d/2$ compositions.

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\mathcal{M}_3$	$\mathcal{M}_4$	$\mathcal{M}_5$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
t	T	F	T	F	T	T	$\cdots$	T	2	≥ 1
t'	F	T	T	T	T	F	$\cdots$	F	4	≥ 2

☛ If $t \equiv G_k$, then t needs $\geq 2^k/2$ compositions.

TASK How can we construct such $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$?

ページ概要：

- 証明の方針：G_k が区別する構造 $\mathcal{M}_0, \mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ を作る
- Key Lemma：d 個を区別するには合成 $\geq d/2$ 個 $\rightsquigarrow$ G_k には合成 $\geq 2^{k-1}$ 個

Proof outline (high level)

Goal Lemma Every CoR term equivalent to G_k needs $2^{\Omega(k)}$ compositions.

Approach We construct structures $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ such that

- G_k distinguishes all of $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$:

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
G_k	T	F	F	$\cdots$	F	2^k	$\geq 2^k/2$

- **Key Lemma** If a CoR term t distinguishes d structures among $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ from $\mathcal{M}_0$, then t needs $\geq d/2$ compositions.

	$\mathcal{M}_0$	$\mathcal{M}_1$	$\mathcal{M}_2$	$\mathcal{M}_3$	$\mathcal{M}_4$	$\mathcal{M}_5$	$\cdots$	$\mathcal{M}_{2^k}$	#diff.	#comp.
t	T	F	T	F	T	T	$\cdots$	T	2	≥ 1
t'	F	T	T	T	T	F	$\cdots$	F	4	≥ 2

☛ If $t \equiv G_k$, then t needs $\geq 2^k/2$ compositions.

TASK How can we construct such $\mathcal{M}_0$ and $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$?


ページ概要：

- 証明の方針：G_k が区別する構造 $\mathcal{M}_0, \mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$ を作る
- Key Lemma：d 個を区別するには合成 $\geq d/2$ 個 $\rightsquigarrow$ G_k には合成 $\geq 2^{k-1}$ 個

Structures as colored complete graphs

View structures as **complete graphs**, colored with 2^k colors $\mu \subseteq [k]$: $P_i \iff i \in \mu$.

Q_i is used as the complement of P_i (i.e., $i \notin \mu$): exactly one of P_i, Q_i holds on each edge.




for every $i \in [3]$,
the x-y and x-z edges
have labels P_i and Q_i, in either order

☛ $G_k(x, y)$ holds iff some x-z edge has the color μ^c, where μ is the color of the x-y edge.

ページ概要：

- 構造を色付き完全グラフとして見る
- $G_k(x, y)$ は x-y の色 μ に対して、x-z の色が μ^c であるような z が存在することと同値

The randomly colored structure $\mathcal{M}_0$



$\mathcal{M}_0$: randomly colored, $O(k4^k)$ vertices,
cf. [JMT95, Algebra Univers.]

Two-witness property For all $u \neq v$ and all colors $\boldsymbol{\mu}, \boldsymbol{\nu}$, there are at least two **witnesses** w such that the u-w and w-v edges have colors $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$, respectively.


Every CoR term t is color-invariant on $\mathcal{M}_0$: $\mathcal{M}_0 \models t(u,v) \iff \mathcal{M}_0 \models t(u',v')$ whenever the u-v and u'-v' edges have the same color.

Proof by induction on t: the case $t = t_1 \,;\, t_2$ follows from the (two-)witness property.

ページ概要：

- ランダム彩色の $\mathcal{M}_0$ は two-witness property をもつ
- 任意の項で同じ辺の色なら真偽が等しい（color-invariant）

The randomly colored structure $\mathcal{M}_0$



$\mathcal{M}_0$: randomly colored, $O(k\,4^k)$ vertices,
cf. [JMT95, Algebra Univers.]

Two-witness property For all $u \neq v$ and all colors $\boldsymbol{\mu}, \boldsymbol{\nu}$, there are at least two **witnesses** w such that the u-w and w-v edges have colors $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$, respectively.

Every CoR term t is color-invariant on $\mathcal{M}_0$: $\mathcal{M}_0 \models t(u, v) \iff \mathcal{M}_0 \models t(u', v')$ whenever the u-v and u'-v' edges have the same color.


Proof by induction on t: the case $t = t_1 \,;\, t_2$ follows from the (two-)witness property.

ページ概要：

- ランダム彩色の $\mathcal{M}_0$ は two-witness property をもつ
- 任意の項で同じ辺の色なら真偽が等しい（color-invariant）

$\mathcal{M}_\lambda$: flipping at the hub $\hbar$

Hub $\hbar$: a fixed arbitrary vertex of $\mathcal{M}_0$.



$\hbar$ has λ-edges

For v with the $\hbar$-v edge of color λ^c: $\mathcal{M}_0 \models G_k(\hbar, v)$ but $\mathcal{M}_\lambda \not\models G_k(\hbar, v)$.
Flipping each of the 2^k colors λ gives $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$.
Definition r **distinguishes** $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ if $\mathcal{M}_0 \models r(u, v)$ but $\mathcal{M}_\lambda \not\models r(u', v')$ (or vice versa) for some u-v edge in $\mathcal{M}_0$ and u'-v' edge in $\mathcal{M}_\lambda$ of the same color.
$\rightsquigarrow$ Each composition newly distinguishes at most two of them (next slides).

ページ概要：


- hub の λ-edges を flip した $\mathcal{M}_\lambda$ を考えると 2^k 個の構造を得る。
- 各合成が新たに区別するのは高々 2 個（次の 2 ページ）

補足：

- 補・逆・和・積は点対ごとの演算なので、新たに区別することはない（口頭）

$\mathcal{M}_\lambda$: flipping at the hub $\hbar$

Hub $\hbar$: a fixed arbitrary vertex of $\mathcal{M}_0$.



$\hbar$ has λ-edges

$\hbar$ does not have λ-edges

For v with the $\hbar$-v edge of color λ^c: $\mathcal{M}_0 \models G_k(\hbar, v)$ but $\mathcal{M}_\lambda \not\models G_k(\hbar, v)$.
Flipping each of the 2^k colors λ gives $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$.
Definition r **distinguishes** $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ if $\mathcal{M}_0 \models r(u, v)$ but $\mathcal{M}_\lambda \not\models r(u', v')$ (or vice versa) for some u-v edge in $\mathcal{M}_0$ and u'-v' edge in $\mathcal{M}_\lambda$ of the same color.
$\rightsquigarrow$ Each composition newly distinguishes at most two of them (next slides).

ページ概要：


- hub の λ-edges を flip した $\mathcal{M}_\lambda$ を考えると 2^k 個の構造を得る。
- 各合成が新たに区別するのは高々 2 個（次の 2 ページ）

補足：

- 補・逆・和・積は点対ごとの演算なので、新たに区別することはない（口頭）

$\mathcal{M}_\lambda$: flipping at the hub $\hbar$

Hub $\hbar$: a fixed arbitrary vertex of $\mathcal{M}_0$.



$\hbar$ has λ-edges

$\hbar$ does not have λ-edges

For v with the $\hbar$-v edge of color λ^c: $\mathcal{M}_0 \models G_k(\hbar, v)$ but $\mathcal{M}_\lambda \not\models G_k(\hbar, v)$.
Flipping each of the 2^k colors λ gives $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$.
Definition r **distinguishes** $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ if $\mathcal{M}_0 \models r(u, v)$ but $\mathcal{M}_\lambda \not\models r(u', v')$ (or vice versa) for some u-v edge in $\mathcal{M}_0$ and u'-v' edge in $\mathcal{M}_\lambda$ of the same color.
$\rightsquigarrow$ Each composition newly distinguishes at most two of them (next slides).

ページ概要：


- hub の λ-edges を flip した $\mathcal{M}_\lambda$ を考えると 2^k 個の構造を得る。
- 各合成が新たに区別するのは高々 2 個（次の 2 ページ）

補足：

- 補・逆・和・積は点対ごとの演算なので、新たに区別することはない（口頭）

$\mathcal{M}_\lambda$: flipping at the hub $\hbar$

Hub $\hbar$: a fixed arbitrary vertex of $\mathcal{M}_0$.



$\hbar$ has λ-edges

$\hbar$ does not have λ-edges

For v with the $\hbar$-v edge of color λ^c: $\mathcal{M}_0 \models G_k(\hbar, v)$ but $\mathcal{M}_\lambda \not\models G_k(\hbar, v)$.
Flipping each of the 2^k colors λ gives $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$.
Definition r **distinguishes** $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ if $\mathcal{M}_0 \models r(u, v)$ but $\mathcal{M}_\lambda \not\models r(u', v')$ (or vice versa) for some u-v edge in $\mathcal{M}_0$ and u'-v' edge in $\mathcal{M}_\lambda$ of the same color.
$\rightsquigarrow$ Each composition newly distinguishes at most two of them (next slides).

ページ概要：

- hub の λ-edges を flip した $\mathcal{M}_\lambda$ を考えると 2^k 個の構造を得る。
- 各合成が新たに区別するのは高々 2 個（次の 2 ページ）

補足：

- 補・逆・和・積は点対ごとの演算なので、新たに区別することはない（口頭）

$\mathcal{M}_\lambda$: flipping at the hub $\hbar$

Hub $\hbar$: a fixed arbitrary vertex of $\mathcal{M}_0$.



$\hbar$ has λ-edges

$\hbar$ does not have λ-edges

For v with the $\hbar$-v edge of color λ^c: $\mathcal{M}_0 \models G_k(\hbar, v)$ but $\mathcal{M}_\lambda \not\models G_k(\hbar, v)$.
Flipping each of the 2^k colors λ gives $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$.
Definition r **distinguishes** $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ if $\mathcal{M}_0 \models r(u, v)$ but $\mathcal{M}_\lambda \not\models r(u', v')$ (or vice versa) for some u-v edge in $\mathcal{M}_0$ and u'-v' edge in $\mathcal{M}_\lambda$ of the same color.
$\rightsquigarrow$ Each composition newly distinguishes at most two of them (next slides).

ページ概要：

- hub の λ-edges を flip した $\mathcal{M}_\lambda$ を考えると 2^k 個の構造を得る。
- 各合成が新たに区別するのは高々 2 個（次の 2 ページ）

補足：

- 補・逆・和・積は点対ごとの演算なので、新たに区別することはない（口頭）

$\mathcal{M}_\lambda$: flipping at the hub $\hbar$

Hub $\hbar$: a fixed arbitrary vertex of $\mathcal{M}_0$.



$\hbar$ has λ-edges

$\hbar$ does not have λ-edges

For v with the $\hbar$-v edge of color λ^c: $\mathcal{M}_0 \models G_k(\hbar, v)$ but $\mathcal{M}_\lambda \not\models G_k(\hbar, v)$.
Flipping each of the 2^k colors λ gives $\mathcal{M}_1, \ldots, \mathcal{M}_{2^k}$.
Definition r **distinguishes** $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ if $\mathcal{M}_0 \models r(u, v)$ but $\mathcal{M}_\lambda \not\models r(u', v')$ (or vice versa) for some u-v edge in $\mathcal{M}_0$ and u'-v' edge in $\mathcal{M}_\lambda$ of the same color.
$\rightsquigarrow$ Each composition newly distinguishes at most two of them (next slides).

ページ概要：

- hub の λ-edges を flip した $\mathcal{M}_\lambda$ を考えると 2^k 個の構造を得る。
- 各合成が新たに区別するのは高々 2 個（次の 2 ページ）

補足：

- 補・逆・和・積は点対ごとの演算なので、新たに区別することはない（口頭）

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (1/2)

Q. If t and s do not distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$, when can $t\,;s$ newly distinguish $\mathcal{M}_\lambda$?

Case $\hbar \notin \{u, v\}$: Never: $t\,;s$ cannot distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ at such (u, v).

Similarly for the converse direction (from $\mathcal{M}_\lambda$ to $\mathcal{M}_0$).


ページ概要：

- hub を含まない点対では、同じ色の組の witness を別の頂点で取れる（two-witness property）
- よってこの場合では区別できない

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (1/2)

Q. If t and s do not distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$, when can $t\,;s$ newly distinguish $\mathcal{M}_\lambda$?

Case $\hbar \notin \{u, v\}$: Never: $t\,;s$ cannot distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ at such (u, v).



Similarly for the converse direction (from $\mathcal{M}_\lambda$ to $\mathcal{M}_0$).


ページ概要：

- hub を含まない点対では、同じ色の組の witness を別の頂点で取れる（two-witness property）
- よってこの場合では区別できない

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (1/2)

Q. If t and s do not distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$, when can $t\,;s$ newly distinguish $\mathcal{M}_\lambda$?

Case $\hbar \notin \{u, v\}$: Never: $t\,;s$ cannot distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ at such (u, v).



② The **two-witness property** shows there is w with the same colors (μ, ν)

Similarly for the converse direction (from $\mathcal{M}_\lambda$ to $\mathcal{M}_0$).


ページ概要：

- hub を含まない点対では、同じ色の組の witness を別の頂点で取れる（two-witness property）
- よってこの場合では区別できない

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (1/2)

Q. If t and s do not distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$, when can $t\,;s$ newly distinguish $\mathcal{M}_\lambda$?

Case $\hbar \notin \{u, v\}$: Never: $t\,;s$ cannot distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ at such (u, v).



Similarly for the converse direction (from $\mathcal{M}_\lambda$ to $\mathcal{M}_0$).


ページ概要：

- hub を含まない点対では、同じ色の組の witness を別の頂点で取れる（two-witness property）
- よってこの場合では区別できない

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (1/2)

Q. If t and s do not distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$, when can $t\,;s$ newly distinguish $\mathcal{M}_\lambda$?

Case $\hbar \notin \{u, v\}$: Never: $t\,;s$ cannot distinguish $\mathcal{M}_\lambda$ from $\mathcal{M}_0$ at such (u, v).



Similarly for the converse direction (from $\mathcal{M}_\lambda$ to $\mathcal{M}_0$).

ページ概要：

- hub を含まない点対では、同じ色の組の witness を別の頂点で取れる（two-witness property）
- よってこの場合では区別できない

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (2/2)

Case $u = \hbar$ (similarly for $v = \hbar$): the $\hbar$-w edge itself may be flipped.

So the answer changes only if t holds on some λ-edges and only on them ($v = \hbar$: the same for s).
Such λ is uniquely determined for t and for s.
$\rightsquigarrow$ Each composition newly distinguishes at most two $\mathcal{M}_\lambda$.
(The remaining cases: loops (u, u), including $(\hbar, \hbar)$, and witnesses $w \in \{u, v\}$. See the paper for details.)

ページ概要：

- hub を含む点対で区別できるのは、t（または s）が λ-edges だけで成り立つときだけ
- 各合成が新たに区別できる $\mathcal{M}_\lambda$ は高々 2 個

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (2/2)

Case $u = \hbar$ (similarly for $v = \hbar$): the $\hbar$-w edge itself may be flipped.




So the answer changes only if t holds on some λ-edges and only on them ($v = \hbar$: the same for s).
Such λ is uniquely determined for t and for s.
$\rightsquigarrow$ Each composition newly distinguishes at most two $\mathcal{M}_\lambda$.
(The remaining cases: loops (u, u), including $(\hbar, \hbar)$, and witnesses $w \in \{u, v\}$. See the paper for details.)

ページ概要：

- hub を含む点対で区別できるのは、t（または s）が λ-edges だけで成り立つときだけ
- 各合成が新たに区別できる $\mathcal{M}_\lambda$ は高々 2 個

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (2/2)

Case $u = \hbar$ (similarly for $v = \hbar$): the $\hbar$-w edge itself may be flipped.




So the answer changes only if t holds on some λ-edges and only on them ($v = \hbar$: the same for s).
Such λ is uniquely determined for t and for s.
$\rightsquigarrow$ Each composition newly distinguishes at most two $\mathcal{M}_\lambda$.
(The remaining cases: loops (u, u), including $(\hbar, \hbar)$, and witnesses $w \in \{u, v\}$. See the paper for details.)

ページ概要：

- hub を含む点対で区別できるのは、t（または s）が λ-edges だけで成り立つときだけ
- 各合成が新たに区別できる $\mathcal{M}_\lambda$ は高々 2 個

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (2/2)

Case $u = \hbar$ (similarly for $v = \hbar$): the $\hbar$-w edge itself may be flipped.




So the answer changes only if t holds on some λ-edges and only on them ($v = \hbar$: the same for s).
Such λ is uniquely determined for t and for s.
$\rightsquigarrow$ Each composition newly distinguishes at most two $\mathcal{M}_\lambda$.
(The remaining cases: loops (u, u), including $(\hbar, \hbar)$, and witnesses $w \in \{u, v\}$. See the paper for details.)

ページ概要：

- hub を含む点対で区別できるのは、t（または s）が λ-edges だけで成り立つときだけ
- 各合成が新たに区別できる $\mathcal{M}_\lambda$ は高々 2 個

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (2/2)

Case $u = \hbar$ (similarly for $v = \hbar$): the $\hbar$-w edge itself may be flipped.



So the answer changes only if t holds on some λ-edges and only on them ($v = \hbar$: the same for s).
Such λ is uniquely determined for t and for s.

$\rightsquigarrow$ Each composition newly distinguishes at most two $\mathcal{M}_\lambda$.


(The remaining cases: loops (u, u), including $(\hbar, \hbar)$, and witnesses $w \in \{u, v\}$. See the paper for details.)

ページ概要：

- hub を含む点対で区別できるのは、t（または s）が λ-edges だけで成り立つときだけ
- 各合成が新たに区別できる $\mathcal{M}_\lambda$ は高々 2 個

Why a composition newly distinguishes at most two $\mathcal{M}_\lambda$ (2/2)

Case $u = \hbar$ (similarly for $v = \hbar$): the $\hbar$-w edge itself may be flipped.



So the answer changes only if t holds on some λ-edges and only on them ($v = \hbar$: the same for s).
Such λ is uniquely determined for t and for s.
$\rightsquigarrow$ Each composition newly distinguishes at most two $\mathcal{M}_\lambda$.
(The remaining cases: loops (u, u), including $(\hbar, \hbar)$, and witnesses $w \in \{u, v\}$. See the paper for details.)

ページ概要：

- hub を含む点対で区別できるのは、t（または s）が λ-edges だけで成り立つときだけ
- 各合成が新たに区別できる $\mathcal{M}_\lambda$ は高々 2 個

Outline

1. Our results

2. Technical overview

3. Conclusion

Conclusion

Thm.

There are FO^3 *formulas* $\varphi_1, \varphi_2, \ldots$ *with* $|\varphi_k| = O(k)$ *such that every* CoR *term equivalent to* φ_k *needs* $\geq 2^{k-1}$ *compositions.*

Our proof also applies to FO^3 vs. FO^2, matrix query languages, and CoR with RTC.

Key Idea the randomly colored structure $\mathcal{M}_0$ with its 2^k flipped structures.
Each composition can newly distinguish ≤ 2 of them.

Problem (Open Questions)

- *a superpolynomial gap over a* ***fixed*** *finite signature? (e.g., one binary relation)*
- *the exact number of compositions for* G_k*?* *(between* 2^{k-1} *and* 2^k*)*
- *the domain size from which the blowup begins?* *(between* $2^k/(8k)$ *and* $O(k4^k)$*)*

Thank you for your attention!

ページ概要：

- 結果（合成 $\geq 2^{k-1}$）
- 鍵となるアイデア（$\mathcal{M}_0$ の 2^k 通りの flip）
- 未解決問題：固定シグネチャ、G_k の正確な合成の個数、領域サイズ（$2^k/(8k)$ と $O(k4^k)$ の間）

Conclusion

Thm.

There are FO^3 *formulas* $\varphi_1, \varphi_2, \ldots$ *with* $|\varphi_k| = O(k)$ *such that every* CoR *term equivalent to* φ_k *needs* $\geq 2^{k-1}$ *compositions.*

Our proof also applies to FO^3 vs. FO^2, matrix query languages, and CoR with RTC.

Key Idea the randomly colored structure $\mathcal{M}_0$ with its 2^k flipped structures.
Each composition can newly distinguish ≤ 2 of them.

Problem (Open Questions)

- *a superpolynomial gap over a* ***fixed*** *finite signature? (e.g., one binary relation)*
- *the exact number of compositions for* G_k*?* *(between* 2^{k-1} *and* 2^k*)*
- *the domain size from which the blowup begins?* *(between* $2^k/(8k)$ *and* $O(k4^k)$*)*

Thank you for your attention!

ページ概要：

- 結果（合成 $\geq 2^{k-1}$）
- 鍵となるアイデア（$\mathcal{M}_0$ の 2^k 通りの flip）
- 未解決問題：固定シグネチャ、G_k の正確な合成の個数、領域サイズ（$2^k/(8k)$ と $O(k4^k)$ の間）

Conclusion

Thm.

There are FO^3 *formulas* $\varphi_1, \varphi_2, \ldots$ *with* $|\varphi_k| = O(k)$ *such that every* CoR *term equivalent to* φ_k *needs* $\geq 2^{k-1}$ *compositions.*

Our proof also applies to FO^3 vs. FO^2, matrix query languages, and CoR with RTC.

Key Idea the randomly colored structure $\mathcal{M}_0$ with its 2^k flipped structures.
Each composition can newly distinguish ≤ 2 of them.

Problem (Open Questions)

- *a superpolynomial gap over a* ***fixed*** *finite signature? (e.g., one binary relation)*
- *the exact number of compositions for* G_k*?* *(between* 2^{k-1} *and* 2^k*)*
- *the domain size from which the blowup begins?* *(between* $2^k/(8k)$ *and* $O(k4^k)$*)*

Thank you for your attention!

ページ概要：

- 結果（合成 $\geq 2^{k-1}$）
- 鍵となるアイデア（$\mathcal{M}_0$ の 2^k 通りの flip）
- 未解決問題：固定シグネチャ、G_k の正確な合成の個数、領域サイズ（$2^k/(8k)$ と $O(k4^k)$ の間）

Conclusion

Thm.

There are FO^3 *formulas* $\varphi_1, \varphi_2, \ldots$ *with* $|\varphi_k| = O(k)$ *such that every* CoR *term equivalent to* φ_k *needs* $\geq 2^{k-1}$ *compositions.*

Our proof also applies to FO^3 vs. FO^2, matrix query languages, and CoR with RTC.

Key Idea the randomly colored structure $\mathcal{M}_0$ with its 2^k flipped structures.
Each composition can newly distinguish ≤ 2 of them.

Problem (Open Questions)

- *a superpolynomial gap over a* ***fixed*** *finite signature? (e.g., one binary relation)*
- *the exact number of compositions for* G_k*?* *(between* 2^{k-1} *and* 2^k*)*
- *the domain size from which the blowup begins?* *(between* $2^k/(8k)$ *and* $O(k4^k)$*)*

Thank you for your attention!

ページ概要：

- 結果（合成 $\geq 2^{k-1}$）
- 鍵となるアイデア（$\mathcal{M}_0$ の 2^k 通りの flip）
- 未解決問題：固定シグネチャ、G_k の正確な合成の個数、領域サイズ（$2^k/(8k)$ と $O(k4^k)$ の間）

References i

[Var95] M. Y. Vardi. "On the Complexity of Bounded-Variable Queries". In: **Fourteenth ACM SIGACT-SIGMOD-SIGART Symposium on Principles of Database Systems (PODS)**. ACM, 1995, pp. 266–276.

[BS21] K. Bringmann and J. Slusallek. "Current Algorithms for Detecting Subgraphs of Bounded Treewidth Are Probably Optimal". In: **48th International Colloquium on Automata, Languages, and Programming (ICALP)**. Vol. 198. Leibniz International Proceedings in Informatics (LIPIcs). Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2021, 40:1–40:16. DOI: 10.4230/LIPIcs.ICALP.2021.40.

[NU27] Y. Nakamura and Y. Uezato. "A Fine-Grained Dichotomy for Bounded-Variable Query Evaluation: The Calculus of Relations, a Boolean Modal Logic, and One-Variable Counting Logic". In: **Proceedings of the ACM on Management of Data** 4.5 (PODS) (2027), 295:1–295:27. DOI: 10.1145/3850130.

[Wil14] R. Williams. "Faster Decision of First-Order Graph Properties". In: **Joint Meeting of the Twenty-Third EACSL Annual Conference on Computer Science Logic and the Twenty-Ninth Annual ACM/IEEE Symposium on Logic in Computer Science (CSL-LICS)**. ACM, 2014, 80:1–80:6. DOI: 10.1145/2603088.2603121.

References ii

[Tar41] A. Tarski. "On the Calculus of Relations". In: **The Journal of Symbolic Logic** 6.3 (1941), pp. 73–89. DOI: 10.2307/2268577.

[TG87] A. Tarski and S. Givant. **A Formalization of Set Theory without Variables**. Vol. 41. Colloquium Publications. Providence, RI: American Mathematical Society, 1987. DOI: 10.1090/coll/041.

[Nak22] Y. Nakamura. "Expressive Power and Succinctness of the Positive Calculus of Binary Relations". In: **Journal of Logical and Algebraic Methods in Programming** 127 (2022), p. 100760. DOI: 10.1016/j.jlamp.2022.100760.

[FGL+15] G. H. L. Fletcher, M. Gyssens, D. Leinders, D. Surinx, J. Van den Bussche, D. Van Gucht, S. Vansummeren, and Y. Wu. "Relative Expressive Power of Navigational Querying on Graphs". In: **Information Sciences** 298 (2015), pp. 390–406. DOI: 10.1016/j.ins.2014.11.031.

[Nak23] Y. Nakamura. "Existential Calculi of Relations with Transitive Closure: Complexity and Edge Saturations". In: **38th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)**. IEEE, 2023, pp. 1–13. DOI: 10.1109/LICS56636.2023.10175811.

References iii

[VS24] H. Vinall-Smeeth. "From Quantifier Depth to Quantifier Number: Separating Structures with k Variables". In: **39th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)**. Full version: arXiv:2311.15885. ACM, 2024, 71:1–71:14.

[BGV19] R. Brijder, M. Gyssens, and J. Van den Bussche. **On Matrices and K-Relations**. 2019. arXiv: 1904.03934 [cs.DB].

[BGV22] R. Brijder, M. Gyssens, and J. Van den Bussche. "On Matrices and K-Relations". In: **Annals of Mathematics and Artificial Intelligence** 90.2–3 (2022), pp. 181–210. DOI: 10.1007/s10472-021-09760-4.

[JMT95] P. Jipsen, R. D. Maddux, and Z. Tuza. "Small Representations of the Relation Algebra $E_{n+1}(1, 2, 3)$". In: **Algebra Universalis** 33 (1995), pp. 136–139. DOI: 10.1007/BF01190770.