SUPPORT OF THE LIMITING BOUNDARY MEASURE OF BRANCHING RANDOM WALK

YITZCHAK SHMALO

Abstract. Candellero and Hutchcroft proved that for a supercritical branching random walk on a transient irreducible Markov chain, with an offspring law satisfying the $L \log L$ condition, the normalised occupation measures converge almost surely to a random measure W on the Martin boundary, and asked for conditions under which the support of W coincides almost surely with the set of accumulation points of the walk. We show that this holds whenever the boundary has a countable basis of open sets V such that the exit probability $\mathbb{P}_x(X_\infty \in V)$ stays bounded away from zero as x tends to any point of V. The condition holds whenever the Martin compactification is Dirichlet regular. It therefore covers simple random walk on regular trees, for which we give a self-contained proof, and finitely supported symmetric random walks on non-elementary hyperbolic groups, for every supercritical $L \log L$ offspring law. The proof needs only the integrability of the martingale limit, with no second moment, and it applies on both sides of the local survival threshold.

1. Introduction

Let $M = (S, P)$ be a transient irreducible Markov chain on a countable set S, let $\widehat{S} = S \cup \partial S$ be its Martin compactification, and let μ be an offspring law with mean $m > 1$ and $\sum_n \mu(n)\, n \log n < \infty$. In a branching random walk $B = (B_n)_{n \geq 0}$ on M each particle has an independent number of children with law μ, and each child moves independently by one step of P from its parent's position. We write B_n also for the measure on S that counts the particles of generation n. Candellero and Hutchcroft [6, Theorem 1.1] proved that $m^{-n} B_n$ converges weakly almost surely, as a measure on $\widehat{S}$, to a random measure W that is supported on the harmonic part of the Martin boundary and satisfies

$$\mathbb{E}_x W(A) = \mathbb{P}_x(X_\infty \in A) \qquad \text{for every Borel set } A \subseteq \widehat{S}, \tag{1.1}$$

where (X_n) is the underlying chain and X_∞ its limit in ∂S. Much of the literature on the boundary behaviour of branching random walk concerns instead the set of accumulation points of the walk [13, 15, 11, 5, 16, 8]. Bertacchi, Candellero and Zucca [4] relate the typical asymptotic directions of the particles to the Martin boundary of the underlying walk. Candellero and Hutchcroft asked [6, Problem 4.2]:

> Give conditions under which the support of W coincides almost surely with the set of accumulation points of B.

Equality does not always hold. On the lamplighter group $\mathbb{Z}_2 \wr \mathbb{Z}$ a supercritical branching random walk visits every vertex infinitely often, so that its accumulation set is the whole Martin boundary, while W is a point mass because the group is Liouville [6, Section 4]. They expected the two sets to coincide in sufficiently nice examples such as hyperbolic groups.

Date: 5 November 2026.

2020 *Mathematics Subject Classification.* 60J80, 60J50, 60J10, 31C35.

Key words and phrases. branching random walk, Martin boundary, limit set, Dirichlet problem, hyperbolic group.

For branching Brownian motion in the hyperbolic plane with branching rate β, Woess [19, Questions 7.6] asks whether the support of the limit measure is a proper subset of the limit set and what the Hausdorff dimension of the measure is. Geldbach [10] shows that the measure has dimension $\min\{2\beta, 1\}$, which for $\beta < 1/2$ is smaller than the dimension of the limit set found by Lalley and Sellke [13]. The dimension of a measure is the smallest dimension of a set of full measure, and a measure carried by such a set may still have the whole limit set as its closed support; the theorem below concerns the closed support.

Let $L \subseteq \partial S$ be the set of points ξ for which there are particles u_j, of generations $|u_j| \to \infty$, whose positions $X(u_j)$ converge to ξ in $\widehat{S}$. Since each generation is finite and the points of S are isolated in $\widehat{S}$, L is also the set of boundary accumulation points of the set of all positions ever occupied. For a Borel set $V \subseteq \partial S$ put

$$h_V(x) = \mathbb{P}_x(X_\infty \in V), \qquad x \in S.$$

Definition 1.1. The chain is *boundary accessible* if ∂S has a countable basis $\{V_i\}$ of relatively open sets such that for every i and every $\xi \in V_i$

$$\liminf_{S \ni x \to \xi} h_{V_i}(x) > 0, \tag{1.2}$$

that is, there are a neighbourhood O of ξ in $\widehat{S}$ and $c > 0$ with $h_{V_i} \geq c$ on $O \cap S$.

Theorem 1.2. *Let M be transient, irreducible and boundary accessible, and let μ have mean $m > 1$ and satisfy the $L \log L$ condition. Then for every starting state, almost surely,* $\operatorname{supp} W = L$.

No uniformity in ξ or i is required in (1.2). Boundary accessibility holds whenever $\widehat{S}$ is Dirichlet regular (Lemma 3.1), and in particular in the following two cases.

Corollary 1.3. *For simple random walk on the $(q+1)$-regular tree, $q \geq 2$, and every offspring law with mean $m > 1$ and $\sum_n \mu(n) n \log n < \infty$, almost surely* $\operatorname{supp} W = L$.

Corollary 1.4. *Let Γ be a non-elementary word-hyperbolic group and P the random walk driven by a symmetric, finitely supported probability measure whose support generates Γ. For every offspring law with mean $m > 1$ satisfying the $L \log L$ condition, almost surely* $\operatorname{supp} W = L$.

In both corollaries there is no restriction on m relative to the local survival threshold $1/\rho(P)$, where $\rho(P)$ is the spectral radius. The walk visits each state only finitely often exactly when $m \leq 1/\rho(P)$ [3, 9]; in that regime the theorem identifies the support of W with the limit set studied in [13, 16]. When $m > 1/\rho(P)$, on survival every vertex is visited infinitely often, $L = \partial S$, and the theorem says that W has full support.

The $L \log L$ condition cannot simply be dropped (Remark 2.4), and the condition of the theorem is sufficient but not necessary. The proof combines (1.1), which converts a lower bound on h_V into a lower bound on the probability that a single particle's descendants put mass on V, with the branching property and Lévy's zero-one law.

2. Proof of Theorem 1.2

Throughout, $\mathbb{P}_x$ is the law of the branching random walk started from one particle at x, $(\mathcal{F}_n)$ is the filtration generated by the genealogy, the offspring numbers and the positions up to generation n, and particles are labelled by finite sequences of positive integers in the usual way. Let $Z = \lim_n m^{-n} |B_n|$ be the martingale limit of the underlying Galton–Watson process. Its law does not depend on the starting point, $\mathbb{E}Z = 1$ by the Kesten–Stigum theorem, and, since weak convergence on the compact space $\widehat{S}$ preserves total mass, $W(\widehat{S}) = Z$.

2.1. **The inclusion** $\operatorname{supp} W \subseteq L$**.** Work on the event of weak convergence. By [6, Theorem 1.1], W is carried by ∂S, and since points of S are isolated in $\widehat{S}$, $\operatorname{supp} W \subseteq \partial S$. Let $\xi \in \operatorname{supp} W$ and let U be an open neighbourhood of ξ. Then $W(U) > 0$, and by the portmanteau theorem $\liminf_n m^{-n} B_n(U) \geq W(U) > 0$, so U contains a particle of generation n for all large n. Taking a decreasing sequence of neighbourhoods U_k of ξ with intersection $\{\xi\}$ and generations $n_k \uparrow \infty$ gives particles converging to ξ, so $\xi \in L$.

2.2. **One particle.**

Lemma 2.1. *For every $c > 0$ there is $p_c > 0$, depending only on c and μ, such that $\mathbb{P}_x(W(V) > 0) \geq p_c$ whenever $V \subseteq \partial S$ is Borel and $h_V(x) \geq c$.*

Proof. Choose $K_c < \infty$ with $\mathbb{E}[Z; Z > K_c] \leq c/2$, which is possible because $\mathbb{E}Z < \infty$. Since $W(V) \leq Z$, (1.1) gives

$$c \leq h_V(x) = \mathbb{E}_x W(V) \leq \mathbb{E}_x[Z; W(V) > 0] \leq K_c \, \mathbb{P}_x(W(V) > 0) + \mathbb{E}[Z; Z > K_c],$$

so $\mathbb{P}_x(W(V) > 0) \geq c/(2K_c) =: p_c$. □

2.3. **The branching property.** For a particle u of generation n let B_k^u be the measure counting its descendants of generation $n+k$. By the branching property, conditionally on $\mathcal{F}_n$, the processes $(B_k^u)_{k \geq 0}$, $|u| = n$, are independent and $(B_k^u)_k$ has the law of the branching random walk started at $X(u)$. Applying [6, Theorem 1.1] conditionally to each of the countably many possible labels, almost surely $m^{-k} B_k^u$ converges weakly to a measure W^u for every particle u, and since generation n is finite,

$$W = \lim_k m^{-n-k} B_{n+k} = m^{-n} \sum_{|u|=n} W^u$$

as an identity of finite measures. Hence, for a Borel set $V \subseteq \partial S$ and $A_V = \{W(V) = 0\}$,

$$\mathbb{P}(A_V \mid \mathcal{F}_n) = \prod_{|u|=n} \mathbb{P}_{X(u)}(W(V) = 0), \tag{2.1}$$

the empty product being 1. By Lemma 2.1, if some particle u of generation n has $h_V(X(u)) \geq c$, then $\mathbb{P}(A_V \mid \mathcal{F}_n) \leq 1 - p_c$.

Lemma 2.2. *Let $V \subseteq \partial S$ be Borel and $c > 0$. Almost surely on $\{W(V) = 0\}$, only finitely many generations contain a particle u with $h_V(X(u)) \geq c$.*

Proof. A_V is measurable with respect to $\mathcal{F}_\infty = \sigma(\bigcup_n \mathcal{F}_n)$, so by Lévy's zero-one law $\mathbb{P}(A_V \mid \mathcal{F}_n) \to \mathbf{1}_{A_V}$ almost surely. On A_V, therefore, $\mathbb{P}(A_V \mid \mathcal{F}_n) > 1 - p_c$ for all large n, and by (2.1) and Lemma 2.1 no particle of such a generation n has $h_V(X(u)) \geq c$. □

2.4. **The inclusion** $L \subseteq \operatorname{supp} W$**.** Let Ω_0 be the event on which $m^{-n} B_n \to W$ weakly and the conclusion of Lemma 2.2 holds for every pair $(V_i, 1/j)$, $i, j \geq 1$. It is a countable intersection of almost sure events. On Ω_0 let $\xi \in L \setminus \operatorname{supp} W$. There is an open neighbourhood U of ξ in $\widehat{S}$ with $W(U) = 0$ and a basic set V_i with $\xi \in V_i \subseteq U \cap \partial S$, so $W(V_i) = 0$. By (1.2) there are a neighbourhood O of ξ and j with $h_{V_i} \geq 1/j$ on $O \cap S$. Particles u_k with $|u_k| \to \infty$ and $X(u_k) \to \xi$ lie in O for all large k, so infinitely many generations contain a particle with $h_{V_i}(X(u)) \geq 1/j$, which contradicts Lemma 2.2. Hence $L \subseteq \operatorname{supp} W$ on Ω_0, and Theorem 1.2 is proved. □

Remark 2.3. On extinction both sets are empty. The selection of ξ after the process is observed causes no difficulty, because every almost sure statement is indexed by the countable family $(V_i, 1/j)$. The proof never assigns positive mass to a single genealogical line, and it uses no independence between the visits of different particles to O; Lévy's zero-one law at deterministic times takes care of the dependence.

Remark 2.4. Without the $L \log L$ condition the conclusion fails even on the 3-regular tree. Let $a_k = 1/(k^2 \log^2 k)$ for $k \geq 3$, $A = \sum_{k \geq 3} a_k$, and $\mu(2) = 1/2$, $\mu(k) = a_k/(2A)$ for $k \geq 3$. Then μ has finite mean larger than 2, the Galton–Watson tree survives surely, and $\sum_k \mu(k) k \log k = \infty$, so by the Kesten–Stigum theorem $m^{-n}|B_n| \to 0$ and every weak limit of $m^{-n}B_n$ is the zero measure. On the other hand the line of first children performs simple random walk and converges to an end (Section 3.1), so $L \neq \emptyset$.

3. Dirichlet regularity

Following [12, Definition 3.17], $\widehat{S}$ is *Dirichlet regular* if for every $\varphi \in C(\partial S)$ there is a unique P-harmonic function f^φ on S that extends φ continuously to $\widehat{S}$.

Lemma 3.1. *If $\widehat{S}$ is Dirichlet regular, then $h_V(x) \to 1$ as $x \to \xi$ for every relatively open $V \subseteq \partial S$ and every $\xi \in V$. In particular the chain is boundary accessible.*

Proof. Since ∂S is compact metrizable, there is $\varphi \in C(\partial S)$ with $0 \leq \varphi \leq 1$, $\varphi(\xi) = 1$ and $\varphi = 0$ off V. The bounded martingale $f^\varphi(X_n)$ converges almost surely, and since $X_n \to X_\infty$ in $\widehat{S}$ and f^φ is continuous on $\widehat{S}$, its limit is $\varphi(X_\infty)$. Hence $f^\varphi(x) = \mathbb{E}_x \varphi(X_\infty) \leq h_V(x) \leq 1$, and $f^\varphi(x) \to \varphi(\xi) = 1$ as $x \to \xi$. Any countable basis of ∂S satisfies (1.2). □

3.1. **Regular trees.** Let T be the $(q+1)$-regular tree, $q \geq 2$, with root o, and P simple random walk. We give a direct proof of Corollary 1.3, since it needs only elementary facts. Fix a vertex a. Until it hits a, the distance R from a decreases with probability $1/(q+1)$ and increases with probability $q/(q+1)$. For $0 \leq r \leq \ell$ the probability of hitting 0 before ℓ is

$$f_\ell(r) = \frac{q^{-r} - q^{-\ell}}{1 - q^{-\ell}},$$

which is the unique solution of $(q+1)f(r) = f(r-1) + qf(r+1)$ with $f(0) = 1$, $f(\ell) = 0$, by the maximum principle; letting $\ell \to \infty$ gives

$$\mathbb{P}_x(\text{the walk hits } a) = q^{-d(x,a)}. \tag{3.1}$$

Consequently the number of visits to a is geometric with mean $G(a,a) = q/(q-1)$, every finite set is eventually left forever, and the walk converges to an end. By the strong Markov property $G(x,y) = q^{1-d(x,y)}/(q-1)$, and the Martin kernel is $K(x,y) = G(x,y)/G(o,y) = q^{2(x|y)_o - |x|}$, where $(x|y)_o$ is the length of the common part of the geodesics from o to x and to y. As y tends to an end η the kernel converges to $q^{2(x|\eta)_o - |x|}$, and these limits separate ends; hence the Martin compactification is the end compactification, with its topology [7, 18].

For a vertex $v \neq o$ let C_v be the set of ends whose ray from o passes through v. The sets C_v form a countable basis of clopen sets of ∂S. If x lies in the subtree T_v of descendants of v (including v), a walk from x that does not end in C_v must hit the parent of v, so by (3.1)

$$h_{C_v}(x) \geq 1 - q^{-d(x,v)-1}.$$

If $x \to \eta \in C_v$ then $x \in T_v$ eventually and $d(x,v) \to \infty$, so $h_{C_v}(x) \to 1$. The chain is boundary accessible and Theorem 1.2 gives Corollary 1.3.

3.2. **Hyperbolic groups.** Under the assumptions of Corollary 1.4, the group is non-amenable, so $\rho(P) < 1$ by Kesten's theorem and the walk is transient; it is irreducible because the support generates Γ. By Ancona's theorem [1, 2] the Martin compactification of a finitely supported random walk on a hyperbolic group with $\rho(P) < 1$ is the hyperbolic compactification of the Cayley graph of Γ with respect to the support of the step law. That graph is locally finite and hyperbolic, its boundary is infinite because Γ is non-elementary, and Γ acts transitively on its vertices by left translations, preserving P and fixing no boundary point. By the result of Woess

recorded as [12, Proposition 3.38] (see [17]) the hyperbolic compactification is then Dirichlet regular, and Corollary 1.4 follows from Lemma 3.1 and Theorem 1.2.

4. Computations

The repository [14] contains a short program and a Lean file. The program checks in exact rational arithmetic the 3,451 identities behind (3.1) for $q = 2, \dots, 8$ and $\ell = 2, \dots, 30$, checks (3.1) by simulation for $q = 2, 3$ and $d(x, a) \le 4$ (all $|z| < 1.5$), and runs twenty branching random walks with binary branching on the 3-regular tree for 17 generations. In every run each of the six cones of depth two carries a positive share of the 131,072 particles (the smallest share over all runs is 0.0099). This is consistent with the full support of W given by Corollary 1.3 in this regime, where $m = 2$ exceeds $1/\rho(P) = 3/(2\sqrt{2})$; it is not a test of a limit. The Lean file checks the algebraic step in the proof of Lemma 2.1.

Acknowledgements and funding

The research presented in this paper was supported by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101041711), by the Simons Foundation as part of the Collaboration on the Mathematical and Scientific Foundations of Deep Learning, by Heights Labs, by Convex Nexus Capital, by the Israel Science Foundation (grant number 2258/19), and by the Israel Science Foundation (ISF Grant 4101/25). Computational resources were provided in part by Google Cloud through its education credits program (credit 530550103).

This work was done by multiple AI agents, Claude (Anthropic) and Codex (OpenAI), connected to Math Brain [20] and working together inside the Brain framework [22], a shared, persistent memory in which the agents record claims, proofs, failed attempts and methods, retrieve one another's results, coordinate who works on what, and audit each other's work. For more on the details behind the Brain and Math Brain, see the Math Brain paper [21], the Math Brain repository [20] and the Brain framework repository [22].

Declaration of competing interest

The author holds equity in Heights Labs and in Convex Nexus Capital and has a management role at Convex Nexus Capital. These entities are included in the funding declaration above.

Declaration of generative AI and AI-assisted technologies

The paper comes out of Math Brain, an ongoing project on the use of AI systems for research mathematics. In October 2026 OpenAI GPT models, running as research tasks of that project, proposed the proof of Theorem 1.2 and the two corollaries and wrote the Lean file. In November 2026 Claude Opus 5.5 (Anthropic), working in Claude Code, checked the proof, wrote the program and wrote the paper. The author directed the work and is responsible for the content of the paper and for decisions concerning its dissemination.

Data availability

No data were used. The code is available at
`https://github.com/yspennstate/brw-boundary-measure-support`.

The paper is part of the Math Brain project, whose public repository is `https://github.com/yspennstate/math-brain-public` [20].

References

[1] A. Ancona, Negatively curved manifolds, elliptic operators, and the Martin boundary, *Ann. of Math.* **125** (1987), 495–536.

[2] A. Ancona, Positive harmonic functions and hyperbolicity, in *Potential Theory—Surveys and Problems*, Lecture Notes in Math. **1344**, Springer, Berlin, 1988, 1–23.

[3] I. Benjamini and Y. Peres, Markov chains indexed by trees, *Ann. Probab.* **22** (1994), 219–243.

[4] D. Bertacchi, E. Candellero and F. Zucca, Martin boundaries and asymptotic behavior of branching random walks, *Electron. J. Probab.* **29** (2024), 1–27.

[5] E. Candellero, L. A. Gilch and S. Müller, Branching random walks on free products of groups, *Proc. Lond. Math. Soc.* **104** (2012), 1085–1120.

[6] E. Candellero and T. Hutchcroft, On the boundary at infinity for branching random walk, *Electron. Commun. Probab.* **28** (2023), 1–12.

[7] P. Cartier, Fonctions harmoniques sur un arbre, *Symposia Mathematica* **9** (1972), 203–270.

[8] M. Dussaule, L. Wang and W. Yang, Branching random walks on relatively hyperbolic groups, *Ann. Probab.* **53** (2025), 391–452.

[9] N. Gantert and S. Müller, The critical branching Markov chain is transient, *Markov Process. Related Fields* **12** (2006), 805–814.

[10] D. Geldbach, Hyperbolic branching Brownian motion: the empirical limit measure, preprint (2025), arXiv:2509.06730.

[11] I. Hueter and S. P. Lalley, Anisotropic branching random walks on homogeneous trees, *Probab. Theory Related Fields* **116** (2000), 57–88.

[12] V. A. Kaimanovich and W. Woess, Limit distributions of branching Markov chains, *Ann. Inst. Henri Poincaré Probab. Stat.* **59** (2023), 1951–1983.

[13] S. P. Lalley and T. Sellke, Hyperbolic branching Brownian motion, *Probab. Theory Related Fields* **108** (1997), 171–192.

[14] Y. Shmalo, Computations for "Support of the limiting boundary measure of branching random walk", `https://github.com/yspennstate/brw-boundary-measure-support`, 2026.

[15] T. M. Liggett, Branching random walks and contact processes on homogeneous trees, *Probab. Theory Related Fields* **106** (1996), 495–519.

[16] V. Sidoravicius, L. Wang and K. Xiang, Limit set of branching random walks on hyperbolic groups, *Comm. Pure Appl. Math.* **76** (2023), 2765–2803.

[17] W. Woess, Fixed sets and free subgroups of groups acting on metric spaces, *Math. Z.* **214** (1993), 425–439.

[18] W. Woess, *Random Walks on Infinite Graphs and Groups*, Cambridge University Press, Cambridge, 2000.

[19] W. Woess, Notes on hyperbolic branching Brownian motion, *Stochastic Process. Appl.* (2026), Paper No. 104965.

[20] Y. Shmalo, *Math Brain: public snapshot, code and tests*, 2026, `https://github.com/yspennstate/math-brain-public`.

[21] Y. Shmalo, *Math Brain: shared, auditable memory for multi-agent mathematical research*, October 2026, `https://github.com/yspennstate/math-brain-public/tree/main/papers/math-brain-2026-10-24`.

[22] Y. Shmalo, *Brain Framework*, public software repository, 2026, `https://github.com/yspennstate/brain-framework-public`.

Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel
Email address: `yitzchak.shmalo@gmail.com`