# FIXED POINTS OF STRICTLY NON-VOLTERRA QUADRATIC STOCHASTIC OPERATORS IN FOUR OR MORE TYPES

YITZCHAK SHMALO

ABSTRACT. A quadratic stochastic operator on the simplex of probability vectors on $m$ types is strictly non-Volterra when a pair of parents of types $i$ and $j$ never produces an offspring of type $i$ or $j$. For $m = 3$, Rozikov and Zhamilov proved that every such operator has a unique fixed point, that it is not attractive, and that the operator has 2-cycles and 3-cycles depending on the parameters; Problem 11 of the 2009 list of open problems on quadratic stochastic operators by Ganikhodzhaev and Rozikov asks whether this holds for $m \geq 4$. We settle both clauses about the fixed point. Non-attractiveness fails from $m = 4$ on, by the globally attracting equiprobable operators of Jamilov, Ladra and Mukhitdinov. Uniqueness fails from $m = 5$ on: a five-type operator with the two fixed points $(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 0, 0)$ and $(\frac{1}{9}, \frac{1}{9}, \frac{1}{9}, \frac{1}{3}, \frac{1}{3})$ was studied by Jamilov in 2017, and we show that it is the simplest member of a family, obtained from any three-type operator by adjoining two types that swap, in which the number of fixed points is always exactly two; for $m \geq 6$ any operator with two disjoint invariant triples of types has at least two fixed points, and a six-type example carries a whole segment of them. At $m = 4$ the fixed point is unique: two fixed points would differ on exactly two positive and two negative coordinates, and the mass transferred across such a cut is contracted by twice the largest coordinate, which at a fixed point is at most $3 - \sqrt{5} < 1$. So the uniqueness clause of the theorem holds exactly for $m \in \{3, 4\}$ and the non-attractiveness clause exactly for $m = 3$. We explain why the constructions stop at four types, and give a fixed-point index count that yields a one-paragraph proof of uniqueness for three types and shows that the four-type fixed point has positive determinant $\det(I - DV|_T)$. The finite verifications are carried out in exact arithmetic, and both they and the four-type theorem are checked in Lean 4.

## 1. INTRODUCTION

Let $S^{m-1} = \{x \in \mathbb{R}^m : x_i \geq 0,\ \sum_i x_i = 1\}$. A *quadratic stochastic operator* (QSO) on $S^{m-1}$ is a map

$$V_k(x) = \sum_{i,j=1}^{m} P_{ij,k}\, x_i x_j \qquad (k = 1, \ldots, m), \tag{1}$$

where the heredity coefficients satisfy $P_{ij,k} = P_{ji,k} \geq 0$ and $\sum_k P_{ij,k} = 1$ for every pair $(i, j)$; then $V$ maps $S^{m-1}$ into itself. These operators, introduced by Bernstein as models of the evolution of type frequencies in a population under random mating, have a large literature; the survey [1] and the problem list [2] describe what is known and what is not. The operator is *Volterra* when $P_{ij,k} = 0$ unless $k \in \{i, j\}$ (an offspring is of the type of one of its parents), and *strictly non-Volterra* when

$$P_{ij,k} = 0 \qquad \text{whenever } k \in \{i, j\}, \tag{2}$$

an offspring is never of the type of either parent. Strictly non-Volterra operators exist only for $m \geq 3$. For $m = 3$ every one of them has the form

$$x' = \alpha y^2 + cz^2 + 2yz, \qquad y' = ax^2 + dz^2 + 2xz, \qquad z' = bx^2 + \beta y^2 + 2xy,$$

with $a + b = c + d = \alpha + \beta = 1$, and Rozikov and Zhamilov [6] proved:

*Date*: September 15, 2026.

**Theorem 1.1** ([6]; Theorem 6 of [2])**.** *For all values of the parameters a strictly non-Volterra operator on $S^2$ has a unique fixed point, which is not attractive, and it has 2-cycles and 3-cycles depending on the parameters.*

Problem 11 of [2] asks: *is Theorem 6 true for $m \geq 4$?* The two clauses about the fixed point fail at different thresholds, and we determine both.

**Theorem 1.2.**

(i) *For every $m \geq 4$ there is a strictly non-Volterra operator on $S^{m-1}$ whose unique fixed point is globally attracting: the equiprobable operator $P_{ii,k} = \frac{1}{m-1}$ $(k \neq i)$, $P_{ij,k} = \frac{1}{m-2}$ $(i \neq j,\ k \notin \{i,j\})$. So the non-attractiveness clause of Theorem 1.1 fails for every $m \geq 4$.*

(ii) *For every $m \geq 5$ there is a strictly non-Volterra operator on $S^{m-1}$ with at least two fixed points. For $m = 5$, every operator obtained from a three-type operator by the swap extension of Theorem 4.2 has exactly two; the simplest, the operator $W$ of Section 4, has rational coefficients and the fixed points $p = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 0, 0)$ and $q = (\frac{1}{9}, \frac{1}{9}, \frac{1}{9}, \frac{1}{3}, \frac{1}{3})$, neither of them attractive. For $m \geq 6$ every operator with two disjoint invariant triples of types has two fixed points, and one such six-type operator has a segment of fixed points. So the uniqueness clause of Theorem 1.1 fails for every $m \geq 5$.*

(iii) *For $m = 4$ every strictly non-Volterra operator on $S^3$ has exactly one fixed point. So the uniqueness clause of Theorem 1.1 holds exactly for $m \in \{3, 4\}$.*

Part (i) is due to Jamilov, Ladra and Mukhitdinov [5], who studied the equiprobable family and proved that its fixed point is globally attracting for $m \geq 4$; we include the computation of the Jacobian at the barycentre for completeness. The operator $W$ of part (ii) was introduced by Jamilov [4], who determined its two fixed points $p$ and $q$ and the limit set of every orbit; together with Theorem 1.1 this already shows that uniqueness fails at $m = 5$. The remainder of part (ii) and part (iii) are new. The example at $m = 5$ is a face construction: the face spanned by three of the types is invariant and carries the unique fixed point of the three-type operator it inherits, and the cross rules are arranged so that a second fixed point appears in the interior; Theorem 4.2 shows that this works for every three-type operator and every distribution of the cross mass, always with exactly two fixed points. For $m \geq 6$ two invariant triples carry two fixed points, and one six-type operator has infinitely many. The obstruction that stops the constructions at $m = 4$ is exactly counted: two invariant triples must be disjoint, so $m = 5$ admits at most one, and at $m = 4$ an invariant triple forces the remaining coordinate to vanish identically. At $m = 4$ neither mechanism is available, and Section 6 proves part (iii): two fixed points would differ on two positive and two negative coordinates, and a transfer inequality for such a cut, which uses that the complement of a pair of types is again a pair, contracts the difference by twice the largest coordinate. Jamilov's uniqueness theorem for the symmetric four-type operators [3] is the special case. The index count of Section 7 gives a one-paragraph proof of the uniqueness clause of Theorem 1.1 and shows that the determinant $\det(I - DV|_T)$ is positive at the four-type fixed point. The random and structured searches of Section 8 are consistent with all of this. The cycle clause of Theorem 1.1 is not addressed.

Every finite verification in the paper (the coefficient conditions of $W$, its two fixed points, the Jacobians and their spectra, the segment of fixed points of the six-type operator, the general face lemma) is carried out in exact rational arithmetic and, independently, in Lean 4, and the four-type uniqueness theorem is formalised in full (Section 9).

## 2. Preliminaries

We write $V = V_P$ for the operator (1). The map $V$ is homogeneous of degree two on $\mathbb{R}^m$, so Euler's identity gives

$$DV(x)\,x = 2V(x) \qquad (x \in \mathbb{R}^m), \tag{3}$$

and at a fixed point $x^* \in S^{m-1}$ the number 2 is an eigenvalue of $DV(x^*)$ with the radial eigenvector $x^*$. The simplex is invariant, its tangent space is the hyperplane $\{v : \sum_i v_i = 0\}$, and attractiveness of a fixed point is decided by the eigenvalues of $DV(x^*)$ on that hyperplane; we call these the *tangent eigenvalues*. For a strictly non-Volterra operator,

$$\frac{\partial V_k}{\partial x_k} = 2\sum_j P_{kj,k}\,x_j = 0, \qquad \text{so} \qquad \operatorname{tr} DV(x) = 0 \quad \text{for every } x, \tag{4}$$

and the tangent eigenvalues at a fixed point sum to $-2$. At $m = 3$ this alone gives one tangent eigenvalue of modulus at least 1, which is the non-attractiveness of Theorem 1.1; for $m \geq 4$ the bound $2/(m-1)$ is below 1 and excludes nothing.

A subset $F$ of the types is an *invariant face* if pairs inside $F$ produce only types in $F$: $P_{ij,k} = 0$ for $i, j \in F$ and $k \notin F$. Then the face $\{x \in S^{m-1} : \operatorname{supp} x \subset F\}$ is mapped into itself, and the restriction of $V$ to it is the QSO on $F$ with coefficients $(P_{ij,k})_{i,j,k\in F}$, which are again stochastic (the mass $\sum_{k\in F} P_{ij,k} = 1$ for $i, j \in F$) and again strictly non-Volterra.

**Lemma 2.1** (invariant faces)**.** *Let $F$ be an invariant face of $V$ and let $x \in S^{m-1}$ with $\operatorname{supp} x \subset F$. Then $V_k(x) = 0$ for $k \notin F$ and $V_k(x) = \sum_{i,j\in F} P_{ij,k} x_i x_j$ for $k \in F$. In particular a fixed point of the restricted operator on $F$, padded by zeros, is a fixed point of $V$.*

*Proof.* Every term $P_{ij,k}x_ix_j$ with $i \notin F$ or $j \notin F$ vanishes because $x$ is supported on $F$, and for $i, j \in F$ and $k \notin F$ the coefficient vanishes by invariance. □

## 3. The equiprobable family: non-attractiveness fails from $m = 4$

Let $m \geq 4$ and let $V$ be the equiprobable strictly non-Volterra operator of Theorem 1.2(i). Its barycentre $b = (\frac{1}{m}, \dots, \frac{1}{m})$ is a fixed point, and the Jacobian there is explicit: by (1) and the symmetry of $P$, $\partial V_k/\partial x_j = 2\sum_i P_{ij,k}x_i$, and at $b$ the sum $\sum_i P_{ij,k}$ equals $P_{jj,k} + \sum_{i\notin\{j,k\}} P_{ij,k} = \frac{1}{m-1} + 1 = \frac{m}{m-1}$ for $j \neq k$ and 0 for $j = k$. Hence

$$DV(b) = \frac{2}{m-1}\,(J - I), \tag{5}$$

with $J$ the all-ones matrix, so $DV(b)v = -\frac{2}{m-1}v$ for every tangent vector $v$ (those with $Jv = 0$) and $DV(b)b = 2b$, as (3) requires. The tangent spectrum is the single eigenvalue $-\frac{2}{m-1}$, of modulus 1 at $m = 3$ and smaller than 1 for every $m \geq 4$: the fixed point is attractive as soon as $m \geq 4$. It is in fact globally attracting. A direct expansion gives, for $k \neq l$ and $S = \sum_i x_i$,

$$V_k(x) - V_l(x) = \frac{(x_l - x_k)}{m-2}\Big(2S - \frac{m\,(x_k + x_l)}{m-1}\Big), \tag{6}$$

and Jamilov, Ladra and Mukhitdinov [5] show from this that the coordinate range contracts along every orbit, so that every orbit converges to $b$; the numerical companion confirms the identities (5) and (6) symbolically and the convergence on random orbits. Either way the non-attractiveness clause of Theorem 1.1 fails for every $m \geq 4$, and (5) is checked in Lean for $m = 4$.

## 4. Five types: two fixed points

The operator of this section was introduced by Jamilov [4]; we describe it by its coefficient table, which is the form that Theorem 4.2 generalises. Let $F=\{1,2,3\}$ and define the coefficients of $W$ on five types as follows (all coefficients not listed, and their symmetric copies $P_{ji,k}=P_{ij,k}$, are zero):

$$\text{on the face: } P_{ii,k}=\tfrac12\ (i,k\in F,\ k\neq i),\qquad P_{ij,k}=1\ (\{i,j,k\}=F);$$
$$\text{across: } P_{i4,5}=P_{i5,4}=1\ (i\in F),\qquad P_{44,5}=1,\qquad P_{55,4}=1,\qquad P_{45,k}=\tfrac13\ (k\in F).$$

Every pair has total mass one, $P$ is symmetric, and $P_{ij,k}=0$ whenever $k\in\{i,j\}$: $W$ is a strictly non-Volterra operator on $S^4$. Its coordinates are, for $k\in F$ with $\{a,b\}=F\setminus\{k\}$,

$$W_k(x)=\frac{x_a^2+x_b^2}{2}+2x_ax_b+\frac23x_4x_5\qquad(k\in F,\ \{a,b\}=F\setminus\{k\}),\tag{7}$$
$$W_4(x)=x_5\,(2x_1+2x_2+2x_3+x_5),\qquad W_5(x)=x_4\,(2x_1+2x_2+2x_3+x_4).$$

The face $F$ is invariant, and the restriction of $W$ to it is the symmetric strictly non-Volterra operator on three types, whose unique fixed point is $(\frac13,\frac13,\frac13)$.

**Proposition 4.1** ([4, Theorem 3])**.** *$W$ has the two fixed points $p=(\frac13,\frac13,\frac13,0,0)$ and $q=(\frac19,\frac19,\frac19,\frac13,\frac13)$. The tangent eigenvalues of $DW$ are $\{2,-1,-1,-2\}$ at $p$ and $\{-\frac43,-\frac13,-\frac13,0\}$ at $q$; neither fixed point is attractive. These are the only fixed points of $W$ in $S^4$.*

Jamilov [4] obtains the count by subtracting pairs of coordinate equations, which gives $x_1=x_2=x_3$ and $x_4=x_5$; this is the argument of Theorem 4.2 below in the present special case. We record the eigenvectors, and a second proof of the count that determines the whole solution set of the polynomial system.

*Proof.* At $p$: $W_k(p)=\frac{1/9+1/9}{2}+\frac29=\frac13$ for $k\in F$ and $W_4(p)=W_5(p)=0$. At $q$: $W_k(q)=\frac{1/81+1/81}{2}+\frac{2}{81}+\frac23\cdot\frac19=\frac{1}{27}+\frac{2}{27}=\frac19$ for $k\in F$, and $W_4(q)=\frac13\,(2\cdot\frac13+\frac13)=\frac13=W_5(q)$. The Jacobians are

$$DW(p)=\begin{pmatrix}0&1&1&0&0\\1&0&1&0&0\\1&1&0&0&0\\0&0&0&0&2\\0&0&0&2&0\end{pmatrix},\qquad DW(q)=\begin{pmatrix}0&\frac13&\frac13&\frac29&\frac29\\\frac13&0&\frac13&\frac29&\frac29\\\frac13&\frac13&0&\frac29&\frac29\\\frac23&\frac23&\frac23&0&\frac43\\\frac23&\frac23&\frac23&\frac43&0\end{pmatrix}.$$

At $p$ the eigenvectors $(1,1,1,0,0)$ and $(0,0,0,1,1)$ have eigenvalue $2$, the vectors $(-1,1,0,0,0)$ and $(-1,0,1,0,0)$ have eigenvalue $-1$, and $(0,0,0,-1,1)$ has eigenvalue $-2$; the radial vector $p$ is a multiple of the first, and the combination $(1,1,1,-\frac32,-\frac32)$ carries the eigenvalue $2$ into the tangent space. At $q$ the eigenvectors

$$3q=(\tfrac13,\tfrac13,\tfrac13,1,1),\quad(0,0,0,-1,1),\quad(-1,1,0,0,0),\quad(-1,0,1,0,0),\quad(-\tfrac23,-\tfrac23,-\tfrac23,1,1)$$

have eigenvalues $2,-\frac43,-\frac13,-\frac13,0$, the last four spanning the tangent space. Each set of five eigenvectors is linearly independent (the determinant of the matrix with these columns is nonzero; the Lean file computes it), so the spectra are complete. Since $p$ has a tangent eigenvalue of modulus $2$ and $q$ one of modulus $\frac43$, neither is attractive. Finally, the fixed-point equations $W(x)=x$, $\sum_k x_k=1$ form a zero-dimensional polynomial system: a lexicographic Gröbner basis has last element $x_5(3x_5-1)(x_5^2-3x_5-3)/3$, and solving the system exactly gives sixteen solutions, all real, of which exactly two have all coordinates nonnegative, namely $p$ and $q$ (Section 8). □

The example is one member of a family in which the count is always exactly two; this is the generalisation that the rest of the section is about.

**Theorem 4.2** (swap extension)**.** *Let $U$ be any strictly non-Volterra operator on the three types $F = \{1, 2, 3\}$ and let $\rho$ be any probability vector on $F$. Define $W = W_{U,\rho}$ on five types by: the coefficients of $U$ on pairs inside $F$; $P_{i4,5} = P_{i5,4} = 1$ for $i \in F$; $P_{44,5} = P_{55,4} = 1$; and $P_{45,k} = \rho_k$ for $k \in F$. Then $W$ is a strictly non-Volterra operator with exactly two fixed points: the fixed point of $U$ on the face $F$, and one point with $x_4 = x_5 = \frac{1}{3}$. The face fixed point has the tangent eigenvalue 2 transverse to the face, so it is not attractive.*

*Proof.* The coefficient conditions are immediate. Write $s = x_4$, $t = x_5$ and $\sigma = x_1 + x_2 + x_3 = 1 - s - t$. From the cross rules, $W_4(x) = t(2\sigma + t) = t(2 - 2s - t)$ and $W_5(x) = s(2 - 2t - s)$. At a fixed point, subtracting the two equations $s = t(2 - 2s - t)$ and $t = s(2 - 2t - s)$ gives $s - t = (t - s)(2 - s - t)$, that is $(s - t)(3 - s - t) = 0$, so $s = t$ since $s + t \leq 1$; then $s = s(2 - 3s)$ forces $s = 0$ or $s = \frac{1}{3}$. If $s = t = 0$ the point lies on the face $F$, where $W$ restricts to $U$, and Theorem 1.1 gives exactly one fixed point there; it has $\partial W_4/\partial x_5 = 2\sigma = 2$ and $\partial W_5/\partial x_4 = 2$ at that point, so the block of $DW$ on the coordinates $(x_4, x_5)$ is $\begin{pmatrix} 0 & 2 \\ 2 & 0 \end{pmatrix}$ and 2 is an eigenvalue of that block with eigenvector $e_4 + e_5$; combined with the radial direction this gives a tangent eigenvector for the eigenvalue 2 pointing into the simplex. If $s = t = \frac{1}{3}$, write $u = (x_1, x_2, x_3) = v/3$ with $v$ a probability vector on $F$; then for $k \in F$, $W_k(x) = U_k(u) + 2\rho_k st = \frac{1}{9}U_k(v) + \frac{2}{9}\rho_k$, and the fixed-point equation $u = W(x)$ reads $v = \frac{1}{3}U(v) + \frac{2}{3}\rho$. The map $v \mapsto \frac{1}{3}U(v) + \frac{2}{3}\rho$ sends the simplex on $F$ to itself and is a contraction for the $\ell^1$ norm: $DU = 2A$ with $A$ column-stochastic, so $\|DU\|_{1\to 1} \leq 2$ and the Lipschitz constant is at most $\frac{2}{3}$. It therefore has exactly one fixed point, which gives exactly one fixed point of $W$ with $x_4 = x_5 = \frac{1}{3}$. □

The operator of Proposition 4.1 is $W_{U,\rho}$ with $U$ the symmetric three-type operator and $\rho$ uniform; there $v = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ and the interior fixed point is $q$. The determinant of $I - DW$ on the tangent space is $-12$ at $p$ and $\frac{112}{27}$ at $q$. For the interior point $q$ this makes its fixed-point index $+1$; the point $p$ lies on the boundary of the simplex with the transverse eigenvalue 2, and its fixed-point index as a fixed point of the self-map of the simplex is 0, so the indices add up to the Lefschetz number 1 as they must (Section 7). The point $p$ is the fixed point of the invariant face, and $q$ is the extra fixed point that the cross rules create in the interior. The dynamics of $W$ are known: Jamilov [4, Theorems 4 and 5] shows that $|x_1 - x_2||x_2 - x_3||x_3 - x_1|$ is non-increasing and $|x_4 - x_5|$ non-decreasing along orbits, and that the limit set of an orbit is $\{p\}$ if $x_4 = x_5 = 0$, $\{q\}$ if $x_4 = x_5 > 0$, and the 2-cycle $\{e_4, e_5\}$ otherwise. Section 8 reports the numerical confirmation.

## 5. Six and more types: two invariant faces

**Proposition 5.1.** *Let $m \geq 6$ and let $V$ be a strictly non-Volterra operator on $S^{m-1}$ with two disjoint invariant triples $F$ and $G$. Then $V$ has at least two fixed points, one supported on $F$ and one on $G$.*

*Proof.* By Lemma 2.1 the restriction of $V$ to the face spanned by $F$ is a strictly non-Volterra operator on three types, which by Theorem 1.1 has a fixed point; padded by zeros it is a fixed point of $V$ supported on $F$. The same holds for $G$, and the two points are distinct since $F \cap G = \emptyset$. □

Such operators exist for every $m \geq 6$: put any three-type strictly non-Volterra operator on each of $F$ and $G$, and let every other pair $(i, j)$ distribute its mass over any types outside $\{i, j\}$; for $m \geq 7$ the remaining types can be treated the same way. One choice is worth recording.

**Example 5.2.** On six types with $F = \{1,2,3\}$ and $G = \{4,5,6\}$, put the symmetric operator ($P_{ii,k} = \frac{1}{2}$, $P_{ij,\text{third}} = 1$) on each triple, and let a cross pair $(i,j)$, $i \in F$, $j \in G$, distribute its mass equally over the four types of $(F \setminus \{i\}) \cup (G \setminus \{j\})$. For $x = (a,a,a,b,b,b)$ with $3a+3b = 1$ one computes $V_k(x) = 3a^2 + 3ab = a(3a+3b) = a$ for $k \in F$ and likewise $V_k(x) = b$ for $k \in G$: the whole segment $\{(a,a,a,\frac{1}{3}-a,\frac{1}{3}-a,\frac{1}{3}-a) : 0 \le a \le \frac{1}{3}\}$ consists of fixed points. So a strictly non-Volterra operator can have infinitely many fixed points.

## 6. Four types: the fixed point is unique

Two invariant triples must be disjoint: if they shared a pair $\{i,j\}$, that pair would have to produce, with probability one, both the third type of the first triple and the third type of the second, which is impossible. So $m = 5$ admits at most one invariant triple, which is why the second fixed point of $W$ has to be created in the interior by the cross rules rather than on a second face. At $m = 4$ an invariant triple $F = \{1,2,3\}$ forces $V_4 \equiv 0$: pairs inside $F$ never produce type 4, and a pair containing type 4 never produces type 4 by (2), so every coefficient $P_{ij,4}$ vanishes. Hence every fixed point of a face-invariant four-type operator lies in the face, and there it is unique by Theorem 1.1. There is no invariant pair at all (a pair inside $\{i,j\}$ has nowhere in $\{i,j\}$ to land). So at $m = 4$ neither mechanism of Sections 4 and 5 is available. In fact there is no counterexample at all.

**Theorem 6.1.** *Every strictly non-Volterra quadratic stochastic operator on four types has exactly one fixed point in $S^3$.*

Two observations are used. The first is a cap on the coordinates of a fixed point: only pairs $(i,j)$ with $i,j \neq k$ contribute to $V_k$, and every coefficient is at most 1, so at a fixed point

$$x_k = V_k(x) \le \sum_{i,j\neq k} x_i x_j = (1-x_k)^2, \qquad \text{hence} \qquad x_k \le c := \frac{3-\sqrt{5}}{2} < \frac{1}{2}. \tag{8}$$

The second is polarisation. For $x, y \in S^{m-1}$ and $u = \frac{1}{2}(x+y)$ let $A(u)$ be the matrix of Section 7, $A(u)_{rb} = \sum_a P_{ab,r} u_a$: nonnegative, column-stochastic, with zero diagonal. From $x_a x_b - y_a y_b = u_a(x_b - y_b) + (x_a - y_a)u_b$ and the symmetry of $P$,

$$V(x) - V(y) = 2A(u)(x-y). \tag{9}$$

**Lemma 6.2** (transfer)**.** *Let $\{i,k\} \cup \{j,\ell\}$ be a partition of the four types into two pairs, $G = \{i,k\}$, and $u \in S^{m-1}$. Then*

$$\sum_{r\in G} \big(A(u)(e_i - e_j)\big)_r \le \max(u_i, u_j).$$

*Proof.* Put $p_{ab} = \sum_{r\in G} P_{ab,r}$, the mass that the pair $(a,b)$ sends into $G$; it is symmetric and lies in $[0,1]$. The pair $\{k,i\} = G$ must produce a type outside $G$, so $p_{ki} = 0$; the pair $\{\ell, j\}$ must produce a type outside $\{j,\ell\}$, that is, in $G$, so $p_{\ell j} = 1$. This is where the number of types enters: the complement of a pair is a pair. Now

$$\begin{aligned}\sum_{r\in G} \big(A(u)(e_i - e_j)\big)_r = \sum_a u_a(p_{ai} - p_{aj}) &= u_i(p_{ii} - p_{ij}) + u_j(p_{ij} - p_{jj}) - u_k p_{kj} + u_\ell(p_{\ell i} - 1) \\ &\le u_i(1 - p_{ij}) + u_j p_{ij} \le \max(u_i, u_j).\end{aligned}$$

□

*Proof of Theorem 6.1.* A fixed point exists by Brouwer's theorem. Suppose $x \neq y$ are fixed points and let $z = x - y$, so $z \neq 0$ and $\sum_k z_k = 0$. If $z$ had exactly one positive coordinate $z_i$, then $x_b \le y_b$ for every $b \neq i$; since $V_i$ is a polynomial with nonnegative coefficients in the coordinates other than $x_i$ (this is the strictly non-Volterra condition), $x_i = V_i(x) \le V_i(y) = y_i$,

a contradiction. Exactly one negative coordinate is excluded in the same way. With four coordinates summing to zero this leaves exactly two positive coordinates, $G = \{i, k\}$, and two negative, $H = \{j, \ell\}$. Let $h = z_i + z_k = -z_j - z_\ell > 0$ and $\theta_{ab} = z_a(-z_b)/h^2$ for $a \in G$, $b \in H$; then $\theta_{ab} \geq 0$, $\sum \theta_{ab} = 1$ and $z = h \sum_{a,b} \theta_{ab}(e_a - e_b)$. By (9) and Lemma 6.2, with $u = \frac{1}{2}(x + y)$,

$$h = \sum_{r \in G} \big(V(x) - V(y)\big)_r = 2 \sum_{r \in G} \big(A(u)z\big)_r = 2h \sum_{a,b} \theta_{ab} \sum_{r \in G} \big(A(u)(e_a - e_b)\big)_r$$
$$\leq 2h \max_a u_a \leq 2ch < h,$$

because every coordinate of $u$ is at most $c$ by (8) and $2c = 3 - \sqrt{5} < 1$. □

The proof includes boundary fixed points and uses no nondegeneracy. It also shows why five types are different: there the complement of a pair is a triple, $p_{\ell j} = 1$ is lost, and the operator $W$ of Section 4 has a second fixed point. The same estimates show that the averaged map $x \mapsto (1 - \lambda)x + \lambda V(x)$ is an $\ell^1$ contraction on $\{x \in S^3 : \max_i x_i \leq q\}$ for $c \leq q < \frac{1}{2}$ and $0 < \lambda \leq q$, so that the fixed point is the limit of a convergent iteration; we do not pursue this here. Theorem 6.1 is formalised in Lean (Section 9).

## 7. An index count

For a QSO $V$ and $x \in S^{m-1}$ let $A(x)$ be the $m \times m$ matrix with entries $A(x)_{kl} = \sum_i P_{il,k} x_i$. Then $DV(x) = 2A(x)$ by the symmetry of $P$, $A(x)x = V(x)$, and $A(x)$ is column-stochastic: $\sum_k A(x)_{kl} = \sum_i x_i \sum_k P_{il,k} = 1$. For a strictly non-Volterra operator its diagonal vanishes. At a fixed point $x$ is a Perron vector of $A(x)$, the sum-zero hyperplane is invariant under $A(x)$ since $A(x)^{\mathsf{T}}\mathbf{1} = \mathbf{1}$, and the tangent eigenvalues of $DV(x)$ are $2\mu$ over the eigenvalues $\mu$ of $A(x)$ other than the Perron eigenvalue 1. Writing $p_A(t) = \det(tI - A)$ and $e_j$ for the elementary symmetric functions of the eigenvalues of $A$ (so $e_1 = \operatorname{tr} A = 0$ and $p_A(1) = 0$), one gets

$$\det\big(I - DV(x)\big|_T\big) = -2^m\, p_A(\tfrac{1}{2}) = \begin{cases} 3 + 4 \det A & (m = 3), \\ 3 + 4e_3(A) - 12 \det A & (m = 4), \end{cases} \tag{10}$$

where $T$ is the tangent space. (Indeed $\prod_{\mu \neq 1}(1 - 2\mu) = 2^{m-1} p_A(\frac{1}{2})/(\frac{1}{2} - 1)$; then $e_2 = e_3 - 1$ at $m = 3$ and $e_2 = e_3 - e_4 - 1$ at $m = 4$ from $p_A(1) = 0$, and the rest is substitution.) These identities were checked symbolically at $m = 3$ and numerically at $m = 4$ to machine precision.

The fixed points of $V$ are the fixed points of a continuous self-map of the compact convex set $S^{m-1}$, so their indices sum to the Lefschetz number 1, and an isolated interior fixed point at which $I - DV(x)|_T$ is invertible has index $\operatorname{sign} \det(I - DV(x)|_T)$. This gives a short proof of the uniqueness clause of Theorem 1.1 and a local form of Theorem 6.1.

**Proposition 7.1.**

(i) *A strictly non-Volterra operator on three types has exactly one fixed point.*

(ii) *Let $V$ be a strictly non-Volterra operator on four types and $x$ its fixed point. Every positive real eigenvalue $\rho$ of $DV(x)|_T$ satisfies $\rho \leq 2 \max_i x_i \leq 3 - \sqrt{5}$, and $\det(I - DV(x)|_T) > 0$, that is, $12 \det A(x) < 3 + 4e_3(A(x))$.*

*Proof.* A fixed point $x$ with support $S$ has $V_k(x) = 0$ for $k \notin S$, and since all $x_i x_j$ with $i, j \in S$ are positive this says that every pair in $S$ lands in $S$: the support of a fixed point is an invariant face. A single type is never invariant ($P_{ii,i} = 0$) and neither is a pair ($P_{ij,k} = 0$ for $k \in \{i, j\}$), so at $m = 3$ every fixed point is interior. (i) At $m = 3$ the matrix $A(x)$ is nonnegative with zero diagonal, so $\det A = a_{12}a_{23}a_{31} + a_{13}a_{32}a_{21} \geq 0$ and (10) gives $\det(I - DV(x)|_T) \geq 3$ at every fixed point; each fixed point is nondegenerate of index $+1$, and the indices sum to 1. (ii) Let $z \in T$ be a real eigenvector, $DV(x)z = \rho z$ with $\rho > 0$. Since $DV(x) = 2A(x)$ is

nonnegative with zero diagonal, $z$ cannot have a single positive coordinate ($(DV(x)z)_i \le 0 < \rho z_i$ there), nor a single negative one, so its signs form two pairs $G$ and $H$, and the decomposition in the proof of Theorem 6.1 applies with $A(x)$ in place of $A(u)$: with $h = \sum_{r\in G} z_r > 0$, $\rho h = \sum_{r\in G}(DV(x)z)_r \le 2h \max_a x_a$, and $\max_a x_a \le c$ by (8). For $0 \le s \le 1$ the matrix $I - sDV(x)|_T$ is therefore invertible (a kernel vector would be an eigenvector with eigenvalue $1/s \ge 1 > 2c$), so $s \mapsto \det(I - sDV(x)|_T)$ is continuous and nonvanishing on $[0,1]$ and equals 1 at $s = 0$; it is positive at $s = 1$. The last form of the statement is (10). □

So the four-type fixed point has index $+1$, as the Lefschetz number requires of a unique nondegenerate fixed point. In constrained searches over four-type operators with $V(x) = x$ imposed, the smallest value of $3 + 4e_3 - 12 \det A$ observed was about 3.14, at a point of the form $(a, a, a, a^2)$ with $a = (\sqrt{13} - 3)/2$, and the largest value of $\det A$ about 0.056, against the 0.25 that a negative index would need with $e_3 = 0$. Without the fixed-point condition the quantity can vanish, so both (ii) and Theorem 6.1 must use $V(x) = x$, which enters through the cap (8).

## 8. Numerical checks

The companion script verifies in exact rational arithmetic the coefficient conditions on $W$, the two fixed points, the Jacobians, their spectra and the identity (6), and it computes the following.

*All fixed points of $W$.* Newton's method on the affine hull of the simplex from three thousand random starts, deduplicated, finds sixteen real solutions of $W(x) = x$, of which two lie in the simplex ($p$ and $q$) and fourteen have a negative coordinate; a lexicographic Gröbner basis of the system confirms that the solution set is finite with last polynomial $x_5(3x_5 - 1)(x_5^2 - 3x_5 - 3)/3$.

*Dynamics of $W$.* Orbits from random starts, iterated four thousand times with the sum renormalised at each step to prevent the compounding of rounding error ($\sum_k W_k(x) = (\sum_k x_k)^2$), were classified by their period, in agreement with [4, Theorem 5]: every one of 300 orbits converged to the 2-cycle $\{e_4, e_5\}$ formed by the two vertices outside the invariant face, which $W$ swaps ($W(e_4) = e_5$ and $W(e_5) = e_4$ by (7); the cycle is superattracting, since $DW$ vanishes on the tangent space at a vertex whose only nonzero coordinate is one of $x_4, x_5$); no orbit approached $p$ or $q$. The point $q$ attracts only the invariant slice $x_4 = x_5$, its transverse eigenvalue being $-\frac{4}{3}$, and $p$ attracts only the face $x_4 = x_5 = 0$, where the symmetric three-type operator has the double eigenvalue $-1$. So this operator has the 2-cycles of clause (2) of Theorem 1.1; no 3-cycle was found.

*Random and vertex operators.* Strictly non-Volterra operators were generated with Dirichlet weights on every admissible target set (concentration 1, and 0.1 for sparse operators), and their fixed points in the simplex counted by Newton's method from 120 starts each. Every one of 500 four-type operators, 300 five-type operators and 60 six-type operators had exactly one fixed point in the simplex. The four-type coefficient polytope has $3^4 \cdot 2^6 = 5184$ vertices (each pair sends its whole mass to one admissible type); every vertex operator has exactly one fixed point in the simplex. The five-type operator $W$ has two, and 200 swap extensions $W_{U,\rho}$ of random three-type operators $U$ with random $\rho$ all had exactly two, both with $x_4 = x_5 \in \{0, \frac{1}{3}\}$ as Theorem 4.2 says; the six-type operator of Example 5.2 has a segment, and a six-type operator with the same invariant triples and random cross rules has exactly the two face fixed points. A finite search proves nothing about the operators it does not reach; the four-type counts are consistent with Theorem 6.1.

## 9. Formal verification

The file `QSOFixedPoints.lean`, available with the code, is checked by Lean 4 (version 4.33.1) against Mathlib. It defines the operator (1) and its Jacobian for an arbitrary number of types and proves, in that generality, Euler's identity (3), the vanishing of the diagonal of the Jacobian and

hence of its trace for strictly non-Volterra operators (4), the invariant-face lemma (Lemma 2.1) as a statement about finite sums, and the four-type observation that a face-invariant operator has $V_4 \equiv 0$. For the five-type operator $W$, given by its coefficient table, it verifies by exact rational computation that $W$ is symmetric, nonnegative, row-stochastic and strictly non-Volterra, that $W(p) = p$ and $W(q) = q$ with $p \neq q$, the two Jacobian matrices, the ten eigen-equations of Proposition 4.1 together with the non-vanishing determinants of the two eigenvector matrices, and the membership of the tangential eigenvectors in the sum-zero hyperplane. It also verifies the equiprobable four-type operator's coefficient conditions and the Jacobian (5) at its barycentre with the tangent eigenvalue $-\frac{2}{3}$, the six-type operator of Example 5.2 with the identity $V(a, a, a, \frac{1}{3} - a, \frac{1}{3} - a, \frac{1}{3} - a) = (a, a, a, \frac{1}{3} - a, \frac{1}{3} - a, \frac{1}{3} - a)$ for every $a$, and the algebraic step of Theorem 4.2 (the two equations for $x_4, x_5$ force $x_4 = x_5 \in \{0, \frac{1}{3}\}$). A second file, `QSOFourTypeUnique.lean`, formalises Theorem 6.1 in full for an arbitrary coefficient table on four types satisfying the hypotheses: the cap (8) in the form $x_k \leq \frac{2}{5}$, the monotonicity of $V_i$ in the other coordinates, the polarisation identity (9), the transfer bound of Lemma 6.2, the count of sign classes, and the final contradiction. `#print axioms` on every theorem in both files reports only `propext`, `Classical.choice` and `Quot.sound`. Not formalised: Theorem 1.1 itself (cited), the Gröbner computation, the contraction step of Theorem 4.2, the global attraction of Section 3, and Proposition 7.1.

## 10. Open questions

**Question 10.1.** For which $m$ and which strictly non-Volterra operators do 2-cycles and 3-cycles exist (clause (2) of Theorem 1.1)? In the five-type example the orbits observed numerically are described in Section 8.

**Question 10.2.** What is the maximal number of isolated fixed points of a strictly non-Volterra operator on $S^{m-1}$? The Gröbner computation bounds it for $W$; a general bound in terms of $m$ is not known to us.

## Funding

The research presented in this paper was supported by the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101041711), by the Simons Foundation as part of the Collaboration on the Mathematical and Scientific Foundations of Deep Learning, by Heights Labs, by Convex Nexus Capital, by the Israel Science Foundation (grant number 2258/19), and by the Israel Science Foundation (ISF Grant 4101/25).

## Declaration of competing interest

The author holds equity in Heights Labs and in Convex Nexus Capital and has a management role at Convex Nexus Capital. These entities are included in the funding declaration above.

## Declaration of generative AI and AI-assisted technologies

This paper is a product of an ongoing project on the use of AI systems for research mathematics, provisionally called Math Brain, which studies how to make such systems better at solving open problems; the project and its tooling will be released publicly in due course. The mathematics, the literature search, the computations, the Lean formalisation and the text were produced by AI systems (Claude Opus 5 and Claude Fable 5.1, Anthropic; Codex with GPT-6 Astra, OpenAI), working from the problem list [2]. Three independent adversarial reviews, also by Claude Fable 5.1, re-derived every claim; the index identities (10), Proposition 7.1(i), the

constrained searches of Section 7 and the statement of Theorem 4.2 originate in those reviews, as do the references [3, 4]. Theorem 6.1 and Proposition 7.1(ii) originate in a Codex run on the project's records. The finite verifications and Theorem 6.1 are machine-checked in Lean 4 (Section 9). The arguments have not been refereed by a human mathematician. The author set the goals of the project, commissioned and directed the work and approved the release; the author did not carry out the mathematics, the verification, the computations or the writing, and takes responsibility for the decision to publish.

## Data and code availability

The numerical code, the Lean file with its build log and axiom report, and the LaTeX source are available at

https://github.com/yspennstate/strictly-nonvolterra-qso-fixed-points

## References

[1] R. N. Ganikhodzhaev, F. M. Mukhamedov and U. A. Rozikov, Quadratic stochastic operators and processes: results and open problems, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* 14 (2011), 279–335.

[2] R. N. Ganikhodzhaev and U. A. Rozikov, Quadratic stochastic operators: results and open problems, arXiv:0902.4207 (2009).

[3] U. U. Jamilov, On symmetric strictly non-Volterra quadratic stochastic operators, *Discontinuity, Nonlinearity, and Complexity* 5 (2016), 263–283.

[4] U. U. Jamilov, Dynamics of a strictly non-Volterra quadratic stochastic operator on $S^4$, *Uzbek Math. J.* 2017, no. 2, 66–75.

[5] U. U. Jamilov, M. Ladra and R. T. Mukhitdinov, On the equiprobable strictly non-Volterra quadratic stochastic operators, *Qual. Theory Dyn. Syst.* 16 (2017), 645–655.

[6] U. A. Rozikov and U. U. Zhamilov, The dynamics of strictly non-Volterra quadratic stochastic operators on the 2-simplex, *Sb. Math.* 200 (2009), 1339–1351.

Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel